

Differentiating *Iconella* from *Surirella* (Bacillariophyceae): typifying four Ehrenberg names and a preliminary checklist of the African taxa

Regine Jahn¹, Wolf-Henning Kusber¹, Christine Cocquyt²

1 Botanischer Garten und Botanisches Museum Dahlem, Freie Universität Berlin, Königin-Luise-Str. 6–8, 14195 Berlin, Germany **2** Botanic Garden Meise, Nieuwelaan 38, 1680, Meise, Belgium

Corresponding author: Regine Jahn (r.jahn@bgbm.org)

Academic editor: K. Manoylov | Received 4 May 2017 | Accepted 31 May 2017 | Published 3 July 2017

Citation: Jahn R, Kusber W-H, Cocquyt C (2017) Differentiating *Iconella* from *Surirella* (Bacillariophyceae): typifying four Ehrenberg names and a preliminary checklist of the African taxa. PhytoKeys 82: 73–112. <https://doi.org/10.3897/phytokeys.82.13542>

Abstract

To comply with the new phylogeny within the Surirellales as supported by molecular and morphological data, re-evaluations and re-combinations of taxa from and within the genera *Surirella*, *Cymatopleura*, and *Stenopterobia* and with the re-established genus *Iconella* are necessary. Since the African diatom flora is rich with taxa from these genera, especially *Iconella*, and the authors have studied these taxa recently, describing also new taxa, a preliminary checklist of African *Iconella* and *Surirella* is here presented. 94 names are contained on this list. 57 taxa have been transferred to *Iconella*; 55 taxa were formerly ranked within *Surirella* and two taxa within *Stenopterobia*. 10 taxa have stayed within *Surirella* and six taxa have been transferred from *Cymatopleura* to *Surirella*. 20 *Surirella* and 1 *Stenopterobia* names are listed which are either unrevised or unrevivable since morphological data is missing. Four names and taxa described by Ehrenberg are here typified. Two had been transferred to *Iconella* already: *Iconella bifrons* (Ehrenb.) Ruck & Nakov and *Iconella splendida* (Ehrenb.) Ruck & Nakov. Two are re-transferred from *Cymatopleura* to *Surirella*: *Surirella librile* (Ehrenb.) Ehrenb. and *Surirella undulata* (Ehrenb.) Ehrenb.; both taxa are currently known by their younger synonyms: *Cymatopleura solea* (Bréb.) W. Smith and *Cymatopleura elliptica* (Bréb. ex Kützing) W. Smith. Lectotypes for *Iconella bifrons*, *I. splendida*, *Surirella librile*, and *S. undulata* were designated.

Keywords

diatoms, Surirellales, types, nomenclatural changes, biodiversity

Introduction

Surirella taxa have been recognized, drawn, and described very early in diatom history since they often have large cells. The genus *Surirella* Bory is within the first published diatom genera which are still in current use: *Bacillaria* by Gmelin in 1791, *Fragilaria* by Lyngbye in 1819, *Achnanthes* and *Navicula* by Bory in 1822, *Diatoma* by Bory in 1824, *Melosira* and *Meridion* by Agardh 1824, *Surirella* by Turpin in 1828; further important genera were later described such as *Cymbella* by C. Agardh in 1830, *Gomphonema* by Ehrenberg in 1832, *Encyonema* by Kützing in 1833, *Eunotia* by Ehrenberg in 1837, *Achnanthidium* by Kützing in 1844, *Campylodiscus* by Ehrenberg ex Kützing in 1844, *Nitzschia* by Hassall in 1845, etc.

The genus name *Surirella* was introduced by P.J.F. Turpin in 1828 who had found it in a collection by the French medical doctor Suriray from brackish waters at the coast of Le Havre in France. He published beautiful drawings which had been enlarged in the microscope by 300x. Ehrenberg also used this 300x enlargement for his research and used this genus name first in 1834 for *Surirella bifrons* and *Surirella splendida*; in his 1838 publication (Ehrenberg 1838) he ranked *Surirella* as a subgenus of *Navicula* and contained in it the species *librile*, *splendida*, *bifrons*, *undulata*, *striatula* (type of the name of the genus *Surirella* introduced by Turpin), and *constricta* (no *Surirella* according to Jahn and Kusber 2004). For each of these he added a ? between the genus and the epithet which meant that he thought that this species might belong to a new genus to be differentiated from *Navicula*; at the end of the text he wrote that they definitely belong to the genus *Surirella* because of their different mode of division in comparison to *Navicula*. By 1845 Ehrenberg (1845a, b) had also recombined *Navicula librile* and *Navicula undulata* with *Surirella* (see typifications below).

Subsequently, more *Surirella* taxa were discovered. W. Smith (1851: 7) explains the morphology of *Surirella*: “Valves concave, with a longitudinal central line and margins produced beyond the suture (winged). ... The concavity of the valves, their winged margins, and the longitudinal central line, which wants the central depression so conspicuous in the *Naviculae*, are characters which sufficiently distinguish *Surirella* from all other genera. I believe a careful examination of the loricae ... would detect the presence of alae in all the species.” In this paper he also described and differentiated his new genus *Cymatopleura* against *Surirella*, the main differences being “the undulated surface of the valves seems to indicate a peculiarity of structure sufficient to constitute a generic difference, and the absence of alae and costae implies a further diversity in the internal character which cannot be regarded as unimportant” (W. Smith 1851: 12). Subsequently, W. Smith recombined *Cymatopleura solea* (= *S. librile* Ehrenb.) and *Cymatopleura elliptica* (= *S. undulata* Ehrenb.). In his *Treatise on the Diatoms*, Van Heurck (1896: 374) reintroduced and validated the genus *Stenopterobia* which had been first mentioned by Brébisson; his short differential diagnosis against *Surirella* is: “Frustules very elongated and very narrow, sometimes sigmoid.”

All the above mentioned genera, *Surirella*, *Cymatopleura*, *Stenopterobia*, *Campylodiscus* (for *C. clypeus* (Ehrenb.) Ehrenb. ex Kütz. see Pouličková and Jahn 2007)

are part of the order Surirellales (sensu Round et al. 1990, Ruck and Kociolek 2004, and Ruck et al. 2016a) which are canal-raphe-bearing diatoms with a circumferential raphe at the entire valve margin. The genera *Epithemia* and *Rhopalodia* which have a canal-raphe-not positioned around the entire valve margin, had been placed into the order Rhopalodiales (Round et al. 1990) but Ruck et al. (2016a) placed them also into the order Surirellales because their monophyly is strongly supported by molecular data (Ruck and Theriot 2011, Ruck et al. 2016a). However, the publications of Ruck et al. (2016a, 2016b), performed with morphology and molecular markers on those Surirellales, strongly reject the monophyly of several genera in the current classification (Round et al. 1990), especially concerning the genera *Surirella* and *Campylodiscus*. In order to provide a home to taxa which do not fit into their strict genus definition, Ruck et al. (2016b) reintroduced the genus *Iconella* which had been established by Jurilij in 1949 and *Coronia* which had been established as a subgenus by Ehrenberg, validated by Grunow and raised to genus rank by Ruck and Guiry (2016).

In the tropical African aquatic ecosystems, taxa from the genera *Surirella* and *Cymatopleura*, as traditionally known, play an important role (Ross 1983, Cocquyt and Vyverman 1994, Cocquyt 2000). In typifying historical material from African waters as described by Otto Müller (Cocquyt and Jahn 2005, 2007a, 2007b, 2007c, 2007d, 2014), by Cholnoky (Cocquyt et al. 2017), by Foged (Cocquyt and Kusber 2010), by Woodhead and Tweed (Cocquyt et al. 2013), we have tried to reevaluate earlier findings of these taxa as well as their endemism. In order to help researchers to name their taxa correctly, we are providing a list of African taxa which have been recombined with a different genus; we are also listing those taxa whose names did not change. Since some of Ehrenberg's species have been the basis for varieties of African taxa, we are including the typification of four taxa originally described by Ehrenberg and synonymizing two younger taxa.

Material and methods

From the Ehrenberg Collection at BHUPM (Museum für Naturkunde, Berlin), the following materials (for details of the collection see Jahn and Kusber 2004) were investigated:

- 540128-6 (*Iconella bifrons*)
- 540178-1 (*Iconella splendida*)
- 540177-3 (*Surirella librile*)
- 540177-4 (*Surirella librile*)
- 540138-6 (*Surirella undulata*)
- Zeichenblatt No 1130 (*Iconella bifrons*)
- Zeichenblatt No 1160 (*Iconella splendida*)
- Zeichenblatt No 1151 (*Surirella librile*)
- Zeichenblatt No 1163 (*Surirella undulata*)

New names and typifications are registered in PhycoBank (Kusber et al. 2017), a registration system for nomenclatural acts (see Barkworth et al. 2016) which is currently in the trial phase. Stable http identifiers are linking to the prototype portal. When possible, we are using long-term stable and semantic web compatible identifiers for specimens according to Güntsch et al. (2017).

Two specimens at BR (Botanic Garden Meise) have been reinvestigated and documented. For specimens not seen at BRM (Alfred-Wegener-Institut für Polar- und Meeresforschung, Hustedt Diatom Study Centre, Bremerhaven), Simonsen (1987) was consulted. Author names are standardized according to IPNI database (The International Plant Names Index 2017). For several nomenclatural details the Index Nominum Algarum (1988+) and the AlgaTerra database (Jahn and Kusber 2005+) have been used.

Results and discussion

Typification of species described by Ehrenberg

Iconella bifrons (Ehrenb.) Ruck & Nakov in *Notulae algarum* 10: 1. 2016.

≡ *Navicula bifrons* Ehrenb. in Abh. Königl. Akad. Wiss. Berlin 1833: 259. 1834.
≡ *Surirella bifrons* (Ehrenb.) Ehrenb. in Abh. Königl. Akad. Wiss. Berlin 1841: 388.
1843.

Lectotype (designated here). BHUPM 540128-6 “Trockenpräparate CXXVIII 6”.
(The valve representing the lectotype is reproduced here as Fig. 1A).

<http://phycobank.org/100029>

Comments. The combination in Ehrenberg (1843) has been accepted by Kützing (1844: 61). The specimen of the lectotype was misprinted as “547806-3” in Cocquyt and Jahn (2007b) (McNeill et al. 2012, Art. 7.10). *Iconella bifrons* was introduced in Ruck et al. (2016a) and validated in Ruck et al. (2016b).

Iconella splendida (Ehrenb.) Ruck & Nakov in *Notulae algarum* 10: 2. 2016.

≡ *Navicula splendida* Ehrenb. in Abh. Königl. Akad. Wiss. Berlin 1831: 81. 1832.
≡ *Surirella splendida* (Ehrenb.) Ehrenb. in Abh. Königl. Akad. Wiss. Berlin 1841: 389.
1843.

Lectotype (designated here). [icon!] Drawing BHUPM 1160. (The cell representing the lectotype is reproduced here as Fig. 2A “2-4”).

<http://phycobank.org/100030>




Figure 1. *Iconella bifrons* **A–B** Lectotype: BHUPM 540128-6 **C–D** Ehrenberg's drawing BHUPM 1130 showing different shapes of the same species in Ehrenberg's concept. Scale bar for **A** = 10 µm.




Figure 2. *Iconella splendida*. Lectotype: Drawing BHUPM 1160. **A** The alive cell representing the lectotype in three views (hand written numbers 2–4), length 188 µm **B** Later documentation of valve details by Ehrenberg, hand written numbers 5–6 represent a 226 µm long cell.

Further material. Mica preparation BHUPM 540178-1 shows a girdle view with dark inclusions and is not informative for identification.

Comment. *Iconella splendida* was introduced in Ruck et al. (2016a) and validated in Ruck et al. (2016b).

Surirella librile (Ehrenb.) Ehrenb. in Ber. Bekanntm. Verh. Königl. Preuss. Akad. Wiss. Berlin 1845: 139 table. 1845.

≡ *Navicula librile* Ehrenb. in Abh. Königl. Akad. Wiss. Berlin 1831: 81. 1832.

Lectotype (designated here). [icon!] BHUPM 1151c, d. (The cell representing the lectotype is reproduced here as Fig. 3A–B).

<http://phycobank.org/100031>

Further material. BHUPM 540177-3 „Trockenpräparate CLXXVII 3“ (Fig. 3D), BHUPM 540177-4“ Trockenpräparate CLXXVII 4“ (Fig. 3C).

Synonyms.

= *Cymbella solea* Bréb. in Brébisson & Godey, Alg. Falaise: 51, pl. VII, p.p. 1835.

≡ *Surirella solea* (Bréb.) Bréb., Consid. Diat.: 17. 1838.

≡ *Cymatopleura solea* (Bréb.) W. Sm. in Ann. Mag. Nat. Hist. ser. 2. 7: 12. 1851.

Nomenclatural comment. Ehrenberg (1845a) introduced and used the name *Surirella librile*. In this publication (1845a) he described all species new to science formally with a Latin diagnosis. Because he did not mark the species as new to science, Ehrenberg introduced the name *Surirella librile* as a new combination of *Navicula librile* under the then accepted genus name *Surirella*. This combination can be verified




Figure 3. *Surirella librile*. Lectotype: Drawing BHUPM 1151. **A** Girdle view representing the lectotype (corresponding to preparation BHUPM 540177-3 in 3D) **B** Valvar view representing the lectotype; Ehrenberg indicated two views of one cell with dots between the undulated girdle view in A and the valvar view in B. **C** Corresponding preparation BHUPM 540177-4 **D** Corresponding preparation BHUPM 540177-3 **E** Documentation of Ehrenberg's observations in 1826 **F** Small cells in girdle view not corresponding to the published protologue. Scale bar for **C–D** = 10 µm.

by the images Ehrenberg (1854) provided e.g. for Berlin material “Brakisches, strichweis lebendes, Erdlager unter Berlin” (Ehrenberg 1854: pl. 14: fig. 38).

Taxonomical comment. Ehrenberg (1832) published *Navicula librile* by a description which included the length of 1/10 Paris Line which is 225.6 µm. But this measurement does not correspond to the first observations he made in Berlin 1826 drawn on a small piece of paper (Fig. 3E) and glued onto the drawing sheet BHUPM 1151. Nevertheless, the published measurement corresponds perfectly to two of his specimens on his drawing sheet BHUPM 1151 showing a living cell in valvar and girdle view (Fig. 3A–B). Therefore, Ehrenberg (1832) was the first who described the species which was some years later described again as *Cymbella solea* Bréb. & Godey (1835) which was later recombined as *Cymatopleura solea* (Bréb.) W. Sm. (1851) and became type of the name of the genus *Cymatopleura* (Smith 1851). Ehrenberg’s specimens, probably deposited in 1835 or 1836 (see Ehrenberg 1838) give proof (Fig. 3C–D) of his earlier findings (Ehrenberg 1832). In addition, Ehrenberg apparently also observed the form which is identified today as “*Cymatopleura solea* var. *apiculata*” (cf. Fig. 3F, e.g. Krammer & Lange-Bertalot 1988, Hofmann et al. 2013). Schoeman and Archibald (1979) had accepted Ehrenberg’s taxon as having priority under *Cymatopleura*. Later *Cymatopleura* was conserved against *Sphinctocystis* Hassall with *Cymatopleura solea* as its type (see Wiersema et al. 2015). Since *Cymatopleura* is here not accepted at the rank of a genus, this conservation is not applicable to our taxonomic treatment.

***Surirella undulata* (Ehrenb.) Ehrenb. in Ber. Bekanntm. Verh. Königl. Preuss. Akad. Wiss. Berlin 1845: 307. 1845.**

≡ *Navicula? undulata* Ehrenb., Infusionstherchen, 187, pl. XXI: fig. XVI. 1838.

Lectotype (designated here). BHUPM 540138-6 “Trockenpräparate CXXXVIII 6” (The valve representing the lectotype is reproduced here as Fig. 4D).

<http://phycobank.org/100032>

Further original material. Drawing BHUPM 1163.

Synonyms.

Surirella elliptica Bréb. ex Kütz., Kieselschal. Bacill., 61, pl. 28: fig. 28. 1844.

≡ *Cymatopleura elliptica* (Bréb. ex Kütz.) W. Sm. in Ann. Mag. Nat. Hist. ser. 2, 7: 13. 1851.

Comment. Ehrenberg (1845b) introduced and used the name *Surirella undulata*. In this publication he described all species new to science formally with a Latin diagnosis. Because he did not mark the species as new to science, Ehrenberg introduced the name *Surirella undulata* as a new combination of *Navicula undulata* under the then accepted genus name *Surirella*. This combination can be verified by the drawing Ehrenberg (1854) provided e.g. for Berlin material “Brakisches, strichweis lebendes, Erdlager unter Berlin” (Ehrenberg 1854: pl. 14: fig. 39). Since Ehrenberg published this taxon name already in 1838, his name has priority over *Surirella elliptica*.




Figure 4. *Surirella undulata*. **A–C** Drawing BHUPM 1163 **D** Lectotype: BHUPM 540138-6, Trock-enräparate CXXXVIII 6. Scale bar for **D** = 10 µm.

Autapomorphies

In the Surirellaceae the raphe canal runs marginally at the edge of the valve. This canal is interrupted on the external valve face at the poles of the valve while internally the raphe is continuous at the head pole, and interrupted at the base pole. Differences between the three genera had been defined as (according to Hofmann et al 2011):

- *Cymatopleura*: valves are crossed by several large undulations which are not interrupted near the median line (= axial area). The raphe is located within a shallow keel (Spaulding and Edlund 2008).
- *Stenopterobia*: valves are elongated or curved sigmoid-like with equally sized poles. The canal raphe is raised above the valve onto a keel (Spaulding and Edlund 2010).
- *Surirella*: valves are iso- or heteropolar, transapical undulations are finely structured and interrupted near the median line.
 - *Pinnatae* group: raphe canal sits directly at the valve mantle; the raphe is interrupted at both poles. Supporting elements are the fibulae which project from the valve mantle more or less into the center of the valve face.
 - *Robustae*: raphe canal rises above valve face and mantle and is located on a wing. Where the canals of the wings, the alar canals, meet the valve face, in LM appears an apically running wavy line which has been named a loop (Schleifenbildung). Between the alar canals lie fenestrae.

These traditional differentiations based on outline, undulations and median line (formerly named pseudoraphe or axial area) were not supported by the molecular data (Ruck et al. 2016a). Ruck et al. (2016a) therefore proposed morphological autapo-

morphies for the differentiation of genera. As a true autapomorphy they accepted only the morphological differentiation between the *Pinnatae* and the *Robustae* group within *Surirella* which means the raphe canal is located either directly on the mantle (*Pinnatae*) or rises above the valve and mantle and has alar canals with fenestral openings occluded by fenestral bars (*Robustae*).

Since the type of the name of the genus *Surirella*, *S. striatula*, belongs to the *Pinnatae* group, the *Pinnatae* make up the true *Surirella* genus including also the taxa from the *Cymatopleura* genera because their raphe canal also is located on the valve mantle. Taxa from the *Robustae* group as well as *Stenopterobia* taxa – and a few *Campylodiscus* taxa i.e. *C. hibernicus* – belong to the reinstated genus *Iconella*. Since alar canals have also been found in marine *Campylodiscus* sensu lato (now *Coronia* (Ehrenb. ex Kütz.) Ruck & Guiry), an additional autapomorphy for *Iconella* besides the occluded fenestral openings are the internally rimmed pores.

This means that the above list of features for identifying the genera needs to be revised (according to Ruck et al. 2016):

- *Campylodiscus* s.s. (*C. clypeus* only plus formerly *Surirella Fastuosae*; most of its marine taxa are now *Coronia*, the freshwater taxa *Iconella*): communication between the raphe canal and interior through a funnel- or chalice-shaped structure.
- *Coronia* (formerly marine *Campylodiscus*): raphe canal rises above the valve and mantle; it has alar canals with fenestral openings often unoccluded and with simple unrimmed pores.
- *Surirella* s.s. (restricted to the *Surirella Pinnatae* plus *Cymatopleura*): the raphe canal is located directly on the mantle.
- *Iconella* (formerly *Surirella Robustae*, *Stenopterobia* plus formerly *Campylodiscus Robusti*): raphe canal rises above the valve face and mantle and has alar canals with fenestral openings occluded by fenestral bars with internally rimmed pores.

Campylodiscus taxa reported from tropical Africa are few. Beside the more common *C. clypeus* and *C. clypeus* var. *bicostata* (W. Sm. ex Roper) Hust. the only endemic species is *Campylodiscus tanganicae* Hust., reported from Lake Tanganyika. Since we cannot determine currently to which genus the African taxa associated historically with *Campylodiscus* belong, we have excluded them from this study. Marine *Coronia* taxa are also not part of this study.

The African *Rhopalodia* and *Epithemia* taxa as described in O. Müllers papers are currently being studied by us and will be published elsewhere.

African *Iconella* Taxa

Iconella aculeata (Hust.) Cocquyt & R. Jahn, comb. nov.

≡ *Surirella aculeata* Hust. in Huber-Pestalozzi, Phytoplankt. Süßwass. vol. 2 (2), 503, fig. 609. 1942.

Lectotype (designated by Simonsen 1987). BRM X1/1 Lake Tanganyika “Tanganikasee”.

<http://phycobank.org/100033>

- *Surirella aculeata* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 354: fig. 9; pl. 355: fig. 1. 1922, nom. inval.

***Iconella acuminata* (Hust.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella acuminata* Hust. in Huber-Pestalozzi, Phytoplankt. Süßwass. vol. 2 (2), 501, fig. 606. 1942.

Lectotype. (designated by Simonsen 1987). BRM X1/7 Lake Tanganyika “Tanganyika See. 6”.

<http://phycobank.org/100034>

- *Surirella acuminata* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 355: 5 - 6. 1922, nom. inval.

***Iconella anassae* (Cholnoky) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella anassae* Cholnoky in Oesterr. Bot. Z. 104: 84, fig. 278–279. 1957.

Lectotype (designated by Cocquyt et al. 2017). UNWH NIWR 186/3707 “Tugela Village, Nkunzini”.

<http://phycobank.org/100035>

***Iconella africana-orientalis* (Cocquyt & R. Jahn) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella africana-orientalis* Cocquyt & R. Jahn in Willdenowia 35: 364. 2005.

≡ *Surirella constricta* var. *africana* O. Müller in Bot. Jahrb. Syst. 34: 32, pl. 2: fig. 1. 1903.

≡ *Surirella muelleri* Hust. [non Forti] in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 355: fig. 2 (caption). 1922, nom. illeg.

Lectotype (designated by Cocquyt and Jahn 2005a). [icon] Müller 1903, pl. 2, fig. 1; reproduced as fig. 8 in Cocquyt and Jahn (2005) “Plankton of Lake Malombe [Malawi]”.

<http://phycobank.org/100036>

= *Surirella constricta* var. *maxima* O. Müll. in Bot. Jahrb. Syst. 34: 32, pl. 2: fig. 2. 1903.

Lectotype (designated by Cocquyt and Jahn 2005a). [icon] Müller 1903, pl. 2, fig. 2; reproduced as fig. 7 in Cocquyt and Jahn (2005a) “Plankton of Lake Malawi, northern part, Tanzania”.

***Iconella agonaensis* (Foged) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella agonaensis* Foged in Biol. Skr. 15 (1): 123, 151, pl. 25: fig. 3. 1966.

Holotype. C Ghana 141/1961. “Southwest Ghana. Fresh water (a small stream in bamboo thicket between the villages Agona and Nsuaem, Loc. No. 12). 9.III.1961.”

<http://phycobank.org/100037>

***Iconella approximata* (Woodhead & Tweed ex Cocquyt, Jüttner & Kusber) Cocquyt, Jüttner & Kusber, comb. nov.**

≡ *Surirella approximata* Woodhead & Tweed ex Cocquyt, Jüttner & Kusber in Diatom Res. 28: 122. 2013.

Holotype. NMW C90.12.179 “River Chigara, Sierra Leone”.

<http://phycobank.org/100038>

- *Surirella approximata* Woodhead & Tweed in Hydrobiologia 12 (2/3): 202, pl. 6 figs 71, 73. 1958, nom. inval.

***Iconella bonsaensis* (Foged) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella bonsaensis* Foged in Biol. Skr. 15 (1): 124, 151, pl. 25: fig. 1. 1966.

Holotype. C Ghana 151/1961. “Southwest Ghana. Fresh water (the Bonsa river, a tributary to the Ankobra river; Loc. No. 14). 9.III.1961.”

<http://phycobank.org/100039>

***Iconella brevicostata* (O. Müll.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella brevicostata* O. Müll. in Bot. Jahrb. Syst. 34: 34-35, pl. 2, fig. 9. 1903.

Lectotype (designated by Cocquyt and Jahn 2005a). B 40 0040181 [<http://herbarium.bgbm.org/object/B400040181>] “Lake Malombe after discharge of Lake Nyassa [Lake Malawi, Malawi] (sample B 52 0000039 [<http://herbarium.bgbm.org/object/B520000039>]”).

<http://phycobank.org/100040>

= *Surirella tanganyikae* G.S. West in J. Linn. Soc., London. Bot. 38: 166, pl. 8: fig. 6. 1907.

Localities. “Tanganyika – In plankton, Kituta Bay (25 Aug. 1904; no. 77), near Mbete (28 Sept. 1904; no. 105, and near Kala (19 Nov. 1904; no. 170)”.

***Iconella brevicostata* var. *constricta* (Hust.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella brevicostata* var. *constricta* Hust. in Huber-Pestalozzi, Phytoplankt. Süsswass. vol. 2 (2), 505, fig. 615. 1942.

Lectotype (designated by Simonsen 1987). BRM 220/39 Lake Tanganyika “Tanganyika - G.S. West, Exp.”.

<http://phycobank.org/100041>

- *Surirella brevicostata* var. *constricta* in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 309: fig. 2. 1914, nom. inval.

***Iconella brevicostata* var. *elongata* (Hust. ex Simonsen) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella brevicostata* var. *elongata* Hust. ex Simonsen, Atlas and Catalogue of the Diatom Types of F. Hustedt 1: 50. 1987.

Holotype. BRM X1/59 Lake Tanganyika “Tanganyika-See.”.

<http://phycobank.org/100042>

- *Surirella brevicostata* var. *elongata* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 309: fig. 1. 1914, nom. inval.

***Iconella chasei* (Cholnoky) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella chasei* Cholnoky in Portugaliae Acta Biol. Sér. B 4: 225, fig. 118–119, 1954.

Lectotype (designated by Cocquyt et al. 2017). UNWH NWU 07–172 “Eastlands, Umtali District, Southern Rhodesia (now Zimbabwe). Stream bank fully exposed to sunlight, source of mountain ravine on a fern hill on border of Eastlands”.

<http://phycobank.org/100043>

***Iconella cataractarum* (Cocquyt & J.C. Taylor) Cocquyt & J.C. Taylor, comb. nov.**

≡ *Stenopterobia cataractarum* Cocquyt & J.C. Taylor in Phytotaxa 158: 78, figs 1–38. 2014.

Holotype. BR 4345. “Zambia, Luapula Province, Ntumbachushi Falls, 09.853736° S, 28.944683° E, leg. J.C. Taylor 12-349”.

<http://phycobank.org/100044>

***Iconella chepurnovii* (Cocquyt & R. Jahn) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella chepurnovii* Cocquyt & R. Jahn in Nova Hedwigia 84: 542, figs 45–47. 2007.

Holotype (in Cocquyt and Jahn 2007d). BR 4099 (ACBUA 576) Lake Tanganyika “Lacus Tanganyika, Gatororongo (Burundi), Africa centralis”.

Isotype (in Cocquyt and Jahn 2007d). B 40 0040243 [<http://herbarium.bgbm.org/object/B400040243>].

<http://phycobank.org/100045>

***Iconella coei* (Cholnoky ex Cocquyt, J.C. Taylor & Kusber) Cocquyt, J.C. Taylor & Kusber, comb. nov.**

≡ *Surirella coei* Cholnoky ex Cocquyt, J.C. Taylor & Kusber in Fottea 17(1): 39, figs 30–39. 2017.

Holotype. UNWH NIWR 332/6627 “Mount Kenya”.

<http://phycobank.org/100046>

- *Surirella coei* Cholnoky in Oesterr. Bot. Z. 107: 362, fig. 25–26, 1960, nom. inval.

***Iconella congolensis* (Cocquyt & J.C. Taylor) Cocquyt & J.C. Taylor, comb. nov.**

≡ *Surirella congolensis* Cocquyt & J.C. Taylor in Eur. J. Taxon. 133: 8, figs 6–9. 2015.

Holotype. BR 4399 “Oriental Province, DR Congo, Lomami River (0.49339° N and 24.16960° E). Epiphyton on dead submerged wood”.

<http://phycobank.org/100047>

***Iconella crawfordii* (Cocquyt & R. Jahn) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella crawfordii* Cocquyt & R. Jahn in Syst. Geogr. Pl. 77: 218, fig. 3C. 2007.

≡ *Surirella fuellebornii* var. *tumida* Hust. in Huber-Pestalozzi, Phytoplankt. Süsswass. vol. 2 (2), 495, fig. 596. 1942.

Lectotype (cited as holotype but in fact designated by Simonsen 1987). BRM X6/63 Lake Tanganyika “Tanganyika See”.

<http://phycobank.org/100048>

- *Surirella fuellebornii* var. *tumida* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 355: fig. 10. 1922, nom. inval.

***Iconella debesii* (Hust.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella debesii* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 356: figs 3, 4. 1922.

Lectotype (designated by Simonsen 1987). BRM X7/59 Lake Tanganyika “Tangani-kasee” of the plate.

<http://phycobank.org/100049>

Comment. Description in the caption in Hustedt (1922).

***Iconella delicatissima* var. *ghanaensis* (Foged) Cocquyt & Kusber, comb. nov.**

≡ *Surirella delicatissima* var. *ghanaensis* Foged in Biol. Skr. 15 (1): 124, 151, pl. 25: fig. 9. 1966.

≡ *Stenopterobia delicatissima* var. *ghanensis* (Foged) Cocquyt & Kusber in Nova Hedwigia 91: 126. 2010.

Holotype. C Ghana 204/1961 “West Ghana. Fresh water (a small river north of the village Dwinyana; Loc. No. 30). 12.III.1961.”

<http://phycobank.org/100050>

- *Surirella delicatissima* var. *africana* Cholnoky 1959, nom. inval.

Comment. Variety of *Iconella delicatissima* Ruck & Nakov in Notulae algarum 10: 3. 2016.

***Iconella dodowaensis* (Foged) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella dodowaensis* Foged in Biol. Skr. 15 (1): 124, 151, pl. 25: fig. 6. 1966.

Holotype. C Ghana 151/1961 “Southeast Ghana. Fresh water (a river near the village Dodowa, Loc. No. 62). 1.III.1961”.

<http://phycobank.org/100051>

***Iconella dumae* (Hust.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella dumae* Hust. in Hedwigia 63: 169. 1921.

Lectotype (designated by Simonsen 1987). BRM 222/72 “D.O. Afrika. Regenpfütze im Dumagebiet” German East Africa, rain barrel.

<http://phycobank.org/100052>

- *Surirella dumae* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 295: fig. 5, 6. 1913, nom. inval.

***Iconella ebalensis* (Cocquyt & J.C. Taylor) Cocquyt & J.C. Taylor, comb. nov.**

Surirella ebalensis Cocquyt & J.C. Taylor in Eur. J. Taxon. 133: 3, figs 1–5. 2015.

Holotype. BR 4398 “Oriental Province, DR Congo, Lomami River (0.49339° N and 24.16960° E). Epiphyton on *Nymphaea lotus*; collected by François Darchambeau and Ernest Tambwe on 24 Nov. 2012”.

<http://phycobank.org/100053>

***Iconella effusa* (Hust.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella effusa* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 357: figs 1, 2. 1925.

Lectotype (designated by Simonsen 1987). BRM X2/9 Lake Tanganyika “Tanganika See. 6”.

<http://phycobank.org/100054>

Comment. Although Hustedt (in Huber-Pestalozzi 1942) reported this taxon as “nicht selten” it was never observed by other investigators.

***Iconella engleri* (O. Müll.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella engleri* O. Müll. in Bot. Jahrb. Syst. 34: 28, pl. 1, fig. 4. 1903.

≡ *Surirella nyassae* var. *engleri* (O. Müll.) Ostenf. in Bull. Mus. Comp. Zool. Harvard Coll. 52: 178. 1909.

Lectotype (designated by Cocquyt and Jahn 2007a). B 40 0040240 [<http://herbarium.bgbm.org/object/B400040240>] (the valve representing the lectotype was published as fig. 1 in Cocquyt and Jahn 2007a) “Lake Malombe after discharge of Lake Nyassa (Lake Malawi), Malawi”.

<http://phycobank.org/100055>

= *Surirella engleri* f. *angustior* O. Müll. in Bot. Jahrb. Syst. 34: 28, pl. 1: fig. 5. 1903.

Lectotype (designated by Cocquyt and Jahn 2007a). slide B 40 0040241 [<http://herbarium.bgbm.org/object/B400040241>] (the valve representing the lectotype was published as fig. 2 in Cocquyt and Jahn 2007a) “Lake Malombe after discharge of Lake Nyassa (Lake Malawi), Malawi”.

= *Surirella engleri* f. *subconstricta* O. Müll. in Bot. Jahrb. Syst. 34: 28-29, pl. 1, fig. 6. 1903.

Lectotype (designated by Cocquyt and Jahn 2007a). B 40 0040239 [<http://herbarium.bgbm.org/object/B400040239>] “Lake Malombe after discharge of Lake Nyassa (Lake Malawi), Malawi”.

= *Surirella engleri* var. *constricta* O. Müll. in Bot. Jahrb. Syst. 34: 29, pl. 1, figs 7, 8. 1903.

Lectotype (designated by Cocquyt and Jahn 2007a). B 40 0040238 [<http://herbarium.bgbm.org/object/B400040238>] (the valve representing the lectotype was published as fig. 4 in Cocquyt and Jahn 2007a) “Lake Malombe after discharge of Lake Nyassa (Lake Malawi), Malawi”.

= *Surirella engleri* [var. *constricta*] f. *minor* Woodhead & Tweed ex Cocquyt, Jüttner & Kusber in Diatom Res. 28:124, fig. 3. 2013.

Holotype. NMW C90.12.229 “Rokupr, site E, Sierra”.

= *Surirella engleri* [var. *constricta*] f. *sublaevis* O. Müll. in Bot. Jahrb. Syst. 34: 29, pl. I, fig. 9. 1903.

Lectotype (designated by Cocquyt and Jahn 2007a). B 40 0040238 [<http://herbarium.bgbm.org/object/B400040238>] (the valve representing the lectotype was published as fig. 5 in Cocquyt and Jahn 2007a) “Lake Malombe after discharge of Lake Nyassa (Lake Malawi), Malawi”.

***Iconella esamangensis* (Foged) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella esamangensis* Foged in Biol. Skr. 15 (1): 125, 151, pl. 25: fig. 2. 1966.

Holotype. C Ghana 144/1961 “Southwest Ghana. Fresh water (a small river in the rain forest near the village Esamang, Loc. No. 12). 9.III.1961”.

<http://phycobank.org/100056>

***Iconella friedelbinziae* (Cocquyt & R. Jahn) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella friedelbinziae* Cocquyt & R. Jahn in Syst. Geogr. Pl. 77: 218. 2007.

≡ *Surirella fuelebornii* var. *elliptica* O. Müll. In Bot. Jahrb. Syst. 34: 31, pl. 1: fig. 13. 1903.

Lectotype (designated in Cocquyt and Jahn 2007b). [icon] Müller 1903, pl. 1, fig. 13 “Lake Tanganyika”. 1D in Cocquyt & Jahn (2007).

Epitype (designated in Cocquyt and Jahn 2007b). BR 4101 “Lake Tanganyika, Burundi; near Kibwe 105 km south of Bujumbura, sandy, stony beach with abundant tufts of *Vossia cuspidata* Griff. (Poaceae)”.

Isoepitype (designated in Cocquyt and Jahn 2007b). B 40 0040242 [<http://herbarium.bgbm.org/object/B400040242>] (ACBUA 660/2).

<http://phycobank.org/100057>

***Iconella fuelebornii* (O. Müll.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella fuelebornii* O. Müll. in Bot. Jahrb. Syst. 34: 30. 1903.

Lectotype (designated in Cocquyt and Jahn 2007b). B 40 0040236 [<http://herbarium.bgbm.org/object/B400040236>] “Lake Malombe”.

<http://phycobank.org/100058>

= *Surirella fuellebornii* var. *constricta* O. Müll. Bot. Jahrb. Syst. 34: 30-31, pl. 1, fig. 12.1903.

Lectotype (designated in Cocquyt and Jahn 2007b). [icon] Müller 1903, pl. 1, fig. 12. “Lake Malombe” according to Cocquyt and Jahn (2007b).

= *Surirella fuellebornii* f. *subconstricta* O. Müll. Bot. Jahrb. Syst. 34: 30, pl. 1, fig. 11 1903.

Iconella gradifera (Hust.) Cocquyt & R. Jahn, comb. nov.

≡ *Surirella gradifera* Hust. in Huber-Pestalozzi, Phytoplankt. Süßwass. vol. 2 (2), 501, fig. 605. 1942.

Lectotype (designated by Simonsen 1987). X2/57 Lake Tanganyika “Tanganikasee. 6”.

<http://phycobank.org/100059>

- *Surirella gradifera* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 353: fig. 8, 9. 1922, nom. inval.

Iconella heidenii (Hust.) Cocquyt & R. Jahn, comb. nov.

≡ *Surirella heidenii* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 355: fig. 2-4. 1922.

Lectotype (designated by Simonsen 1987). BRM X2/58 Lake Tanganyika “Tanganyika See”.

<http://phycobank.org/100060>

Iconella kusberi (Cocquyt & R. Jahn) Cocquyt & R. Jahn, comb. nov.

≡ *Surirella kusberi* Cocquyt & R. Jahn in Syst. Geogr. Pl. 77: 221. 2007.

≡ *Surirella bifrons* var. *intermedia* O. Müll. in Bot. Jahrb. Syst. 34: 27, pl. 1: fig. 1. 1903.

Lectotype (designated by Cocquyt and Jahn 2007b). [icon] Pl. 1. fig. 1 in Müller (1903) reproduced as fig. 1A in Cocquyt and Jahn (2007b) “unknown” locality.

Epitype (designated by Cocquyt and Jahn 2007b). Slide B 40 0040235 [<http://herbarium.bgbm.org/object/B400040235>], from Müller's material B 52 0000058 [<http://herbarium.bgbm.org/object/B520000058>] (the valve representing the epitype in Cocquyt and Jahn 2007b as fig. 7C “The River Olunga (Tanzania)”.

<http://phycobank.org/100061>

***Iconella lancettula* (Hust.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella lancettula* Hust. in Huber-Pestalozzi, Phytoplankt. Süßwass. vol. 2 (2), 505, fig. 613. 1942.

Lectotype (cited as holotype but in fact designated by Simonsen 1987). BRM X7/58 Lake Tanganyika “Tanganikasee”.

<http://phycobank.org/100062>

- *Surirella lancettula* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 354: figs 1, 2. 1922, nom. inval.

***Iconella latecostata* (Hust.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella latecostata* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 353: figs 5–7. 1922.

Lectotype (designated by Simonsen 1987). BRM X2/70 Lake Tanganyika “Tanganyika See.”

<http://phycobank.org/100063>

***Iconella likomensis* (Cocquyt & R. Jahn) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella likomensis* Cocquyt & R. Jahn in Willdenowia 35: 361. 2005.

≡ *Surirella bifrons* [var. *tumida*] f. *minor* O. Müll. in Bot. Jahrb. Syst. 34: 28, pl. 1, fig. 3. 1903.

Lectotype (designated by Cocquyt and Jahn 2005a). [icon] Müller 1903: t. 1, fig. 3; reproduced as Fig. 4 in Cocquyt and Jahn (2005) Lake Nyassa [Lake Malawi] near Likoma on the bottom.

Epitype (designated by Cocquyt and Jahn 2005a). B 40 0040180 [<http://herbarium.bgbm.org/object/B400040180>] Lake Malombe after discharge of Lake Malawi, Malawi (sample B 52 0000039 [<http://herbarium.bgbm.org/object/B520000039>]).

<http://phycobank.org/100064>

Taxonomical remark. According to Cocquyt and Jahn (2005a), the taxonomic concept *Surirella biseriata* var. *bifrons* (Ehrenb.) Hust. sec. Hustedt in Schmidt (1912) pro parte falls into synonymy.

***Iconella linearis* var. *elliptica* (O. Müll.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella linearis* var. *elliptica* O. Müll. in Bot. Jahrb. Syst. 34: 30, pl. 1: fig. 10. 1903.

Lectotype (designated by Cocquyt and Jahn 2005a). B 40 0040182 [<http://herbarium.bgbm.org/object/B400040182>] Lake Malombe after discharge of Lake Malawi, Malawi (sample B 52 0000039 [<http://herbarium.bgbm.org/object/B520000039>]).

<http://phycobank.org/100065>

Comment. Variety of *Iconella linearis* (W. Sm.) Ruck & Nakov in Notulae algarum 10: 2. 2016.

***Iconella linearis* var. *elongata* (Compère) Cocquyt & R. Jahn, comb. nov. et stat. nov.**

Surirella linearis f. *elongata* Compère in Bull. Jard. Bot. Nat. Belg. 45: 380, figs 11, 23. 1975.

Holotype. BR 982 “Chad, Lake Chad”.

<http://phycobank.org/100066>

***Iconella malombae* (O. Müll.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella malombae* O. Müll. in Bot. Jahrb. Syst. 34: 34, pl. 2: figs 5–6. 1903.

≡ *Surirella nyassae* var. *malombae* (O. Müll.) Ostenf. in Bull. Mus. Comp. Zool. Harvard Coll. 52: 178. 1909.

Lectotype (designated by Cocquyt and Jahn 2007d). B 40 0040230 [<http://herbarium.bgbm.org/object/B400040230>] “Lake Malombe after discharge of Lake Nyasa (Lake Malawi), Malawi” (sample B 52 0000039 [<http://herbarium.bgbm.org/object/B520000039>]”).

= *Surirella malombae* f. *acuta* O. Müll. in Bot. Jahrb. Syst. 34: 34, pl. 2, fig. 7. 1903.

Lectotype (designated by Cocquyt and Jahn 2007d). [icon] Pl. 2: fig. 7 in Müller (1903) “Lake Malombe, after discharge of Lake Malawi, Malawi”.

Epitype (designated by Cocquyt and Jahn 2007d). B 40 0040231 [<http://herbarium.bgbm.org/object/B400040231>] “Lake Victoria near the isle of Djuma” (sample B 52 0000100 [<http://herbarium.bgbm.org/object/B520000100>]).

<http://phycobank.org/100067>

= *Surirella malombae* var. *tumida* Ostenf. in Bot. Jahrb. Syst. 41: 343. 1908.

***Iconella margaritifera* (Hust.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella margaritifera* Hust. in Huber-Pestalozzi, Phytoplankt. Süsswass. vol. 2 (2), 501, fig. 607. 1942.

Lectotype (designated by Simonsen 1987). BRM X2/85 Lake Tanganyika “Tanganyika See. 6”.

<http://phycobank.org/100068>

- *Surirella margaritifera* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 354: figs 3–5. 1922, nom. inval.

***Iconella margaritacea* (O. Müll.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella margaritacea* O. Müll. in Bot. Jahrb. Syst. 34: 37, pl. 2: fig. 12. 1903.

Lectotype (designated by Cocquyt and Jahn 2005a). slide B 40 0040183 [<http://herbarium.bgbm.org/object/B400040183>], river Songwe [Tanzania] (sample B 52 0000036 [<http://herbarium.bgbm.org/object/B520000036>]).

<http://phycobank.org/100069>

***Iconella muelleri* (Forti) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella muelleri* Forti in Atti R. Ist. Veneto Sc. Lett. Ed Arti 69(2): 1284, 1294, pl. 3: fig. 9, 10. 1910.

Type locality. Ethiopia, lago Zulay. Coll. Giov. Negri.

<http://phycobank.org/100070>

***Iconella murielae* (Compère) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella murielae* Compère in Bull. Jard. Bot. Nat. Belg. 45: 381, figs 12, 26. 1975.

Holotype. BR 984, Compère 3875. “Lac Tchad, à 10 km au N du delta du Chari, plancton”.

<http://phycobank.org/100071>

Comment. This species is illustrated by LM and SEM in Bogaerts et al. (2014), additional illustrations are given here in Fig. 5.

***Iconella nagbogensis* (Foged) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella nagbogensis* Foged in Biol. Skr. 15 (1): 125, 151, pl. 25: fig. 7. 1966.

Holotype. C Ghana 279/1961 “Northeast Ghana. Fresh water (a small river near the village Nagbog, Loc. No. 53). 21.III.1961”.

<http://phycobank.org/100072>

***Iconella nervosa* (A.W.F. Schmidt) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella tenera* var. *nervosa* A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 23: fig. 15. 1875.
≡ *Surirella nervosa* (A.W.F. Schmidt) Ant. Mayer in Ber. Naturwiss. Vereins Regensburg 14: 341. 1913.

Lectotype (here designated). [icon!] A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 23: fig. 15. "Whatabevot"

<http://phycobank.org/100073>

Taxonomical comment. From two different localities A.W.F. Schmidt (1875) depicted three valves with an axial area including a central line and spines at both ends of this line. The valve depicted as pl. 23: fig. 15 fits the criterium "illustration with analyses" (McNeill et al. 2012, Art. 38.10) because many small spinules on the valve surface are clearly shown. Therefore we have chosen pl. 23: fig. 15 as the lectotype. fig. 16 is less detailed. We exclude the depicted specimen collected at Khayenmatay (fig. 17) from the species because with its denser costae and less distinct wing projection it probably belongs to a different species.

***Iconella nyassae* (O. Müll.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella nyassae* O. Müll. in Bot. Jahrb. Syst. 34: 33, pl. 2: fig. 3. 1903.

Lectotype (designated by Cocquyt and Jahn 2007d). B 40 0040228 "Lake Malawi near Langenburg, Tanzania, between 40-70 m depth (sample B 52 000014 [<http://herbarium.bgbm.org/object/B520000014>]).

<http://phycobank.org/100074>

= *Surirella nyassae* var. *sagitta* O. Müll. in Bot. Jahrb. Syst. 34: 33, pl. 2: fig. 4. 1903.

Lectotype (designated by Cocquyt and Jahn 2007d). B 40 0040229 "Lake Malawi near Langenburg, Tanzania, between 40-70 m depth" (sample B 52 000013 [<http://herbarium.bgbm.org/object/B520000013>]).

***Iconella obtusiuscula* (G.S. West) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella obtusiuscula* G.S. West in J. Linn. Soc., London Bot. 38: 165, pl. 8: fig. 7. 1907.

Comment. Type specimen not studied but specimens from Lake Tanganyika observed (Cocquyt 1998).

Localities. "Tanganyika – In plankton, Komba Bay (11 Oct. 1904; no. 135) and near Kala (19 Nov. 1904; no. 170)."

<http://phycobank.org/100075>




Figure 5. *Iconella murielae*. Valve from the holotype slide BR 984. **A, C** Overview of the entire valve at different foci **B, D** Detail of the middle part of the valve at different foci **B** is showing the fenestral openings below the raphe canal and **D** the striae and the transapical undulations. Scale bar for **A, C** = 20 µm; scale bar for **B, D** = 10 µm.

***Iconella oliffii* (Cholnoky) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella oliffii* Cholnoky in Oesterr. Bot. Z. 103: 90, fig. 134, 1956.

Lectotype (designated by Cocquyt et al. 2017). [icon]. Fig. 134 in Cholnoky (1956) “Umgeni river by Albert Falls. 14.X.1954”, leg. W.D. Oliff.

Epitype (designated by Cocquyt et al. 2017). UNWH NIWR 193/3860 “Kwa-Zulu Natal, Umgeni River at Albert Falls, Umgeni, South Africa”.

<http://phycobank.org/100076>

***Iconella pangaensis* (O. Müll.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella pangaensis* O. Müll. in Bot. Jahrb. Syst. 34: 257-258, figs 3–4. 1904.

Lectotype (designated by Cocquyt and Jahn 2005). [icon] Müller 1904, fig. 3; reproduced as Fig. 22 in Cocquyt and Jahn (2005a), Rufidji (Usambara-Usagara region) Pangani rapids.

<http://phycobank.org/100077>

***Iconella plana* (G.S. West) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella plana* G.S. West in J. Linn. Soc., London Bot. 38: 165, pl. 8: fig. 5. 1907.

Locality. “Tanganyika – In plankton, near Ndauvie (7 Feb. 1905; no. 227).”

<http://phycobank.org/100078>

Comment. Type specimen not studied but specimens from Lake Tanganyika observed (Cocquyt 1998).

***Iconella propinqua* (Hust.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella propinqua* Hust. in Exploration du Parc National Albert, Mission H. Damas 8: 153, pl. 14: fig. 5, 6. 1949.

Lectotype (designated by Simonsen 1987), BRM 242/6 DR Congo “Belg. Kongo. 39”.

<http://phycobank.org/100079>

***Iconella pseudothienemannii* (Cholnoky) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella pseudothienemannii* Cholnoky in Beih. Nova Hedwigia 21: 72–73, fig. 184, 185, 1966.

Holotype. UNWH NIWR 169/336 „Uferwasser des Kunene-Flusses bei Swart Boois Drift. Stille Bucht am Südufer, 8.8.1961“.

<http://phycobank.org/100080>

***Iconella reicheltii* (Hust.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella reicheltii* Hust. in Huber-Pestalozzi, Phytoplankt. Süsswass. vol. 2 (2), 501, fig. 607. 1942.

Lectotype (designated by Simonsen 1987). BRM X3/69 Lake Tanganyika “Tanganyika See”.

<http://phycobank.org/100081>

- *Surirella reicheltii* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 354: figs 3–5. 1922, nom. inval.

***Iconella sorriensis* (Foged) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella sorriensis* Foged in Biol. Skr. 15 (1): 125, 152, pl. 25: fig. 8. 1966.

Holotype. C Ghana 223/1961 “North Ghana. Fresh water (the Sorri river, the White Volta river system, Loc. No. 35). 16.III.1961”.

<http://phycobank.org/100082>

***Iconella spiralooides* (Hust.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella spiralooides* Hust. in Huber-Pestalozzi, Phytoplankt. Süßwass. vol. 2 (2), 507, fig. 617. 1942.

Lectotype (designated by Simonsen 1987). BRM X4/45 Lake Tanganyika “Tanganika See. 6”.

<http://phycobank.org/100083>

- *Surirella spiralooides* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 353: fig. 2, 3. 1922, nom. inval.

***Iconella subcontorta* (Hust.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella subcontorta* Hust. in Huber-Pestalozzi, Phytoplankt. Süßwass. vol. 2 (2), 518, fig. 633. 1942.

Lectotype (designated by Simonsen 1987). BRM X4/57 Lake Tanganyika “Tanganyika See”.

<http://phycobank.org/100084>

- *Surirella subcontorta* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 356: fig. 1, 2. 1922, nom. inval.

***Iconella takoradiensis* (Foged) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella takoradiensis* Foged in Biol. Skr. 15 (1): 126, 152, pl. 25: fig. 4. 1966.

Holotype. C Ghana 119/1961 “Southeast Ghana. Fresh water (a small river in the rain forest west of Takoradi; Loc. No. 8) 8.III.1961”.

<http://phycobank.org/100085>

= *Surirella takoradiensis* var. *suhinensis* Foged in Biol. Skr. 15 (1): 126, 152, pl. 25: fig. 5.

Holotype. C Ghana 218/1961 “West Ghana. Fresh water (the Suhin river, the Black Volta river system; Loc. No. 33. 13.III.1961”.

***Iconella tchadensis* (Compère) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella tchadensis* Compère in Bull. Jard. Bot. Nat. Belg. 45: 380, figs 11, 23. 1975.

Holotype. BR 987 (see also Bogaerts et al. 2014), Compère 3880, Tchad.

Valves from the holotype slide are given in Fig. 6

<http://phycobank.org/100086>

***Iconella tumida* (O. Müll.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella bifrons* var. *tumida* O. Müll. in Bot. Jahrb. Syst. 34: 27, t. 1, fig. 2. 1903.

≡ *Surirella tumida* (O. Müll.) Cocquyt & R. Jahn in Willdenowia 35: 361. 2005.

Lectotype (designated by Cocquyt and Jahn 2005). [icon] Müller 1903: t. 1, fig. 2; reproduced as fig. 1 in Cocquyt and Jahn (2005a) “Lake Malombe after discharge of Lake Nyassa [Lake Malawi, Malawi]”.

Epitope (designated by Cocquyt and Jahn 2005a). B 40 0040179 [<http://herbarium.bgbm.org/object/B400040179>] “Lake Malombe after discharge of Lake Malawi, Malawi (sample B 52 0000038 [<http://herbarium.bgbm.org/object/B520000038>])”.

<http://phycobank.org/100087>

Taxonomical remark. According to Cocquyt and Jahn (2005a), the taxonomic concept *Surirella biseriata* var. *bifrons* (Ehrenb.) Hust. sec. Hustedt (in Schmidt 1911) pro parte falls into synonymy.

***Iconella turbo* (O. Müll.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella turbo* O. Müll. in Bot. Jahrb. Syst. 34: 34, pl. 2, fig. 8. 1903.

Lectotype (designated by Cocquyt and Jahn 2005a). [icon] Müller 1903, pl. 2, fig. 8; reproduced as fig. 24 in Cocquyt and Jahn (2005a) “Lake Malawi, near Island of Likoma (sample B 52 0000025 [<http://herbarium.bgbm.org/object/B520000025>])”.

<http://phycobank.org/100088>

***Iconella vasta* (Hust.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella vasta* Hust. in Huber-Pestalozzi, Phytoplankt. Süsswass. vol. 2 (2), 503, fig. 611. 1942.

Lectotype (designated by Simonsen 1987). BRM X4/89 Lake Tanganyika “Tanganyika See. 6”.

<http://phycobank.org/100089>

- *Surirella vasta* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 354: fig. 6, 7. 1922, nom. inval.

***Iconella vasta* var. *linearis* (Hust.) Cocquyt & R. Jahn, comb. nov.**

≡ *Surirella vasta* var. *linearis* Hust. in Huber-Pestalozzi, Phytoplankt. Süsswass. vol. 2 (2), 504. 1942.

Lectotype (cited as holotype but in fact designated by Simonsen 1987). X4/9589 Lake Tanganyika “Tanganyika See. 6”.

<http://phycobank.org/100090>

Updated taxonomy of African *Surirella* taxa

***Surirella afrocalcarata* Cocquyt & R. Jahn, nom. nov.**

≡ *Cymatopleura calcarata* Hust. in Huber-Pestalozzi, Phytoplankt. Süsswass. vol. 2 (2), 480, fig. 579. 1942.

Lectotype (designated by Simonsen 1987). BRM Xa/20 Lake Tanganyika “Tanganika See”.

<http://phycobank.org/100091>

- *Cymatopleura calcarata* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 367: figs 1–2. 1927, nom. inval.

Comment. The epithet name “afrocalcarata” was chosen because of *Suriraya calcarata* Pfitzer in Bot. Abh. Morphol. Physiol. 2: 107. 1871. *Suriraya* is a homotypic synonym of *Surirella* Turpin.

***Surirella clavata* (O. Müll.) Cocquyt & R. Jahn, comb. nov.**

≡ *Cymatopleura solea* var. *clavata* O. Müll. in Bot. Jahrb. Syst. 34: 22, fig. 1. 1904.

≡ *Cymatopleura clavata* (O. Müll.) Cocquyt & R. Jahn in Pl. Ecol. Evol. 147 (3): 413. 2014.

Lectotype (designated by Cocquyt and Jahn 2014). B 40 0040250 [<http://herbarium.bgbm.org/object/B400040250>] (the valve representing the lectotype was published as fig. 1D in Cocquyt and Jahn (2014) “Malawi, Lake Malombe, after the discharge of Lake Malawi”.

<http://phycobank.org/100092>




Figure 6. *Iconella tchadensis*. **A–D** Valves from the holotype slide BR 987 **A–C** valvar views showing the size range **D** girdle view. Scale bar = 10 µm.

***Surirella comperei* (Cocquyt & R. Jahn) Cocquyt & R. Jahn, comb. nov.**

≡ *Cymatopleura comperei* Cocquyt & R. Jahn in Pl. Ecol. Evol. 147 (3): 419, figs 6–8. 2014.

Holotype. B 40 0040184 [<http://herbarium.bgbm.org/object/B400040184>]; the valve representing the holotype was published as fig. 6E in Cocquyt and Jahn (2014) “Malawi, Lake Malawi near Langenburg”.

<http://phycobank.org/100093>

- *Cymatopleura solea* var. *subconstricta* O. Müll. in Bot. Jahrb. Syst. 34: 23. 1904, nom. inval.

- *Cymatopleura solea* var. [*subconstricta*] f. *major* O. Müll. in Bot. Jahrb. Syst. 34: 23. 1904, nom. inval.

- *Cymatopleura solea* var. [*subconstricta*] f. *minor* O. Müll. in Bot. Jahrb. Syst. 34: 23. 1904, nom. inval.

- *Cymatopleura solea* var. [*subconstricta*] f. *minor* O. Müll. in A.W.F.Schmidt, Atlas Diatom.-Kunde, pl. 245: fig. 3. 1904, nom. inval.

***Surirella distinguenda* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 283: fig. 5. 1912.**

Lectotype (cited as holotype but in fact designated by Simonsen 1987). BRM 218/56. “Togo, Westafrika 1912, Lagunenschlick”.

***Surirella fasiculata* O. Müll. in Bot. Jahrb. Syst. 34: 36, pl. 1: fig. 14. 1903.**

Lectotype (designated by Cocquyt and Jahn 2007c). [icon] Müller (1903): pl. 1: fig. 14 “Lake Nogzi, a brackish water lake in the crater of the mountain Nogzi on the northern edge of Kondeland, at 2000 m asl, Tanzania”.

Epitype (designated by Cocquyt and Jahn 2007c). B 40 0040234 [<http://herbarium.bgbm.org/object/B400040234>] (the valve representing the epitype was illustrated as fig. 4 in Cocquyt and Jahn 2007c) “basin near the hot spring of Utengule, Beya Mountain (Tanzania)”.

***Surirella laticeps* (O. Müll.) Cocquyt & R. Jahn, comb. nov.**

≡ *Cymatopleura solea* var. *laticeps* O. Müll. in Bot. Jahrb. Syst. 34: 22-23, fig. 2. 1904.

≡ *Cymatopleura laticeps* (O. Müll.) Cocquyt & R. Jahn in Pl. Ecol. Evol. 147 (3): 418. 2014.

Lectotype (designated in Cocquyt and Jahn 2014). B 40 0040251 [<http://herbarium.bgbm.org/object/B400040251>] (the valve representing the lectotype was published as fig. 5 B in Cocquyt and Jahn (2014) “Nyassaland”, Tanzania, Lake Malawi near Langenburg”.

<http://phycobank.org/100094>

***Surirella modesta* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 357: fig. 8, 9. 1925.**

Lectotype (designated by Simonsen 1987). BRM X2/87 Cameroon “Kamerun, Lagune”.

***Surirella nyansae* (G.S. West) Cocquyt & R. Jahn, comb. nov.**

≡ *Cymatopleura nyansae* G.S. West in J. Linn. Soc. Bot. 38: 167, pl. 8: fig. 8. 1907.

Lectotype (designated in Cocquyt and Jahn 2014). BM 34183 “Tanganyika – In plankton, near Kala (19 Nov. 1904; no. 170).”

<http://phycobank.org/100095>

***Surirella olungensis* Cocquyt & R. Jahn in Cryptog. Algol. 28: 111, figs 7–12, 18–21. 2007.**

Holotype. B 40 0040232 [<http://herbarium.bgbm.org/object/B400040232>] (the valve representing the holotype is illustrated as fig. 7 in Cocquyt and Jahn 2007c)

“Olunga River (Ohmga) in Ussangu northern Mount Kinga, Tanzania (sample B 52 0000058 [<http://herbarium.bgbm.org/object/B520000058>]”).

***Surirella ostentata* Cholnoky in Hydrobiologia 19: 106, 1962.**

≡ *Surirella ovata* var. *africana* Cholnoky in Ber. Deutsch. Bot. Ges. 68: 21–22, fig. 46. 1955.

Lectotype (designated by Cocquyt et al. 2017). UNWH NIWR 191/3802 “Bewässerungskanal bei Vredendal near Olifantsriver”, leg. A.H.P. Engelbrecht.

<http://phycobank.org/100096>

***Surirella ovalis* var. *apiculata* O. Müll. in Bot. Jahrb. Syst. 34: 36, pl. 2: fig. 10. 1903.**

Lectotype (designated by Cocquyt and Jahn 2007c). [icon] Müller (1903): pl. 2: fig. 10 “basin near the hot spring at Utengule, Tanzania”.

= *Surirella ovalis* [var. *apiculata*] f. *minor* O. Müll. in Bot. Jahrb. Syst. 34: 36, pl. 2: fig. 11. 1903.

Lectotype (designated by Cocquyt and Jahn 2007c). [icon] Müller (1903): pl. 2: fig. 11 “Lake Rukwa”.

***Surirella pseudotenuis* Cholnoky in Portugaliae Acta Biol. Sér. B. 4: 226, fig. 120, 1954.**

Lectotype (designated by Cocquyt et al. 2017). UNWH NWU 07–138 “Moss growing on rocks at edge of stram in full sunshine in gully South of road to Vumba, Umtali – 27.7.1952” leg. N.C. Chase.

<http://phycobank.org/100097>

***Surirella rудis* Hust. in Arch. Hydrobiol. Suppl. 15: 505. 1938.**

Lectotype (cited as holotype but in fact designated by Simonsen 1987). BRM X4/3 Lake Tanganyika “Tanganyika See”.

- *Surirella rудis* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 356: fig. 5, 6. 1922, nom. inval.

***Surirella sparsipunctata* Hust. in Huber-Pestalozzi, Phytoplankt. Süßwass. vol. 2 (2): 516, fig. 631. 1942.**

Lectotype (cited as holotype but designated by Simonsen 1987). BRM X4/30 Lake Tanganyika “Tanganikasee. 3rd Tang. Exp., G.S. West”.

- *Surirella sparsipunctata* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 309: fig. 15. 1914, nom. inval.

= *Surirella sparsipunctata* var. *laevis* Hust. in Huber-Pestalozzi, Phytoplankt. Süsswass. vol. 2 (2), 517, fig. 631A. 1942.

Lectotype (designated by Simonsen 1987). BRM X4/34 Lake Tanganyika “Tanganikasee 6”.

Comment. For taxonomical results and discussion see Cocquyt and Vyverman (1993).

Surirella striolata Hust. in Arch. Hydrobiol. 18: 249. 1927.

Lectotype (designated by Simonsen 1987). BRM 224/92 Lake Tanganyika “Tanganika See. Grund. 6, 2”.

- *Surirella striolata* Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 356: fig. 7. 1922, nom. inval.

Surirella subrugosa Cocquyt & R. Jahn, nom. nov.

≡ *Cymatopleura solea* var. *rugosa* O. Müll. in Bot. Jahrb. Syst. 34: 23, fig. 3. 1904.

≡ *Cymatopleura rugosa* (O. Müll.) Cocquyt & R. Jahn in Pl. Ecol. Evol. 147 (3): 416. 2014.

Lectotype (designated in Cocquyt and Jahn 2014). B 40 0040252 [<http://herbarium.bgbm.org/object/B400040252>] (the valve representing the lectotype was published as fig. 3D–E in Cocquyt and Jahn 2014) “Malawi, Lake Malombe, after discharge of Lake Malawi”.

<http://phycobank.org/100098>

Nomenclatural comment. The new epithet was chosen because of *Surirella rugosa* Bramb. & P.B. Ham.

Unrevisable and unrevised taxa

Surirella acanthophora Giffen in Beih. Nova Hedwigia 21: 145, pl. 4: figs 92–95. 1966. Holotype. Giffen collection 30/6 “Fort Hare, Cape Province” South Africa.

Surirella asperrima f. *rokuprensis* Woodhead & Tweed in Rev. Algol. N. S. 5: 144, fig. 4. 1960, nom. inval.

Locality. “Sierra Leone, Rokupr” (Several localities are cited but no type is indicated McNeill et al. 2012, Art. 40.1).

Comment. Taxon unrevisable according to Cocquyt et al. (2013).

Surirella capensis Ehrenb. ex Cocquyt & R. Jahn in Cryptog. Algol. 26: 150. 2005.

Lectotype (designated by Cocquyt and Jahn 2005b). BHUPM 130715 b “Lacus in monte Camdebo Graaf Reinet proximo, Provincia Capensis, Africa Meridionalis”. - *Surirella capensis* Ehrenb., Mikrogeologie 245, 254. 1854, nom. inval.

Taxonomical comment. Species closely related to *Surirella sparsipunctata* and *Iconella anassae*. Further studies are needed, including SEM to evaluate its taxonomic position, which is only possible if material from the type locality can be obtained; otherwise unrevisable.

Surirella cuspidata f. *constricta* Hust. in Explor. Parc Natl. Albert. Mission H. Damas 8: 155, pl. 15: fig. 11. 1949.

Lectotype (cited as holotype but in fact designated by Simonsen 1987). BRM 244/34a DR Congo “Belg. Kongo. 39. Karisimbi-See. +3800 m”.

Comment. Sampling site is located very probably on the Rwandan side of the border with DR Congo.

Taxonomic comment: *S. cuspidata* Hust. in Int. Rev. Hydrobiol. Hydrogr. 42: 156, figs 391–393. 1942, described from Indonesia was transferred to *Stenopterobia cuspidata* (Hust.) Vyverman in Bull. Soc. Bot. Belgique 122: 74. 1989. Further studies are needed to evaluate Hustedt's forma and its taxonomic position.

Surirella engleri f. *densecostata* R. Maillard in Bull. Mus. Natl. Hist. Nat. [Paris], Bot. 30: 39, 43 Pl. 3: fig. 1. 1977, nom. inval.

Localities. Mali “Congo et Mozambique” (Several localities are cited but no type is indicated McNeill et al. 2012, Art. 40.1).

Surirella engleri f. *sierra-leonensis* Woodhead & Tweed in Hydrobiologia 12: 202. 1958, nom. inval.

Locality. Sierra Leone: R. Makoko at Maranda (see Cocquyt et al. 2013).

Surirella fuellebornii var. *worthingtonii* H.Bachm. in Ber. Schweiz. Bot. Ges. 42: 707, 709, pl. 26: fig. 7, 8. 1933.

Locality. Lake Victoria “Victoria Nyanza”.

Surirella gracilis var. *africana* Cholnoky in Hydrobiologia 7: 184, fig. 82, 83. 1955.

Syntype localities. Rayton-vlei 30 km E of Pretoria, South Africa, “Tümpelchen” and “Bächlein” leg. Cholnoky.

Surirella gracilis f. *constricta* Cholnoky in Hydrobiologia 7: 184. 1955.

Type indicated. 10–12 km N of Rayton, 30–35 km NE of Pretoria, South Africa “Leeufonstein Quellen”

Surirella ignota Cholnoky in Nova Hedwigia 2: 118, figs 342, 343. 1960.

Type indicated. “Port Shepstone 362, Kleiner, sickernder Seitenbach des Unzimkulwana-Flusses nahe dem Paddock-Eingange im Oribi Gorge. 22.7.1958.”

Surirella pseudospinifera Iltis in Rev. Algol. 10 (4): 334. 1972, nom. inval.

- *Surirella acanthophora* Iltis in Rev. Algol. 10 (2) 174, figs 10–12, pl. 2: fig. 3, 4. 1971, nom. inval. et nom. illeg. [non Giffen 1966].

Localities. Chad, Mali “Mare du 3e barrage à Bol. 13°30'N, 14°43'30"E. Puits près de la mare de Latir. 13°36'N, 14°44'E” (Two localities are cited but no type is indicated McNeill et al. 2012, Art. 40.1).

Comment. *Surirella pseudospinifera* Iltis was the intended substitute name for *Surirella acanthophora* Iltis [non Giffen 1966] but based on an invalid name (McNeill et al. 2012, Art. 40.1).

Stenopterobia recta Woodhead & Tweed in Hydrobiologia 12: 202, fig. 72. 1958.

Locality. Sierra Leone, Lake Sofon.

Comment. Taxon unrevisable according to Cocquyt et al. (2013).

Surirella rokuprensis Woodhead & Tweed in Rev. Algol. 5: 145, fig. 5. 1960, nom. inval.

Locality. “Sierra Leone, Rokupr” (Several localities are cited but no type is indicated McNeill et al. 2012, Art. 40.1).

Comment. Taxon unrevisable according to Cocquyt et al. (2013).

Surirella rufa var. *sierra-leonensis* Woodhead & Tweed in Rev. Algol. 5: 146, fig. 9. 1960

Type. Mambolo (2352).

Comment. Taxon unrevisable according to Cocquyt et al. (2013).

Surirella rufa [var. *sierra-leonensis*] f. *constricta* Woodhead & Tweed in Rev. Algol. 5: 146, fig. 7. 1960, nom. inval.

Locality. Sierra-Leone (Several localities are cited but no type is indicated McNeill et al. 2012, Art. 40.1).

Comment. Taxon unrevisable according to Cocquyt et al. (2013).

Surirella scutum Reichelt in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 295: fig. 4. 1913, nom. inval.

Locality. Kalahari.

Comment. Type and description of the depicted species are missing.

Surirella subrobusta Hust. in A.W.F. Schmidt, Atlas Diatom.-Kunde, pl. 353: fig. 1. 1922.

Lectotype (cited as holotype but in fact designated by Simonsen 1987). BRM 4/59 “Lafirio-Fluß. Deutsch-O-Afrika” (Simonsen 1987).

Comment. Description in the caption of the plate.

Surirella schweickerdtii Cholnoky in Bot. Not. 1954: 290, figs 95, 96. 1954.

≡ *Stenopterobia schweickerdtii* (Cholnoky) Brassac, T.Ludwig & Torgan in Diatom Research 18: 186. 2003.

Locality. “Moosrasen auf einer kleinen Insel zwischen Gras. Debegeni” South Africa.

Surirella taiaensis J.R. Carter & Denny, Beih. Nova Hedwigia 73: 325, pl. 8: fig. 274. 1982.
Holotype. BM 78108 “Sierra Leone, River Jong (Taia) at Njala”.

Surirella tenera var. *minor* Cholnoky in Portugaliae Acta Biol. Sér. B, 6: 140 fig. 168. 1958.
Holotype. FR 118 “In rivulo apud Modderpoort prope oppidum Nylstroom (Transvaal)”.

Surirella welshii Cholnoky 1962 in Hydrobiologia 20: 337, fig. 45. 1962.
Type indicated. “Unnamed mountain stream between Piggs Peak and Mbabane,
3.7.1961, leg. H. Welsh”, “Swaziland”.

Conclusion

55 taxa – formerly ranked within *Surirella* - have been transferred to *Iconella*; most of these have been shown to be endemic (Ross 1983, Cocquyt et al. 1993, Cocquyt and Vyverman 1994, Cocquyt 1998, 2000,) and many of them, especially the large species, have become planktonic in the East African great lakes (Müller 1905, Hustedt in Huber-Pestalozzi 1942, Cocquyt 1998). In addition, two taxa – formerly ranked within *Stenopterobia* – have been transferred to *Iconella*. 10 taxa have stayed within *Surirella*, (although the position of *S. sparsipunctata* has to be genetically verified), and six taxa have been transferred from *Cymatopleura* to *Surirella*. For completeness sake, 21 taxa have been listed which are either unrevised or unrevisable because missing morphological data do not allow us to decide if the raphe is raised on a keel.

When more taxa from the genera *Iconella* and *Surirella* have been studied molecularly, especially the endemic species from Africa and other tropical regions, further autapomorphies might be discovered which might support the differentiation into further groups. With the currently available data, the solution by Ruck et al (2016a, b) clarifies their phylogeny and presents a very workable approach.

Acknowledgements

We wish to thank Ingrid Jüttner (Cardiff, United Kingdom) and Jonathan C. Taylor (Potchefstroom and Grahamstown, South Africa) for contributing to the respective new combinations. The authors gratefully acknowledge the Andrew Mellon Foundation for funding the “Pilot project on the digitization of African micro-algae types and typical specimens: the diatom family Surirellaceae”, and the EU project SYNTHESYS for financial support to CC for two research award at the BGBM in 2005 and 2015. Marco Cantonati supplied us with a missing reference, and RJ and WHK wish to thank Dr. David Lazarus for providing access to the Ehrenberg Collection (BHUPM). Additionally, they acknowledge DFG Grant JA 874/8-1 for Names Registration PhycoBank.

References

- Agardh CA (1824) *Systema algarum, Literis Berlingianis.* 312 pp.
- Agardh CA (1830) *Conspectus criticus diatomacearum.* Lund. 1-2: 1–38.
- Bachmann H (1933) Phytoplankton von Victoria Nayza-, Albert Nyanza- und Kiogasee gesammelt von E. B. Worthington. Berichte der Schweizerischen Botanischen Gesellschaft. 42: 705–717, pl. 23–26.
- Barkworth ME, Watson M, Barrie FR, Belyaeva IV, Chung RCK, Dašková J, Davidse G, Dönmez AA, Doweld AB, Dressler S, Flann C, Gandhi K, Geltman D, Glen HF, Greuter W, Head MJ, Jahn R, Janarthanam MK, Katinas L, Kirk PM, Klazenga N, Kusber WH, Kvaček J, Malécot V, Mann DG, Marhold K, Nagamasu H, Nicolson N, Paton A, Patterson DJ, Price MJ, Prud'homme van Reine W, Schneider CW, Sennikov A, Smith GF, Stevens PF, Yang ZL, Zhang XC, Zuccarello GC (2016) Report of the Special Committee on Registration of Algal and Plant Names (including fossils). *Taxon* 65 (3): 670–672. Report of the Special Committee on Registration of Algal and Plant Names (including fossils). *Taxon* 65(3): 670–672. <https://doi.org/10.12705/653.43>
- Bogaerts A, de Haan M, Van de Vijver B, Cocquyt C (2014) A complete catalogue of algal taxa described by Pierre Compère. *Plant Ecology and Evolution* 147(3): 311–324. <https://doi.org/10.5091/plecevo.2014.993>
- Bory de Saint Vincent JBM (1822–1831) *Dictionnaire classique d'Histoire Naturelle.* Paris.
- Carter JR, Denny P (1982) Freshwater algae of Sierra Leone III. Bacillariophyceae. Beihefte zur Nova Hedwigia 73: 281–331.
- Cholnoky BJ (1954) Diatomeen aus Süd-Rhodesien. *Portugaliae Acta Biologica (Série B)* 4: 197–227.
- Cholnoky BJ (1954) Diatomeen und einige andere Algen aus dem „de Hoek“-Reservat in Nord-Transvaal. *Botaniska Notiser* 1954: 269–296.
- Cholnoky BJ (1955) Hydrobiologische Untersuchungen in Transvaal I. Vergleichung der herbstlichen Algengemeinschaften in Rayton-vlei und Leeufontein. *Hydrobiologia* 7: 137–209. <https://doi.org/10.1007/BF00046690>
- Cholnoky BJ (1956) Neue und seltene Diatomeen aus Afrika II. Diatomeen aus dem Tugela-Gebiete in Natal. *Österreichische Botanische Zeitschrift* 103: 53–97. <https://doi.org/10.1007/BF01287833>
- Cholnoky BJ (1957) Neue und seltene Diatomeen aus Afrika III. Diatomeen aus dem Tugela-Flussystem, hauptsächlich aus den Drakensbergen in Natal. *Österreichische Botanische Zeitschrift* 104: 25–99. <https://doi.org/10.1007/BF01289120>
- Cholnoky BJ (1958) Beiträge zur Kenntnis der Südafrikanischen Diatomeenflora II. Einige Gewässer im Waterberg-Gebiet, Transvall. *Portugaliae Acta Biologica (Série B)* 6: 99–160.
- Cholnoky BJ (1960) Beiträge zur Kenntnis der Diatomeenflora von Natal (Südafrika). *Nova Hedwigia* 2: 1–128, pl. 1–9.
- Cholnoky BJ (1960) Diatomeen aus einem Teiche am Mt. Kenya in Mittelafrika. *Österreichische Botanische Zeitschrift* 107: 351–365. <https://doi.org/10.1007/BF01289759>
- Cocquyt C (1998) Diatoms from the northern basin of Lake Tanganyika. *Bibliotheca Diatomologica* 39: 1–276.

- Cocquyt C (2000) Biogeography and Species Diversity of Diatoms in the Northern Basin of Lake Tanganyika. Advances in Ecological Research 31: 125–150. [https://doi.org/10.1016/S0065-2504\(00\)31010-8](https://doi.org/10.1016/S0065-2504(00)31010-8)
- Cocquyt C, Jahn R (2005a) Rare *Surirella* taxa (Bacillariophyta) from East Africa described by Otto Müller: typifications, recombinations, new names, annotations and distributions. Willdenowia 35: 359–371. <https://doi.org/10.3372/wi.35.35218>
- Cocquyt C, Jahn R (2005b) *Surirella capensis* Ehrenberg ex Cocquyt & R.Jahn, spec. nov., a rare and historical diatom species from South Africa. Cryptogamie Algologie 26: 149–153.
- Cocquyt C, Jahn R (2007a) *Surirella engleri* O. Müller – a study of its types, infraspecific variability and distribution. Diatom Research 22: 1–16. <https://doi.org/10.1080/0269249X.2007.9705691>
- Cocquyt C, Jahn R (2007b) *Surirella fuellebornii* (Bacillariophyta) and related taxa: lectotypification and distribution. Systematics and Geography of Plants 77(2): 213–228.
- Cocquyt C, Jahn R (2007c) Taxa related to *Surirella ovalis* from Tanzania, East Africa: Typification of O. Müller's taxa and description of a new species. Cryptogamie Algologie 28: 107–116.
- Cocquyt C, Jahn R (2007d) *Surirella nyassae* O. Müller, *S. malombae* O. Müller, *S. chepurnovii* Cocquyt & R. Jahn sp. nov. (Bacillariophyta) – typification and variability of three closely related East African diatoms. Nova Hedwigia 84: 529–548. <https://doi.org/10.1127/0029-5035/2007/0084-0529>
- Cocquyt C, Jahn R (2014) A re-investigation of Otto Müller's *Cymatopleura* taxa (Bacillariophyta) from East Africa. Plant Ecology and Evolution 147(3): 412–425. <http://dx.doi.org/10.5091/plecevo.2014.992>
- Cocquyt C, Jüttner I, Kusber WH (2013) Reinvestigation of West African Surirellaceae (Bacillariophyta) described by Woodhead & Tweed from Sierra Leone (West Africa). Diatom Research 28: 121–129. <https://doi.org/10.1080/0269249X.2012.752411>
- Cocquyt C, Kusber WH (2010) Reinvestigation of West African Surirellaceae (Bacillariophyta) described by Foged from Ghana. Nova Hedwigia 91: 111–136. <https://doi.org/10.1127/0029-5035/2010/0091-0111>
- Cocquyt C, Taylor JC (2015) New and interesting *Surirella* taxa (Surirellaceae, Bacillariophyta) from the Congo Basin (DR Congo). European Journal of Taxonomy 133: 1–15. <https://doi.org/10.5852/ejt.2015.133>
- Cocquyt C, Taylor JC, Kusber WH (2017) Reinvestigation of African *Surirella* taxa (Bacillariophyta) described by B.J. Cholnoky with some remarks on digitization of diatom types. Fottea 17(1): 34–56. <https://doi.org/10.5507/fot.2016.012>
- Cocquyt C, Taylor JC, Wetzel CE (2014) *Stenopterobia cataractarum* sp. nov. (Bacillariophyta), a new benthic diatom from a waterfall in Zambia, Africa. Phytotaxa 158(1): 76–84. <https://doi.org/10.11646/phytotaxa.158.1.5>
- Cocquyt C, Vyverman W (1993) *Surirella sparsipunctata* Hustedt and *S. sparsipunctata* var. *laevis* Hustedt, a light and electron microscopical study. Hydrobiologia 269/270: 97–101. <https://doi.org/10.1007/BF00028009>
- Cocquyt C, Vyverman W (1994) Composition and diversity of the algal flora in the East African Great Lakes: a comparative survey of lakes Tanganyika, Malawi (Nyasa) and Victoria. Archiv für Hydrobiologie Beiheft Ergebn. Limnol. 44: 161–172.

- Cocquyt C, Vyverman W, Compère P (1993) A checklist of the algal flora of the East African Great Lakes: Lake Malawi, Lake Tanganyika and Lake Victoria. *Scripta Botanica Belgica* 8: 1–56.
- Compère P (1975) Diatomées de la région du lac Tchad, taxons, combinaisons et noms nouveaux. *Bulletin du Jardin Botanique National de Belgique* 45: 373–382, 26 figs.
- Ehrenberg CG (1832) Über die Entwicklung und Lebensdauer der Infusionsthiere nebst ferneren Beiträgen zu einer Vergleichung ihrer organischen Systeme. *Abhandlungen der Königlichen Akademie der Wissenschaften in Berlin* 1831: 1–154, Tafel 1–4.
- Ehrenberg CG (1834) Dritter Beitrag zur Erkenntniss grosser Organisation in der Richtung des kleinsten Raumes. *Abhandlungen der Königlichen Akademie der Wissenschaften in Berlin* 1833: 145–336, Tafel 1–11. 1834.
- Ehrenberg CG (1837) Über ein aus fossilen Infusorien bestehendes, 1832 zu Brod verbackenes Bergmehl von den Grenzen Lapplands in Schweden. Bericht über die zur Bekanntmachung geeigneten Verhandlungen der Königlich Preussischen Akademie der Wissenschaften zu Berlin 1837: 43–45.
- Ehrenberg CG (1838) Die Infusionsthierchen als vollkommene Organismen. Leipzig.
- Ehrenberg CG (1840) Characteristik von 274 neuen Arten von Infusorien. Bericht über die zur Bekanntmachung geeigneten Verhandlungen der Königlich Preussischen Akademie der Wissenschaften zu Berlin. 1840: 197–190.
- Ehrenberg CG (1842) Über fossile Infusorien aus Irland. Bericht über die zur Bekanntmachung geeigneten Verhandlungen der Königlich Preussischen Akademie der Wissenschaften zu Berlin, 1842: 321–339.
- Ehrenberg CG (1845a) Über vulkanischen Infusorien-Tuff am Rhein, auf der Insel Ascension, in Patagonien und im Phonolith von Wisterschan, und über einen anscheinend organischen Einschluss im Trachyt vom Zimspan. Bericht über die zur Bekanntmachung geeigneten Verhandlungen der Königlich Preussischen Akademie der Wissenschaften zu Berlin, 1845: 133–149.
- Ehrenberg CG (1845b) Über die kleinsten Lebensformen in Portugal und Spanien. Bericht über die zur Bekanntmachung geeigneten Verhandlungen der Königlich Preussischen Akademie der Wissenschaften zu Berlin. 1845: 304–308.
- Ehrenberg CG (1854) Mikrogeologie. Das Erden und Felsen schaffende Wirken des unsichtbar kleinen selbstständigen Lebens auf der Erde. Atlas. Leipzig.
- Foged N (1966) Freshwater diatoms from Ghana. *Biologiske Skrifter Det Kongelige Danske Videnskarbernes Selskab* 15: 1–169, 25 pls.
- Forti A (1910) Contribuzioni diatomologiche. X. Diatomaceae quaternariae e subfossili d'acqua dolce raccolte in Etiopia dal dott. Giovanni Negri. *Atti del Reale Istituto Veneto Scienze, Lettere & Arti* 69: 1284, 1273–1303.
- Giffen MH (1966) Contributions to the diatom flora of Southern Africa. II. Diatoms from the Hog's Back Region of the Amatola Mountains, Eastern Cape Province, South Africa. *Nova Hedwigia Beiheft* 21: 123–150, pl. 1–4.
- Gmelin JF (1791) In *Systema Naturae*. Edition 13 1(6). Lipsiae.
- Güntsch A, Hyam R, Hagedorn G, Chagnoux S, Röpert D, Casino A, Droege G, Glöckler F, Gödderz K, Groom Q, Hoffmann J, Holleman A, Kempa M, Koivula H, Marhold K, Nicolson N, Smith VS, Triebel D (2017) Actionable, long-term stable and semantic web

- compatible identifiers for access to biological collection objects. Database (Oxford) 2017 (1): bax003. <https://doi.org/10.1093/database/bax003>
- Hassall AH (1845) A history of the British freshwater algae. 1–2: 1–464, 103 pl.
- Heurck H (1896) A treatise on the Diatomaceae. London, William Wesley & Son, 558 pp.
- Hofmann G, Lange-Bertalot H, Werum M (2013) Diatomeen im Süßwasser-Benthos von Mitteleuropa. Bestimmungsflora Kieselalgen für die ökologische Praxis. Über 700 der häufigsten Arten und ihre Ökologie. Königstein, Koeltz Scientific Books.
- Huber-Pestalozzi G (1942) Das Phytoplankton des Süßwassers. In: Thienemann A (ed.): Die Binnengewässer, 16(2/2). Schweizerbart'sche Verlag, Stuttgart.
- Hustedt F (1912) Tafeln 281–284 in A. Schmidts Atlas Diatomaceenkundekunde 71. Leipzig, Reisland.
- Hustedt F (1913) Tafeln 293–300 in A. Schmidts Atlas Diatomaceenkundekunde 74–75. Leipzig, Reisland.
- Hustedt F (1914) Tafeln 305–312 in A. Schmidts Atlas Diatomaceenkundekunde 77–78. Leipzig, Reisland.
- Hustedt F (1921) VI. Bacillariales in Schröder B: Zellpflanzen Ostafrikas, gesammelt auf der Akademischen Studienfahrt 1910. *Hedwigia* 63: 117–173, pl. 1.
- Hustedt F (1922) Tafeln 353–356 in A. Schmidts Atlas Diatomaceenkundekunde 89. Leipzig, Reisland.
- Hustedt F (1925) Tafeln 357–360 in A. Schmidts Atlas Diatomaceenkundekunde 90. Leipzig, Reisland.
- Hustedt F (1927) Tafeln 365–368 in A. Schmidts Atlas Diatomaceenkundekunde 92. Leipzig, Reisland.
- Hustedt F (1938) Systematische und ökologische Untersuchungen über die Diatomeenflora von Java, Bali und Sumatra nach dem Material der Deutschen Limnologischen Sunda-Expedition. Teil I. Systematischer Teil, Schluß. Archiv für Hydrobiologie, Supplement 15: 393–506.
- Hustedt F (1949) Süßwasser-Diatomeen aus dem Albert-Nationalpark in Belgisch-Kongo. In: Exploration du Parc National Albert, Mission H. Damas 8: 1–199 +16 pls.
- Iltis A (1971) Algues nouvelles des mares du Kanem (Tchad). Revue Algologique 10(2): 172–176, pl. 2.
- Iltis A (1972) Remarques taxinomiques sur les algues des Mares du Kanem (Tchad). Revue Algologique 10(4): 334.
- Index Nominum Algarum (1988+) University Herbarium, University of California, Berkeley. Compiled by Paul Silva. Available online at <http://ucjeps.berkeley.edu/CPD/> [accessed 2017-03-10]
- Jahn R, Kusber WH (2004) Algae of the Ehrenberg Collection - 1. Typification of 32 names of diatom taxa described by C.G. Ehrenberg. Willdenowia 34: 577–595. <https://doi.org/10.3372/wi.34.34219>
- Jahn R, Kusber WH (2005+) AlgaTerra Information System. <http://www.algaterra.org> [accessed 2017-03-10]
- Jurilij A (1949) Nove dijatomeje – Surirellaceae – iz Ohridskog jezera i njihovo filogenetsko značenje. Prirodoslovna Istraživanja 24: 171–260.

- Krammer K, Lange-Bertalot H (1988) Bacillariophyceae. 2. Teil: Bacillariaceae, Epithemiaceae, Surirellaceae. In: Ettl H, Gerloff J, Heynig H, Mollenhauer D (Eds) Süßwasserflora von Mitteleuropa 2/2. Gustav Fischer Verlag, Stuttgart, New York.
- Kützing FT (1844) Die Kieselchaligen Bacillarien oder Diatomeen. W. Köhne, Nordhausen, [i-vii], [1]-152, pls 1–30. <https://doi.org/10.5962/bhl.title.64360>
- Kützing FT (1833) Synopsis Diatomacearum oder Versuch einer systematischen Zusammenstellung der Diatomeen. Linnaea 8: 529–620, pl. 13–19.
- Kusber WH, Kohlbecker A, Güntsch A, Berendsohn WG, Jahn R (2017) PhycoBank - Registration of nomenclatural acts for diatomists and diatom research. In: Kopálová K, Bishop J, Van de Vijver B (Eds) Program & Abstracts, 11th Central European Diatom meeting, Prague, Czech Republic, 22–25 March 2017. Charles University, Prague, 40–41. <https://www.ced2017.eu/files/2017/03/ABSTRACT-book.pdf>
- Lyngbye HC (1819) Tentamen hydrophytologiae Danicae continens omnia hydrophyta cryptogama Daniae, Holsatiae, Foeroeae, Islandiae, Groenlandiae hucusque cognita, systematae disposita, descripta et iconibus illustrata, adjectis simul speciebus Norvegicis. Hafniae, 248 pp, 70 pl.
- Maillard R (1977) Diatomées d'eau douce du Mali, Afrique. Bulletin du Muséum National d'Histoire Naturelle [Paris], Botanique 30: 17–45.
- Mayer A (1913) Die Bacillarien der Regensburger Gewässer. Berichte des Naturwissenschaftlichen Vereins zu Regensburg 14: 412 pp, 30 pl.
- McNeill J, Barrie FR, Buck WR, Demoulin V, Greuter W, Hawksworth DL, Herendeen PS, Knapp S, Marhold K, Prado J, Prud'homme van Reine WF, Smith GF, Wiersema JH, Turland NJ (2012) International Code of Nomenclature for algae, fungi and plants (Melbourne Code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011. *Regnum Vegetabile* 154: 1–208.
- Müller O (1904a) Bacillariaceen aus dem Nyassalande und einigen benachbarten Gebieten. I. Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie 34: 9–38.
- Müller O (1904b) Bacillariaceen aus dem Nyassalande und einigen benachbarten Gebieten. II. Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie 34: 256–301.
- Müller O (1905) Bacillariaceen aus dem Nyassalande und einigen benachbarten Gebieten. III. Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie 36: 137–205.
- Müller O (1910) Bacillariaceen aus dem Nyassalande und einigen benachbarten Gebieten. IV. Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie 45: 69–122.
- Ostenfeld CH (1908) Phytoplankton aus dem Victoria Nyanza. Sammelausbeute von A. Borgert, 1904–1905. VIII. Abhandlung. Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie 41: 330–350.
- Ostenfeld CH (1909) Notes on the Phytoplankton of Victoria Nyanza, East Africa. Bull. Museum of Compar. Zoology Harvard Coll. 52(10): 171–181.

- Pouličková A, Jahn R (2007) *Campylodiscus clypeus* (Ehrenberg) Ehrenberg ex Kützing: Typification, morphology and distribution. Diatom Research 22: 135–146. <https://doi.org/10.1080/0269249X.2007.9705701>
- Ross R (1983) Endemism and cosmopolitanism in the diatom flora of the East African Great Lakes. In: Sims RW, Price JH, Whalley PES (Eds) Evolution, time and space. Syst. Assoc. Special 23, 40–41.
- Round FE, Crawford RM, Mann DG (1990) The diatoms. Biology, and morphology of the genera. Cambridge University Press, 747 pp.
- Ruck EC, Guiry MD (2016) Validation of the diatom genus name *Coronia* (Surirellaceae, Bacillariophyta). Notulae algarum 11: 1–3.
- Ruck EC, Kociolek JP (2004) Preliminary phylogeny of the family Surirellaceae (Bacillariophyta). Bibliotheca Diatomologica 50: 1–236.
- Ruck EC, Nakov T, Alverson AJ, Theriot EC (2016a) Phylogeny, ecology, morphological evolution, and reclassification of the diatom orders Surirellales and Rhopalodiales. Molecular Phylogenetics and Evolution 103: 155–171. <http://dx.doi.org/10.1016/j.ympev.2016.07.023>
- Ruck EC, Nakov T, Alverson AJ, Theriot EC (2016b) Nomenclatural transfers associated with the phylogenetic reclassification of the Surirellales and Rhopalodiales. Notulae Algarum 10: 1–4.
- Ruck EC, Theriot EC (2011) Origin and Evolution of the Canal Raphe System in Diatoms. Protist 162: 723–737. <https://doi.org/10.1016/j.protis.2011.02.003>
- Schoeman FR, Archibald REM (1979) The diatom flora of Southern Africa. CSIR Special Report WAT 50(5). Pretoria.
- Simonsen R (1987) Atlas and catalogue of the diatom types of Friedrich Hustedt. 1–3. J. Cramer, Berlin & Stuttgart.
- Smith W (1851) Notes on the Diatomaceae, with descriptions of British species included in the genera *Campylodiscus*, *Surirella*, *Cymatopleura*. Annals and magazine of natural history, including zoology, botany, and geology. London, 7: 1–14, 3 pl.
- Spaulding S, Edlund M (2008) *Cymatopleura*. In Diatoms of the United States. <http://westerndiatoms.colorado.edu/taxa/genus/Cymatopleura> [accessed 2017-03-06]
- Spaulding S, Edlund M (2010) *Stenopterobia*. In Diatoms of the United States. <http://westerndiatoms.colorado.edu/taxa/genus/stenopterobia> [accessed 2017-03-06]
- The International Plant Names Index (2017) <http://www.ipni.org> [accessed 2017-03-10]
- Turpin PJF (1828) Observations sur le nouveau Genre *Surirella*. Mémoires du muséum d'histoire naturelle, Paris 16: 361–368.
- Vyverman W (1989) Diatoms (Bacillariophyta) from Mount Giluwe (Southern Highlands Province, Papua New Guinea). Bulletin de la Société Royale de Botanique de Belgique 122: 61–80, 2 pl.
- West GS (1907) Report on the freshwater algae, including phytoplankton, of the Third Tanganyika Expedition conducted by Dr. W.A. Cunnington, 1901–1905. Journal of the Linnean Society, Botany 38: 81–197. <http://dx.doi.org/10.1111/j.1095-8339.1907.tb00848.x>

- Wiersema JH, McNeill J, Turland NJ, Barrie FR, Buck WR, Demoulin V, Greuter W, Hawksworth DL, Herendeen PS, Knapp S, Marhold K, Prado J, Prud'homme Van Reine WF, Smith GF (2015) International Code of Nomenclature for algae, fungi and plants (Melbourne Code). Appendices II-VIII. *Regnum Vegetabile* 157: 1–492.
- Woodhead N, Tweed TD (1958) Freshwater algae of Sierra Leone. I. New and unusual algae from the Sula Hills. *Hydrobiologia* 12: 181–225. <https://doi.org/10.1007/BF00034148>
- Woodhead N, Tweed TD (1960) Freshwater algae of Sierra Leone 3. The algae of Rokupr and Great Scarcies River. *Revue Algologique* 5: 116–150, pls 14–15.