

Synopsis of *Acalypha* (Euphorbiaceae) of continental Ecuador

José María Cardiel Sanz¹, Pablo Muñoz Rodríguez¹

¹ Departamento de Biología, Facultad de Ciencias, Universidad Autónoma de Madrid. Ciudad Universitaria de Cantoblanco, Postal Code 28049, Madrid, Spain

Corresponding author: José María Cardiel Sanz (jm.cardiel@uam.es)

Academic editor: Dmitry Geltman | Received 4 April 2012 | Accepted 20 August 2012 | Published 11 September 2012

Citation: Cardiel JM, Muñoz P (2012) Synopsis of *Acalypha* (Euphorbiaceae) of continental Ecuador. PhytoKeys 17: 1–17. doi: 10.3897/phytokeys.17.3190

Abstract

A critical review of the Ecuadorian species of *Acalypha* L. (Euphorbiaceae) is presented; 20 of the 38 previously recognized species are accepted, 9 are considered synonyms and 9 are based on misidentifications. Comprehensive nomenclatural information is supplied and 13 lectotypes are designated. An identification key is also provided.

Resumen

Se presenta una revisión crítica de las especies ecuatorianas de *Acalypha* L. (Euphorbiaceae); se aceptan 20 de las 38 especies previamente reconocidas, 9 se consideran sinónimos y otras 9 están basadas en identificaciones erróneas. Se aporta una exhaustiva información nomenclatural y se designan 13 lectótipos. Se incluye también una clave de identificación.

Keywords

Acalypha, Ecuador, Euphorbiaceae, lectotypification, species identification

Introduction

Acalypha L. is the third largest genus in the Euphorbiaceae sensu stricto (after *Euphorbia* L. and *Croton* L.). It comprises c. 500 species found mostly in the tropics worldwide, although some reach temperate regions. The Americas are home to two thirds of the species, from southeastern Canada and United States to Uruguay and northern Argentina. They thrive in a wide variety of habitats, from tropical rainforests to subde-

sertic areas, and from sea level up to 4000 meters of altitude. *Acalypha* belongs to subfamily Acalyphoideae, the most diverse and complex in the Euphorbiaceae (Hayden and Hayden 2000). This subfamily appears to be paraphyletic, but the central group of taxa, Acalyphoideae Beilschm. sensu stricto, is clearly monophyletic. Molecular analysis support the monophyly of *Acalypha* (Tokuoka 2007, Wurdack and Davis 2009).

Despite its great diversity, *Acalypha* is one of the lesser known genera of the Euphorbiaceae. The last treatment of the whole genus was made by Pax and Hoffman (1924), on which 18 species were recorded in Ecuador (Table 1). The state of the knowledge of the genus in South America includes updated national floristic treatments (Cardiel 1995, 1999, Levin 2001), and checklists (Bacigalupo and Mulgura 1999, Berry et al. 2007, Brako 1993, Cardiel 2007, 2010, Levin 2008). Regarding Ecuador, the only complete work on *Acalypha* is the treatment of Euphorbiaceae for the Catalogue of the Vascular Plants of Ecuador (Webster 1999), which recognized 34 species for the continental land of Ecuador (and four other from Galapagos Island). This work was updated by Ulloa Ulloa and Neil (2005), who added another four species of *Acalypha*, but none in the second update (Neil and Ulloa Ulloa 2011). There are also several regional or thematic floras mentioning *Acalypha* species (Bonifaz and Cornejo 2004, Cerón et al. 2007, De la Torre et al. 2008, León-Yánez et al. 2011, Madsen et al. 2001, Valencia et al. 2000). In addition Cardiel (2000) described two new species and proposed several new synonyms for Ecuadorean *Acalypha*.

This work presents a reviewed critical synopsis of the species of *Acalypha* for continental Ecuador and provides a key to help identification.

Materials and methods

We studied 987 Ecuadorian collections of *Acalypha* from the following herbaria: A, AAU, B, BM, COL, DAV, F, G, GB, GH, HAL, HBG, JE, K, L, M, MA, MO, NY, P, PR, QCA, QCNE, S, SEL, U, UPS, US, W, WRSL and Z (acronyms according to Thiers 2011). We reviewed all the collections cited by Webster (1999) and Ulloa Ulloa and Neill (2005), solving the doubts raised by some names. We also found a large number of type specimens, clarifying the identity of many names. Typifications were made after a carefully review of the original literature on the taxa, and examination of the nomenclatural types. Where no holotype was indicated, or it has been lost or destroyed, a lectotype is designated according International Code of Botanical Nomenclature rules and recommendations (McNeill et al. 2006).

The structure of the checklist follows, in general terms, those of Brako and Zarucchi (1993) and Webster (1999). The accepted species are cited in alphabetical order, including original publications, homotypic synonyms and nomenclatural synonyms based on Ecuadorian collections. For each name, the information concerning the type collections is included, with studied specimens indicated with an exclamation mark (!). Then, we summarize information about habit, habitat and altitudinal range in 250 meters intervals; this information was obtained exclusively from the studied specimens. We follow the geographic regions proposed by Jørgensen and León-Yánez (1999).

Table 1. Species of *Acalypha* cited for Ecuador by Pax and Hoffmann (1924), Webster (1999), and in this work; the vouchers refers to cited by Webster (1999).

| Pax and Hoffmann (1924) | Webster (1999) | In this work |
|--|--|--|
| <i>A. alopecuroides</i> Jacq. | accepted | |
| <i>A. amentacea</i> Roxb. subsp. <i>wilkesiana</i> (Fosb.) Müll. Arg. accepted | = <i>Acalypha wilkesiana</i> Müll. Arg. (fide Sagun et al. 2010) = <i>A. padifolia</i> Kunth (fide Cardiel 2000) | The voucher cited (P. Jørgensen 1551) corresponds to <i>A. padifolia</i> Kunth. <i>A. argomuelleri</i> has not been found in Ecuador. |
| <i>A. andina</i> Müll. Arg. | | |
| <i>A. argomuelleri</i> Briq. | | The voucher cited (P. Jørgensen 1449) corresponds to <i>A. padifolia</i> Kunth. <i>A. aronioides</i> has not been found in Ecuador. |
| <i>A. aronioides</i> Pax & K. Hoffm. accepted | accepted | The voucher cited (Berg & Akkermans 1016) corresponds to <i>A. stachyura</i> Pax. <i>A. benensis</i> is a synonym of <i>A. stricta</i> Poepp., not found in Ecuador (fide Cardiel 2007) |
| <i>A. arvensis</i> Poepp. | accepted | Doubtful taxon not found in Ecuador (fide Cardiel 2007) |
| <i>A. benensis</i> Britton | | |
| <i>A. brachyclada</i> Müll. Arg. | accepted | |
| <i>A. cuneata</i> Poepp. | accepted | |
| <i>A. cuspidata</i> Poepp. | accepted | |
| <i>A. dicynoneura</i> Müll. Arg. accepted | accepted | |
| <i>A. diversifolia</i> Jacq. accepted | accepted | |
| <i>A. ecuadorica</i> Pax & K. Hoffm. accepted | | = <i>A. schiediana</i> Schlechl. (fide Cardiel 2000) |
| <i>A. eggersii</i> Pax & K. Hoffm. accepted | | = <i>A. cuneata</i> Poepp. (fide Cardiel 1995) |
| <i>A. heterolepis</i> Müll. Arg. = <i>A. macrostachya</i> Jacq. | | = <i>A. macrostachya</i> Jacq. (fide Cardiel 1995) |
| <i>A. hispida</i> Burm. f. accepted | accepted | |
| <i>A. infesta</i> Poepp. not cited | accepted | |
| <i>A. macbridei</i> I.M. Johnst. | | = <i>A. salicifolia</i> Müll. Arg. (fide Cardiel 2000) |
| <i>A. macrodonia</i> Müll. Arg. accepted | | = <i>A. padifolia</i> Kunth (fide Cardiel 2007) |
| <i>A. macrostachya</i> Jacq. | accepted | accepted |
| <i>A. mapirensis</i> Pax | | The voucher cited (C. Cerón & C. Igúago 5461) correspond to <i>A. stachyura</i> Pax. <i>A. mapirensis</i> is a synonym of <i>A. stricta</i> Poepp., not found in Ecuador (fide Cardiel 2007) |
| <i>A. ostryfolia</i> Riddell | | The voucher cited (Dodson & al. 7109) corresponds to <i>A. subastrata</i> Aresch. <i>A. ostryfolia</i> has not been found in Ecuador |

| Pax and Hoffmann (1924) | Webster (1999) | In this work |
|---------------------------------------|--|---|
| <i>A. platyphylla</i> Müll. Arg. | <i>A. padifolia</i> Kunth accepted | accepted |
| <i>A. ruiziana</i> Müll. Arg. | accepted | The voucher cited (Sodiro 151) corresponds to <i>A. dicynoneura</i> Pax. <i>A. ruiziana</i> is a synonym of <i>A. padifolia</i> Kunth (fide Cardiel 2007) |
| <i>A. salicifolia</i> Müll. Arg. | accepted | accepted |
| | <i>A. scandens</i> Benth. | accepted |
| | <i>A. schimpffii</i> Diels | <i>A. schiediana</i> Schleidl. (cited first time in Ecuador by Cardiel (2000)) = <i>Acalypha padifolia</i> Kunth (fide Cardiel 2000) |
| <i>A. setosa</i> A. Rich. | accepted | The voucher cited (Madsen 63051) corresponds to <i>A. subcaerulea</i> Aresch. <i>A. setosa</i> has not been found in Ecuador |
| | <i>A. stachyura</i> Pax. | accepted |
| | | <i>A. stellata</i> Cardiel (fide Cardiel 2000) |
| <i>A. stellipila</i> Pax & K. Hoffm. | = <i>Acalypha dicynoneura</i> Müll. Arg. | = <i>Acalypha dicynoneura</i> Müll. Arg. |
| | <i>A. stenoloba</i> Müll. Arg. | The voucher cited (Van der Werff & W. Palacios 10355) corresponds to <i>A. stachyura</i> Pax. <i>A. stenoloba</i> has not been found in Ecuador |
| | <i>A. subandina</i> Ule | = <i>A. platyphylla</i> Müll. Arg. (fide Cardiel 1995) |
| | accepted | accepted |
| <i>A. subcaerulea</i> Aresch. | | |
| <i>A. renipes</i> Pax & K. Hoffm. | accepted | = <i>A. cuspidata</i> Jacq. (fide Cardiel 2000) |
| <i>A. ninguraguae</i> Pax & K. Hoffm. | accepted | = <i>A. cuspidata</i> Jacq. (fide Cardiel 2000) |
| <i>A. villosa</i> Jacq | accepted | accepted |
| | | <i>A. websteri</i> Cardiel (fide Cardiel 2000) |

They defined three regions for continental Ecuador: Coastal, Andean and Amazonian. The Coastal region is defined as below 1000 meters elevation, from west of the Andes to the coast, while the Amazonian region is defined in the same altitudinal range to the east of the Andes. The Andean region is defined for lands above 1000 meters elevation. Ecuadorian provinces where the species are recorded are cited in accordance with the studied collections, following the Ecuadorian provinces after the 2007 reorganization (i.e., including the new provinces of Santa Elena and Santo Domingo de los Tsáchilas, as well as Orellana, created in 1998). We indicate the total number of collections reviewed per taxa and one representative specimen (voucher), indicating the herbaria acronym where it is deposited. Finally we indicate post-Webster (1999) bibliographic sources which offer updated information about the species. In the “notes” section we include, when needed, any other relevant information, including justifications for nomenclatural decisions.

Data resources

All the information gathered as part of this work is available online in the regularly updated “*Acalypha* Taxonomic Information System” Website (Cardiel J.M., P. Muñoz, E. Dorda & M. Pardo de Santallana. *Acalypha* Taxonomic Information System. <http://www.acalypha.es>). In addition, the information of the studied specimens has been also uploaded to the Global Biodiversity Information Facility (GBIF) (<http://data.gbif.org/datasets/resource/12046/>).

Results

Our work records 20 accepted species of *Acalypha* for continental Ecuador. Two of them are endemic: *Acalypha stellata* and *A. websterii*, and two others are allochthonous: *A. hispida* and *A. wilkesiana*. An identification key is provided. Of the 34 species recognized by Webster (1999) for this territory, nine are considered as synonyms and nine are based on misidentifications (Table 1). The four species added by Ulloa Ulloa and Neill (2005) are accepted. We identify 17 synonyms based on Ecuadorian collections, including the new one *A. pilocardia* Gilli. We indicate the type specimens of almost all the treated names, and 13 lectotypes are designated.

Key to the species of *Acalypha* of continental Ecuador

Key to the subgenera

- 1 Female flowers pedicellate; calyx with 4 or 5 sepals, the subtending bracts inconspicuous, not becoming foliaceous in fruit **Subgen. *Linostachys***

- 2 Female flowers sessile; calyx with 3 sepals, the subtending bracts becoming foliaceous and accrescent in fruit (except in *A. hispida*) **Subgen. *Acalypha***

Key to species

Subgenus *Linostachys*

- 1a Leaf blade palmately nerved, brightly colored minute resinous droplets present, mainly on lower surface **18. *A. villosa***
 1b Leaf blade pinnately nerved, brightly colored minute resinous droplets absent..... **2**
 2a Female inflorescences paniculate. Leaf blade with 10–17 veins per side; stipules generally more than 5 mm long. Petioles more than 1 cm long..... **11. *A. platyphylla***
 2b Female inflorescences racemose. Leaf blade with 9–13 veins per side; stipules inconspicuous, ca. 1 mm long. Petioles less than 1 cm long ... **12. *A. salicifolia***

Subgenus *Acalypha*

- 1a Herb or suffrutex **2**
 1b Trees or shrubs..... **6**
 2a Female inflorescences ellipsoid or cylindrical, densely flowered, with the axis completely covered by the flowers **3**
 2b Female inflorescences loosely flowered, with the axis conspicuously visible... **5**
 3a Female bract with long awned lobes..... **4**
 3b Female bract with triangular awnless lobes..... **5. *A. infesta***
 4a Young branches and leaves without glandular hairs; leaf blade acute; styles branched..... **2. *A. arvensis***
 4b Young branches and leaves with glandular hairs; leaf blade acuminate; styles unbranched..... **1. *A. alopecuroidaea***
 5a Female inflorescences terminal, bracts with filiform lobes cut more than $\frac{1}{2}$ length to the base..... **17. *A. subcastrata***
 5b Female inflorescences axillary, bracts with triangular lobes cut ca. $\frac{1}{4}$ length to the base..... **4. *A. cuspidata***
 6a Leaves with indumentum of stellate hairs..... **7**
 6b Leaves without indumentum of stellate hairs **8**
 7a Female inflorescences terminal, subtending bracts and styles with stellate hairs..... **16. *A. stellata***
 7b Female inflorescences generally axillary, rarely terminal, subtending bracts and styles without stellate hairs **5. *A. dictyoneura***
 8a Female or bisexual inflorescences terminal **9**
 8b Female or bisexual inflorescences axillary **12**

| | | |
|-----|---|----------------------------------|
| 9a | Female bracts without glandular hairs | 18. <i>A. stachyura</i> |
| 9b | Female bracts with glandular hairs | 10 |
| 10a | Female bracts subtriangular at maturity, with the central tooth prominent, lanceolate, acuminate..... | 19. <i>A. websteri</i> |
| 10b | Female bracts suborbicular at maturity, with the central tooth not or slightly prominent..... | 11 |
| 11a | Leaf blade generally broadly ovate-lanceolate; accrescent bracts with glandular hairs ca. 0.3–0.5 mm long; styles 4–5 mm long | 14. <i>A. schiedeana</i> |
| 11b | Leaf blade generally narrowly ovate-lanceolate; accrescent bracts with glandular hairs ca. 1 mm long; styles 7–8 mm long | 10. <i>A. padifolia</i> |
| 12a | Plants with both unisexual and bisexual inflorescences..... | 13 |
| 12b | Plants with all the inflorescences unisexual..... | 14 |
| 13a | Leaf blade generally triangular-lanceolate, palmately nerved.. | 4. <i>A. cuspidata</i> |
| 13b | Leaf blade elliptic-lanceolate or oblong-lanceolate, pinnately nerved..... | 6. <i>A. diversifolia</i> |
| 14a | Leaf blade pinnately nerved | 15 |
| 14b | Leaf blade palmately nerved | 16 |
| 15a | Leaf blade generally obovate, the base subcuneate; female inflorescences 7–15 cm long | 3. <i>A. cuneata</i> |
| 15b | Leaf blade ovate to oblong-lanceolate, the base rounded to subcordate; female inflorescences 25–40 cm long | 12. <i>A. scandens</i> |
| 16a | Female inflorescences extremely densely flowered, with the axis hidden; bracts non-accrescent | 9. <i>A. hispida</i> |
| 16b | Female inflorescences more or less densely flowered, with the axis visible; bracts conspicuously accrescent..... | 17 |
| 17a | Leaf blade generally variegated; female inflorescences up to 10 cm long..... | 20. <i>A. wilkesiana</i> |
| 17b | Leaf blade not variegated; female inflorescences more than 15 cm long..... | 9. <i>A. macrostachya</i> |

Catalogue to the species of *Acalypha* of continental Ecuador

01. *Acalypha alopecuroides* Jacq., Collectanea 3: 196. 1789[1791].

Type. Crescit in Venezuela, tab. 620 in Jacq., Ic. Pl. Rar. 3 (1792), lectotype designated by Cardiel, Anales Jard. Bot. Madrid 54: 233 (1995[1996]).

≡ *Ricinocarpus alopecuroides* (Jacq.) Kuntze, Revis. Gen. Pl. 2: 617. 1891.

Herb. Coastal, 0–250 m. – Provincial distribution: Guayas (2 collections examined). – Voucher: F.M.Valverde 334 (COL, US).

References. Levin (2001), Madsen et al. (2001), Cardiel (2007).

Note. We found only two Ecuadorian collection of this species, which is widely distributed in Central America, Venezuela and Colombia.

02. *Acalypha arvensis* Poepp. in Poepp. Endl., Nov. Gen. Sp. Pl. 3: 21. 1841.

Type. [PERU] Crescit in cultis et ruderalis provinciae Maynas ad Yurimaguas, toto anno florens, E.Poeppig 2215[2115] (**lectotype:** W!, **designated here**; isolectotypes G[2 sheets]!, F[fragment ex W]!, W!). Other type collection: [BRAZIL] ad Serpa in provincia Paranaensi, E.Poeppig s.n. (W!).

≡ *Ricinocarpus arvensis* (Poepp.) Kuntze, Revis. Gen. Pl. 2: 617. 1891.

Annual herb or small suffrutex. Coastal, 0–1000 m. Roadsides and disturbed vegetation. – Provincial distribution: Cotopaxi, Guayas, Los Ríos, Manabí, Napo, Pichincha, Sucumbíos (14 collections examined). – Voucher: O.Haught 2994 (GH, US).

References. Levin (2001), Cardiel (2007).

Note. *A. arvensis* was described based on two collections: E.Poeppig 2215 and E.Poeppig s.n., from Peru and Brazil respectively. We select one of the two specimens of the collection E.Poeppig 2215 found in the W herbarium, as lectotype. Some duplicates of this collection from G and W herbaria also show an original label with the number 2115, in addition of 2215.

03. *Acalypha cuneata* Poepp., Nov. Gen. Sp. Pl. 3: 22. 1841.

Type. [PERU] Crescit in fruticetis maynensibus ad Yurimaguas. Martio lecta, E.Poeppig 2230 (**lectotype:** W[113778]!, **designated here**; isolectotypes, B[destroyed, photo 5288], F!, W!). Other type collections: E.Poeppig 2317 (B[destroyed, photo 5288], F!, G!, P[2 sheets]!, W!), 2330 (F!, G[4 sheets]!, P[2 sheets]!, W!), 2807 (W!).

≡ *Ricinocarpus cuneatus* (Poepp.) Kuntze, Revis. Gen. Pl. 2: 617. 1891; *Acalypha obovata* Benth. var. *cuneata* (Poepp.) J.F. Macbr., Candolea 6: 26. 1940.

= *Acalypha obovata* Benth., Bot. Voy. Sulphur. 163, tab.53. 1846; *Acalypha cuneata* Poepp. var. *obovata* (Benth.) Müll. Arg., Linnaea 34: 14. 1865. **Type:** [ECUADOR, Esmeraldas] Atacames, tab. 53 in Benth., loc. cit. 1846.

= *Acalypha eggersii* Pax, Bot. Jarhb. Syst. 26: 205. 1899. **Type:** Ecuador: prov. Manabí, prope Hacienda El Recreo, H.Eggers 15007 (**lectotype:** Sl!, **designated here**; isolectotypes: B[destroyed, photo 5291], C!, F[2 sheets]!, MA!, MO!, NY!, US!).

Shrub or small tree. Mainly Amazonian and Coastal, excepcionally Andean, (0-) 250–1000(-1500) m. Generally associated with primary rainforests. – Provincial distribution: Esmeraldas, Guayas, Los Ríos, Manabí, Morona-Santiago, Napo, Orellana, Pastaza, Santa Elena, Santo Domingo de los Tsáchilas, Sucumbíos, Tungurahua, Zamora-Chinchipe (222 collections examined). – Voucher: E. Gudiño 61 (GB, DAV, MO, QCNE, U).

References. Bonifaz and Cornejo (2004), Cardiel (2007), De la Torre et al. (2008), Santiana and Cerón (2000).

Notes. Poeppig described *A. cuneata* based on four Peruvian collections: E.Poeppig 2230, 2317, 2330 and 2807. We selected the best preserved specimen, in the W herbarium, as lectotype. The synonym *A. eggersii* Pax was described from

a single collection (F.A.Eggers 15007), which was distributed to several herbaria; due to the destruction of Berlin specimen, we designate as lectotype the specimen from the S herbarium.

04. *Acalypha cuspidata* Jacq., Pl. Hort. Schoenbr. 2: 63, tab. 243. 1797.

Type. [VENEZUELA] Crescit ad Caracas, tab. 243 in Jacq., loc. cit. 1797, lectotype designated by Cardiel, Anales Jard. Bot. Madrid 54: 233 (1995[1996]).

= *Acalypha vestita* Benth., Bot. Voy. Sulpuhur. 164. 1844. **Type:** [ECUADOR] Guayaquil, Sinclair s.n. (K[in hb. Hook.]).

= *Acalypha tenuipes* Pax & K. Hoffm. in Engl., Pflanzenr. 147, 16 (heft. 85): 122. 1924.

Type: Ecuador trockene Gebüsche bei Agua Amarga, Elrecreo, H.Eggers 15833 (**lectotype:** US!, **designated here**; isolectotypes: B[destroyed, photo 5325], F!, K! [2 sheets], L!, M!).

Shrub or suffrutex. Coastal, 0–250 m. Associated with dry deciduous forest, savanna and thickets. – Provincial distribution: El Oro, Guayas, Manabí, Santa Elena (8 collections examined). – Voucher: J.E. Madsen 63938 (AAU, MA, QCA, QCNE).

References. Cardiel (2000, 2007), Madsen et al. (2001), De la Torre et al. (2008).

Notes. *A. cuspidata* Jacq. is often confused with *A. plicata* Müll. Arg., which has a conspicuous glandular indumentum on the young branches, leaves and inflorescences. *A. plicata* is frequent in the Andean zones of Colombia and Peru, but has not been found in Ecuador where it is also likely to be present. The synonym *A. tenuipes* Pax & K. Hoffm. was described from a single collection (H.Eggers 15833), which was distributed to several herbaria; due to the destruction of Berlin specimen, we select as lectotype the best preserved and most complete specimen found in the US herbarium. See comments about this synonym in Cardiel (2000).

05. *Acalypha dictyoneura* Müll. Arg., Linnaea 34: 12. 1865.

Type. [PERU] In Peruvia prope Chachapoyas, A.Matthews s.n. (holotype: G!; isotype: K!).

= *Acalypha pilocardia* Gilli, Feddes Repert. Spec. Nov. Regni Veg. 92: 678. 1981.

Syn. nov. Type: [ECUADOR] Schlucht des Angamarca-Flusses bei El Corazón, 1300 m, 1.7.75, 301, fl., A.Gilli 301 (holotype: W!).

= *Acalypha stellipila* Pax & K. Hoffm. in Engl., Pflanzenr. 147, 16 (heft. 85): 49. 1924. **Type:** Ecuador, Gualea, Sodiro 151/26^a (**lectotype:** F[644679]!, **designated here**; isolectotype B[destroyed, photo 71872]). Other type collections: Ecuador, Puente de Chimbo, Sodiro 151/26^b (B[destroyed, photo 5321]).

Shrub or small tree. Andean, (1300)1750–2500 m. Montane rainforests. – Provincial distribution: Bolívar, Chimborazo, Imbabura, Napo, Orellana, Pichincha, Santo Domingo de los Tsáchilas, Tungurahua (52 collections examined). – Voucher: M.Balslev & E.Madsen 10362 (AAU, COL, C, F, MO, NY, QCA, SEL, US).

References. Cardiel (2000, 2007), Santiana and Cerón (2000).

Note. The synonym *A. stellipila* Pax & K. Hoffm. was described from two collections (Sodiro 151/26^a and 151/26^b), both destroyed in B herbarium. We designate as lectotype a fragment of the first one, preserved in F herbarium.

06. *Acalypha diversifolia* Jacq., Pl. Hort. Schoenbr. 2: 63, tab. 244. 1797.

Type. [VENEZUELA] ex Caracas, tab. 244 in Jacq., Pl. Hort. Schoenbr. 2 (1797), lectotype designated by Cardiel, Anales Jard. Bot. Madrid 54: 233 (1995[1996]).

Shrub or small tree. Amazonian, Andean and Coastal, 0–2000 m. Lowland rain-forests, deciduous and semi-deciduous forests (often along river banks) and disturbed areas. – Provincial distribution: Bolívar, Carchi, Chimborazo, Cotopaxi, El Oro, Esmeraldas, Guayas, Imbabura, Loja, Los Ríos, Manabí, Morona-Santiago, Napo, Orellana, Pastaza, Pichincha, Santo Domingo de los Tsáchilas, Sucumbíos, Tungurahua, Zamora-Chinchipe (310 collections examined). – Voucher: V.Zak & J.Jaramillo 2822 (DAV, F, GB, GH, K, MO, NY, QCNE, SEL).

References. Levin (2001), Cardiel (2007), De la Torre et al. (2008).

07. *Acalypha hispida* Burm. f., Fl. Ind. 302, tab. 61, f. 1. 1768.

Type. Habitat in India, tab. 61 in Burm., loc. cit. 302. 1768.

Shrub. Coastal, 0–250 m. Cultivated and naturalized. – Provincial distribution: Guayas, Santo Domingo de los Tsáchilas (4 collections examined). – Voucher: L.P.Kvist & E.Asanza 40725 (AAU, QCA, QCNE).

References. Levin (2001), Cardiel (2007), De la Torre et al. (2008).

Note. Native to Malaysia or Melanesia, this species is grown in gardens throughout the tropics, and sometimes appears naturalized.

08. *Acalypha infesta* Poepp. in Poepp. & Endl., Nov. Gen. Sp. Pl. 3: 22. 1845.

Type. [PERU] Crescit in cultis ad Cuchero, E.Poeppig 1701 (**lectotype:** W[103476]!, **designated here**; isolectotypes: F!, Pl!, US!, W!).

Annual herb or suffrutex. Andean, 1000–2500 m. Disturbed areas. – Provincial distribution: Bolívar, Cañar, Chimborazo, Pichincha (6 collections examined). – Voucher: F.R.Fosberg & M.A.Giler 22627 (NY, US).

References. Cardiel (2001, 2007), Ulloa Ulloa and Neill (2005).

Notes. This species is often confused with *A. poiretii* Spreng., present in South America (mostly in Argentina and Bolivia), and not found in Ecuador. *A. poiretii* has bisexual inflorescences, the calyx of female flowers with four sepals and female bracts with smaller teeth. *A. infesta* Poepp. was described from a single collection (E.Poeppig 1701), which was distributed to several herbaria. We select as lectotype one of the two sheets conserved in the W herbarium.

09. *Acalypha macrostachya* Jacq., Pl. Hort. Schoenbr. 2: 63, t. 245. 1797.

Type. [VENEZUELA] crescit ad Caracas, tab. 245 in Jacq., Pl. Hort. Schoenbr.: 2 (1797), lectotype designated by Cardiel, Anales Jard. Bot. Madrid 54: 233 (1995[1996]).

= *Acalypha heterodonta* Müll. Arg. var *hirsuta* Müll. Arg., Linnaea 34: 12. 1865.

Type: [Ecuador] Ad pedem montis Chimborazo, R.Spruce 6147 (holotype: K!; isotypes: BM!, W!).

Shrub or small tree. Amazonian, (250-)500–2000 m. In lowland and lower montane rainforests, frequent in disturbed areas. – Provincial distribution: Carchi, Chimborazo, El Oro, Esmeraldas, Morona-Santiago, Napo, Orellana, Pastaza, Pichincha, Santo Domingo de los Tsáchilas, Sucumbíos, Tungurahua, Zamora-Chinchipe (103 collections examined). – Voucher: H.Balslev & E.Madsen 10462 (AAU, C, COL, F, GB, MO, NY, QCA, SEL, US).

References. Levin (2001), Cardiel (2007), Cerón et al. (2007), De la Torre et al. (2008).

10. *Acalypha padifolia* Kunth in Humb. & Bonpl., Nov. Gen. Sp. (quarto ed.) 2: 97. 1817.

Type. [COLOMBIA] Crescit locis sylvaticis subfrigidis inter Almaguer et Pasto, prope villam Meneses, alt. 1322 hex, A.Humboldt & A.Bonpland 2136 (holotype: P-Bonpl.!; isotype: P!).

= *Acalypha andina* Müll. Arg., Flora 55: 26. 1872. **Type:** [ECUADOR] In andibus orientalibus ecuadorensibus inter Bannas et Rio verde altitudine circ. 5000 ped., M.Wagner s.n. (holotype: M!).

= *Acalypha tunguraguae* Pax & K. Hoffm. in Engl., Pflanzenr. 147, 16 (heft. 85): 66. 1924. **Type:** Ecuador, am vulkan Tunguragua, 1800–2000 m., F.Lehmann 6641 (**lectotype:** K[600517]!, **designated here**; isolectotype: B[destroyed, photo 5327]).

= *Acalypha schimpffii* Diels, Biblioth. Bot. 29: 103. 1937. **Type:** Mittel-Ecuador: West-Kordillere: Prov. Chimborazo: Tal des R. Chanchan bei Huigra an buschigen Abnhägen, 1260 m. 16 November, H.J.F.Schimpff 429 (**lectotype:** MO!, **designated here**; isolectotypes: A[3 sheets]!, M!, NY, US!, Z[3 sheets]!). Other type collection: Ecuador, loc. cit. 21 September 1993, Diels 1128 ([B destroyed]).

Shrub or small tree. Andean, 2000–3250(3800) m. Upper and lower montane rainforests, mainly disturbed. – Provincial distribution: Azuay, Bolívar, Cañar, Carchi, Chimborazo, Imbabura, Loja, Pastaza, Pichincha, Tungurahua (83 collections examined). – Voucher: C.E.Cerón & G.Benavides 2551 (F, MO, QCA, QCNE).

References. Cardiel (2000, 2007), De la Torre et al. (2008), Santiana and Cerón (2000).

Notes. The synonym *A. tunguraguae* Pax & K. Hoffm. was described from a single collection (H. Eggers 15833); due to destruction of the Berlin specimen, we select as lectotype the duplicate found in the K herbarium. The synonym *Acalypha schimpffii*

Diels was described based on two collections: Diels 1128, destroyed in B herbarium, and Schimpff 429, poorly preserved, which was distributed to several herbaria; we select as lectotype the specimen preserved in the MO herbarium.

11. *Acalypha platyphylla* Müll. Arg., Linnaea 34: 6. 1865.

Type. In Ecuador Peruviae, L.Fraser s.n. (holotype: G-DC[324093]!).

≡ *Ricinocarpus platyphyllus* (Müll. Arg.) Kuntze, Revis. Gen. Pl. 2: 618. 1891.

Shrub or small tree. Andean, 1250–2500 m. In lower and upper montane rainforests. – Provincial distribution: Azuay, Cotopaxi, Morona-Santiago, Napo, Orellana, Pichincha, Santo Domingo de los Tsáchilas (87 collections examined). – Voucher: J.Jaravillo & V.Zak 7481 (F, MO, NY, QCA, QCNE).

References. Cardiel (2007).

12. *Acalypha salicifolia* Müll. Arg., Flora 47: 438. 1864.

Type. [ECUADOR] in Andibus Ecuadorensibus, R.Spruce 4963 (holotype: W!; isotypes, F, K[2 sheets, fragments ex W!]!).

≡ *Ricinocarpus salicifolius* (Müll. Arg.) Kuntze, Revis. Gen. Pl. 2: 618. 1891.

Shrub or small tree. Amazonian, 500–1250(-1750) m. In lower montane rainforests and lowland rainforests. – Provincial distribution: Morona-Santiago, Napo, Sucumbíos, Zamora-Chinchipe (28collections examined). – Voucher: H.Van der Werff & E.Gudiño 11350 (G, GB, MO, NY, QCNE).

References. Cardiel (2000, 2007), Cerón et al. (2007), De la Torre et al. (2008).

13. *Acalypha scandens* Benth., J. Bot. (Hooker) 6: 329. 1854.

Type. [BRAZIL] On the island of the Amazon opposite Santarem, R.Spruce 1000 (holotype: K!).

Shrub or small tree. Amazonian, 0–250 m. Lowland rainforests, often along river banks. – Provincial distribution: Orellana, Pastaza, Sucumbíos (6 collections examined). – Voucher: C.E.Cerón et N.Gallo 5186 (AAU, MO, QCNE).

References. Cardiel (2007).

14. *Acalypha schiedeana* Schltdl., Linnaea 7: 384. 1832.

Type. [MEXICO] In sylvis umbrosis Jalapae, C.J.W. Schiede 44 (**lectotype:** P[645420]! **designated here**). Other type collections: C.J.W.Schiede 72 (B[destroyed, photo 5319!]!), 247 (HAL[072241]!).

= *Acalypha ecuadorica* Pax & K. Hoffm. in Engl., Pflanzenr. 147, 16 (heft 85): 68. 1924. **Type:** Ecuador, Agua Amarga bei El recreo, trockene Gebüsche, H.Eggers 15535 (**lectotype:** S[S-R-7770]!, **designated here**; isolectotypes: B[destroyed, photo 5290], BM, Cl, F!, GH, K[2 sheets]!, L!, M!, MA!, NY!, PR, US!).

Shrub or small tree. Coastal, 0–500(-750) m. Generally associated with dry forests and thickets. – Provincial distribution: Guayas, Loja, Los Ríos, Manabí (19 collections examined). – Voucher: E.Asplund 15240 (S)

References. Cardiel (1999, 2000), Santiana and Cerón (2000), Ulloa Ulloa and Neill (2005).

Notes. *A. schiedeana* Schldl. was described from a single collection, Schiede 72. The specimen deposited in the B herbarium was destroyed. We select as lectotype the duplicate found in the K herbarium. The synonym *A. ecuadorica* Pax & K. Hoffm. was described from a single collection, H.Eggers 15535, distributed to many herbaria. It represents young branches with immature inflorescences. Because of the destruction of the Berlin specimen, we select as lectotype the duplicate from S herbarium, which has the most developed flowers. See comments about this synonym in Cardiel (2000).

15. *Acalypha stachyura* Pax, Repert. Spec. Nov. Regni Veg. 7: 110. 1909.

Type. [BOLIVIA] Charopampa und San Carlos bei Mapiri, 750 m –August und November 1909. O.Buchtien 1315 (**lectotype:** M!, **designated here**; isolectotype US!). Other type collections: O.Buchtien 1307 (WRSL, US!), 1314 (US!).

Shrub or small tree. Amazonian region, 0–1000(-1250) m. In lowland rainforests. – Provincial distribution: Morona-Santiago, Napo, Orellana, Pastaza, Sucumbíos, Zamora-Chinchipe (136 collections examined). – Voucher: J.Zaruma & al. 126 (F, GB, MO, NY, QCNE).

References. Cardiel (2007), Cerón et al. (2007), De la Torre et al. (2008).

Note. *A. stachyura* Pax was described based on three collections of O.Buchtien: 1307, 1314 and 1315, all from the same locality. We select as lectotype the best preserved collection, from the M herbarium.

16. *Acalypha stellata* Cardiel, Novon 10(4): 362. 2000.

Type. ECUADOR, Bolívar: Limón, estribaciones de la Cordillera Occidental, 880–1100 m, 14 Oct. 1943, M.Acosta Solís 6639 (holotype: F!; isotype: F!). Shrub or small tree. Coastal, 500–1000 m. Lower western slopes of the Andes. – Provincial distribution: Bolívar, Chimborazo, El Oro (5 collections examined).

References. Cardiel (2000), Ulloa Ulloa and Neill (2005).

Note. Ecuadorian endemic.

17. *Acalypha subcastrata* Aresch., Pl. Itin. Eugeniae: 137. 1910.

Type. [ECUADOR] On Puna i Guayaquil viken, N.J.Andersson 160 (**lectotype:** S[S-R-7773]!, **designated here**; isolectotype: S[08-1622]!).

Annual herb. Coastal, 0–500(-1200) m. Generally associated with dry forest and open sites. – Provincial distribution: Guayas, Loja, Los Ríos, Manabí, Santa Elena (24 collections examined). – Voucher: C.M.Dodson & P.M.Dodson 13680 (MO, QCNE, SEL).

References. Cardiel (2007).

Notes. *A. subcastrata* Aresch. is often confused with *A. setosa* A. Rich, which has the ovary hispid and female bracts without glandular hairs (vs. ovary glabrous and bracts with glandular hairs). *A. setosa* occurs in Venezuela and Colombia, and has not been found in Ecuador. *A. subcastrata* was described based on a single collection, N.J.Andersson 160, of which we found two sheets in S herbarium. We select one of them as the lectotype.

18. *Acalypha villosa* Jacq., Enum. Syst. Pl. 32.1760.

Type. [COLOMBIA] Habitat Carthagena in silvis & sepibus, tab. 183, fig. 16 in Jacq., Select. Stirp. Amer. Hist. (1763). Lectotype designated by R.A. Howard (1776). Epitype: tab. 47 in Jacq., Hort. Bot. Vindov. (1776), designated by Cardiel, Anales J. Bot. Madrid (1995[1996]).

= *Acalypha villosa* Jacq. var. *latiuscula* Pax. & K. Hoffm. in Engl., Pflanzenr. 147, 16 (heft. 85): 17. 1924. **Type:** Ecuador, Manabí, bei Hacienda El Recreo, H.Eggers 15616 (**lectotype:** K[600533]!, **designated here**; isolectotype: L!). Other type collections: Ecuador, Manabí, bei Hacienda El Recreo, H.Eggers 15047(F!, K!, L!, M!, US!); Brasilien, Alto Amazonas, Rio Acre, Seringal S. Francisco, E.Ule 9537 (K!); [BRAZIL] Matto Grosso, Malme 2543; [BRAZIL] Madeira Fälle, H.Rusby 1272 (K!, NY!, US!); [BOLIVIA] Bolivien, am Zusammenfluß des Beni und Madre de Dios, H.Rusby 1271 (BM!, MA!, K!, MO!, NY!, US!).

Shrub or small tree. Coastal, 0–250 m. Generally associated to secondary woods and thickets, also with savanna and deciduous forest, mainly on sandy and humid soils. – Provincial distribution: El Oro, Esmeraldas, Guayas, Los Ríos, Manabí, Pichincha, Santo Domingo de los Tsáchilas (54 collections examined). – Voucher: L.P.Kvist 40466 (AAU, MO, NY, QCA).

References. Levin (2001), Cardiel (2007), De la Torre et al. (2008).

Note. The synonym *A. villosa* var. *latiuscula* Pax & K. Hoffm. was described based on six different collections from Bolivia, Brazil and Ecuador. We select as lectotype the quite well preserved Ecuadorian collection H.Eggers 15616, from the K herbarium.

19. *Acalypha websteri* Cardiel, Novon 10(4): 360. 2000.

Type. ECUADOR, Prov. Chimborazo, Huigra, ca. 1200 m, E.Asplund 15427 (holotype S!, isotype Sl!). Shrub. Andean, 1200 m. – Provincial distribution: Chimborazo (3 collections examined). – Voucher: J.E.Madsen 36847 (AAU).

References. Cardiel (2000), Ulloa Ulloa and Neill (2005).

Note. Ecuadorian endemic.

20. *Acalypha wilkesiana* Müll. Arg. in DC., Prodr. 15(2): 817. 1866.

Type. [FIJI] In insulis Fidji (U.S. Expl. Exped. Under. Capt. Wilkes), B.C.Seeman 22 (holotype, G-DC; isotypes, GH!, K[2]!, US[2]!).

≡ *Acalypha amentacea* Roxb. subsp. *wilkesiana* (Müll. Arg.) Fosberg, Smithsonian Contr. Bot. 45: 10. 1980.

Shrub. Cultivated and naturalized. Coastal, Andean and Amazonian, 0–1500 m. – Provincial distribution: Guayas, Imbabura, Manabí, Napo, Santo Domingo de los Tsáchilas (6 collections examined). – Voucher: M.Acosta-Solís 13076 (F).

References. Levin (2001), Cardiel (2007), Sagun et al. (2010).

Note. Native to the Polynesian island of Fiji, this species is widely used as an ornamental in the tropics. It has been treated as a subspecies of *A. amentacea* Roxb. because of its morphological similarities (Fosberg and Sachet 1980), but a molecular study by Sagun et al. (2010) showed that it was closer to *A. hispida* Burm. than to *A. amentacea*, and therefore should be the species rank.

Acknowledgements

This work was partly financed by the Spanish Government, through the research project EUI 2008-0388. Support was also received from the Universidad Autónoma de Madrid (Spain) and the Regional Government (Comunidad de Madrid), through the research project CCG07-UAM/AMB-1453. We kindly thank the curators and staff of the aforementioned herbaria for facilitating the study of their specimens. Francisco Pando, manager of GBIF-Spain, kindly helped us in everything relating to biodiversity databases. The authors are also grateful to Grady Webster for his advice in the early studies on Ecuadorean *Acalypha*; rest in peace.

References

- Bacigalupo NM, Mulgura ME (1999) *Acalypha*. In: Zuloaga O, Morrone O (Eds) Catálogo de las plantas vasculares de la República Argentina. Missouri Botanical Garden Press. Missouri, vol. 2, 590–592.

- Berry PE, Caruzo MBR, Cordeiro I, Esser HJ, Fernández-Casas FJ, Levin GA, Lima LR, Riina R, Wurdack KJ (2007) *Acalypha*. In: Zuloaga O, Morrone O, Belgrano MJ (Eds) Catálogo de las plantas vasculares del Cono Sur (Argentina, Sur de Brasil, Chile, Paraguay y Uruguay). Missouri Botanical Garden Press, Missouri, 528–535.
- Bonifaz C, Cornejo X (2004) Flora del Bosque de Garúa (árboles y epífitas) de la comuna Loma Alta, cordillera Chongón Colonche, provincia del Guayas, Ecuador. Universidad de Guayaquil, Missouri Botanical Garden, Fundación GAIA.
- Brako L (1993) *Acalypha*. In: Brako L, Zarucchi JL (Eds) Catálogo de las Angiospermas y Gimnospermas del Perú. Monographs in Systematic Botany from the Missouri Botanical Garden, St. Louis, 45: 428–429.
- Brako L, Zarucchi JL (1993) Catálogo de las Angiospermas y Gimnospermas del Perú. Monographs in Systematic Botany from the Missouri Botanical Garden, St. Louis, 45.
- Cardiel JM (1995) *Acalypha* (Euphorbiaceae). In: Flora de Colombia, monografía nº15. Universidad Nacional de Colombia y Real Jardín Botánico de Madrid-CSIC.
- Cardiel JM (1995 [1996]) Tipificación de las especies de *Acalypha* (Euphorbiaceae) descritas por Jacquin. Anales Jard. Bot. Madrid 54: 230–233.
- Cardiel JM (1999) Contribuciones a la Flora de Venezuela: Revisión del género *Acalypha* L. (Euphorbiaceae). Acta Botánica Venezolana 22(2): 255–324.
- Cardiel JM (2000) Nuevas especies y sinónimos de *Acalypha* L. (Euphorbiaceae) de Ecuador. Novon 10: 123–126. doi: 10.2307/3392986
- Cardiel JM (2001) *Acalypha infesta* Poepp. (Euphorbiaceae), novedad para la flora colombiana. Revista de la Academia Colombiana de Ciencias Exactas 25(97): 463–465.
- Cardiel JM (2007) Catálogo de las especies peruanas de *Acalypha* Linnaeus (Euphorbiaceae). Fontqueria 55(50): 405–414.
- Cardiel JM (2010) *Acalypha*. In: Campostrini R et al. (Eds) Lista de espécies da Flora do Brasil. Jardim Botânico do Rio de Janeiro: 963–964.
- Cerón CE, Reyes CI, Montalvo C, Vargas LM (2007) La cuenta alta del río Oglán, Pastaza-Ecuador, diversidad, ecología y flora. Editorial Universitaria, Quito.
- De la Torre L, Navarrete H, Muriel MP, Macía MJ, Balslev H (Eds) (2008) Enciclopedia de las Plantas Útiles del Ecuador. Herbario QCA de la Escuela de Ciencias Biológicas de la Pontificia Universidad Católica del Ecuador, Herbario AAU del Departamento de Ciencias Biológicas de la Universidad de Aarhus, Quito & Aarhus.
- Fosberg FR, Sachet MH (1980) Systematic studies of Micronesian plants. Smithsonian Contributions to Botany 45: 1:40.
- Hayden WJ, Hayden SM (2000) Wood anatomy of Acalyphoideae (Euphorbiaceae). International Association of Wood Anatomists Journal 21: 213–235.
- Howard RA (1989) *Acalypha*. In: Flora of the Lesser Antilles, Leeward and Windward Island. Arnold Arboretum, Harvard University Jamaica Plain, Massachusetts. 5(2): 5–11.
- Jørgensen PM, León-Yáñez S (Eds) (1999) Catalogue of the Vascular Plants of Ecuador. Monographs in Systematic Botany from the Missouri Botanical Garden, St. Louis.
- León-Yáñez S, Valencia R, Pitman N, Endara L, Ulloa Ulloa C, Navarrete H (Eds) (2011) Libro rojo de las plantas endémicas del Ecuador, 2^a edición. Publicaciones del Herbario QCA, Pontificia Universidad Católica del Ecuador, Quito.

- Levin GA (2001) *Acalypha*. In: Flora of the Venezuelan Guayana. Missouri: Botanical Garden Press, 5: 81–85.
- Levin GA (2008) *Acalypha*. In: Hokche O et al. (Eds) Nuevo Catálogo de la Flora Vascular de Venezuela. Fundación Instituto Botánico de Venezuela Dr. Tobías Lasser, Caracas, 6.
- Madsen JE, Mix RL, Balslev H (2001) Flora of Puná Island. Plant resources on a Neotropical Island. Aarhus University Press.
- McNeill J, Barrie FR, Burdet HM, Demoulin V, Hawksworth DL, Marhold K, Nicolson DH, Prado J, Silva PC, Skog JE, Wiersema JH, Turland NJ (Eds) (2006) International Code of Botanical Nomenclature (Vienna Code): Adopted by the Seventeenth International Botanical Congress, Vienna, Austria, July 2005. Regnum Vegetabile 146. Gantner, Ruggell.
- Pax FA, Hoffmann K (1924) *Acalypha*. In: Engler A (Ed) Das Pflanzenreich. Verlag von Wilhelm Engelmann, Leipzig, IV, 147–16 (heft 85): 1–231.
- Sagun VG, Levin GA, van Welzen PC (2010) Revision and phylogeny of *Acalypha* (Euphorbiaceae) in Malesia. Blumea 55: 21–60. doi: 10.3767/000651910X499141
- Santiana J, Cerón C (2000) Euphorbiaceae. In: Valencia R, Pitman N, León-Yanez S, Jørgensen PM (Eds) Libro Rojo de las Plantas Endémicas del Ecuador. Herbario QCA, Pontificia Universidad Católica del Ecuador, Quito, 190–195.
- Thiers B (2011) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium: <http://www.sweetgum.nybg.org/ih/>
- Tokuoka T (2007) Molecular phylogenetic analysis of Euphorbiaceae sensu stricto based on plastid and nuclear DNA sequences and ovule and seed character evolution. Journal of Plant Research 120: 511–522. doi: 10.1007/s10265-007-0090-3
- Ulloa Ulloa C, Neill DA (2005) Cinco años de adiciones a la flora del Ecuador: 1999–2004. Universidad Técnica Particular de Loja, Loja.
- Neill DA, Ulloa Ulloa C (2011) Adiciones a la Flora del Ecuador: Segundo Suplemento, 2005–2010. Fundación Jatun Sacha. Quito.
- Valencia R, Pitman N, León-Yanez S, Jørgensen PM (Eds) (2000) Libro Rojo de las Plantas Endémicas del Ecuador. Herbario QCA, Pontificia Universidad Católica del Ecuador, Quito. Ecuador.
- Webster GL (1999) *Acalypha*. In: Jørgensen PM, León-Yáñez S (Eds) Catalogue of the Vascular Plants of Ecuador. Monographs in Systematic Botany from the Missouri Botanical Garden, St. Louis, 75: 455–457.
- Wurdack KJ, Davis CC (2009) Malpighiales phylogenetics: Gaining ground on one of the most recalcitrant clades in the angiosperm tree of life. American Journal of Botany 96: 1551–1570. doi: 10.3732/ajb.0800207