

# Evolutionary relationships and taxonomy of *Microtea* (*Microteaceae*), a basal lineage in the core Caryophyllales

Alexander P. Sukhorukov<sup>1</sup>, Alexander N. Sennikov<sup>2,3</sup>, Maya V. Nilova<sup>1</sup>, Yuri Mazei<sup>4</sup>,  
Maria Kushunina<sup>5</sup>, Maria Salete Marchioretti<sup>6</sup>, Pavel Hanáček<sup>7</sup>

**1** Department of Higher Plants, Biological Faculty, Lomonosov Moscow State University, 119234, Moscow, Russia **2** Botanical Museum, Finnish Museum of Natural History, P.O. Box 7, 00014 University of Helsinki, Finland **3** Herbarium, Komarov Botanical Institute of Russian Academy of Sciences, Prof. Popov St. 2, 197376 St. Petersburg, Russia **4** Department of Hydrobiology, Biological Faculty, Lomonosov Moscow State University, 119234, Moscow, Russia **5** Department of Plant Physiology, Biological Faculty, Lomonosov Moscow State University, 119234, Moscow, Russia **6** Instituto Anholtano de Pesquisas/UNISINOS, São Leopoldo, RS, Brazil **7** Department of Plant Biology, Mendel University in Brno, Zemědělská 1, 613 00 Brno, Czech Republic

Corresponding author: Alexander P. Sukhorukov (sukhor@mail.ru)

---

Academic editor: Eric Roalson | Received 12 August 2018 | Accepted 7 December 2018 | Published 9 January 2019

---

**Citation:** Sukhorukov AP, Sennikov AN, Nilova MV, Mazei Y, Kushunina M, Marchioretti MS, Hanáček P (2019) Evolutionary relationships and taxonomy of *Microtea* (*Microteaceae*), a basal lineage in the core Caryophyllales. *PhytoKeys* 115: 1–50. <https://doi.org/10.3897/phytokeys.115.29041>

---

## Abstract

The basal position of the small American genus *Microtea* within the core Caryophyllales was suggested only recently in accordance with molecular phylogeny. However, the specific relationships within the genus were not traced. The results of our phylogenetic analysis based on the *matK* chloroplast gene suggest the monophyly of *Microtea*, and *Ancistrocarpus* and other related genera should be included in it. *Microtea* is divided into two major sister clades: clade A consisting of *M. glochidiata*, *M. maypurensis* and *M. tenuifolia*, and clade B comprising *M. debilis*, *M. sulcicaulis*, *M. scabrida*, *M. celosioides*, and *M. papillosa*. The nrDNA dataset (ITS), although containing only a limited number of accessions, shows the same species number in clade A, and the remaining species studied (*M. debilis*, *M. scabrida* and *M. celosioides*) form clade B. Subgeneric status is assigned to clades A and B corresponding with the names *Microtea* subgen. *Ancistrocarpus* subgen. nov. and *Microtea* subgen. *Microtea*, respectively. The diagnostic characters at the subgeneric level are as follows: length of pedicels, number of flowers at each node, number of stamens and styles. A multivariate analysis of 13 distinguishing morphological characters supports the results of phylogenetic analysis. All species have similar pericarp and seed ultrasculpture and anatomy, and they share the reticulate pericarp surface (independent of presence or absence of finger-shaped outgrowths on its surface) and rugose or slightly alveolate seed ultrasculpture. On the basis of morphological characters, we

accept 10 *Microtea* species. A checklist includes a new diagnostic key, morphological descriptions and distribution patterns of each species. *Galenia celosioides* is the oldest legitimate name available for the plants previously known as *Microtea paniculata*, for which the combination *Microtea celosioides* is validated here. The neotypes of *Galenia celosioides* and *Microtea sprengelii* were designated from the collections of Prinz Wied at BR. The name *M. foliosa* is discussed and finally synonymized with *M. scabrida*. The lectotypes of *Ancistrocarpus maypurensis* ( $\equiv$ *Microtea maypurensis*), *Microtea debilis* var. *ovata* ( $=$ *M. debilis*), *M. glochidiata*, *M. maypurensis* var. *angustifolia* ( $=$ *M. tenuifolia*), *M. glochidiata* f. *lanceolata* ( $=$ *M. maypurensis*), *M. longebracteata* ( $=$ *M. celosioides*), *M. paniculata* var. *latifolia* ( $=$ *M. scabrida*), *M. portoricensis*, *M. scabrida*, *M. sulcicaulis*, and *Potamophila parviflora* ( $=$ *M. maypurensis*) are designated. *Microtea sulcicaulis* is reported for the first time as native to Bolivia, and *M. maypurensis* is reported from Indonesia (Java), where it is found as an alien plant with an unclear invasion status.

## Keywords

Caryophyllales, *Microtea*, molecular phylogeny, reproductive characters, taxonomy

## Introduction

The genus *Microtea* Sw. was described by Swartz (1788) with one species, *M. debilis* Sw., native to the Lesser Antilles (the Caribbean). The author placed it within the group “Pentandria–Digynia” due to the pentaphylloous perianth, five stamens, and two styles. A pericarp with distinct echinate outgrowths was reported as another indicative character of *Microtea* (Swartz 1788). Kunth (1817) described a related genus *Ancistrocarpus* Kunth with the type species *A. maypurensis* Kunth, and he pointed out that the main differences between *Microtea* and *Ancistrocarpus* belong to the reproductive characters such as the different numbers of stamens (five vs eight, respectively) and styles (two vs four or five) and the shape of the pericarp outgrowths (echinate vs apically hooked). Further new generic and species names were mostly based on the same reproductive characters (Roemer and Schultes 1820, Schrank 1821, Link 1821), but none of the genera allied to *Microtea* have been commonly accepted, and currently *Microtea* has been considered a single genus that includes all closely related taxa (e.g., Steudel 1841, Moquin-Tandon 1849, Urban 1885, Walter 1909, Rohwer 1993, Marchioretto and de Siqueira 1998). According to the latest studies (Marchioretto and de Siqueira 1998, Hernández-Ledesma et al. 2015), *Microtea* comprises 10–12 species distributed in Central and South America. They can be distinguished and classified by life history, presence of bracteoles by each flower, and morphology of the pericarp (e.g., Walter 1909, Marchioretto and de Siqueira 1998). However, many important reproductive traits are still poorly studied in this genus, including the fruit and seed anatomy that has been depicted schematically only for *M. debilis* (Melikian 1993).

Traditionally, *Microtea* occupied a provisional position within the core Caryophyllales and has been considered as part of the Chenopodiaceae (Kunth 1817, Walter 1906, Takhtajan 2009), Petiveriaceae (Brown and Varadarajan 1985) or Phytolaccaceae (Moquin-Tandon 1849, Walter 1909, Hutchinson 1926, Nowicke 1968, Behnke 1993, Rohwer 1993, Atha 2004, Zhu and Sanderson 2017). Friedrich (1956) suggested that *Microtea* may be a connecting link between the Phytolaccaceae and Che-

nopodiaceae. Also, Behnke (1993) and Behnke and Mabry (1994) reported that the structures of sieve-element plastids and of the pollen grains deviate from those of the other Phytolaccaceae. The recently combined molecular phylogeny based on the *petD* and *matK* regions revealed a distant position of *Microtea* from both Chenopodiaceae and Phytolaccaceae (Schäferhoff et al. 2009). However, only two *Microtea* species – *M. debilis* and *M. scabrida* Urb. – were included in this molecular analysis (Schäferhoff et al. 2009). Currently, the monophyly of *Microtea* and the relationships between its species have not been confirmed.

The aims of the present paper are (1) to include more species of *Microtea* in the molecular analysis in order to clarify the relationships between the species of the genus and to confirm the monophyly of *Microtea*, (2) to provide new data on the carpological characters as the most diverse and taxonomically important traits, and (3) to provide a new taxonomic description of the genus and better determination of the range of each species.

## Methods

### Field studies and revision of the herbarium material

Field work was done by the first author (AS) in Grenada in November 2016 (Main Island, Carriacou, and Petit Martinique) and in March 2018 in the Dutch Caribbean (Curaçao); however, no *Microtea* species were found. The field investigations in Brazil were provided by Maria Salete Marchioretto. The revision of herbarium specimens was undertaken in B, BM, BR, E, G, H, K, L (incl. U & WAG), LE, LY, M, MEXU, MHA, MSB, MW, P, and PACA. The Virtual database of the Brazilian herbaria (<http://reflora.jbrj.gov.br/reflora/PrincipalUC/PrincipalUC.do>), National Herbarium of Colombia (<http://ciencias.bogota.unal.edu.co/icn/colecciones-cientificas/herbario/>) and the Tropicos database (<http://tropicos.org/Name/24800059?tab=specimens>) were used as references for some specimens kept in ASE, CEN, COL, FURB, GB, HUFS, NY, NYBG, RB, SJRP, and US if their identification was possible using the digital images.

### Carpological studies

Several fruits of all species were taken from the herbaria vouchers deposited in herbaria with the permission of the curators. Seed ornamentation was examined using a scanning electron microscope (SEM) JSM–6380 (JEOL Ltd., Japan) at 15 kV after sputter coating with gold-palladium in the laboratory of Electron Microscopy at the Lomonosov Moscow State University. To restore the soft pericarp tissue prior to scanning electron microscopy, the fruits were dehydrated in aqueous ethyl alcohol solutions of increasing concentration, followed by alcohol-acetone solutions and pure acetone. The seeds did not require a complicated treatment prior to SEM due to the presence of the

hard seed coat. The cross-sections of the fruits and seeds were prepared using a rotary microtome Microm HM 355S (Thermo Fisher Scientific, USA). Before sectioning, the seeds were soaked in water:alcohol:glycerin (1:1:1) solution, dehydrated in an ethanol dilution series and embedded in Technovit 7100 resin (Heraeus Kulzer, Germany). The cross-sections were observed using a Nikon Eclipse Ci microscope and photographed with a Nikon DS-Vi1 camera (Nikon Corporation, Japan) at the Department of Higher Plants (Moscow State University).

## Phylogenetic analysis

The list of vouchers and their accession numbers is provided in Table 1.

**Table 1.** Voucher information and GenBank accession numbers for *Microtea* species and outgroups included in the phylogenetic analysis. Sample codes are provided only for the newly sequenced samples.

Sample code	Species	Voucher	GenBank accession number	
			matK	ITS
Mi02	<i>Microtea debilis</i>	French Guiana, Eau Claire, 15 Aug 1993, S. Mori <i>et al.</i> 23295 (P05197089)	MH678599	—
—	<i>M. debilis</i>	USA, California, cultivated, Yuncker <i>et al.</i> (UC851834)	—	JX232577
Mi14	<i>M. celosioides</i>	Brazil, Bahia, Milagres, 6 Mar 1977, R.M. Harley 19451 (U 1473444)	MH678600	MH726167
Mi07	<i>M. celosioides</i>	Brazil, Piaui, Caracol, 25 Feb 2011, Melo <i>et al.</i> 9216 (PACA 115982, sub <i>M. longebracteata</i> )	MH678601	—
Mi06	<i>M. glochidiata</i>	Brazil, Tucano Mun., 20 Feb 1992, A.M. de Carvalho & D.J.N. Hind 3841 (G)	MH678602	MH726168
Mi10	<i>M. maypurensis</i>	Brazil, Bahia, Rio Jacurici, 16 Jan 1997, M.M. Arbo <i>et al.</i> 7276 (G)	MH678603	MH726169
Mi22	<i>M. papillosa</i>	Brazil, Minas Gerais, Diamantina, 11 May 1982, L. Rossi <i>et al.</i> 76279 (PACA 76279)	MH678604	—
Mi19	<i>M. scabrida</i>	Paraguay, Concepcion, Paso Horqueta, 18 Nov 1993, E. Zardini & T. Tilleria 37460 (MW0581802)	MH678605	—
Mi26	<i>M. scabrida</i>	Paraguay, Cordillera Dept., Cerro Tobati, 28 Oct 1987, R. Degun & E. Zardini 447 (G)	MH678606	MH726170
Mi27	<i>M. scabrida</i>	Paraguay, Guaira, Melgarejo, 13 Mar 1989, E. Zardini & C. Velasquez 11391 (MW)	MH678607	MH726171
Mi28	<i>M. scabrida</i>	Paraguay, National Park Ybicu'i, 11 Nov 1989, E. Zardini & U. Velasquez s.n. (G)	MH678608	—
Mi47	<i>M. sulcicaulis</i>	Paraguay, dept. Cordillera, Colonia Rosado, 26 Oct 1986, A. Schinini & E. Bordas 24850 (G)	MH678609	—
Mi34	<i>M. tenuifolia</i>	Brazil, Bahia, 23 Mar 1974, Belmonte 17305 (U1473428)	MH678610	MH726172
Mi36	<i>Griselinia littoralis</i>	Switzerland, Botanical Garden of Geneva, Sep 2017, A. Konstantinova (living collection)	MH678611	MH726173
Mi41	<i>G. scandens</i>	Chile, Santiago, 1925, A. Marillo 923 (LE)	MH678612	MH726174
—	<i>Macarthuria australis</i>	Australia, Lepisci & Brims 1943 (K)	FN825765	—
—	<i>M. neocambrica</i>	Australia, Coveny & Wilson 11674 (K)	FN825766	—
—	<i>Stegnosperma halimifolium</i>	no data	HQ878442	—
—	<i>Simmondsia chinensis</i>	no data	AF204863	—

### DNA extraction and PCR amplification

The nuclear (ITS) and chloroplast (*matK*) regions of genomic DNA were used for the phylogenetic analysis. Total DNA was isolated from dried leaves using Invisorb<sup>®</sup> Spin Plant Mini Kit (Stratec Molecular GmbH, Berlin, Germany). ITS-A (Blattner and Kaderer 1999) and ITS4 (White et al. 1990) primers were used for ITS region amplification, and MatK-1RKIM-f and MatK-3FKIM-r were used for *matK* region ([http://botany.si.edu/projects/dnabarcodes/matK\\_PCR\\_&\\_Sequencing\\_Proocols.pdf](http://botany.si.edu/projects/dnabarcodes/matK_PCR_&_Sequencing_Proocols.pdf)). PCRs were performed using 0.75 units of MyTaq Red DNA polymerase (Bioline, London, UK) in 15 µl of original buffer containing MgCl<sub>2</sub> and dNTPs, with 0.3 µM of each primer and 1 µl of unquantified DNA template. Thermocycling was carried out in TProfessional Basic Thermocycler (Biometra, Göttingen, Germany) using the thermal and cycling conditions as described in Shaw et al. (2007): initial denaturation at 80 °C for 5 min; 30 cycles of 95 °C for 1 min, 50 °C for 1 min, a ramp of 0.3 °C/s to 65 °C and incubation at 65 °C for 4 min, with a final extension step of 65 °C for 5 min. A clean-up reaction with exonuclease I and alkaline phosphatase (Thermo Fisher Scientific, Waltham, Massachusetts, USA) was used to remove unincorporated primers and nucleotides before sequencing. The PCR products were sent to Macrogen Europe (Netherlands) for automated sequencing. The primers used for amplification were also used for the sequencing reactions.

### Sequence alignment and phylogenetic reconstruction

Data files were assembled, edited and evaluated using Geneious 8.1 software (Biomatters Ltd, Auckland, New Zealand). Regions of ambiguous alignment were excluded from all analyses. After exclusion of these regions we used 754 characters in the nuclear (ITS) and 828 characters in the chloroplast (*matK*) analysis. The nuclear and chloroplast data were analyzed separately with MEGA7 software (Kumar et al. 2016) which delivered a maximum likelihood (ML) tree based on the Kimura 2-parameter model (Kimura 1980) with support for nodes measured by bootstrap percentages (N70% considered significant). The percentage of trees in which the associated taxa clustered together is shown next to the branches. Initial tree(s) for the heuristic search were obtained automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using the Maximum Composite Likelihood (MCL) approach, and then selecting the topology with superior log likelihood value. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. Evolutionary analyses were conducted in MEGA7 (Kumar et al. 2016).

### Multivariate analysis


Different *Microtea* species were classified by group average linkage algorithm of cluster analysis constructed on a Gower similarity matrix (Gower 1971) based on thirteen characters including general morphology (life history, pubescence, leaves) and reproductive

traits. This approach recognizes the species grouping based on similar characters, but does not provide a true phylogenetic context. The reliability of grouping was assessed at the level  $p<0.05$  using SIMPROF algorithm (Clarke 1993, Clarke and Warwick 2001). Calculations were performed using PRIMER 6.1.6 statistical software (Clarke and Gorley 2006).


## Results

### Phylogenetic analysis of matK region

The most representative phylogenetic analysis based on cpDNA (*matK*) dataset shows the monophyly of the genus *Microtea*, which is divided into two well-supported clades (Fig. 1): clade A consisting of *M. glochidiata* + *M. maypurensis* / *M. tenuifolia*, and clade B comprising the remaining species included in the analysis (*M. debilis*, *M. celosioides*, *M. scabrida*, *M. sulcicaulis*, and *M. papillosa*). In clade B, *M. debilis* is a sister to the remaining species. The position of all *Microtea* species is considered to be close to *Macarthuria* (core Caryophyllales). Three clades – *Macarthuria*, *Stegnosperma* and *Microtea*, even if not fully represented in the trees based on different phylogenetic markers – occupy a



**Figure 1.** The phylogenetic tree from maximum likelihood analysis of *matK* region sequences. The tree with the highest log likelihood (-2983.53) is shown. The analysis involved 18 nucleotide sequences. Codon positions included were 1<sup>st</sup>+2<sup>nd</sup>+3<sup>rd</sup>+Noncoding. There were a total of 828 positions in the final dataset.



**Figure 2.** The phylogenetic tree from maximum likelihood analysis of ITS region sequences. The tree with the highest log likelihood (-2848.78) is shown. The analysis involved 9 nucleotide sequences. There were a total of 754 positions in the final dataset.

basal position within the core Caryophyllales (Brockington et al. 2009, Schäferhoff et al. 2009, Sukhorukov et al. 2015). The tree based on the *matK* region that includes these three basal lineages suggests the monophyly of *Microtea* (Fig. 1).


### Phylogenetic analysis of ITS region

Based on the nrDNA tree (Fig. 2), the clade A (*M. glochidiata* + *M. maypurensis* / *M. tenuifolia*) is a sister to the clade B comprising the rest of the species studied (*M. debilis* + *M. celosioides* / *M. scabrida*) with good statistical support. However, only a limited number of species were included in the ITS dataset.

Based on both *matK* and ITS phylogenetic analyses, subgeneric status was assigned to two major clades: clade A (*M. glochidiata* + *M. maypurensis* / *M. tenuifolia*) represents *M. subgen. Ancistrocarpus* stat. nov., and clade B represents the type subgenus (*M. subgen. Microtea*).

### Carpological investigations


In all the species studied the fruit is one-seeded (Fig. 3), and the pericarp surface is reticulate (Figs 4–8, A, B, E, F). The finger-shaped (echinate) outgrowths over the




**Figure 3.** Cross-section of the fruit of *Microteca glochidiata*, showing the embryo and perisperm covered by the tegmen (Brazil, Bahia, Tucano Mun., 20 Feb 1992, A.M. de Carvalho & D.J.N. Hind 3841, PACA). Abbreviations: P – pericarp with plumose outgrowths, T – testa of the seed coat (tegmen covers perisperm and embryo), PE – perisperm, E – embryo. Magnification – 50×.

entire fruit surface are present in almost all species (Figs 3; 4A, B; 5–6A, B, E, F; 7E, F; 8 A, B) except *M. portoricensis* (Fig. 7A–B), but they are usually scattered or even obscure in *M. tenuifolia* (Fig. 8E–F) and in some specimens of *M. celosioides* (Fig. 4E–F). In *M. glochidiata* they can reach 0.65 mm in length (Fig. 7E–F). The echinate outgrowths may be covered by large horizontal unicellular papillae (plumose outgrowths: *M. glochidiata*, Fig. 7E–F), or each outgrowth is terminated by a group of 2–4 recurved (hooked) hairs (*M. maypurensis*, Fig. 8A–B). Such plumose or hooked outgrowths clearly assist epizoochorous dispersal. Two thick styles are characteristic for the species forming clade B (*M. bahiensis*, *M. celosioides*, *M. debilis*, *M. papillosa*, *M. portoricensis*, *M. scabrida*, *M. sulcicaulis*), and those in the remainder of the genus (clade A: *M. glochidiata*, *M. maypurensis*, *M. tenuifolia*) possess three to five thin styles. The pericarp consists of several layers; the cells of the innermost layers are usually filled with tannins (Fig. 9 A, B). The pericarp outgrowths emerge from the mesophyll and consist of several prosenchymatous cell layers.


The seeds are spherical and black, with a rugose (Figs 4–8C, D, G, H) or slightly alveolate (*M. sulcicaulis*: Fig. 6D) surface, and are basally inserted on the fruit wall. The seed coat of all species consists of a thick (40–50  $\mu\text{m}$ ) exotestal layer, with outer cell




**Figure 4.** Fruits and seeds of *Microtea debilis* and *M. celosioides*: **A, B** fruit of *M. debilis*, enclosed in the perianth (St. Lucia, Soufrière, 1958, G.R. Proctor 17789, BM000019256) **C, D** seed of *M. debilis* (Honduras, nr Cangrejal river, foothills of Ceiba, 29 Jul 1938, T.G. Yuncker et al. 8674, G) **E, F** fruit of *M. celosioides* (Retiro das Pedras, Brumadinho, 14 Dec 1998, J.R. Stehmann & C.E.S. Ferreira 2399, PACA) **G, H** seed of *M. celosioides* (Retiro das Pedras, Brumadinho, 14 Dec 1998, J.R. Stehmann & C.E.S. Ferreira 2399, PACA). Magnification: **A, E** – 30x, **B, F** – 100x, **C, G** – 50x, **D, H** – 300x.




**Figure 5.** Fruits and seeds of *Microtea papillosa* and *M. scabrida*: **A, B** fruit of *M. papillosa*, enclosed in the perianth (Diamantina Mun., Estrada Conselheiro Mata, 11 Apr 1982, L. Rossi et al. 3322, PACA) **C, D** seed of *M. papillosa* (Diamantina Mun., Estrada Conselheiro Mata, 11 Apr 1982, L. Rossi et al. 3322, PACA) **E, F** fruit of *M. scabrida*, enclosed in the perianth (Paraguay, Alto Paraná, 1909, K. Fiebrig 5468, M) **G, H** seed of *M. scabrida* (Paraguay, Alto Paraná, 1909, K. Fiebrig 5468, M). Magnification: **A, E** – 30×, **B, F** – 100×, **C, G** – 50×, **D, H** – 300×.




**Figure 6.** Fruits and seeds of *Microtea sulcicaulis* and *M. bahiensis*: **A, B** fruit of *M. sulcicaulis*, enclosed in the perianth (Paraguay, Caazapá Dept., Tavai, 7 Dec 1988, F. Mereles 2122, G) **C, D** seed of *M. sulcicaulis* (Paraguay, Caazapá Dept., Tavai, 7 Dec 1988, F. Mereles 2122, G) **E, F** fruit of *M. bahiensis* (Brazil, Bahia state, Salvador, Dunas de Itapuá, nr Hotel Stella Maris, N from Condomínio Alamedas da Praia, 8 Jun 1993, P. de Queiroz 3211, PACA) **G, H** seed of *M. bahiensis* (Brazil, Bahia state, Salvador, Dunas de Itapuá, nr Hotel Stella Maris, N from Condomínio Alamedas da Praia, 8 Jun 1993, P. de Queiroz 3211, PACA). Magnification: **A, E** – 30x, **B, F** – 100x, **C, G** – 50x, **D, H** – 300x.



**Figure 7.** Fruits and seeds of *Microtea portoricensis* and *M. glochidiata*: **A, B** fruit of *M. portoricensis*, enclosed in the perianth (Puerto Rico, Cabo Rojo, 1864, Grosourdy 13, P04598159) **C, D** seed of *M. portoricensis* (Puerto Rico, Cabo Rojo, 1864, Grosourdy 13, P04598159) **E, F** fruit of *M. glochidiata* (Brazil, Maranhão, Barao do Grajau, 21 Jan 2012, R.M. Harley et al. 56455, K) **G, H** seed of *M. glochidiata* (Brazil, Maranhão, Barao do Grajau, 21 Jan 2012, R.M. Harley et al. 56455, K). Magnification: **A, E** – 30 $\times$ , **B, F** – 100 $\times$ , **C, G** – 50 $\times$ , **D, H** – 300 $\times$ .



**Figure 8.** Fruits and seeds of *Microtea maypurensis* and *M. tenuifolia*: **A, B** fruit of *M. maypurensis*, enclosed in the perianth (Bolivia, La Paz Dept., Beni river, Jul 1886, H.H. Rusby 1379, LE) **C, D** seed of *M. maypurensis* (Bolivia, La Paz Dept., Beni river, Jul 1886, H.H. Rusby 1379, LE) **E, F** fruit of *M. tenuifolia* enclosed in the perianth (Brazil, Minas Gerais, Serra das Vertentes, Jun 1893, A. Glaziou 20437, B) **G, H** seed of *M. tenuifolia* (Brazil, Jacobina Mountains in Bahia, 1836, Blanchet 2588, P00798998). Magnification: **A, E** – 30×, **B, F** – 100×, **C, G** – 50×, **D, H** – 300×.



**Figure 9.** Cross-section of *Microtea glochidiata* fruit (Brazil, Bahia, Tucano Mun., 20 Feb 1992, A.M. de Carvalho & D.J.N. Hind 3841, PACA). **A** image of the cross-section **B** schematic representation. Abbreviations: P – pericarp; T – testa; TE – tegmen.


walls much thicker than the arch-like protoplast (Figs 3; 9), and 1–2 barely visible layers of tegmen with bar-thickenings of its cell walls. The cells of the exotesta (Fig. 9) are dark brown due to the presence of unstructured tannin-like substances, but without additional stalactite-shaped deposits (vertical or oblique depositions of tannins originating from the outer cell walls). The annular embryo occupies a peripheral position in the seed, and is located vertically. The perisperm is abundant.

All *Microtea* species share the reticulate pericarp surface (regardless of the presence or absence of echinate outgrowths) and the rugose or slightly alveolate seed surface. The fruit and seed structure of *Microtea*, namely the homocellular pericarp consisting of several layers, seed coat with much thicker testa and barely noticeable tegmen with bar-thickenings of the cell walls, vertical embryo position in one-seeded fruits, and abundant perisperm, is typical for the core Caryophyllales (Sukhorukov et al. 2015, 2018).

#### Diagnostic characters in *Microtea* and multivariate analysis

All characters discovered in *Microtea* species are summarized in Table 2.

The results of cluster analysis of the characters suggest the existence of five significantly different groups within the *Microtea*, these branches being highlighted in black colour (Fig. 10): (1) *M. glochidiata*–*M. maypurensis*–*M. tenuifolia*, (2) *M. debilis*–*M. portoricensis*, (3) *M. scabrida*–*M. sulcicaulis*, (4) *M. celosiooides*–*M. bahiensis*, and (5)



**Figure 10.** Classification of *Microtea* species by group average linkage algorithm of cluster analysis based on 13 characters. Black branches connect significantly ( $P < 0.05$ ) different groups, red branches – insignificantly different groups.

*M. papillosa*. The groups are significantly ( $p<0.05$ ) distinguished on different levels of Gower's index. Clusters (2–5) correspond with clade B in the phylogenetic analysis. The most prominent distinctions are observed between three major clusters consisting of (1) *M. glochidiata*, *M. maypurensis*, *M. tenuifolia*; (2) *M. debilis*–*M. portoricensis*; and (3) *M. papillosa*, *M. bahiensis*–*M. celosioides*, *M. scabrida*–*M. sulcicaulis*. Cluster (1) corresponds with clade A in the phylogenetic analyses (Figs 1, 2), and this group is clearly distant from the remaining species due to character sets 4, 5, 9 and 10. Cluster (2) comprises the species without bracteoles (character 3, state 0), and cluster 3 unites the rest of the genus with the similar sets of characters 3, 4, 5, 8, 9, 10, and 12.

**Table 2.** Characters of *Microtea* species.

Species/character	<i>M. bahiensis</i>	<i>M. debilis</i>	<i>M. celosioides</i>	<i>M. glochidiata</i>	<i>M. maypurensis</i>	<i>M. papillosa</i>	<i>M. portoricensis</i>	<i>M. scabrida</i>	<i>M. sulcicaulis</i>	<i>M. tenuifolia</i>
1 Life history: 0 – annual or rarely biennial; 1 – upright perennials with a taproot or caudex; 2 – perennial lianas	1	0	0	0	0	1	0	2	1	0
2 Pubescence: 0 – (almost) glabrous; 1 – papillate	0	0	0	0	0	1	0	0	0	0
3 Bracteoles. 0: absent; 1: present	1	0	1	1	1	1	0	1	1	1
4 Flower arrangement. 0: solitary (inflorescence is a spike); 1: two or three in clusters (thyrsoid inflorescence)	0	0	0	1	0	0	0	0	0	0
5 Presence of pedicel. 0: flowers sessile or subsessile (pedicel up to 1.3 mm); 1: pedicel 1.35–3.0 mm	0	0	0	1	1	0	0	0	0	1
6 Perianth segments. 0: oblong or ovoid; 1: roundish	0	0	0	1	0	0	0	0	0	0
7 Number of perianth segments. 0: always five; 1: four or five (varying)	0	0	0	0	0	0	1	0	0	0
8 Number of stamens. 0: four or five; 1: more than five (usually seven or eight)	1	0	1	1	1	1	0	1	1	1
9 Number of stigmas. 0: two, rarely three; 1: three to five	0	0	0	1	1	0	0	0	0	1
10 Shape of stigmas. 0: thick; 1: filiform	0	0	0	1	1	0	0	0	0	1
11 Diameter of fruit body (without stigmas and outgrowths if the latter are present). 0: 0.9–1.1 mm; 1: 1.1–1.3 mm; 2: 1.4–2.0 mm	1	1	1	0	0	1	0	2	2	0
12 Fruit/perianth ratio. 0: fruit longer than perianth (protruding); 1: fruit equal to perianth (fruit not protruding)	0	0	0	0	0	0	0	0	0	1
13 Pericarp. 0: without any projections, reticulate; 1: with scattered, thick and simple outgrowths; 2: with abundant thick outgrowths; 3: with fimbriate (plumose) projections; 4: with projections terminating in hooks	1	2	1	3	4	1	0	1	1	0

## Taxonomy

### Artificial key to the *Microtea* species

- 1 Leaning or twining perennial herb up to 150 cm; leaf blades basally truncate ..... **4. *M. scabrida***
- Smaller herbs or dwarf subshrubs up to 100 cm; leaves cuneate ..... **2**
- 2 Stems decumbent; each flower supported by a bract; bracteoles not present; stamens 4–5; annuals with obovate leaves ..... **3**
- Stems usually erect; each flower supported by a bract and two transverse bracteoles; stamens 5–8; annuals or perennials with leaves not obovate ..... **4**
- 3 Pericarp with echinate (finger-shaped) outgrowths ..... **1. *M. debilis***
- Pericarp not echinate (its surface reticulate) ..... **7. *M. portoricensis***
- 4(2) Perennials or annuals with shortly but densely papillate stem and leaves ..... **3. *M. papillosa***
- Perennials or annuals with glabrous stems, or with leaves papillate at margins and mid-ribs ..... **5**
- 5 Perennial herb or dwarf subshrub with well-expressed caudex and rosulate leaves; cauline leaves short (up to 2.0 cm) ..... **6. *M. bahiensis***
- Annuals or perennial herbs (in latter case without a caudex); cauline leaves usually larger (2.5–12.0 cm) ..... **6**

- 6 Flowers 1–6 per node; pericarp outgrowths plumose..... **8. *M. glochidiata***  
 – Flowers 1(2) per node; pericarp outgrowths (if present) glabrous or hooked at apices..... **7**  
 7 Flowers 1(2) per node; pericarp outgrowths hooked at apices.....  
 ..... **9. *M. maypurensis***  
 – Flowers always one per node; pericarp outgrowths (if present) not hooked ..... **8**  
 8 Perennial herb; leaves lanceolate or narrowly oblong, usually appressed to the stem, stiff; fruit 1.75–2.0 mm long..... **5. *M. sulcicaulis***  
 – Annuals, biennials; leaves filiform to oblong, not appressed to the stem, not stiff; fruit less than 1.5 mm long ..... **9**  
 9 Pedicels 1.35–1.7(2.5) mm long; fruit not protruding or slightly protruding from the perianth; pericarp smooth, verrucous or with barely visible outgrowths; leaves usually filiform or lanceolate ..... **10. *M. tenuifolia***  
 – Pedicels up to 1 mm long; fruit twice the length of the perianth; pericarp with short finger-shaped outgrowths; leaves narrowly lanceolate to oblong (rarely ovoid) ..... **2. *M. celosioides***

**Gen. *Microtea* Sw., Prodr. [O.P.Swartz]: 53 (1788).**

*Microtea debilis* Sw. (type species)

=*Schollera* Rohr, Skr. Naturhist.-Selsk. 2: 210 (1792), nom. illegit., non Roth (1788).

≡*Microtea* subgen. *Schollera* (Rohr) H.Walter, Pflanzenr. (Engler) 39: 127 (1909), nom. inval. (Art. 22.2). Type species: *M. debilis* Sw. Note: Vahl (1792) established that a new generic name, *Schollera* was based on the plant that he considered conspecific with *Microtea debilis*. The type of *Schollera* Rohr is therefore that of *Microtea debilis* (Turland et al. 2018: Art. 10.2).

=*Ancistrocarpus* Kunth, Nov. Gen. Sp. [quarto] 2: 186 (1817). Type species: *A. maypurensis* Kunth (≡*Microtea maypurensis* (Kunth) G.Don).

=*Potamophila* Schrank, Pl. Rar. Hort. Monac. 2: tab. 63 (1821) nom. illegit., non R.Br. (1810). Type species: *P. parviflora* Schrank (=*Microtea maypurensis* (Kunth) G.Don).

=*Ceratococca* Willd. ex Roem. & Schult., Syst. Veg., ed. 15, 6: LXX (1820). Type species: *C. maypurensis* Humb. & Bonpl. ex Roem. & Schult. (=*Microtea maypurensis* (Kunth) G.Don).

=*Aphananthe* Link, Enum. Hort. Berol. Alt. 1: 383 (1821), nom. rej. Type species: *A. celosioides* (Spreng.) Link (≡*Microtea celosioides* (Spreng.) Moq. ex Sennikov & Sukhor.). Note: The generic name *Aphananthe* Link is rejected in favour of its later homonym, *Aphananthe* Planch. (Cannabaceae).

**Description of the genus.** Annuals, perennial herbs, rarely dwarf subshrubs; stems angulate, glabrous or papillate; leaves alternate, sessile or pedunculate, entire, filiform to ovate or obovate, cuneate or truncate, apically mostly acuminate, a persistent leaf rosette usually present, cauline leaves resembling the rosulate leaves or much shorter;

inflorescence a spike or thyrsoid; pedicel inconspicuous or up to 3 mm long; flowers actinomorphic, bisexual, subtended by a hyaline bract and two similar bracteoles, sometimes bracteoles absent; perianth of (4)5 glabrous segments or lobes, green, white or yellowish; stamens (4)5 in alternisepalous position, or 6–8 (in both antesepalous and alternisepalous positions), anthers 0.15–0.30 mm, introrse, thecae globose, pollen grains pantoporate; ovary roundish; style not present or very short, stigmas 2–5; fruit nut-like, single-seeded, dry; pericarp projections (if present) not evident in flowering condition, pericarp at fruiting stage reticulate, mostly having finger-shaped outgrowths (emergences) that can be plumose (with additional smaller hair-like projections) or hooked at their apices; seeds spherical, black, with rugose or alveolate surface, with annular embryo located vertically and abundant perisperm.

Ten species distributed in the (sub)tropics of the Americas; two – *M. debilis* and *M. maypurensis* – are considered as aliens in the humid tropics of Africa (Cameroon) and Asia (Indonesia), respectively.

### **Taxonomic synopsis of *Microtea***

#### ***Microtea* subgen. *Microtea***

=*Microtea* subgen. *Moquinia* Nowicke in Ann. Missouri Bot. Gard. 55: 349 (1968).

Type: *M. paniculata* Moq. (=*M. celosioides* (Spreng.) Moq. ex Sennikov & Sukhor.).

**Description of the subgenus.** Annuals, perennial herbs or dwarf subshrubs; bracteoles present or absent; pedicels inconspicuous or very short (up to 1.3 mm long); flowers single per node (inflorescence a spike); stigmas 2(3), thick. The species are distributed across the (sub)tropical South America, in Central America and Antilles.

#### **1. *M. debilis* Sw., Prodr. [O.P.Swartz]: 53 (1788).**

=*M. debilis* Sw. var. *ovata* Delile ex Moq. in DC., Prodr. 13(2): 17 (1849). Lectotype (Sukhorukov, designated here): Overseas territories of France. Guadeloupe, herb. Desfontaines (P00798994!).

=*M. debilis* var. *rhombifolia* Moq. in DC., Prodr. 13(2): 17 (1849). Holotype: [without locality data] “Herb. Poiret in Herb. Moquin-Tandon” (P00798997!). Note: One specimen at K contains both varieties (*M. debilis* var. *ovata* and *M. debilis* var. *rhombifolia*) mounted on one sheet and identified as such by Moquin-Tandon. It contains several plant fragments with different labels. The varieties can barely be distinguished from one another.


**Lectotype.** (designated by Howard and Howard 1982: 76): OVERSEAS TERRITORIES OF THE NETHERLANDS. St. Eustatius, *F. Masson* s.n. in Herb. Banks (BM000019252!). Note. The species was described from the West Indies, and Saint

Christopher Island (also known as Saint Kitts Island, St. Kitts & Nevis) was reported as the only locality in the protologue. Although the personal herbarium of Swartz is incorporated in S, he noted for some species in his book (Swartz 1788) that in such cases he used the herbarium collections in Banks's possession (Stearn 1961, Howard and Howard 1982). The collections used by Swartz (1788) are listed in Swartz's subsequent publication, *Flora Indiae Occidentalis*, in which the original locality of *M. debilis* was stated more accurately as the island of "St. Eustathii", and "Masson" was indicated as the collector (Swartz 1797: 543). This specimen is the only element associated with the taxon by the original author, and consequently it was designated as the lectotype of the name (Howard and Howard 1982: 76). Although this was not indicated in the protologue, probably a further collection was available to Swartz by that time: a specimen of *M. debilis* collected by H. de Ponthieu (reportedly in Grenada) was acquired by Swartz from Banks and subsequently given to L.J. Montin, a Swedish collector of herbarium material, whose private herbarium became part of S along with that of Swartz himself (Lindman 1916).


**Description.** Annuals, glabrous; stems decumbent, up to 30 cm (Fig. 11A); rosulate leaves up to 9 (10–12) cm, long-petiolate, obovate or oblong, mostly persistent; caudine leaves rhombic or ovate, cuneate; inflorescence a spike (Fig. 11B); flowers sessile or very shortly pedicellate (pedicels at fruiting ~1 mm); bracteoles absent; perianth segments 5, greenish, lanceolate or oblong; stamens (4)5; stigmas 2, thick; fruit roundish, 1.1–1.25 mm long and 1.0–1.2 mm wide (Fig. 4A, B), with finger-shaped outgrowths (up to 0.4 mm long); seed ~1.0 mm, with rough surface (Fig. 4D).

**Habitat.** Sands, forest margins, or as a weed; altitudes up to 1000(1200) m a.s.l.

**Distribution.** Native to American tropics (Fig. 12).



**Figure 11.** *Microtea debilis*: **A** general view of the plants (La Selva Biological Station, Costa Rica, 2001) **B** close-up of the inflorescence (La Selva Biological Station, Costa Rica, 2001). Photographs by Orlando Vargas Ramírez. See also [https://sura.ots.ac.cr/florula4/find\\_sp3.php?key\\_species\\_code=LS001515](https://sura.ots.ac.cr/florula4/find_sp3.php?key_species_code=LS001515).




**Figure 12.** Distribution map of *Microtea debilis* in its native range.

**Specimens examined.** **ANTIGUA & BARBUDA:** Dimsdale, 29 Aug 1937, H.E. Box 1005 (BM000019212); **BELIZE:** Manatee Lagoon, 1905, M.E. Peck 42 (K); 22 mile of Stann Creek river, 250 ft, 3 July 1932, W.A. Schipp 286 (K); Bright Lookout Bank, Sibun river, 12 Jan 1935, P.H. Gentle 1487 (K); Toledo distr., 28 Oct 2005, C. Whitefoord & V. Quiroz 106292 (BM000895516); **BOLIVIA:** La Paz, Upper Rio Beni, Dec 1906, J.W. Evans 54 (BM); Rio Acre, Jan 1912, E. Ule 9363 (G, K); Beni Dept., provinces Ballivián & Yacuma, 27 May 1985, I. Guareco 477 (M); Beni Dept., provinces Ballivian & Yacuma, comunidad Charaton, 11 Mar 1995, E. Rivero 293 (M); **BRAZIL:** Acre: Sena Madureira Mun., 10 Oct 1968, G.T. Prance & al. 7958 (P05197144, U1473418); Serra do Moa, 24 Apr 1971, G.T. Prance et al. 12403 (K); Sena Madureira to Rio Branco, 10 Oct 1968, G.T. Prance et al. 7958 (K); Cruzeiro do Sul, 18 Nov 2001, T.B. Croat 85419 (NY01187675 – image!); **Amazonas:** Barra do Rio Negro [Manaus], 1850, R. Spruce 1078 (BM000019274, M); Mun. Boca do Acre, Rio Acre, Jan 1912, E. Ule 9363 (L1673158); Barcelos, 9 Sep 1962, A.P. Duarte 7284 (RB00272328 – image!); Manaus, 20 Oct 1971, P.J.M. Maas & H. Maas 498

(U1473424); Manaus Mun., Rio Cuieiras, 50 km upstream, farm of Sr. Nemerio, 2 Apr 1974, *J.C. Ongley & J.F. Ramos* P21790 (K, U1473416); Rio Mapari, ca 30 km E of Borba, 24 Jun 1983, *C. Todzia et al.* 2246 (K); Maués Mun., ca. 20 km E of Maués, 23 Jul 1983, *S.R. Hill* 13146 (K); **Maranhão**: Rio Perizes, 12 Dec 1976, *B.G.S. Ribeiro & G.S. Pinheiro* 1752 (IAN154119 – image!); **Pará**: 1908, *C.F. Baker* 29 (K, M); Rio Tapajos, 29 Feb 1920, *anonym* 14684 (RB00272338 – image!); Belém, 10 Nov 1942, *W.A. Archer* 7794 (K); **Pernambuco**: Aliança, Rio Branco, Apr 1913, *J.G. Kuhlmann* 3139 (RB00272342 – image!); **Rio Grande do Sul**: Porto Alegre, Jan 1898, *Reinedly s.n.* (E). Note: The isolated record from Rio Grande do Sul may be locally alien, since the specimen was collected near the port. **COLOMBIA**: **Cundinamarca Dept.**: Bogotá, 1851–1857, *J. Triana* 5260 (P04598044); **Guainía Dept.**: Rio Negro, San Felipe, 28 Sep 1921, *Ph.v. Luetzelburg* 22249 (M); **Magdalena Dept.**: Santa Marta, 1898–1901, *H.H. Smith* 1246 (BM000019264, G, E, K, L1673160, LE, P04598033); **Antioquia Dept.**: Vuelta de Acuña, 14 Jan 1918, *E.W. Pennell* 3815 (K); **Bolívar Dept.**: nr Turbaco, 9 Nov 1926, *E.P. Killip & A.C. Smith* 14359 (LE); **Amazonas Dept.**: Rio Putumayo, Sep/Oct 1930, *G. Klug* 1647 (BM000019246, K); Rio Igara Paraná, Puerto Buenaventura, 12 Oct 1973, *C. Sastre* 2484 (P04448856); **Santander Dept.**: Puerto Berrio, 6 Jun 1935, *O. Haught* 1752 (BR, BM000019237); nr Barrancabermeja, 5 Mar 1967, *J. de Bruijin* 1593 (WAG1166513); **Tolima Dept.**: San Sebastián de Mariquita, 25 Sep 2001, *G.C. Bernal et al.* 1096 (COL000007803 – image!); **COSTA RICA**: Talamanca, Mar 1894, *A. Tonduz* 8712 (BM000019277, P03321182); Sipurio, Apr 1894, *H. Pittier & Th. Durand* 8712 (M, P04598123); Alajuela prov., Los Chiles, 4 Aug 1949, *R.W. Holm & H.H. Iltis* 933 (G, P04598124); Limón prov., Cordillera de Talamanca, 9°40'25"N, 83°01'35"W, 24 Feb 1989, *G. Herrera* 2439 (BM000019239); **DOMINICA**: 1838, *Murray* 344 (K); Oct 1881, *H. Eggers* 563 (BR, G, L1673154, LE, LY, M, P04597992); Dubuc, 21 Jul 1983, *Assi & Portecop* 16503 (P06806988); St. Patrick, 8 Feb 1986, *C. Whitefoord* 5386 (BM000019281); **DOMINICAN REPUBLIC**: Higüey, Nov 1946, *R.A. Howard & E.S. Howard* 9731 (P04598162); **ECUADOR**: **Esmeraldas prov.**: Timbre, 2 Jun 1955, *E. Asplund* 16558 (B, G, K, P04598007); **Guayas prov.**: Balao, [without date] *Hamilton* 373 (K, BM000019242); Balao, Dec 1891, *Eggers* 14122 (B); Guayaquil, 1846–1849, *W. Jameson* 373 (BM); nr Naranjito, Jun 1945, *W.H. Camp* 3575 (P04598008); **Los Ríos prov.**: Clementina on Rio Pita, 12 Mar 1939, *E. Asplund* 5253 (BR, G, K, P04598009); **Manabi prov.**: El Recreo, 1897, *H. Eggers* 14927 (LE); Naranjapata, 21 Nov 1933, *H.J.F. Schimpff* 505 (G, M); **Napo prov.**: Puerto Misahualli, May 1983, *W. Palacios et al.* 366 (K); **FRENCH GUIANA**: St Georges, 14 May 1983, *M.F. Prévost* 1356 (P04598030); Eau Claire, 3°37'N, 53°12'W, 15 Aug 1993, *S. Mori et al.* 23295 (P05197089); Elahé, Abati Ti Wan, 6 Mar 1999, *M. Pignal* 927 (P00176753); Cayenne, 15 May 1999, *M.-F. Prévost* 3657 (P5197094, U1473414); **grenada** [Main Island]: 1844, *J. Goudot s.n.* (P04598036); Victoria [city], 20 Nov 1957, *G.R. Proctor* 17140 (BM000019255); St. George's, 11 May 1905, *Broadway* 4366 (BR); **GUADELOUPE** (selected specimens): see type of *M. debilis* var. *ovata*; [without exact location and date] *Bertero*, herb. *J. Gay* 1820 (K); [without exact location and date] coll. *Bertero* s.n. [herb. De Candolle] (G00676754);

[without exact location] 1907, *P. Duss* 41 (LY); Rivage de Capesterre, 1 Apr 1943, *Questel* 5064 (P04597987); **GUATEMALA**: Izabal, Aug 1870, *G. Bernoulli* 877 (G, K); **GUYANA** (selected specimens): [without location] 1868, *Schomburgk* 229 (BR, L1673157, P04598083); Mazaruni river, Sep 1880, *G.S. Jenman* 748 (K); Mazaruni river, Aug 1889, *Jenman* 5277 (BM000019244, K); Rockstone, Sep 1905, *A. W. Bartlett* 8559 (K); Aruka, NW district, Nov 1915, *C.K. Bancroft* & *N. Persaud s.n.* (K); Tumatumari village, Jun 1921, *H.A. Gleason* 18 (K); Amakura river, Northwest district, 8°10'N, 60°W, Mar 1923, *J.S. de la Cruz* 3548 (K); Pomeroon district, Moruka river, Jul 1927, *J.S. de la Cruz* 4601 (K); Cuyuni river, near Lower Camaria landing, 23 Nov 1929, *N.Y. Sadwith* 666 (K); Mabaruma compound, 11 Jul 1934, *W.A. Archer* 2256 (K); Waranama ranch, intermediate Savannahs, Berbice river, 9 Jun 1958, *Harrison* 1073 (K); Potaro river, near Amatuk, 2000 ft, 10 Sep 1959, *B.A. Whitton* 344 (K); [West Demerara Region] Essequibo, Henrietta, 4 Feb 1960, *J.A. Harris* 143 (MHA); Royal Island, 5 May 1985, *C. Feuillet* 2181 (B, P04598031); Cuyuni-Mazaruni Region, near Eping river, 5°58'N, 60°13'W, 396 m, 10 Feb 1991, *T. McDowell* 3985 (K, U1473392); Pomeroon-Supenaam Region, Kabakaburi river, 8 Sep 1992, *B. Hofmann* & *L. Roberts* 2443 (U1473417); Rupununi distr., Manari, 2 Aug 1995, *M.J. Jansen-Jacobs* et al. 4733 (U1473413); North-West distr., Moruka river, 9 Oct 1997, *T. van Andel* 1959 (U1473415); **HAITI**: Ile La Tortue [Tortuga Island], La Vallée, 21 May 1925, *E.L. Ekman* 4063 (G, K); **HONDURAS**: Roatan Island, Aug 1886, *G.F. Gaumer* 27 (K); Santa Barbara Dept., San Pedro Sula, May 1888, *C. Thieme* 5427 (K); nr Cangrejal river, foothills of Ceiba, 29 Jul 1938, *T.G. Yuncker* et al. 8674 (BM000019283, G, K); Gracias a Dios dept., Cocobila, 10 Feb 1981, *G.R. Proctor* 38968 (BM000019258); Gracias a Dios dept., Rio Platano Biosphere Reserve, 30 May 1985, *S.G. Knees* 2821 (BM000019248); Atlántida, La Ceiba, 26 Sep 1991, *M. Chorley* 439 (BM); Gracias a Dios dept., Tawahka Asangni Biosphere Reserve, 1994, *P. House* 99 (BM000833935); **JAMAICA**: Negril, 9 Mar 1908, *W. Harris* 10214 (BM000019231, K, P04598156); **MARTINIQUE**: [year] 1833, *Sieber* 93 (BM000019251, BR, G, M, P04597991); [without exact location] 1839, *Rivoire s.n.* (P04597989); [without exact location] 1868–1869, *Hahn* 1026 (B); nr St Pierre, Jun 1879, *Cosson* 811 (K); June 1913, *M. Mouret* 128 (P04597990); Tivoli, 12 Dec 1944, *H. Stehle* 5552 (U1473388); **MONT-SERRAT**: Plymouth, 5 Feb 1959, *G.R. Proctor* 19021 (BM000019257); **NETHERLANDS ANTILLES**: **St. Eustatius Island**, [without exact location] 1908, *I. Boldsingh* 89 (U1473381); **Saba Island**, Bottom city, 29 Aug 1947, *F. Arnoldo* 892 (U1473385); **St. Martin**: Philipsburg to Belvedere, 18 Aug 1908, *I. Boldsingh* 2588 (U1473387); **NICARAGUA**: Omotepe Island, Oct 1869, *P. Lévy* 241 (G, P04598125); Puerto Cabezas, 5 Oct 1978, *W.D. Stevens* 10563 (MO-20480120 – image!); Wiwili Mun., 20 Jan 2006, *I. Coronado* et al. 30888 (MEXU); **PANAMÁ**: Isthmus of Panama, Feb 1850, *A. Fendler* 109 (K); [no exact location] Jul 1861, *J. Hayn* 196 (BM000019238, K); Isla Brava, 8 Jun 1909, *E. André* 350 (K); El Real, 16 Jun 1959, *W.L. Stern* et al. 621 (LE); Fort San Lorenzo, 17 Dec 1966, *D. Burch* et al. 1025 (K); Ailigandi, 8 Oct 1978, *B. Hammel* 5041 (MEXU); **PERU**: **Huánuco Region**: Prov. Pachitea, Dept. Huanuco, Bosque National de Iparia, 6 Dec 1966, *J. Schunke* 1322 (G); Panguana, Nov/Dec

2008, *G. Gerlach* 136 (M); **Puno Region:** [San] Gaban, Aug 1854, *R.F. Hohenacker* 2443 (G, M); **Loreto Region:** Maucallacta, Rio Paranapura, Jan 1935, *G. Klug* 3959 (BM000019247, K); Loreto prov., 30 Oct 1940, *E. Asplund* 14151 (G, K); Loreto dept., Rio Nanya, 7 Aug 1972, *T.B. Croat* 18874 (E); Dept. Iquitos, Maynas, 19 May 1986, *M. Rimachi* 8194 (BR); **San Martín Region:** Prov. Mariscal Cáceres, Dept. Tocache Nuevo, 6 Nov 1969, *J. Schunke* 3585 (G); **PUERTO RICO:** nr Dorado, Mar 1922, *N.L. Britton et al.* 6650 (NY00992870 – image!); Yabacoa, 11 Oct 1968, *R.J. Wagner* 1687 (U1473389); Mun. de Patillas, 15 May 1988, *C.M. Taylor & J. Druitt* 8097 (NY00992874 – image!); **ST. KITTS & NEVIS:** St. Kitts, nr Canada Estate, Sep/Oct 1901, *N.L. Britton & J.F. Cowell* 275 (NY01509872 – image!); **ST. LUCIA:** St Lucia, Jun 1879, *H.B. Murray s.n.* (K); Soufrière, 1958, *G.R. Proctor* 17789 (BM000019256); Anse Mamin, 21 Nov 1938, *H.E. Box* 1999 (BM); **ST. VINCENT & GRENADINES:** St. Vincent [Island], 1822, *L. Guilding s.n.* (K); St. Vincent Island, 1826, *Lambert s.n.* (BR); [St. Vincent Island] pastures at Petit Bordell estate, 14 Oct 1949, *I. Velez* 3341 (K); St. Vincent Island, [without date and collector] 178 (BM000019219); **SURINAME** (selected specimens): 1843, *Hostmann s.n.* (K); Paramaribo, 1851, *Wullschlaegel* 445 (BR); Groningen, 10 May 1916, *J.A. Samuels* 115 (K, L1673164, P04598019); Corantijnpolder nr Nieuw Nickerie, 27 Aug 1933, *J. Lanjouw* 634 (K); Landsboerderij, 11 Feb 1955, *J.C. Lindeman* 521 (U1473398); Zuid river, 3°20'N, 56°49'W, 30 Sep 1963, *H.S. Irwin et al.* 57701 (M, P04598018); distr. Suriname, Kalpoeweg, 13 Apr 1981, *Ch. Kalpoe* 16588 (U1473404); Marowijne, Bigiston, 13 Apr 2006, *T.R. van Andel & L. McIntosh* 5180 (L0842725); **TRINIDAD & TOBAGO** (selected specimens): Trinidad [Island], 1826, *Sieber* 134 (L1673148, LE, M, P04598179); Tobago [Island], Nov 1889, *H. Eggers* 5826 (P04597985); Tobago, Roxborough, 16 Oct 1912, *W.E. Broadway* 4642 (G, K); Tobago, 15 May 1913, *anon. s.n.* (P04597986); Trinidad, Imperial College of Tropical Agriculture [St Augustine], New Farm, 80 ft, 20 Feb 1958, *J.W. Purseglove* 6112 (K); Trinidad, Curepe, 8 miles E of Port-of-Spain, 25 May 1975, *A. Raynal* 15549 (K); Tobago, Castara, 18 Jul 1910, *W.R. Broadway* 4068 (E); **VENEZUELA** (selected specimens): **Amazonas:** Dept. Casiquiare, Maroa, 25 Aug 1978, *O. Huber* 2571 (K); Alto Orinoco Mun., 1 May 2005, *A. Fernández et al.* 21234 (NYBG03142590 – image!); **Aragua State:** Maracay, [without date] *P.C. Vogl* 20 (BR, M); **Barinas State:** Barinas to San Cristobal, 13 Mar 1964, *F.J. Breteler* 3692 (WAG1166512); **Capital District:** Caracas, 1864, *Grosourdy* 19 (P04598045); Cordillera de la Costa, 19 Aug 2000, *W. Meier et al.* 7418 (M); Cordillera de la Costa, 1 May 2002, *W. Meier et al.* 8153 (G); **Carabobo State:** Companero, 1843, *J. Linden* 1338 (LE, P04598050); San Esteban, 1893–1894, *Mocquerys s.n.* (H1332163); Chirgua, 700 m, 1 Jan 1939, *A.H.G. Alston* 5960 (BM000019204, LE, U1473454); Distr. Bejuma, 16 Apr 2000, *W. Meier & N. Flauger* 6878 (G); **Delta Amacuro State:** Delta of Orinoco, San Antonio, 16 Feb 1911, *F.E. Bond et al.* 140 (K); **Falcón State:** Rio Tucuyo, 26 Jan 1966, *J.A. Steyermark & A. Braun* 94504 (M); **Mérida State:** El Vigia, [without date] *Mocquerys* 981 (P05197141); **Monagas State:** Lower Orinoco, Sacupana, Apr 1896, *H.H. Rusby & R.W. Squires* 77 (E, G, K, M); **Sucre State:** Cumaná, 1893–1894, *Mocquerys* 814 (K, P04598035); **Trujillo State:**



**Figure 13.** Records of *Microtea debilis* as an alien plant in Africa.

Trujillo city, 22 May 1971, C. Emilin & B. de Rojas 964 (U1473456); **Vargas State:** road from La Suagra to Macuto, Apr 1854, *anonym s.n.* (K); **Yaracuy State:** Chivacoa, 6 Mar 2004, W. Meier & S. Nehlin 10101 (G); San Felipe distr., 23 Mar 2004, W. Meier & J.L. Escalona 10219 (B 10 0455316); **Zulia State:** Maracaibo, 1826, *Plée s.n.* (P04598048); [without exact location] 1893–1894, *Mocquerys* 931 (P04598034); **VIRGIN ISLANDS (US):** St Croix, Jolly Hill, 20 Jan 1906, C. Raunkier *s.n.* (BR, P04598163); St. Thomas, Charlotte Amalie, 9 Feb 1913, N.L. Britton *et al.* 470 (NYBG01509850 – image!); Note: Data were not available on the presence of the species in Cuba, in agreement with the recent treatment of *Microtea* in this country (Greuter 2002). As alien found in tropical Africa (Bamps 1974; see also Fig. 13); **CAMEROON:** Douala, sea shore, Oct 1938, H. Jacques-Félix 2203 (P04621264); Douala, 50 m, 21 Mar 1967, A. Meurillon 661 (K); Douala, 8 Oct 1969, *anonym* 1721 (P04621263, WAG1166511); West province, Moliwe, 3 miles N of Victoria, 400 ft, 16 Aug 1969, H. Chuml 301 (K); South-West Province, NE of Muyuka, at foot of Cameroon Mt., 26 Aug 1983, D. Thomas 2538 (K, P05156267, P05156282, WAG1166510).

## 2. *M. celosioides* (Spreng.) Moq. ex Sennikov & Sukhor., comb. nov. urn:lsid:ipni.org:names:77193071-1

≡*Galenia celosioides* Spreng., Nov. Prov.: 18 (1818). ≡*Aphananthe celosioides* (Spreng.) Link, Enum. Hort. Berol. Alt. 1: 383 (1821). Neotype (designated here by Sennikov & Sukhorukov): BRAZIL. Bahia, Mucuri, März [March] 1816, Maximilian, Prinz zu Wied 53 (BR000005575398! Fig. 14). Notes. Although *Galenia celosioides* was synonymized with *Microtea maypurensis* (Moquin-Tandon 1849,

Schmidt 1872, Walter 1909, Nowicke 1968) or *M. debilis* (Steudel 1841), the protologue (Sprengel 1818) contains the phrases “flosculi ... brevissime pedicellati... styli duo” [flowers very shortly pedicellate, styles two], the characters found in *M. celosioides*. The description did not include reference to the presence of hooked outgrowths, a peculiar feature of *M. maypurensis*, and the short pedicels and two styles also exclude this species. Nevertheless, Moquin-Tandon (1849) reported his former intention to transfer Sprengel’s name to *Microtea*, apparently on the basis of his analysis of its protologue. He changed his mind probably because he did not examine the original material of the name. *Galenia celosioides* was introduced anonymously to the Berlin Botanical Garden, from which it was described as new to science by the end of 1818 (Sprengel 1818). Although the source of introduction was not recorded, among European plant collectors in Brazil it was only Alexander Philipp Maximilian, Prince of Wied-Neuwied (1782–1867), who collected specimens early enough to bring the material to Berlin before the protologue of *G. celosioides* had been published. Prince Maximilian travelled in Brazil during 1815–1817, in the States of Rio de Janeiro, Bahia and Espírito Santo, mostly along the coast. He left for Europe in May 1817 and arrived at Lisbon on 1<sup>st</sup> July. Very shortly after the arrival he started to distribute dried specimens, seeds and even living collections which he donated to botanical experts and gardens in Germany and Belgium. Large amounts of seeds were sent to Ghent, Antwerp and Enghien, but it is unknown which seeds were sent to Germany (Moraes 2009). A herbarium voucher of *M. paniculata* numbered 53 by C.G.D. Nees von Esenbeck, a German botanist who received a large set of specimens from Prince Maximilian for collaborative work, can be found in the Nees herbarium at BR. This number was cited by Walter (1909) as referable to *M. paniculata* (specimen destroyed at B). Along with the characters stated in the protologue of *G. celosioides*, this fact provides indirect evidence that Prince Maximilian’s collections were the likely source of the introduction of *G. celosioides* to the Botanical Garden in Berlin. There are no specimens of the original material of *G. celosioides* in existence, which probably was acquired by B and then destroyed (Stafleu and Cowan 1985). Since the characters of the Prince Maximilian’s specimen at BR are in good agreement with the protologue of *G. celosioides*, we designate this specimen as a neotype of the species name. This name is therefore the earliest one available for the species also known as *M. paniculata*.

=*M. paniculata* Moq. in DC., Prodr. 13(2): 18 (1849), **syn. nov.** Lectotype (designated by Nowicke 1968: 350): BRAZIL. Bahia, “Serra Jacobina” [Villa do Barra, according to Moquin-Tandon (1849)], [1837], Blanchet 2709 (K000601204! islectotype P00743956!).

=*M. longebracteata* H. Walter, Pflanzenr. (Engler) 39: 129 (1909), **syn. nov.** Lectotype (Senikov & Sukhorukov, designated here): BRAZIL. Prov. Bahia, zwischen den Campos und Vittoria, [1815], Sellow 359 (B 10 0250568! islectotypes – B 10 0250569! B 10 0250570!). Note: The main character distinguishing *M. longebracteata* from *M. paniculata* is the length of the bracts that is equal to those of the flowers (Walter 1909). However, the length of the bracts in both taxa is equal to that of the flower buds, and the length of fully opened flowers is greater than the length of the subtending bract.




**Figure 14.** Neotype of *Galenia celosioides* (BR0000005575398).

**Description.** Annuals or biennials; stems erect, up to 100 cm; leaves petiolate (petioles up to 2.0 cm), blades 3.0–10.0(12.0) cm long, 0.2–2.0 cm wide, cuneate, lanceolate to oblong, rarely ovate, glabrous or their margins and mid-rib below covered with papillae; inflorescence a spike, long and spreading, whip-like; flowers with a bract and two filiform bracteoles (often not well-visible), bracts longer than flowers at the beginning, then equal to the perianth segments; pedicels up to 1.0 mm at fruiting, perianth segments 5, oblong or ovoid; stamens 5–8; stigmas 2, thick; fruit with scattered short outgrowths, fruit body (1.0)1.1–1.4 × 0.9–1.1 mm, 1.5–2 times as long as the perianth (Fig. 4E, F); pericarp readily scraped off the seed; seed ~1.0 mm, with rough surface (Fig. 4H).

**Habitat.** Forest margins, roadsides, river banks, on sandy and rocky substrates; altitudes up to 1000 m a.s.l.

**Distribution.** (Fig. 15) Eastern South America.

**Specimens examined.** BRAZIL (selected specimens): **Alagoas:** Maceió, 4 Sep 1987, S. Tsugaru & Y. Sano B-1470 (NYBG01014516 – image!); **Bahia:** Gloria Mun., Barra do Tarrachil, 17 Jul 1962, G. Eiten & L.T. Eiten 4966 (K); Victoria da Conquista, 17 Jan 1965, E. Pereira & G. Pabst 34875 (M); Rio Cumbuca, Mucugê, 4 Feb 1974, R.M. Harley 15971 (K, P04598066, U1473440); Serra de Itiuba, Itiuba, 19 Feb 1974, R.M. Harley 16204



**Figure 15.** Distribution map of *Microtea celosioides*.

(P04598067, U1473439); Milagres, 6 Mar 1977, *R.M. Harley* 19451 (U1473444); Morro do Chapéu Mun., 980 m, 1 Jun 1980, *R.M. Harley* 22970 (E, K, U1473442); Cachoeira do ferro Doido, Morro do Chapéu, 30 Nov 1980, *A. Furlan & al. s.n.* (PACA 76292); **Ceará:** Cedro, May 1933, *Ph. von Luetzelburg* 26522 (M); Jaguaribe Mun., Maciço do Pereiro, 12 Apr 2011, *A.M. Miranda & K. Manso* 6327 (HUEFS178925 – image!); **Distr. Federal:** Brasilia, 27 Oct 1965, *H.S. Irwin et al.* 9600 (LE); Paranoá, 13 Dec 1965, *H.S. Irwin & al.* 11258 (IAN122990 – image!); **Espírito Santo:** Linhares, 20 Apr 2011, *J. Meirelles et al.* 624 (HUEFS193997 – image!); **Goiás:** Serra Dourada, 19 Jan 1966, *H.S. Irwin & al.* 11812 (NYBG00864091 – image!); **Maranhão:** Loreto Mun., Loreto city, 11 Feb 1970, *G. Eiten & L.T. Eiten* 10581 (K); Carolina Mun., Rodovia, 24 Apr 2008, *G. Pereira-Silva et al.* 13260 (CEN00091428 – image!); **Mato Grosso:** Barra do Garças, 17 Jun 1966, *D.R. Hunt* 6050 (K); Vale dos Sonhos, Barra do Garças Mun., 93 km S of Xavantina, 9 Nov 1968, *R.M. Harley & al.* 10980 (K); **Mato Grosso do Sul:** Estrada Forte Coimbra, Lagoa do Jacadigo, Corumbá, 1 May 1989, *A. Pott et al.* 4744 (PACA); Faz. Retirinho, Aquiduana, 23 Nov 1989, *A. Pott et al.* 5433 (PACA); Maracaju Mun., Serra de Maracaju, 12 Nov 1993, *G. Hatschbach et al.* 58937 (H1679509, K); **Minas Gerais:** Serra de Belo Horizonte, Belo Horizonte, 10 Feb 1927, *W. Hoebne s.n.* (PACA 76836); Diamantina, 3 Jun 1955, *E. Pereira* 1680 (PACA); Estrada Diamantina–Milho Verde, 33 km from Diamantina, 3 Dec 1981, *N. Hensold et al. s.n.* (PACA 76282); Caeté Mun., Serra da Piedade, 10 Jan 1982, *N. Hensold et al. s.n.* (PACA 76294); Subida ao Pico do Itambé, Santo Antonio do Itambé, 5 Apr 1982, *A. Furlan et al. s.n.* (PACA 76296); Tiradentes, 6 Jun 1984 (PACA 76835); São Thomé das Letras, 30 Oct 1984, *I. Cordeiro et al. s.n.* (PACA 76291); Serra do Cabral, Joaquim Felício, 21 Nov 1984, *R.M. Harley et al. s.n.* (PACA 76295); N of Grão Mogol, 27 Nov 1984, *R.M. Harley et al. s.n.* (PACA 76283); Estrada Serra-Diamantina, Trinta Réis, 27 Jan 1986, *N.L. Monezes et al. s.n.* (PACA 76284); Itacambira, Rodovia to Juramento, 14 Feb 1988, *J.R. Pirani et al.* 2272 (PACA); Caeté Mun., Morro da Piedade, Serra da Moenda, 24 Apr 1990, *J.A. Paula & S.B. Velten s.n.* (PACA 76847); Pico de Itabirito, Itabirito, 21 May 1994, *W.A. Teixeira s.n.* (PACA 76841); Retiro das Pedras, Brumadinho, 14 Dec 1998, *J. R. Stehmann & C.E.S. Ferreira* 2399 (PACA); Faz. Santana, Salto da Divisa, 21 Aug 2003, *J.A. Lombardi et al.* 5333 (PACA); Vilarejo do Funil, Rio Preto, 21 May 2004, *F.R.G. Salimena et al.* 1284 (PACA); Parque Estadual de Grão Mogol, Grão Mogol, 13 Jun 2006, *C.V. Vidal* 187 (PACA); Lima Duarte, 20 Nov 2006, *F.M. Ferreira et al.* 1139 (K); Santana do Riacho Mun., Cachoeira Véu de Noiva, 15 Mar 2007, *M.S. Marchiorotto* 353 (PACA); Serra de Antônio Pereira, Samarco, Ouro Preto, 4 May 2007, *M. Messias et al.* 1296 (PACA); Serra do Lenheiro, São João Del Rei, 25 Dec 2012, *M. Sobral* 15294 (PACA); **Paraíba:** Paraíba do Norte, Serra Borborema, 20 Mar 1913, *Ph. v. Luetzelburg* 12490 (M); Varzea, 23 Mar 1936, *Luetzelburg* 27000 (M); Areia, 17 Jun 1953, *J.C. de Moraes* 973 (IAN082599 – image!); Alagoa Nova Mun., Brejo Paraibano Reg., 5 Mar 2012, *E. Melo et al.* 10917 (PACA); **Pernambuco:** Floresta, 26 Aug 1994, *M. Sales* 319 (K); Buique, 20 May 1995, *K. Andrade et al.* 62 (K); **Piauí:** São Raimundo Nonato, 1 May 1978, *E. Laure* 195 (P06806987); Serra Branca, São Raimundo Nonato, 7 Feb 1986, *L. Emperaire* 2817 (P05197106); São Joao do Piaui Mun., Porfirio,

Aug 1995, *F.G. Alcoforado* Folho 480 (K); Caracol, 25 Feb 2011, *Melo et al.* 9216 (PACA 115982) as *M. longebracteata*; Caracol Mun., Serra das Confusões, Serra Grande, 18 Jul 2011, *A.A. Conceição et al.* 4037 (PACA); **Rio de Janeiro:** Rio de Janeiro [city], 1879, *A. Glaziou* 11440 (LE); Guanabara, 7 Feb 1964, *W. Hoehne* 5581 (P05197105); Corcovado, 5 Feb 1940, *B. Rambo* 3581 (PACA); Próximo de Recreio dos Bandeirantes, Guanabara, 4 Apr 1964, *W. Hoehne* 5708 (PACA); Est. da Guanabara, Pedra de Estauna, 31 Jan 1965, *Newton Santos* 5409 (M); **Rio Grande do Norte:** Grossos, Salina Salmar, 27 Jun 2007, *A.A. Roque* 137 (UFRN00004903 – image!); Nísia Floresta, Sete Lagoas, 18 Aug 2016, *V.F. Sousa* 456 (UFRN00022373 – image!); **Sergipe:** Canindé de São Francisco, 9 Sep 2014, *K.M. Pimenta* 589 (RB00956693 – image!); Pirambú Mun., [without date] *M. Ramos & E. Santos* 98 (ASE0017438 – image!); **Tocantins:** Goias, Araguaina, 9 Mar 1982, *A. Krapovickas et al.* 37841 (G); Rodovia, 1 Feb 2012, *R.M. Harley et al.* 56656 (K); **PARAGUAY:** Valenzuela, 18 Mar 1884, *B. Balansa* 4571 (B, P04598095); Tobati, [without date] *E. Hassler* 3981 (P03321381); Dept. Cordillera, Cordillera de Altos, 6 Mar 1984, *A. Schinini* 23957 (G); Dept. Cordillera, Tobati, 8 Feb 1991, *E. Zardini & C. Velázquez* 26244 (B); Paraguay, Concepcion, Paso Horqueta, 18 Nov 1993, *E. Zardini & T. Tilleria* 37460 (MW0581802).

### **3. *M. papillosa* M.S.Marchioretto & J.C.de Siqueira, Pesquisas, Botânica 48: 30 (1998).**


**Holotype.** BRAZIL. Minas Gerais, Estrada Conselheiro Mata, a 2 km do asfalto, Diamantina, 11 April 1982, *L. Rossi, A. Furlan, N.L. Menezes, N. Hensold, H.L. Wagner & E.M. Isejima* 3317 (PACA!).

**Description.** Perennial with a taproot or rarely annual (?) herb with caudex; stems erect, densely covered with short papillae; leaves papillate, oblong or lanceolate, appressed or somewhat spreading, cuneate, 2.0–4.0 cm long and 0.2–0.7 cm wide; inflorescence a spike; flowers subsessile, with the pedicels 0.25–0.5 mm long, with a bract and two bracteoles; perianth segments 5, oblong, greenish; stamens 6–8; stigmas 2, thick; fruit 1.1–1.25 × 0.9–1.1 mm, with short finger-shaped outgrowths (Fig. 5A, B); seed 0.9–1.0 mm, with rough surface (Fig. 5D). Morphologically, this species is most similar to *M. celosioides* and is distinguished by the papillate stems and leaves.

**Habitat.** Sandy substrates at altitudes 500–1400 m a.s.l.

**Distribution.** (Fig. 16) Endemic to Minas Gerais (Diamantina Mun.), Brazil.

**Specimens examined.** BRAZIL. Minas Gerais: Diamantina Mun., Estrada Conselheiro Mata, a 2 km do asfalto, 11 Apr 1982, *L. Rossi et al.* 3322 (PACA); N of Grão Mogol, 900–1000 m, 27 Nov 1984, *R.M. Harley et al.* 36121 & 37090 (K); 15 km from Diamantina towards Mendanha, 2 Dec 1984, *B. Stannard et al.* 36280 (K); Itacambira, 1220 m, 11 Nov 1988, *J.R. Pirani et al.* 2272 (K); 5 km W de Diamantina, 1200 m, 16 Feb 1991, *M.M. Arbo et al.* 5220 (K); Diamantina Mun., 14 km from Diamantina, 30 Jan 2000, *R.C. Forzza & R. Mello-Silva* 1486 (G, K).



**Figure 16.** Distribution map of *Microtea papillosa*.

#### 4. *M. scabrida* Urb., Ber. Deutsch. Bot. Ges. 3: 325 (1885).


- ≡*M. paniculata* Moq. var. *scabrida* (Urb.) Kuntze, Revis. Gen. Pl. 3(3): 268 (1898). Lectotype (Sennikov & Sukhorukov, designated here): BRAZIL. [Without exact location and date] *F. Sellow s.n.* (F0BN005735, image!). Note: The lectotype is chosen according to the protologue (Urban 1885). Urban's collections of *M. scabrida* in B have probably been missing since 1945.
- =*M. paniculata* var. *latifolia* Kuntze, Revis. Gen. Pl. 3(3): 268 (1898). Lectotype (Sennikov & Sukhorukov, designated here): BOLIVIA. Rio Yapacani, June 1892, *O. Kuntze s.n.* (B!).
- =*M. foliosa* Chodat, Bull. Herb. Boissier, ser. 2, 3: 418 (1903). Lectotype (designated by Nowicke 1968: 351): PARAGUAY. In regione collium “Cerros de Tobaty”, September 1900, *E. Hassler 6254* (MO216419, image!) isolectotypes – P00743942! K000601209!. Note: This species was synonymized with *M. paniculata* (=*M. celosiooides*) by Marchioretto and de Siqueira (1998). We have seen the original specimens cited in the protologue (Chodat and Hassler 1903): the specimens with the numbers 6254 (MO216419 – image! K000601209! P00743942!), 1649 (P03321197), 1988 (P00743946!) and 1988b (B! P00743944!) are *M. scabrida*, and the specimens with the number 7605 (K000601208! P00743941!) belong to *M. sulcicaulis*, a species described in the same article by Chodat and Hassler (1903). The authors mentioned in the publication three *Microtea* species:

*M. foliosa*, *M. paniculata* and *M. sulcicaulis* (Chodat and Hassler 1903), and *M. scabrida* was omitted in the species list. The protologue of *M. foliosa* combines the characters of both *M. scabrida* and *M. sulcicaulis*, but the epithet “foliosa” belongs to *M. scabrida* with inflorescence leafy in its lower part. Based on the typification made by Nowicke (1968), and in agreement with Walter (1909), we synonymize *M. foliosa* with *M. scabrida*.

=*M. scandens* Rusby, Mem. New York Bot. Gard. 7: 239 (1927). Holotype: BO-LIVIA. La Paz, Iturralde, Ixiamas, alt. 244 m, 18 Dec 1921, M. Cárdenas 1942 (NYBG01163848, image!).

**Description.** Leaning or twining perennial herb up to 150 cm, glabrous or slightly scabrid; leaves ovate or oblong (Fig. 17A), long-petiolate (petioles 1.0–4.0 cm), blades 5.0–10.0 cm long and 2.0–4.0 cm wide, basally truncate, apically acuminate; inflorescence lax, a spike (Fig. 17B); flowers almost sessile (pedicels up to 1.0 mm), with a bract and two bracteoles; perianth segments 5, whitish or green, oblong or ovoid, stamens 6–7, stigmas 2–3; fruit 1.75–2.0 mm long and 1.6 mm wide, with finger-shaped outgrowths, some of them basally concrescent (Fig. 5E, F); seed 1.3–1.5 mm, with verrucous surface (Fig. 5H).

**Habitat.** Forests, shrub thickets; alt. up to 1000 m.



**Figure 17.** *Microtea scabrida*: **A** a fragment of the shoot (São Bento do Sul, Santa Catarina, Brazil, 23 Sep 2016) **B** inflorescence (São Bento do Sul, Santa Catarina, Brazil, 14 Dec 2013). Photographs by Paulo Schwirckowski.



**Figure 18.** Distribution map of *Microtea scabrida*.

**Distribution.** (Sub)tropical parts of South America (Fig. 18).

**Specimens examined.** ARGENTINA: Chaco prov.: Isla del Cerrito, 11 Dec 1971, A.G. Schulz 17927 (G); Corrientes prov.: Itati, 6 Feb 1964, T.M. Pedersen 68 (P04598111); Santo Tome Dep., Potrero Luna, 8 Dec 1981, S.G. Tressens et al. 1612 (G); Ituzaingö Dept., Puerto Mora, 11 Dec 1973, A. Krapovickas et al. 24329 (RB00272365 – image!); Entre Ríos prov.: Concepción del Uruguay, Dec 1917, L. Hauman s.n. (BR); Dep. Uruguay, 31 Mar 1967, T.M. Pedersen 8202 (E00621962, P04598112); Concordia Dept., Nueva Escócia, 24 Nov 1988, N.M. Bacigalupo et al. 863 (B); Misiones prov.: Dept. San Ignacio, El Colorado, 24 Jun 1946, G.J. Schwarz 2845 (L1678246); Dept. San Pedro, El Alcazar, 30 Mar 1949, E. Schwindt 1428 (BR); Cainguás Dept., Oro Verde, 29 Apr 1949, G.J. Schwarz 7789 (G); Candelaria Dept., Santa Ana, 10 Mar 1951, J.E. Montes 15101 (P05197102); Leandro N. Alem Dept., Paso Carreta, 9 Mar 1969, A. Krapovickas et al. 15003 (G); BOLIVIA: see lectotype of *M. paniculata* var. *latifolia* (B); San Rafael, 27 Mar 1902, R.S. Williams 222 (BM000019282, K); La Paz, Upper Rio Beni, Dec 1906, J.W. Evans 53 (BM000019216); Santa Cruz Dept., Rio Yapacani, 8 Mar 1926, J. Steinbach 7498 (BM000019275, E, K, U1473457); Prov. Sara, Dept. Santa Cruz, Rio Yapacani, 8 Mar 1926, J. Steinbach 7498 (G); La Paz, Ixiamas, 18 Dec 1921, M. Cárdenas 1942

(NYBG – image!); **BRAZIL: Acre:** Santa Rosa Mun., Rio Chambuiacu, 14 Mar 2002, *D.C. Daly et al.* 11320 (NY00865147 – image!); **Amazonas:** Seringal São Francisco, Aug 1911, *E. Ule* 9361 (G, K, L1678244); **Bahia:** [without exact location and year] *Blanchet s.n.* (LE); Rio Grongogy, Nov 1915, *H.M. Curran* 148 (US01344703 – image!); Milagres, 6 Mar 1977, *R.M. Harley* 19451 (E); Abaira Mun., 31 Jan 1992, *J.R. Pirani et al.* 51372 (E00324053); **Minas Gerais:** [without exact location] 1816–1821, *A. Saint-Hilaire* 43 (P04598127); **Paraná:** Tomazina, Barra Grande, 29 Jan 1911, *P. Dusén* 11265 (K); Cerre Azul Mun., Ribeirao do Veado, 9 Feb 1960, *G. Hatschbach* 6725 (L1678240); Foz do Iguaçu Mun., Parque Nacional, 18 Feb 1963, *G. Hatschbach* 9737 (U1473447); Xambre Mun., 10 Dec 1965, *G. Hatschbach* 13304 (B, P04598113); Icaraima & Porto Camargo, 20 Jan 1967, *G. Hatschbach et al.* 4285 (U1473449); Icaraima Mun., Porto Camargo, 20 Jan 1967, *G. Hatschbach* 15765 (B, P04598114); Altônia Mun., Porto Byington, 23 Jan 1967, *J.C. Lindeman & J.Y. de Haas* 4396 (K); Cerro Azul Mun., Cabeceiras do Riberao do Tigre, 16 Dec 1992, *G. Hatschbach & O.S. Ribas* 58457 (G); Cerro Azul Mun., Rua Serra da Paranapiacaba, 16 May 1997, *G. Hatschbach et al.* 66536 (SJRP00009310 – image!); **Paraíba:** Areia, 15 May 1944, *J.M. Vasconcellos* 240 (RB00272395 – image!); **Pernambuco:** [without exact location] 1838, *M. Gardner* 1738 (G); Recife Mun., Caxangá, 29 Jul 1887, *Ridley et al. s.n.* (BM000019259); Recife, Mar 1936, *S. Vasconcellos* 4097 (US01344756 – image!); **Rio de Janeiro:** [without exact location] 1816–1821, *A. Saint-Hilaire* 376 (P04598132); Maricá Mun., Itaipuáçu, 27 Jan 1935, *Brade* 29310 (B); Niterói, 6 Mar 1998, *M.C.F. dos Santos* 153 (RB00766894 – image!); **Rio Grande do Sul:** [without exact location] 1816–1821, *A. Saint-Hilaire* 2711 (P04598128); Iraí, Nov 1949, *K. Emrich s.n.* (PACA 48169); **Santa Catarina:** [without exact location] Jul 1840, *F. Müller* 460 (K); Itapiranga ad fl. Uruguai superius, 16 Feb 1934, *B. Rambo* 1788 (PACA), São Bento do Sul, Rio Natal, 14 Dec 2013, *P. Schwirkowski* 128 (FURB36981 – image!); **São Paulo:** [without exact location] 25 Feb 1874, *A. Glaziou* 2007 (P04598040); Serra de São Pedro, São Pedro, 22 Dec 1965, *J. Mattos & N. Mattos* 13028 (PACA); **PARAGUAY: Asunción:** 13 Sep 1874, *B. Balansa* 1988 (G, K, P04459470); **Caa-guazú:** Caaguazú city, Nov 1874, *B. Balansa* 1988b (B, P00743944); **Caazapá:** San Juan Nepomuceno, 12 Dec 1989, *I. Basualdo* 2794 (G); National Park Caaguazu, 24 Nov 1997, *E.M. Zardini & A. Benitez* 47391 (P05197101); **Cordillera:** Tobati, 10 Jan 1903, *K. Fiebrig* 677 (E, G, L, M); Tobati, 14 Jan 1903, *F. Fiebrig* 737 (E, G, K, L1678243, LY, M) sub *M. foliosa*; Cerro Tobati, 28 Oct 1987, *R. Degun & E. Zardini* 447 (G); Cerro Ybitu Silla, Tobati, 28 May 1988, *E. Zardini* 4319 (G); **Guairá:** Villarica, 13 Sep 1874, *B. Balansa* 1988 (P00743946); Villarica, 13 Nov 1945, *G.W. Teague* 444 (BM000019276); Colonia Independencia, 30 Mar 1972, *T.M. Pedersen* 10122 (K, P04598115); Tebicuary, 17 Nov 1978, *L. Bernardi* 18685 (G); Melgarejo, 13 Mar 1989, *E. Zardini & C. Velásquez* 11391 (E00047207, K); Cordillera de Ybytyruzu, 28 May 1989, *E. Zardini & U. Velásquez* 12354 (G); **Paraguári:** 15 Dec 1875, *B. Balansa* 1988 (P04598096); prope Sapucay, 1885–1895, *E. Drake* 1649 (P03321197); Sapucay, Jul 1913, *E. Hassler* 11878 (BM000019235, E, G); La Rosada, 12 Dec 1979, *G. Schmeda* 234 (G); Ybicui National Park, La Po-

sada, 3 Feb 1992, *E. Zardini & P. Aquino* 30230 (G); Estero del Ypoá, Trinchera Cué, 5 Aug 1993, *E.M. Zardini & T. Tilleria* 36838 (B); **Alto Paraná**: [without exact location] 1909, *K. Fiebrig* 5468 (BM000019222, E, G, L1678245, LY, M); **PERU**: reported for the Dept. Madre de Dios (Brako and Zarucchi 1993); **URUGUAY**: [without exact location and year] *J. Tweedie s.n.* (BM).

### **5. *M. sulcicaulis* Chodat, Bull. Herb. Boissier, ser. 2, 3: 419 (1903).**


**Lectotype.** (designated by Nowicke 1968: 351, first-step lectotype; Sennikov & Sukhorukov, second-step lectotype designated here): PARAGUAY. In regione fluminis Tapiroguay, Aug.[ust] [without year], *E. Hassler* 4328 (K000601203! isolectotypes B! G! P00743937! P00634433!). Note: This species was synonymized with *M. celosioides* by Marchiorotto and de Siqueira (1998, sub *M. paniculata*). We reinstate *M. sulcicaulis* to specific rank due to (1) strongly perennial life history, (2) stiff (not spreading) inflorescences, (3) larger fruit diameter, (4) alveolate seed surface (in *M. celosioides* it is rough but without alveolae), and (5) predominant distribution in subtropical South America (Paraguay, South Brazil, and SE Bolivia).

**Description.** Perennial herb with a taproot; stems up to 60 cm, sparsely branched, glabrous; rosulate leaves at least partially persistent, appressed to the stem, lanceolate to narrowly oblong, 3.0–8.0 × 0.5–1.0 cm, tapering into the short petiole; caudine leaves numerous, continuously decreasing in size towards the inflorescence, sessile; inflorescence a spike, branched, with lateral shoots directed upwards and not spreading or whip-like; pedicels up to 0.5 mm, flowers with a bract and two bracteoles; perianth segments 5, greenish with white margin or yellowish, ovoid; stamens 5–8; stigmas 2, thick; fruit 1.5–2.0 × 1.30–1.50 mm (Fig. 6A); pericarp readily scraped off the seed, with small finger-shaped outgrowths up to 0.3 mm (Fig. 6B); seed 1.3–1.5 mm, with alveolate surface (Fig. 6D).

**Habitat.** Rocky and sandy substrates at altitudes up to 1500 m a.s.l.

**Distribution.** Subtropical South America (Fig. 19).

**Specimens examined.** **BOLIVIA** (the first record for the country): **Santa Cruz Dept.**, Fortin Suarez Arana, 19 Oct 1977, *C. Evrard* 8203 (BR) as *Microtea* sp. Previously not reported for this country (Jørgensen et al. 2015); **BRAZIL: Bahia**: RiodeContasMun., PicodasAlmas, 6Nov1988, *R.M. Harley* et al. 25939 (K); Agua QuenteMun., Pico das Almas, 1400m, 13Dec1988, *R.M. Harley* & *D.J.N. Hind* 27232(K); Mun. de Abaira, 13 Dec 1993, *W. Ganey* 2615(K); **Mato Grosso do Sul**: Pacuri, 12 Dec 1982, *G. Hatschbach* 45918 (G); **Rio de Janeiro**: 1857, *A. Glaziou* 16311 (BR); **PARAGUAY: Alto Paraguay Dept.**: Fuerte Olimpo, 24 Oct 1946, *T. Rojas* 13637 (E); **Amambay Dept.**: Pedro Juan Caballero, Bela Vista, 23 Nov 1963, *J. Correa Gomes* 1465 (G); Pedro Juan Caballero, 19 Oct 1986, *T.M. Pedersen* 14669 (G); Estancia 5 Hermanos, 2 Nov 1997, *I. Basualdo* 6437 (G); National Park Cerro Corá, Cerrado, 350 m, 10 Nov 1999, *E.M. Zardini* & *P. Baéz* 52218 (P05197108); **Caazapá Dept.**: Tavai, 7 Dec 1988, *F. Mereles* 2122 (G); **Canindeyú Dept.**: Nanduro Kai, 1 Nov 1978, *L. Bernandi*



**Figure 19.** Distribution map of *Microtea sulcicaulis* (circles) and *M. babiensis* (star).

18347 (G); Lagunita, 23 Sep 1988, T.N. Pedersen 15089 (G); [without exact location] 11 Oct 1996, B. Jiménez & G. Marin 1616 (BM000527219); Mbaracayú Natural Reserve, 31 Oct 1998, E.M. Zardini & I. Chaparro 49429 (B 10 0058099); **Central Dept.:** Ypacarai Lake, Dec 1913, E. Hassler 12395 (BM000019236, E, G, L1678248, LY); **Concepción.:** nr Concepción, Oct 1901, E. Hassler 7605 (G, K000601208, LY, sub *M. foliosa*); **Cordillera Dept.:** Piribebuy, 11 Jan 1877, B. Balansa 2576 (P04598143); Colonia Rosado, 26 Oct 1986, A. Schinini & E. Bordas 24850 (G); **San Pedro Dept.:** Yaguareté forest, 24 Aug 1995, E. Zardini & A. Vargas 43638 (MW).

## 6. *M. babiensis* M.S.Marchioretto & J.C.de Siqueira, Pesquisas, Botânica 48: 11 (1998).

**Holotype.** BRAZIL. Estado Bahia, Município de Salvador, ca. 30 km a N do Centro de Salvador, Estrada para o aeroporto, arredores de Itapuá, dunas [Bahia State, Salvador Municipality, ca. 30 km N from Salvador city, on the way to the airport, surroundings of Itapuá, dunes], 23 May 1981, Carvalho, Mori & Boom 706 (CEPEC! isotypes – ALCB, NY).

**Description.** Perennial herb or dwarf subshrub, glabrous, up to 30 cm tall; caudex well-developed; stems erect or ascending; rosulate leaves up to 9.0(10.0–12.0) cm, obovate or oblong, mostly persistent at fruiting; caudine leaves rhombic, ovate or obovate, cuneate and shorter (up to 2.0 cm) than the rosulate leaves; inflorescence a spike, mostly one-sided; flowers sessile or very shortly pedicellate (pedicels up to 0.5 mm); bracteoles present, very short, perianth segments 5, greenish with white margins, oblong or ovoid; stamens 8; stigmas 2, thick; fruit roundish, 1.1–1.3 × 1.0–1.2 mm, with short finger-shaped outgrowths (Fig. 6E, F); seed 1.1–1.3 mm, with rough surface (Fig. 6H). Note: We were unsuccessful in extracting DNA from the available specimens. However, the characters of this species indicate its position within the type subgenus.

**Habitat.** Sand dunes at altitudes up to 500 m a.s.l.

**Distribution.** Endemic to Bahia state, Brazil (Fig. 19).

**Specimens examined.** BRAZIL. Bahia State, Salvador Mun., Itapuá, 27 Feb 1983, *P. de Queiroz* 496 (ALCB); Itapuá, 20 Apr 1983, *P. de Queiroz* 544 (HUEFS, PACA); Bahia [state], Salvador, Dunas de Itapuá, nr Hotel Stella Maris, N from Condomínio Alamedas da Praia, 8 Jun 1993, *P. de Queiroz* 3211 (K, PACA).


## 7. *M. portoricensis* Urb., Ber. Deutsch. Bot. Ges. 3: 324 (1885).

**Lectotype.** (designated by Nowicke 1968: 348, first-step lectotype; Sennikov & Sukhorukov, second-step lectotype designated here): PUERTO RICO. Cabo Rojo, in campis, 20 January 1885, *P. Sintenis* 717 (S-R-3531 – image! isolectotypes P00743938! B 10 0296282! B 10 0296820! BM000019288! G! K000601210! P00743938! P00743939! L1678241! L1678242! LE!).

**Description.** Annual, glabrous, stems decumbent, up to 30 cm; rosulate leaves up to 8.0 cm, long-petiolate (petioles up to 3.0 cm), obovate or oblong, mostly persistent at fruiting; caudine leaves rhombic or ovate, cuneate; inflorescence a spike, one-sided, flowers sessile or very shortly pedicellate (pedicels at fruiting ~1.0 mm); bracteoles mostly absent or tiny; perianth segments 4–5, greenish, lanceolate or oblong; stamens 4–5; stigmas 2, thick; fruit almost orbicular or broadly ovate, 0.9–1.1 × 0.9–1.0 mm, reticulate, without any projections (Fig. 7A, B); seed 0.9–1.0 mm, with rough surface (Fig. 7D). Note: This species is assigned here to the type subgenus, although it was not included in the molecular analysis. Morphologically, it is closely related to *M. debilis*.

**Distribution.** Endemic to the Greater Antilles (Fig. 20).

**Specimens examined.** DOMINICAN REPUBLIC: reported by Moscoso (1943) and Nowicke (1968 [“herbarium S”] (n.v.); HAITI: reported by Nowicke (1968), herbarium S (n.v.); CUBA: Santiago de las Vegas, 10 May 1904, *H.A. van Hermann* 121 (BM00019279, P04598109); Ramón de la Sagra, [without date] *A. Jamain s.n.* (P04598110); Limonar, 1864, *Angel* 771 (LE); Havana, 3 Nov 1921, *E.L. Ekman* 13408 (G); PUERTO RICO: see type specimens; Cabo Rojo, 1864, *Grosourdy* 13 (P04598159); Laguna Derrumadero, nr Bayaguana, 7 Sep 1981, *T. Zanoni & M. Mejia* 16407 (NY01509884 – image!).



**Figure 20.** Distribution map of *Microtea portoricensis*.

***Microtea* subgen. *Ancistrocarpus* (Kunth) Sukhor. & Sennikov, comb. & stat. nov.  
urn:lsid:ipni.org:names:77193077-1**


≡*Ancistrocarpus* Kunth, Nov. Gen. Sp. [quarto] 2: 186 (1817). ≡*Microtea* subgen. *Eumicrotea* H.Walter, Pflanzenr. (Engler) 39: 127 (1909), nom. inval. (Art. 21.3). Type: *M. maypurensis* (Kunth) G.Don.

**Description of the subgenus.** Annuals; bracteoles present; pedicels conspicuous (1.35–3.0 mm long); flowers single or clustered (2–6 per node); stigmas 3–5, thin. The species are mostly distributed in Brazil, with irradiations to the neighbouring countries.

#### 8. *M. glochidiata* Moq. in DC., Prodr. 13(2): 18 (1849).

**Lectotype.** (Sukhorukov & Sennikov, designated here): BRAZIL. Piauhy [Piauí], near Boa Esperança [-6.809768, -41.380520], February 1839, Gardner 2311 (K000601202! isolectotypes B! P00743948! P00743949! P00743950!). Notes: The lectotype specimen, Gardner 2311, was collected when G. Gardner used the hospitality of Rev. Marcos de Araújo Costa (Gardner 1846), who was a land-owner, amateur botanist and educator in the province of Piauí, Brazil. The other collections cited in the protologue (Moquin-Tandon 1849), Blanchet 2680 from Villa di Barra, belong to *M. maypurensis* (P00798999) and *M. celosioides* (BM000019298).

**Description.** Annual, glabrous, up to 40 cm; stem erect, usually branched from the base with mostly persistent rosulate leaves; leaves linear to oblong, cuneate, 1.0–



**Figure 21.** Distribution map of *Microtea glochidiata*.

3.0 cm long, 0.1–0.4 cm wide; inflorescence thyrsoid, flowers 1–6 at each node, pedicels at fruiting 1.5–2.5(3.0) mm long; each flower with a bract and two bracteoles; perianth segments 5, white or yellowish, broadly ovoid or suborbicular, 0.8–1.0 mm long, imbricate; stamens 5–8; stigmas 3–5, filiform; fruit orbicular, its body 1.0–1.2 mm, covered with plumose outgrowths 0.4–0.7 mm long (Fig. 7E, F); pericarp easily scraped off the seed; seed ~1.0 mm, with slightly alveolate surface (Fig. 7H).

**Habitat.** Forest margins, roadsides, ruderal sites, on sandy substrates at altitudes up to 500 m a.s.l.

**Distribution.** (Fig. 21) Eastern tropical Brazil.

**Specimens examined.** BRAZIL: Bahia: Jacobina, [without date] *Blanchet* 2630 (P04598062); Nova da Rainha, [without date] *Martius* 304/18 (M); Gentio do Ouro, 22 Feb 1977, *R.M. Harley* 18907 (E, K, M, P05197098, U1473427); Gentio do Ouro Mun., nr Santo Inácio, 26 Feb 1977, *R.M. Harley* 19120 (E, P05197099, U1473426); Upper São Francisco river, Caldeirão Grande, 500 m, 18 Apr 1980, *R.M. Harley* 21511 (K); Riacho Grande, 4–5 km NE from Itatim, Santa Terezinha, 16 May 1983, *L. Noblick et al.* 3243 (PACA); Tucano Mun., 20 Feb 1992, *A.M. de Carvalho & D.J.N. Hind* 3841 (G, K, PACA); Paramirim Mun., Lago do Leito, 17 Jan 1997, *G. Hatschbach et al.* 65893 (H1693402); Rio de Contas Mun., 5 Feb 1997, *M.L. Guedes*

*et al.* 5118 (K); Remanso, 27 Feb 2000, *M.R. Fonseca et al.* 46314 (K, P000868555); Remanso, 10 Mar 2005, *L.P. Queiroz et al.* 10070 (HUEFS093066 – image!); **Ceará:** Sobral, 8 Apr 1984, *A. Fernandes* 12451 (ASE0023720 – image!); **Espírito Santo:** between Colatina & Patrimônio, 4 Dec 1971, *A.P. Duarte* 13986 (PACA); **Maranhão:** São Raimundo das Mangabeiras Mun., 15 Mar 1962, *G. Eiten & L.T. Eiten* 3674 (K); Barao do Grajau, 21 Jan 2012, *R.M. Harley et al.* 56455 (K); **Pernambuco:** Petrolina, 22 Jan 1970, *P. Carauta* 1008 (RB169035 – image!); **Piauí:** see lectotype and isolecotypes of *M. glochidiata*; Caracol, 25 Feb 2011, *E. Melo et al.* 9216 (PACA); **Rio Grande do Norte:** Jucurutu, 1 Jun 2008, *A.A. Roque* 570 (HUEFS164628 – image!).

## 9. *M. maypurensis* (Kunth) G.Don, Hort. Brit. [Loudon]: 98 (1830).

≡*Ancistrocarpus maypurensis* Kunth, Nov. Gen. Sp. [quarto] 2: 186 (1817). Lectotype (Sennikov & Sukhorukov, designated here): COLOMBIA. “In pratis Maypure”, ex herb. Bonpland, ex herb. Kunth s.n. (B 10 0296283!). Note: Kunth, who revised the main set of Humboldt's and Bonpland's collections at P, acquired a large part of this herbarium. After his death, the collections were transferred to the Botanical Museum in Berlin (Urban 1881), where they partly survived.

=*Ceratococca maypurensis* Humb. & Bonpl. ex Roem. & Schult., Syst. Veg., ed. 15, 6: 800 (1820). Holotype: COLOMBIA. Maypure, Cataracta, Orinoco, *A.J.A. Bonpland & F.W.H.A. von Humboldt* s.n. (B-W 06266-01!). Notes: *Ceratococca maypurensis* was apparently based on a duplicate of the collection on which *Ancistrocarpus maypurensis* was described earlier by Kunth. However, a specimen in the Herbarium of Willdenow was not accessible to Kunth at the time when the protologue was prepared (McVaugh 1955).

=*Potamophila parviflora* Schrank, Pl. Rar. Horti Monac. 2: tab. 63 (1821), nom. illegit., non R.Br. 1810.

≡*Ancistrocarpus schrankii* Ledeb., Index Seminum Horti Academicii Dorpatensis 1821 (Appendix I): 21 (1821). Described on the basis of plants cultivated in the Botanical Garden in Munich. Lectotype (Sennikov & Sukhorukov, designated here): [icon] fig. 63 in Schrank (1821). (Fig. 22). Note: von Ledebour (1821) intended (Art. 6.11) to introduce his new species name as a replacement name for the later homonym published by Schrank, and both names are therefore necessarily homotypic. Schrank (1821) published his new species names based on seeds sent or brought from Brazil by C.F. von Martius. Original herbarium collections of *Potamophila parviflora* Schrank were not found at M and most likely had never been prepared (H.-J. Esser, pers. comm. 2018), and the illustration published as part of the protologue is the only original element available for lectotypification (Art. 9.4). The protologue of *Potamophila parviflora* indicates that the new species has five styles (although only three are visible in the accompanying illustration) and a pericarp with setae. Coupled with pedicellate flowers and petiolate leaves, easily recognizable in the illustration, these characters are indicative of *Microtea may-*

*purensis*, of which *P. parviflora* (*Ancistrocarpus schrankii*) is a later synonym. This agrees with the conclusions of Moquin-Tandon (1849), Walter (1909) and Novicke (1968). Schrank (1821) stated that the new species was collected in “Brasilia prope sinum Omnium Sanctorum ad flumina locis umbrosis”. There is a gathering, Herb. Martius 2198 (M 0274659, M 0274661, M 0274662), which was collected by Martius in Bahia, Cachoeira, along the Paraguaçu River at the distance of ca. 35 km from Baía de Todos os Santos (Bay of All Saints), in December 1818 (see the route of Martius’ expedition in Tiefenbacher (1983)). This is the likely type locality of *P. parviflora*, and the gathering may be the voucher for the seed collection sent by Martius to the Munich Botanical Garden. The plants of this gathering are unmistakeably referable to *M. maypurensis*. Two other specimens, Herb. Martius 2309 (M 0274665, M 0274666), collected by Martius in Bahia, Monte Santo, in April 1819, were mistakenly labelled as *P. parviflora*. They clearly disagree with the protologue and belong to *M. celosioides*. This curatorial mislabelling probably occurred because of the confusingly similar localities on its label and in the protologue.


=*M. spengelii* Moq. in DC., Prodr. 13(2): 19 (1849). Neotype (Sennikov & Sukhorukov, designated here): BRAZIL. Bahia, Rio Belmonte, [Aug. 1816], *Maximilian, Prinz zu Wied* 53 (BR000005537679, image!). Note: Moquin-Tandon (1849) based the protologue of *Microtea spengelii* entirely on the description of *M. maypurensis* in Sprengel (1820). He stressed the diagnostic character of opposite leaves, which is almost impossible in *Microtea*. However, Sprengel’s description closely matches the characters of *M. maypurensis* except for the opposite leaves, and unlikely belongs to any other species. Moquin-Tandon (1849) assumed a technical error in Sprengel (1820), with which we agree. Sprengel noted specifically that he based the description on plants from Brazil. No relevant specimens survived. As a neotype, we designate a specimen of *M. maypurensis* collected in Brazil by Prinz Maximilian, a contemporary collection that may have been available to Sprengel.

=*M. glochidiata* f. *lanceolata* Chodat & Hassler, Bull. Herb. Boiss., ser. 2, 3: 418 (1903). Lectotype (Sennikov & Sukhorukov, designated here): PARAGUAY. Ad marginem silvae [prope] Caraguatay, Aug.[ust] [1897], E. Hassler 3126 (P00743952! isolectotypes P00743951! P00743953!). Note: Our synonymy confirms the opinion of Heimerl (1912).

**Description.** Annual or biennial, glabrous; stems erect, up to 60 cm, branched; rosulate leaves oblong, usually withered, lower leaves oblong or spatulate, cuneate, petiolate (petioles up to 2.5 cm), 3.0–8.0 cm long and (0.2–0.4)0.5–2.0 cm wide (sometimes narrower), acuminate; inflorescences a spike, not one-sided, often spreading; flowers solitary (rarely two per node), with a bract and two bracteoles, pedicellate (pedicels 1.5–3.0 mm), perianth segments 5, oblong to ovoid, white or yellowish; stamens 5–8, stigmas 3–5, filiform; fruit slightly protruding from the perianth or up to twice its length, fruit body 1.0–1.1 mm across, with outgrowths 0.2–0.5 mm long terminating in a group of 2–4 hooked hairs (Fig. 8A, B); pericarp readily scraped off the seed; seed ~1 mm, with rough surface (Fig. 8D).



**Figure 22.** Lectotype of *Ancistrocarpus schrankii* (fig. 63 in Schrank, 1821). Image provided by the library of Biological Faculty, Lomonosov Moscow State University.




**Figure 23.** Distribution map of *Microtea maypurensis* in its native range.

**Habitat.** Forests or ruderal sites; 0–1500 m.

**Distribution.** Tropical South America (Fig. 23).

**Specimens examined.** **BOLIVIA:** **La Paz Dept.:** Beni river, Jul 1886, *H.H. Rusby* 1379 (BM000019263, E, G, LE, P04598104); nr Guanay, 1892, *M. Bang* 1589 (B, BM000019206, E, G, K); Tipuani, Hacienda Simaco, 1400 m, Jan 1920, *O. Buchtien* 5404 (K, M); Tipuani, 1400 m, 29 Mar 1923, *O. Buchtien* 7290 (E, G); Mapiri, 850 m, 29 Nov 1926, *O. Buchtien* 694 (M); **Santa Cruz Dept.:** Ñuflo de Chávez prov., Pascana Ministro, 12 Jun 1995, *J.R. Abbott* 16976 (K); Ñuflo de Chávez prov., Lomerio, 14 Apr 1995, *F. Mamani & M. Saucedo* 817 (MEXU); **BRAZIL:** **Amazonas:** Rio Juruá, Juruá Miry, May 1901, *E. Ule* 5503 (G, L1678235); Rio Branco, Nov 1913, *J.G. Kuhlmann* 120 (RB00272533 – image!); Rio Negro between Ilha Uabetuba & Ilha da Silva, 14 Oct 1971, *G.T. Prance et al.* 15235 (NY00779040 – image!); Rio Jutai, 17 Nov 1975, *N.A. Rosa & L. Coelho* 563 (IAN151165 – image!); **Bahia:** [without exact location] 1834, *Blanchet s.n.* [herb. De Candolle] (G00687542); Jacobina, [without date] *Blanchet* 2588 (P00743954, a plant in the middle); Andaraí, 500–600 m, 13 Feb 1977, *R.M. Harley* 18632 (E, K, P05197100); Iaçu Mun., Lagedo Alto, 25 Sep 1984, *L.R. Noblick & M.J. Lemos* 3407 (PACA); Ilheus Mun., Rodovia, 10 Apr



**Figure 24.** The record of *Microtea maypurensis* as an alien plant in Indonesia.

1986, *J.L. Hage s.n.* (G); Dom Basilio Mun., 28 Dec 1989, *A. de Carvalho et al.* 2679 (G, PACA); Jacobina Mun., 30 Nov 1992, *M.M. Arbo et al.* 5450 (K); Iaçu Mun., Rio Paraguacu, 10 Apr 1992, *G. Hatschbach et al.* 56968 (G, K); Rio Jecuriçá, 16 Jan 1997, *M.M. Arbo et al.* 7276 (G, K); **Maranhão:** Carolina Mun., National Park Chapada das Mesas, 9 Apr 2016, *A.C. Sevilha et al.* 5723 (CEN00097885 – image!); **Minas Gerais:** Pedra Azul, 12 Dec 1984, *A.M. Giulietti et al.* 36297 (K); towards Jequitinhonha, Pedra Azul, 20 Oct 1988, *R.M. Harley et al.* 25232 (K, PACA); **Pará:** Marabá to Altamira, 16 Jan 1976, *P. Bamps* 5170 (BR); **Paraíba:** Esperança, 14 Sep 1958, *J.C. de Moraes* 1936 (RB00272388 – image!); **Paraná:** Londrina, 17 Nov 1969, *G. Hatschbach* 22898 (NYBG00779051 – image!); **Pernambuco:** [without exact location] 1838, *Gardner* 1138 (P04598076); Petrolina Mun., Tapera, Aug 1930, *D.B. Pickel* 50 (B, BM000019250); **Rio de Janeiro:** Rio de Janeiro [city], 1883, *A. Glaziou* 15355 (LE); **Roraima:** Rio Ajarani, 29 Apr 1974, *J.M. Pires et al.* 14408 (IAN144038 – image!); **São Paulo:** Rio Negro, between Ilha Uabetuba & Ilha da Silva, 14 Oct 1971, *G.T. Prance et al.* 15235 (G, K, M, P04598013, U1473419); **Sergipe:** Cristinapolis Mun., 2 Apr 1976, *G. Davidse et al.* 11810 (U1473430); **Tocantins:** Darcinopolis Mun., Rio Tocantins, 16 Apr 2008, *G. Pereira-Silva et al.* 12933 (CEN00091111 – image!); **COLOMBIA:** Rio Vaupés, 1852, *R. Spruce* 2546 (E, K); **ECUADOR:** El Oro, 18 km on road Huaquillas–Arenillas, [without date] *Harling & Andersson* 18835 (GB – image!); **GUIANA:** [without exact location] 1868, *Schomburgk* 835 (B, E, K, P04598085); Pirara, 1841, *Schomburgk* 325 (BM000019270); Region Upper Takutu, Essequibo,

1 Jun 1996, *D. Clarke* 1874 (U1473431); **PARAGUAY: Amambay Dept.**: between Rio Apa & Rio Aquidabán, 13 Feb 1908, *K. Fiebrig* 4928 (BM000019221, G, E, K, L); **Cordillera Dept.**: see lectotype of *M. glochidiata* f. *lanceolata*; Caraguatay, Aug 1900, *E. Hassler* 3126 (B, BM000019232, G, P00743953, P00743951, P00743952); **San Pedro Dept.**: [without exact location] 26 Oct 1953, *A.L. Woolston* 210 (K); **PERU: Cusco Region**: Paucartambo prov., Cusco dept., Pillcopata, 20 Jun 1959, *J. Infantes* 5913 (B); **Puno Region**: Carabaya prov., San Gaban distr., [without date] ex herb. Steudel 2288 (G, P04598108); **SURINAME**: nr Paramaribo, 12 Feb 1904, *van Hell* 150 (U1473435); Paramaribo, 10 April 1916, *J.A. Samuels* 8 (K); Paramaribo, 12 Apr 1916, *J.A. Samuels* 65 (BM000019265, L1678237, P04598014); Found as alien in Southeast Asia (Fig. 24); **INDONESIA**: Java, Pasuruan, 1924, *anonymous s.n.* (L1678238). Previously not reported from this region (Backer 1954).

## 10. *M. tenuifolia* Moq. in DC., Prodr. 13(2): 18 (1849).


=*M. maypurensis* var. *angustifolia* Moq. in DC., Prodr. 13(2): 18 (1849). Lectotype (Sennikov & Sukhorukov, designated here): BRAZIL. Jacobina Mountains in Bahia, 1836, *Blanchet* 2588 (P00798998!).

**Lectotype.** (designated by Nowicke 1968: 352): BRAZIL. Espírito Santo State, Pico d'Habira [Pico do Itabira], 1843, *Claussen* 392 (P00743993!).

**Description.** Annual or short-lived perennial herb with several or numerous stems 10–40 cm high; leaves sessile (sometimes rosulate leaves shortly pedunculate), cuneate, 10–30 mm long and 0.3–3.0(5.0) mm wide; inflorescence a spike; pedicels 1.35–1.7(2.5) mm; flowers with a bract and two bracteoles, perianth segments 5, white, oblong; stamens 6–8, stigmas 3–5, thin; fruit 0.9–1.1 × 1.0–1.1 mm (Fig. 8E), equal to perianth or slightly protruding; pericarp smooth or with small and scattered tubercles (Fig. 8F), more or less reticulate, readily scraped off the seed; seed 0.9–1.1 mm, with rough surface (Fig. 8H).

We report for the first time that the perianth/fruit ratio is a useful distinguishing character for this species. Also, the number of stigmas can be useful in delimiting *M. tenuifolia* and similar forms of *M. celosioides* with narrower leaves. The character set of *M. tenuifolia* supports its close relationship to *M. maypurensis*, especially the forms with reduced pericarp outgrowths. Remarkably, Moquin-Tandon (1849) described a new variety of *M. maypurensis* (var. *angustifolia* Moq.) represented by two specimens of *M. tenuifolia* (P00743955! and P00798998!) and one specimen containing two individuals, *M. tenuifolia* and a narrow-leaved *M. maypurensis* (leg. *Blanchet* 2588, P00743954!). The figure of *M. tenuifolia* in Calió and Pirani (2006) showing the tuberculate perianth is rather an exception, and the individuals with pericarp lacking the outgrowths have so far been collected more frequently.

**Habitat.** Forest margins, rocky places; 0–1000 m.



**Figure 25.** Distribution map of *Microtea tenuifolia*.

**Distribution.** Endemic to Eastern Brazil, found only in Bahia, Espírito Santo, Minas Gerais and Rio de Janeiro States (Fig. 25).

**Specimens examined.** **BRAZIL: Bahia:** 4 km SW of Belmonte, 23 Mar 1974, R.M. Harley 17305 (K, P04598081, U17305, U1473428); see lectotype of *M. maypurensis* var. *angustifolia*; Jacobina, [without date], Blanchet 153 (BM000019209); Morro do Chapeu, 1000 m, 1 Jun 1980, R.M. Harley 22924 (U1473452); **Espírito Santo:** see lectotype of *M. tenuifolia*; **Minas Gerais:** [without exact location] Feb 1839, Riedel 48 (P007990000); [without exact location] 1841, Claussen 4 (P00743934); [without exact location] 1845, Widgren s.n. (M); Caldas, 5 May 1870, A. Glaziou 11 (P04598116); Caldas, 15 May 1870, A.T. Regnell s.n. (B); Serra das Vertentes, Jun 1893, A. Glaziou 20437 (B, K, P04598119); Turvo, 24 Apr 1926, W. Hohne & A. Gehrt s.n. (PACA 76840); Buenópolis Mun., Curimatai, [without date] A. Glaziou 13127 (P04598093); Buenópolis Mun., Curimatai, [without date] A. Glaziou 19399 (P04598074); Serra do Espinhaco, 15 Feb 1969, H.S. Irwin et al. 23315 (G); Bacia do Córrego Escurona, Grão Mogol, 2 Nov 1987, M.C. Assis & al. s.n. (PACA 76297); Vale do Rio Itacambiruçu, Grão Mogol, 10 Dec 1989, A. Freire-Fierres et al. s.n. (PACA

76298); Vale do Córrego Escurona, Grão Mogol, 13 Jun 1990, A.A. Oliveira et al. s.n. (PACA 76299); Serra dos Inconfidentes, Pico do Itabirito, Itabirito, 4 Jan 1994, W.A. Teixeira s.n. (PACA); Estação Ecológica da Mata do Cedro, Carmópolis, 13 Jul 2004, L. Echternacht & T. Dornas 573 (PACA); Estação Ecológica da Mata do Cedro, Carmópolis, 23 Jan 2005, L. Echternacht & T. Dornas 830 (PACA); Grão Mogol Mun., Parque Estadual de Grão Mogol, 13 Apr 2006, C.V. Vidal 177 (PACA); Santana do Riacho Mun., Entre a Rodovia BR 251 e Grão Mogol, Cachoeira Véu de Noiva, 15 Mar 2007, M.S. Marchioretto 352 & 355 (PACA); **Rio de Janeiro:** nr Rio de Janeiro, Feb 1882, A. Glaziou 13127 (G, K); Rio de Janeiro, 1887, A. Glaziou 17748 (LE).

### Species excluded

All species cited below under *Microtea* belong to the South African genus *Lophiocarpus* Turcz., with a distinct position within Caryophyllales (Cuénoud et al. 2002, Schäferhoff et al. 2009) and with a different seed anatomy (Sukhorukov et al. 2015). The transfers of *Lophiocarpus* to *Microtea* were undertaken due to the morphological similarity of their members (Brown 1909), which is a case of homoplasy between phylogenetically distant Caryophyllales genera (Schäferhoff et al. 2009, Brockington et al. 2009).

*Microtea burchellii* (Hook.f.) N.E.Br., Bull. Misc. Inf. Kew 1909(3): 135 [1909]  
 $\equiv$  *Lophiocarpus burchellii* Hook.f. in Bentham & Hooker f., Gen. Pl. 3(1): 50 (1880);

*Microtea gracilis* A.W.Hill, Bull. Misc. Inf. Kew 1910(2): 56 [1910]  
 $=$  *Lophiocarpus polystachyus* Turcz., Bull. Soc. Imp. Naturalistes Moscou 16: 56 (1843);

*Microtea polystachya* (Turcz.) N.E.Br., Bull. Misc. Inf. Kew 1909(3): 135 [1909]  
 $\equiv$  *Lophiocarpus polystachyus* Turcz., Bull. Soc. Imp. Naturalistes Moscou 16: 56 (1843);

*Microtea tenuissima* (Hook.f.) N.E.Br., Bull. Misc. Inf. Kew 1909(3): 134 [1909]  
 $\equiv$  *Lophiocarpus tenuissimus* Hook.f., Hooker's Icon. Pl. 15: 50, tab. 1463 (1883).

### Acknowledgments

We thank Hajo Esser (Botanische Staatssammlung, Munich, Germany), Filip Verloove (Meise, Belgium) and Denis Melnikov (Komarov Botanical Institute, St.-Petersburg, Russia) for communication and discussion of the specimens kept in the herbaria M and LE, respectively; Geoffrey Harper for comments on the text; reviewers Walter Judd and Lucas Majure, editor Eric H. Roalson for their comments to the first draft of the manuscript, Orlando Vargas Ramírez and Paulo Schwirkowski for the excellent images of *M. debilis*

and *M. scabrida*, respectively, reproduced with permissions; staff of the library of Biological Faculty, Lomonosov Moscow State University (Moscow, Russia) for a scan of *Ancistrocarpus schrankii* illustration. The study was supported by the Russian Science Foundation (project 14-50-00029: carpological research by APS, MK and MN) and Scientific program AAAA-A16-116021660045-2 of the Department of Higher Plants, Lomonosov Moscow State University (revision of the herbaria in Moscow and St.-Petersburg).

## References

- Atha D (2004) Phytolaccaceae. In: Smith N, Mori SA, Henderson A, Stevenson DW, Heald SV (Eds) Flowering plants of the Neotropics. Princeton University Press, Princeton, 292–294.
- Backer CA (1954) Phytolaccaceae. In: van Steenis CGGJ (Ed.) Flora Malesiana, Vol. 4. Noordhoff-Kolff, Djakarta, 228–232.
- Bamps P (1974) Présence d'une Phytolaccacée américaine, *Microtea debilis* Swartz, sur la côte du Cameroun. Bulletin du Jardin Botanique National de Belgique 44(3/4): 439. <https://doi.org/10.2307/3667682>
- Behnke HD (1993) Further studies of the sieve-element plastids of the Caryophyllales including *Barbeuia*, *Corrigiola*, *Lyallia*, *Microtea*, *Sarcobatus*, and *Telephium*. Plant Systematics and Evolution 186(3–4): 231–243. <https://doi.org/10.1007/BF00940800>
- Behnke HD, Mabry TJ (1994) Caryophyllales: Evolution and Systematics. Springer Verlag, Berlin – Heidelberg – New York. <https://doi.org/10.1007/978-3-642-78220-6>
- Blattner FR, Kadereit JW (1999) Morphological evolution and ecological diversification of the forest-dwelling poppies (Papaveraceae: Chelidonioideae) as deduced from a molecular phylogeny of the ITS region. Plant Systematics and Evolution 219(3–4): 181–197. <https://doi.org/10.1007/BF00985578>
- Brako L, Zarucchi JL (1993) Catalogue of the flowering plants and gymnosperms of Peru. Missouri Botanical Garden, Missouri.
- Brockington S, Dos Santos P, Glover B, Ronse de Craene L (2009) Androecial evolution in Caryophyllales in light of a paraphyletic Molluginaceae. American Journal of Botany 100: 1757–1778. <https://doi.org/10.3732/ajb.1300083>
- Brown NE (1909) List of plants collected in Ngamiland and the northern part of the Kalahari desert, chiefly in the neighbourhood of Kwebe and along the Botlele and Lake Rivers. Bulletin of Miscellaneous Information Kew 1909: 89–148.
- Brown GK, Varadarajan GS (1985) Studies in Caryophyllales I: Re-evaluation of classification of Phytolaccaceae s.l. Systematic Botany 10(1): 49–63. <https://doi.org/10.2307/2418434>
- Calió MF, Pirani JR (2006) Flora de Grão Mogol, Minas Gerais: Phytolaccaceae. Boletim de Botânica, Departamento de Botânica. Instituto de Biociências. Universidade de São Paulo 24: 15–16.
- Chodat RH, Hassler E (1903) Plantæ Hasslerianæ – Soit énumération des plantes récoltées au Paraguay par le Dr. Émile Hassler, d'Aarau (Suisse) de 1885 à 1902. Bulletin de l'Herbier Boissier, ser. 2 3: 387–421.
- Clarke K (1993) Non-parametric multivariate analyses of changes in community structure. Australian Journal of Ecology 18(1): 117–143. <https://doi.org/10.1111/j.1442-9993.1993.tb00438.x>

- Clarke K, Gorley R (2006) PRIMER version 6: user manual/tutorial PRIMER-E. Plymouth.
- Clarke K, Warwick R (2001) Change in marine communities: An approach to statistical analysis and interpretation, 2<sup>nd</sup> ed. PRIMER-E, Plymouth.
- Cuénoud P, Savolainen V, Chatrou LW, Powell M, Grayer RJ, Chase MW (2002) Molecular phylogenetics of Caryophyllales based on nuclear 18S rDNA and plastid rbcL, atpB, and matK DNA sequences. *American Journal of Botany* 89(1): 132–144. <https://doi.org/10.3732/ajb.89.1.132>
- Friedrich HC (1956) Studien über die natürliche Verwandtschaft der Plumbaginales und Centrospermae. *Phyton* 6: 220–263.
- Gardner G (1846) Travels in the interior of Brazil, principally through the northern provinces, and the gold and diamond districts, during the years 1836–1841. Reeve Brothers, London.
- Gower JC (1971) A general coefficient of similarity and some its properties. *Biometrics* 27(4): 857–871. <https://doi.org/10.2307/2528823>
- Greuter W (2002) Phytolaccaceae. In: Greuter W, Rodríguez RR, Manitz H (Eds) *Flora de Cuba*, Fasc. 6. Gantner Verlag, Ruggell, 3–35.
- Heimerl A (1912) Die Nyctaginaceen und Phytolaccaceen des Herbarium Hassler. *Verhandlungen der Kaiserlich-Königlichen Zoologisch-botanischen Gesellschaft in Wien* 62: 1–17.
- Hernández-Ledesma P, Berendsohn WG, Borsch T, von Mering S, Akhani H, Arias S, Castañeda-Noa I, Eggli U, Eriksson R, Flores-Olvera H, Fuentes-Bazán S, Kadereit G, Klak C, Korotkova N, Nyffeler R, Ocampo G, Ochoterena H, Oxelman B, Rabeler RK, Sanchez A, Schlumpberger BO, Uotila P (2015) A taxonomic backbone for the global synthesis of species diversity in the angiosperm order Caryophyllales. *Willdenowia* 39(2): 281–383. <https://doi.org/10.3372/wi.45.45301>
- Howard RA, Howard ES (1982) The West Indian taxa in Solander's "Florula Indiae occidentalis". *Journal of the Arnold Arboretum* 63: 63–81.
- Hutchinson J (1926) The families of flowering plants, Vol. 1. (Dicotyledones). Macmillan & Co, London.
- Jørgensen PM, Nee MH, Beck SG (2015) Catálogo de las Plantas Vasculares de Bolivia. Systematic Botany Monographs 127: 1–1741.
- Kimura M (1980) A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* 16(2): 111–120. <https://doi.org/10.1007/BF01731581>
- Kumar S, Stecher G, Tamura K (2016) MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution* 33: 1870–1874. <https://doi.org/10.1093/molbev/msw054>
- Kunth KS (1817) Nova genera et species plantarum, Vol. 2. Sumtibus Librariae Graeco-Latino-Germanico, Paris.
- Lindman CAM (1916) Botaniska afdelning. In: Anonymous (Ed) *Naturhistoriska Riksmuseets historia: Dess uppkomst och utveckling*. Almqvist & Wiksell, Stockholm, 93–129.
- Link JHF (1821) *Enumeratio plantarum Horti Regii Berolinensis altera*, Vol. 1. G. Reimer, Berlin.
- Marchioretto MS, de Siqueira JC (1998) O gênero *Microtea* Sw. (Phytolaccaceae) no Brasil. *Pesquisas Botânica* 48: 5–51.

- McVaugh R (1955) The American collections of Humboldt and Bonpland, as described in the *Systema Vegetabilium* of Roemer and Schultes. *Taxon* 4(4): 78–86. <https://doi.org/10.2307/1217774>
- Melikian AP (1993) Phytolaccaceae. In: Takhtajan AL (Ed.) *Anatomia seminum comparativa*, Vol. 3. Nauka, Leningrad, 8–9.
- Moquin-Tandon A (1849) Phytolaccaceae. In: De Candolle A (Ed.) *Prodromus systematis universalis regni vegetabilis*, Vol. 13, pt. 2. V. Masson, Paris, 3–40.
- Moraes PLR (2009) The Brazilian herbarium of Maximilian. Prince of Wied. *Neodiversity* 4(2): 16–51. <https://doi.org/10.13102/neod.42.1>
- Moscoso RM (1943) Catalogus Flora Domingensis, Part 1. Printing Co., New York.
- Nowicke WH (1968) Palynotaxonomic study of the Phytolaccaceae. *Annals of the Missouri Botanical Garden* 55(3): 294–364. <https://doi.org/10.2307/2395128>
- Roemer JJ, Schultes JA (1820) Caroli a Linné ... *Systema vegetabilium: secundum classes, ordines, genera, species, cum characteribus, differentiis et synonymiis*, ed. 15, Vol. 6. Sumtibus J.G. Cottae, Stuttgart.
- Rohwer JG (1993) Phytolaccaceae. In: Kubitzki K, Rohwer JG, Bittrich V (Eds) *The families and genera of vascular plants*, Vol. 2. Springer, Berlin–Heidelberg–New York, 506–515. [https://doi.org/10.1007/978-3-662-02899-5\\_59](https://doi.org/10.1007/978-3-662-02899-5_59)
- Schäferhoff B, Müller KF, Borsch T (2009) Caryophyllales phylogenetics: Disentangling Phytolaccaceae and Molluginaceae and description of Microteaceae as a new isolated family. *Willdenowia* 39(2): 209–228. <https://doi.org/10.3372/wi.39.39201>
- Schmidt JA (1872) Phytolaccaceae. In: Martius KFP, Hooker JD, Eichler AG, Engler A, Baker JG, Rohrbach P, Schmidt JA (Eds) *Flora Brasiliensis*, Vol. 14, pt. 2. Fleischner, Leipzig, 326–344.
- Schrink F (1821) *Plantae rariores Horti academici Monacensis*, Vol. 2. Venditur in Instituto lithographico Scholae festivalis, Berlin.
- Shaw J, Lickey EB, Schilling EE, Small RL (2007) Comparison of whole chloroplast genome sequences to choose noncoding regions for phylogenetic studies in angiosperms: The tortoise and the hare III. *American Journal of Botany* 94(3): 275–288. <https://doi.org/10.3732/ajb.94.3.275>
- Sprengel K (1818) *Novi proventus hortorum academicorum Halensis et Berolinensis*. Geibauer, Halle.
- Sprengel K (1820) *Neue Entdeckungen im ganzen Umfang der Pflanzenkunde*, Vol. 1. F. Fleischer, Leipzig, 1–154.
- Stafleu FA, Cowan RS (1985) *Taxonomic literature. A selective guide to botanical publications and collections with dates, commentaries and types*, Vol. 5. Bohn, Scheltema & Holkema, Utrecht & Antwerpen; dr. W. Junk b.v., Publishers, The Hague & Boston. *Regnum Vegetabile*: 112.
- Stearn WT (1961) Botanical gardens and botanical literature in the eighteenth century. In: Stevenson A (Ed.) *Catalogue of botanical books in the collection of Rachel McMasters Miller Hunt*, Vol. 2(1). The Hunt Botanical Library, Pittsburgh, vii–ccxlv.
- Steudel ET (1841) *Nomenclator botanicus seu synonymia plantarum universalis*, Vol. 2. J.G. Cottae, Stuttgart & Tübingen.

- Sukhorukov AP, Mavrodiev EV, Struwig M, Nilova MV, Dzhalilova KK, Balandin SA, Erst A, Krinityna AA (2015) One-seeded fruits in the core Caryophyllales: Their origin and structural diversity. PLoS One 10(2): e0117974. <https://doi.org/10.1371/journal.pone.0117974>
- Sukhorukov AP, Nilova MV, Erst AS, Kushunina M, Baider C, Verlooove F, Salas-Pascual M, Belyaeva IV, Krinityna AA, Bruyns PV, Klak C (2018) Diagnostics, taxonomy, nomenclature and distribution of perennial *Sesuvium* (Aizoaceae) in Africa. PhytoKeys 92: 45–88. <https://doi.org/10.3897/phytokeys.92.22205>
- Swartz OP (1788) Nova genera and species plantarum seu prodromus descriptionum vegetabilium, maximam partem incognitorum quae sub itinere in Indiam Occidentalem annis 1783–[17]87. Bibliopoliis Acad. M. Swederi, Holmiae.
- Swartz OP (1797) Flora Indiae occidentalis, Vol. 1. J. Palmius, Erlangen & B. White et fil., London.
- Takhtajan A (2009) Flowering plants. Springer Science & Business Media, New Delhi. <https://doi.org/10.1007/978-1-4020-9609-9>
- Tiefenbacher L (1983) Die Brasilienexpedition von J. B. v. Spix und C. F. Ph. v. Martius in den Jahren 1817 bis 1820. Spixiana 9(Supplement): 35–42.
- Turland NJ, Wiersema JH, Barrie FR, Greuter W, Hawksworth DL, Herendeen PS, Knapp S, Kusber W-H, Li D-Z, Marhold K, May TW, McNeill J, Monro AM, Prado J, Price MJ, Smith GF (Eds) (2018) International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code) adopted by the Nineteenth International Botanical Congress Shenzhen, China, July 2017. Glashütten, Koeltz Botanical Books (Regnum Vegetabile 159).
- Urban I (1881) Geschichte des Königlichen botanischen Gartens und des Königlichen Herbariums zu Berlin, nebst einer Darstellung des augenblicklichen Zustandes dieser Institute. Jahrbuch des Königlichen botanischen Gartens und des botanischen Museums zu Berlin 1: 1–164.
- Urban I (1885) Über den Blüthenbau der Phytolaccaceen-Gattung *Microtea*. Berichte der Deutschen Botanischen Gesellschaft 3: 324–332.
- Vahl M (1792) [Comments on *Schollera*]. Skrifter af Naturhistorie-Selskabet i Kjøbenhavn 2: 210.
- von Ledebour CF (1821) Indicis seminum Horti Academicii Dorpatensis (Appendix I). J.C.Schünmann, Dorpat.
- Walter H (1906) Die Diagramme der Phytolaccaceen. Botanische Jahrbücher für Systematik. Pflanzengeschichte und Pflanzengeographie (Beiblatt) 37(4): 1–57.
- Walter H (1909) Phytolaccaceae. In: Engler A (Ed.) Das Pflanzenreich, Heft 39. Wilhelm Engelmann, Leipzig.
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds) PCR Protocols: a guide to methods and applications. Academic Press, New York, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Zhu GL, Sanderson SC (2017) Genera and a new evolutionary system of World Chenopodiaceae. Science Press, Beijing.