

The Cuito catchment of the Okavango system: a vascular plant checklist for the Angolan headwaters

David J. Goyder^{1,2}, Nigel Barker³, Stoffel P. Bester^{4,5}, Arnold Frisby³,
Matt Janks⁶, Francisco M.P. Gonçalves^{7,8}

1 Royal Botanic Gardens, Kew, TW9 3AE, UK **2** National Geographic Okavango Wilderness Project, Wild Bird Trust, South Africa **3** Department of Plant and Soil Sciences, University of Pretoria, South Africa **4** South African National Biodiversity Institute, Pretoria, South Africa **5** Unit for Environmental Sciences and Management, North-West University, Potchefstroom, South Africa **6** GroundTruth, 9 Quarry Road, Hilton 3245, KwaZulu-Natal, South Africa **7** Herbarium of Lubango, ISCED-Huila, Sarmento Rodrigues s/n, Lubango, Angola **8** University of Hamburg, Institute for Plant Science and Microbiology, Hamburg, Germany

Corresponding author: David J. Goyder (d.goyder@kew.org)

Academic editor: T. Couvreur | Received 9 October 2018 | Accepted 1 November 2018 | Published 27 November 2018

Citation: Goyder DJ, Barker N, Bester SP, Frisby A, Janks M, Gonçalves FMP (2018) The Cuito catchment of the Okavango system: a vascular plant checklist for the Angolan headwaters. *PhytoKeys* 113: 1–31. <https://doi.org/10.3897/phytokeys.113.30439>

Abstract

This paper aims to provide a baseline for conservation planning by documenting patterns of plant diversity and vegetation in the upper catchment of the Cuito River. 417 species are recorded from this region. Nine of these are species potentially new to science. Ten species are newly recorded from Angola, with an additional species only recorded previously within Angola from the northern enclave of Cabinda. The 108 new provincial records for Moxico clearly indicate the lack of collections from Angola's largest province. We note the existence of extensive peat deposits in the Cuito river system for the first time and suggest that one of Barbosa's vegetation types in the area needs to be reassessed.

Keywords

Angola, Botswana, Cuando Cubango, Moxico, peat deposits, Namibia

Introduction

Internationally famous for its wildlife, the Okavango Delta wetland in northern Botswana was the 1000th World Heritage Site to be designated by UNESCO and is surrounded by desert. The hydrology and ecology of the Delta are dependent entirely on rainfall in the highlands of central Angola, and the flow of water south and east through the Okavango's two principal tributaries, the Cuito and Cubango rivers. The Cubango system has been studied extensively in recent years (Oldeman et al. 2013), but little attention has been paid to biodiversity or conservation of the Cuito drainage.

Central and eastern Angola is overlain by deep Kalahari sands formed from uplifted and reworked deposits of an ancient palaeo-lake. The upper catchment of the Cuito and Cuanavale rivers falls mostly within Moxico Province where the plateau lies at an altitude of around 1500m, and the rivers have cut down to an elevation of around 1350 m. The landscape receives rainfall of approximately 1250 mm a year in the headwater lakes region, dropping to around 750 mm at the southern limits of the core study area which is marked by the Menongue – Longa – Cuito Cuanavale road in Cuando Cubango Province. The rainy season lasts from November to April and soils are highly leached. In consequence, they support very little agriculture (Diniz 1973).

Barbosa (1970) assigned the vegetation of the region stretching from just east of Camacupa [General Machado] to Luena [Luso] and south to Longa to vegetation type 17A. This he described as dense, high, mixed (Zambesian and Congolian) miombo woodland with “chanas” or geoxylic-rich grasslands. According to Barbosa, these woodlands comprise *Brachystegia* species (*B. spiciformis* Benth. and *B. longifolia* Benth.) and *Julbernardia paniculata* (Benth.) Troupin, with some *Guibourtia* species, *Cryptosepalum* species and *Marquesia* species. Around Longa, the vegetation transitions into Barbosa's vegetation type 24, which he describes as a mosaic of savanna, woodland and dry forest with characteristic woody vegetation containing *Brachystegia bakeriana* Hutch. & Burtt Davy and *Burkea africana* Hook.

White (1977) drew attention to the high rainfall highly leached Kalahari sand system and its peculiar flora in a seminal paper on the underground forests of Africa, extrapolating from his knowledge of similar habitats in north-west Zambia. But detailed surveys of the flora of this region are lacking.

Angola is the least intensively inventoried country in southern Africa for plants (Goyder and Gonçalves in press) – this can be seen graphically in the paucity of plant distribution records for the country (Fig. 1) at the start of the National Geographic Okavango Wilderness Project series of expeditions in 2015. Not only is the whole country under-recorded in terms of plants, but the eastern half of the country has very little geo-referenced specimen data (Marshall et al. 2016, Stropp et al. 2016, Sosef et al. 2017). Early collectors such as the Austrian botanist Friedrich Welwitsch collected along the coast, and along routes into the interior as far as Malange Province in the north and the Huíla Plateau in the south, but no further east (Albuquerque 2008, Goyder and Gonçalves in press). Swiss botanist John Gossweiler collected in all of Angola's provinces over the course of fifty years but spent very little time in central and




Figure 1. Plant distribution records for southern Africa. Raw data from GBIF (<https://www.gbif.org>). Note the absence of records for the Upper Cuito River.

eastern parts of the country apart from surveys of the Dundo area, Lunda Norte, in 1946 and 1948 funded by the diamond concession DIAMANG (Cavaco 1959, Figueiredo and Smith 2008). In addition, many of Gossweiler's collections are difficult to localise with outdated place names, and duplicates in herbaria accessible to the authors frequently omit locality data altogether. Slowly, as Angola has become more accessible following the end of the civil conflict in 2002, botanical surveys have resumed in areas of high endemism or conservation concern along the western escarpment (Hind and Goyder 2014, Gonçalves and Goyder 2016, Gonçalves et al. 2016), but the large eastern provinces of Moxico and Cuando Cubango remain poorly documented.

The only major expedition to study parts of the Cuito catchment botanically was the Kunene-Sambesi Expedition led by Pieter van der Kellen, and that covered only the area either side of the present-day Menongue – Longa – Cuito Cuanavale road. The expedition was in the Cuito system from 17th December 1899 to around 4th March 1900,

and again between 4–18th April 1900. The botany of the expedition was written up by Warburg (1903) and summarised by Figueiredo et al. (2009) who included notes on the botanist Hugo Baum and on the itinerary. Collections which form the basis of the many species described by predominantly Berlin-based botanists in Warburg (1903) and by subsequent authors were made from the Longa, Cuiriri and Cuito rivers. The area was revisited by Mendes whose 1959–1960 expedition covered the area between Kuvango [Artur de Paiva], Menongue [Serpa Pinto] and Cuito Cuanavale. Prior to the start of the Okavango Wilderness Project many species were known only from this area, and the surveys offered the chance to see if they occurred more widely.

Material and methods

The core study area is located to the south of Munhango (Figs 2, 3), and fieldwork was centred initially around the source lakes of the Cuito and Cuanavale rivers (Fig. 4), with excursions radiating from these points to the area south of Tempue and to nearby headwater lakes of other river catchments. In addition, more southerly tributaries such as the Longa (Fig. 5), Luassingua and Cuiriri river valleys were accessed from the Menongue – Cuito Cuanavale road. The darker green area towards the top left of Fig. 2 corresponds with the elevated and dissected plateau covered with moist miombo woodland which formed our core study area.

Botanical surveys were conducted at four different seasons to maximise recording of plant diversity – May–June 2015 (dry season), February–March 2016 (height of the rainy season), October–November 2016 (early rainy season) and April 2018 (late rains/early dry season). DG took part in all four surveys and focussed principally on the higher rainfall zones of the catchment between the headwaters and the Menongue – Cuito Cuanavale road (Barbosa vegetation type 17A and its transition to vegetation type 24). FG participated in the third of these surveys, and AF focussed on the Longa and Cuiriri river valleys (transition zone between Barbosa 17A and 24 vegetation types), which were the core of Baum's study in 1899 and 1900, and which had proved to be of particular botanical interest in earlier surveys. NB, SB and MJ surveyed the Longa area and the catchment south of the Menongue – Cuito Cuanavale road in June 2015.

Plant diversity was mostly assessed through walk-over surveys of each habitat in turn. But for grasses specifically, plots were set up in February–March 2016 following the methodology of Vorontsova et al. (2016) in order to feed into wider continental assessments of natural and anthropogenic grassland diversity. One plot was set up in undisturbed valley grassland near Tempué, a second in grassland possibly cleared from plateau woodland, but apparently long established, above the Cuito source lake, and the third plot was placed in open miombo woodland on the slope immediately adjacent to the Cuito source lake.

The major vegetation types generally form discrete, readily observable units in different parts of the landscape and were categorised informally.

Herbarium collections were made in sets of four where possible and deposited in two Angolan institutions (the National Biodiversity Institute of the Ministério do Am-




Figure 2. The Okavango Basin and its two principal tributaries the Cuito and Cubango rivers. The core study area is in the more elevated darker green zone of the upper Cuito river.

biente in Luanda and the Lubango Herbarium (LUBA) at ISCED-Huila), one in the UK (Royal Botanic Gardens, Kew (K)) and one in South Africa (the SANBI Herbarium in Pretoria (PRE)). Plants covered by CITES regulations (*Aloe*, succulent *Euphorbia*, Orchidaceae) were deposited only in Angolan institutions, and identified from photographs. Plants were dried on a frame over a gas burner, using aluminium corrugates to transmit heat and dry air through the press. Collections were identified principally by DG at Kew by reference to the unrivalled tropical African collections and literature held there. Expert opinion was sought from specialists in particular plant groups: Gill Challen – Euphorbiaceae, Phyllanthaceae; Phillip Cribb – Orchidaceae; Iain Darbyshire – Acanthaceae, Linderniaceae, Orobanchaceae; Sebsebe Demissew – *Asparagus*; Peter Goldblatt – *Gladiolus*; Nicholas Hind – Compositae; Isabel Larridon – Cyperaceae; Gwylim Lewis – Leguminosae; Mike Lock – Xyridaceae, Zingiberaceae; Inger Nordal – *Crinum*; Jorge Paiva – *Polygala*; Alan Paton – Lamiaceae; Sylvia Phillips – Eriocaul-




Figure 3. Locations visited during 2015 and 2016 surveys.

aceae; Brian Schrire – *Indigofera*; Andre Schuiteman – Orchidaceae; Maria Vorontsova – Gramineae; Kaj Vollesen – Acanthaceae; Martin Xanthos – Cyperaceae, Gramineae.

Angiosperm classification and nomenclature follows APG IV (2016) at family level, and the African Plant Database (version 3.4.0) or the World Checklist of Selected Plant Families (WCSP 2016) in most cases at lower taxonomic levels. Fern and lycopod names follow Roux (2009). On occasion, accepted names diverge from these resources where expert opinion suggests otherwise. Where new country or provincial records are reported, Figueiredo and Smith (2008), recent taxonomic revisions, and searchable online herbarium catalogues (principally Kew (K), the Natural History Museum, London (BM) and the Tropical Institute, Lisbon (LISC)) have been used as the baselines for comparison.

Local usage of plants was documented on 5th and 9th March 2016 thanks to the inhabitants of Samenunga village (12°56'00"S, 018°48'54"E) who explained which plants had medicinal properties, and which were used to make items such as fish traps



Figure 4. Cuito River source lake, Moxico Province. Extensive moist miombo woodland on the plateau with a few partially cleared areas on the slopes, peaty marsh surrounding the source lake and a narrow strip of fire-maintained grassland between the marsh and the miombo. Photograph D. Goyder.

and beehives. Several cultural artefacts were purchased and deposited in the Economic Botany collections at Kew, where some have since been put on public display. Vouchers of the relevant plants were taken for verification at Kew.

Results

Approximately 1100 plant collections were made over the course of the four expeditions, with a further 40+ site-based observations recorded.

The principal vegetation types of the core study area are outlined below.

Vegetation

Moist miombo woodlands

Vast swathes of central and eastern Angola are covered in this vegetation. The most common trees we observed were *Brachystegia bakeriana*, *B. longifolia*, *Cryptosepalum exfoliatum*



Figure 5. Upper Longa River valley at the southern end of the study area, Cuando Cubango Province. Moist miombo woodland on the plateau with a much broader valley containing more extensive peaty wetlands and fire-maintained grassland zones. The river is fast-flowing in deep sinuous channels with bare sandy bottoms. Photograph D. Goyder.

De Wild. subsp. *pseudotaxus* (Baker f.) P.A.Duvign. & Brenan, *Julbernardia paniculata*, with frequent *Pterocarpus angolensis* DC., *Erythrophleum africanum* (Welw. ex Benth.) Harms, *Baphia massaiensis* Taub. subsp. *ovovata* (Schinz) Brummitt var. *ovovata*, *Bobgunnia madagascariensis* (Desv.) J.H. Kirkbr. & Wiersema, *Guibourtia coleosperma* (Benth.) J.Léonard, *Monotes dasyanthus* Gilg., *M. glaber* Sprague, and *Englerophytum magalismontanum* (Sond.) T.D.Penn. Shrubs include *Bauhinia mendoncae* Torre & Hillc., *Bauhinia urbaniana* Schinz and *Copaifera baumiana* Harms. Rainfall is generally between 750–1250 mm a year in the upper Cuito catchment. Where the rainfall drops below this, to the south (lower Longa valley and Cuito Cuanavale southwards), other elements such as *Baikiaea plurijuga* Harms come in, and by M'Pupo Falls, all elements of miombo are replaced by dry thorn-scrub.

Isoberlinia angolensis (Benth.) Hoyle & Brenan var. *lasiocalyx* Hoyle & Brenan and *B. spiciformis* are essentially absent from the Cuito catchment, occurring instead on richer substrate to the west. We only noted a single occurrence of *B. spiciformis* in plateau woodland in the Cuito system.

Brachystegia bakeriana is most common near the outer margins of Cuito miombo woodland, and where the miombo patches are very small, as in the “fairy forests” near

the Cuanavale source, these are dominated by this species. More extensive miombo is on the slopes is usually dominated by *Julbernardia paniculata*, and some plateau miombo (presumably with different soil composition) by *Cryptosepalum exfoliatum* subsp. *pseudotaxus*, which can form dense, closed canopy stands of miombo forest rather than woodland. Forest lacks the flammable grass layer that is present in woodland and under *Cryptosepalum* we frequently observed the presence of a hummock-forming moss not generally found elsewhere. *Julbernardia paniculata* was seen as the principle nectar source for honey bees during our 2016 surveys.

Swamp forest

We spent a short time in a small patch of swamp forest at the source of the Rio Cuiva (Kwanza drainage). Swamp forest appears to be rare and highly localised in Moxico, unlike in Lunda Norte where extensive formations occur along tributaries of the Kasai River (Congo drainage). The Cuiva swamp forest contained species of Guineo-Congolian affinity such as *Zanthoxylum gilletii* (De Wild.) P.G.Waterman and *Syzygium owariense* (P.Beauv.) Benth.

Seasonally burned savannas

These high rainfall grasslands receive 750–1250 mm of rain a year in the upper Cuito catchment, and are on highly leached Kalahari sand. Eastern Angola contains probably 80% of this habitat, which also extends into parts of NW Zambia and western parts of the DR Congo. This habitat is fire-adapted, and is dominated by grasses or by geoxyllic suffrutescences, plants with large underground woody biomass and seasonal above-ground shoots. Factors governing whether grasses dominated, or geoxyllic suffrutescences dominated these areas were not clear. Maurin et al. (2014) argue that across Africa, fire is the evolutionary driver of such lifeforms, whereas Finckh et al. (2016) provide convincing evidence that in upland central and eastern Angola, frost also plays a principal role, with cold air pooling in valley bottoms in the winter dry season and “burning” new shoots. Proximity to the water table limits growth of trees also.

The 2016 surveys took us to several sites with significant expanses of natural or little disturbed grasslands. They were particularly extensive near the confluence of the Cuito and Calua rivers downstream of the Cuito source lake, and the equivalent confluence downstream of the Cuanavale source lake. The third notable site was the Tempué valley grasslands. Grassland diversity plots were placed at three sites – one on the plateau above the Cuito source lake, one in the nearby miombo, and one in the Tempué valley grassland. *Loudetia* species dominated – *L. simplex* (Nees) C.E.Hubb. in open areas and *L. lanata* (Stent & J.M.Rattray) C.E.Hubb. in the woodland. Five to seven grass species were found in each plot. Total grass diversity in the upper Cuito-Cuanavale system was 27 species, the majority (18) occurring in open grassland. Grassland diversity appears significantly higher than in the lower altitude plateau grasslands of Lunda Norte, also dominated by *Loudetia simplex* (Darbyshire et al. 2011, 2014). *Polygala*

robusta Gürke seems to be associated with diverse natural grassland and could perhaps be considered an indicator of good quality habitat. Another rare species encountered in this environment was the Angolan endemic *Blepharis flava* Vollesen, known from just eight earlier collections. Both of these species are newly recorded from Moxico. A spectacular blue-flowered *Barleria* is new to science and was collected at the Cuito-Calua confluence. Also new to science is a geoxyllic species of *Baphia* (Leguminosae), a genus of around 50 species of tree and shrub – the “underground forest” life form had not been recorded in *Baphia* before. This taxon was only seen in one area of the upper Lungué-Bungo catchment, in plains with a rich flora of geoxyllic legume species.

Burkea africana was a common tree in savanna vegetation at the Cuanavale source lake. This was encountered much less frequently in the Cuito source region.

Further south, the upper Longa valley, despite large-scale conversion to rice cultivation, has extensive areas of burned savanna, with some extremely rare species. *Orthonanthera gossweileri* C.Norman was known only from the type, but we recollected it in the Longa valley in March 2016, and at the Cuanavale source in October 2016 extending its range some 200 km to the north.

Wetland

Wetlands tend not to be very diverse botanically, nor to have local endemics. They are however, poorly sampled in Angola.

The extensive peaty wetlands of the Cuito have a much more diverse flora than the rather limited equivalent on the Cubango, which is a much faster flowing river running through a rocky valley. Clump- or tussock-forming plants such as Eriocaulaceae and Xyridaceae are common, while plants such as Droseraceae and Lentibulariaceae are able to supplement the limited nutrients available to other plants by trapping and digesting insects or aquatic invertebrates. Sedges (Cyperaceae) are present but are not as common as preliminary palynological records might suggest (unpublished preliminary results).

The headwater lakes of the Cuito system support a wider range of open water aquatics (true water lilies (Nymphaeaceae) and other aquatics such as *Nymphoides* and *Brasenia*) than is present on the Cubango. One unusual aquatic plant encountered in the fast-flowing upper Longa river was *Mayaca baumii* Gürke (Mayacaceae), a near-endemic and the only old-world representative of this otherwise entirely neotropical family.

Conversely, rocky rapid specialists such as *Hydrostachys triaxialis* Engl. & Gilg (Hydrostachyaceae) and *Inversodicraea warmingiana* (Gilg) Engl. (Podostemaceae) which are present on suitable portions of the Cubango (Cheek et al. 2017) are completely absent from the Cuito.

Robust river-margin plants include *Gardenia imperialis* K.Schum. (Rubiaceae) are present throughout both river catchments, while plants such as *Tacazzea rosmarinifolia* Oliv. (Apocynaceae) with rheophytic adaptations and requiring a rocky footing are found only on the Cubango.

Many wetland species have their known distributions extended dramatically. *Genlisea angolensis* R.D.Good, for example, was formerly known in Angola from just Cuando Cubango and from one collection in the DR Congo – collections in both 2015 and 2016 demonstrate this species occurs throughout the catchment of the Cuito and Cuanavale rivers (Goyder 2016). Wetland species of *Polygala* and Eriocaulaceae show similar distributions. The photographic record of *Crinum binghamii* Nordal & Kwembeya from just N of Cuito Cuanavale demonstrates this also, as it was formerly known only from western Zambia (Nordal and Kwembeya 2004, Zimudzi et al. 2008). While extending the known distributions, the new limits reflect the high rainfall, low nutrient Kalahari sand ecology.

The source lakes generally have deep accumulations of unconsolidated peat at their margins. We measured these to a depth of at least five metres at the Cuito source lake. The valleys also have more consolidated peat deposits. Such deposits are rare in tropical Africa. Reiley and Page (2016), in a recent volume on tropical peatland, state that the only significant peat deposits in Angola are on the lower Cuanza River 50 km from Luanda. The upper Cuito and Cuanavale lakes and wetlands seem to have been overlooked, despite reference in the same volume to peaty deposits in the Okavango Delta in Botswana. Analysis of peat cores from these deposits in ongoing at the University of the Witwatersrand in South Africa – pollen trapped in different strata has the potential to shed light on changes in vegetation in the region over thousands of years.

Plant diversity

417 species of vascular plant were recorded from the core study area of the high-rainfall upper Cuito and Cuanavale drainage system. The Checklist was compiled principally from our own collections from the high-rainfall zone, but with some additional collection made by Hugo Baum in the transition zone to the south. The majority of Baum's collections from the Cuito drainage system were, however, made in Barbosa's drier vegetation type 24 even further to the south and are not included in this checklist. Note that Baum's specimens citing Longa as the locality refer to the river, not to the village currently known as Longa, which is at the southern limit of our core study area, nor to Baixo Longa 100 km to the S, and outside the core study area. A further point of confusion is Warburg's (1903) map showing the route of the Kunene-Sambesi Expedition places "Hadjon Longa" close to the confluence of the Longa and Cuito rivers even further south in the region of the present-day village of Nankova.

We report nine species from the core study area which are potentially new to science (Table 1). Ten species are newly recorded for Angola with an additional species which had only been recorded within Angola from the northerly enclave of Cabinda. *Orthochilus* is a new generic record for the country (Table 2). But it is the new provincial records that give the clearest indication of how poorly studied the core project area has been to date – we recorded ten new records for Bié Province, ten for Cuando Cubango, and 108 for Moxico – the largest province in Angola.

Table 1. Species potentially new to science.

Family	Species	Notes
Acanthaceae	<i>Barleria</i> sp. nov.	Grassland at the Cuito-Calua confluence. Also in grasslands of upper Langué-Bungo tributary
Compositae	<i>Vernonia</i> sp. nov.	Growing in the floating peaty mat at Cuanavale source lake
Euphorbiaceae	<i>Acalypha</i> sp. not matched	Similar to <i>A. angustissima</i> but dioecious. Pyrophytic grassland at head of Rio Cuanavale valley and N of Tempué
Gramineae	<i>Loudetia</i> sp. nov.	Closest to <i>L. densispica</i> . Grassland in Longa river valley
Lamiaceae	<i>Endostemon</i> sp. nov.	Grassland at the Cuito-Calua confluence, Moxico province
Leguminosae	<i>Baphia</i> sp. nov.	Found at a single locality in upper Langué-Bungo catchment
Linderniaceae	<i>Crepidorhopalon</i> sp. nov.	Open sand in upper Langué-Bungo catchment
Orchidaceae	possibly sp. nov.	Same site as the <i>Barleria</i> sp. nov. A eulophioid orchid, but generic affinities uncertain
Orobanchaceae	<i>Buchnera</i> sp. not matched at Kew	May be undescribed, or a species from DR Congo. Awaiting comment from expert

Table 2. Species newly recorded from Angola.

Family	Species	Notes
Acanthaceae	<i>Justicia subsessilis</i> Oliv.	Westerly range extension
Amaryllidaceae	<i>Crinum binghamii</i> Nordal & Kwembaya	Cuanavale River N of Cuito-Cuanavale. Known also from similar habitats in western Zambia
Apocynaceae	<i>Landolphia cuneifolia</i> Pichon	Known from NW Zambia and DR Congo
Apocynaceae	<i>Secamone dewevrei</i> De Wild. subsp. <i>elliptica</i> Goyder	Only known previously from western Zambia.
Cyperaceae	<i>Cyperus unioloides</i> R.Br.	Widely distributed across tropical and subtropical Africa
Gramineae	<i>Schizachyrium claudopus</i> (Chiov.) Chiov.	Known from Tanzania, DR Congo and Zambia
Loranthaceae	<i>Englerina gabonensis</i> (Engl.) Balle	Congolian element, near Cuanavale source. New record for Angola excluding Cabinda
Orchidaceae	<i>Brachycorythis congoensis</i> Kraenzl.	Marsh in the Longa and Cuiriri valleys
Orchidaceae	<i>Bulbophyllum josephi</i> (Kuntze) Summerh.	Moist miombo woodland in Moxico
Orchidaceae	<i>Orthochilus aurantiacus</i> (Rolfe) Bytebier	New generic record for Angola
Rubiaceae	<i>Gardenia resiniflua</i> Hiern subsp. <i>resiniflua</i>	Suffrutescent form – Longa valley

Botanically, the pyrophytic grassland zone between the marsh and the miombo woodland contains most of the new and interesting species. Over 40 underground forest species (whose nearest relatives are forest trees or shrubs) were recorded from this zone and as part of the ground flora of neighbouring miombo woodlands. They include *Napoleonaea gossweileri* Baker f. (Lecythidaceae), *Trichilia quadrivalvis* C.DC (Meliaceae), and an undescribed species of *Baphia* (Leguminosae). The *Baphia* was flowering profusely at ground level in the upper Langué-Bungo catchment, where it occurred in an assemblage of other underground forest species. *Baphia* is a genus of 50 species of trees and shrubs in Africa and Madagascar – this is the first record of a pyrophytic underground forest species in the genus, and it appears to be a species new to science. The diversity of rubber-producing Apocynaceae species in the grassland zone was not-

ed – *Landolphia lanceolata* (K.Schum.) Pichon, *L. thollonii* Dewèvre, *Chamaelitandra henriquesiana* (Hallier f.) Pichon, and *Raphionacme michelii* De Wild. were common elements and have been used as sources of natural rubber in earlier times. Other much rarer species of Apocynaceae were also recorded from this zone, including *Orthanthera gossweileri* C.Norman, which we found at the source of the Cuanavale river, 200 km north of its earlier known distribution. The new species of *Baphia* will be described separately, along with a more detailed discussion of the geoxyllic suffrutex flora of the region.

Thirty-nine legume species were recorded from the upper catchment of the Cuito Cuanavale system and were found in both open and woodland habitats. Most of the miombo trees belong to this family, but there were many herbs also. Other significant elements of the flora include Rubiaceae (26 spp.), Apocynaceae (19 spp.), Lamiaceae (20 spp.) and the genus *Polygala* (Polygalaceae) with 14 species recorded – each habitat had its own suite of *Polygala* species. Monocot diversity was also substantial, with 31 grass species recorded, 17 orchids – mostly in the marsh and grassland zones – and seven species of *Gladiolus* (Iridaceae).

A flame lily species, *Gloriosa sessiliflora* Nordal & M.G.Bingham, was recorded from Angola for the second and third times ever, by the headwater team and the Longa/Cuiriri team respectively. It was described from similar marshland habitats in western Zambia in 1998.

Discussion

Miombo woodland is generally regarded as Zambesian floristically. Nevertheless, we encountered a small but significant element of moist-miombo species with Guineo-Congolian affinities. These include several species of Apocynaceae, *Uvaria angolensis* Welw. ex Oliv. in the Annonaceae, *Paropsia brazzaeana* Baill. in the Passifloraceae and *Englerina gabonensis* (Engl.) Balle in the Loranthaceae. The small patch of swamp forest at the head of the Río Cuiva is also Guineo-Congolian in affinity with *Syzygium owariense* (Myrtaceae) and *Zanthoxylum gilletii* (Rutaceae) widely distributed in the Congo Basin and West Africa. Phylogenetically, *Crinum binghamii* (Amaryllidaceae), a wetland species, is closer to Congolian members of the genus than to Zambesian species (Nordal and Kwembeya 2004).

Cape elements in the flora were restricted to savanna or grassland habitats, sometimes where rocky substrate was encountered locally. *Protea*, *Cliffortia* and *Erica* are three genera with predominantly Cape affinities and species radiations.

Floristic links outside of Africa are demonstrated by a couple of wetland taxa. *Mayaca baumii* (Mayacaceae) is the only African species of an otherwise entirely neotropical genus and family. *Mesanthemum glabrum* Kimpouni (Eriocaulaceae) is allied phylogenetically to a species from Ecuador (Larridon pers. comm., unpublished work in progress). These distributions probably reflect historic transatlantic dispersal events involving birds.

Human population in the region is low, and the few villages we passed are far apart. The low-nutrient landscape does not support much agriculture. Nevertheless, one or two

villages grew a diverse range of crops, and neighbouring miombo woodland was cleared for shifting maize and cassava cultivation. Habitat conversion is local but increasing in what is otherwise a remarkably intact ecosystem. Major human impact on the vegetation of the Cuito-Cuanavale system was only really apparent around the town of Cuito Cuanavale, and the section of the Longa valley affected by the large-scale rice project, although many of the grasslands are burned more frequently than they would be without human presence. Also, timber in the upper Lungué-Bungo valley is increasingly targeted as this area is closer to the provincial capital Luena than the rest of the core project area.

Many native plant species are used as medicines or for construction. It is mostly the most common species that are used.

The most frequent miombo tree, *Julbernardia paniculata*, is not only the principle source of nectar for honey bees but is also the preferred tree for the construction of beehives (Fig. 6). A cylinder of bark is removed from the tree (killing the individual), stapled together with stakes made from another legume timber (*Bobgunnia madagascariensis*), and tied together with stringy underbark from a third (*Julbernardia paniculata*, *Brachystegia bakeriana* or *Cryptosepalum exfoliatum* subsp. *pseudotaxus*). Internal bracing hoops come from flexible young shoots of either *Diplorhynchus condylocarpon* (Müll.Arg.) Pichon or *Baphia massaiensis* subsp. *obovata*, the permanent wooden cap at one end is made from *Parinari curatellifolia* Planch. ex Benth., while the removable cap at the other end is of woodland grasses, mostly *Loudetia* spp. Such traditional methods of construction are destructive but sustainable when population levels are low. However, harvesting of honey and production of beehives is becoming an industry, with some villages boasting of 300+ hives in active use.

Large fish traps were constructed from saplings of *Englerophytum magalismontanum*, tied together with fine bark string as above. The small fishtrap was constructed from the grass *Loudetia densispica* (Rendle) C.E.Hubb.

Locally made bark canoes were present in most lakes and major watercourses we visited (Fig. 7). These were generally made from bark of the legume tree *Erythrophleum africanum* and stitched together as above with strips of stringy underbark from *Brachystegia* or *Cryptosepalum* spp.

It was noted that local people have a detailed understanding of plants with active biological properties in their immediate environment and know how to use these to treat a variety of conditions in the absence of accessible western medicine.

Conclusions

Over 1100 plant collections were made during the course of the National Geographic Okavango Wilderness Project, the majority from the core project area of the upper Cuito and Cuanavale river catchments. These form the basis of what is undoubtedly the most detailed specimen-based assessment of the vegetation and plant diversity of this region.

The flora of the upper Cuito and Cuanavale system is diverse and endemism is high, although the latter has not been analysed in detail for this study. New records extend the known geographic range of many species 200 km to the north, to the



Figure 6. Beehive made in Samenunga village, Moxico Province, now displayed in the Economic Botany collection of the Royal Botanic Gardens, Kew, UK. The body of the beehive is a cylinder of bark from the locally dominant legume tree *Julbernardia paniculata*. Other species are used to provide stakes, bracing hoops and endcaps. Photograph D. Goyder.

headwaters of the Cuito and Cuanavale rivers. They also underline the need for further surveys in Moxico Province where 108 new provincial records were reported, and provide evidence that the absence of plant records for eastern Angola revealed on the GBIF data map of southern Africa is real, and not a data artefact. All four *Protea* species collected in Moxico had never been recorded there before.

Barbosa (1970)'s vegetation type 17A needs to be critically reconsidered in the light of our findings in this area – we observed a fundamental change in composition of the miombo woodlands east of Cuemba once we moved onto the deep white sands, where several woody species drop out – no *Isoberlinia angolensis* (Benth.) Hoyle & Brenan was seen east of this point, and *Brachystegia spiciformis* occurred exceptionally rarely. Both *Burkea africana* and *Brachystegia bakeriana* are significant elements of the landscape in the headwater lakes region, not just in the transition zone around Longa. We saw no *Marquesia* species in the headwaters zone, but *Monotes* is common. *Baphia massaiensis* subsp. *obovata*, more commonly associated with dry *Baikiaea*-dominated woodland, was a common element of the miombo right up into the headwater region.

We also highlight the existence of extensive peat deposits in the Cuito river system. These are not as extensive as those recently reported from the Congo Basin (Dargie et al. 2017), but must be significant in terms of carbon storage nevertheless.



Figure 7. Bark canoe (foreground) made from *Erythrophleum africanum* (Leguminosae) alongside a fibre-glass “dugout” brought in by the National Geographic Okavango Wilderness Project from the Okavango Delta. Cuito source lake. Photograph D. Goyder.

Checklist

An annotated checklist of the upper Cuito & Cuanavale drainage system – the flora of high rainfall (annual precipitation more than c. 750 mm), highly leached Kalahari sand deposits from the headwaters to c. 15°S, based principally on 2015, 2016 and 2018 field surveys (Barbosa vegetation type 17A and transition to vegetation type 24).

Table Checklist

Family	Species	Habitat	Vouchers	New Records
LYCOPODIOPHYTA				
Lycopodiaceae	<i>Lycopodiella affinis</i> (Bory) Pic.Serm.	Wetland	Frisby 3027; Goyder 8261	
	<i>Lycopodiella cernua</i> (L.) Pic.Serm.	Wetland	sight record 38	
	<i>Lycopodiella sarcocaulon</i> (A.Braun & Welw. ex Kuhn) Pic.Serm.	Wetland	Goyder 8298	
PTERIDOPHYTA				
Aspleniaceae	<i>Asplenium aethiopicum</i> (Burm.f.) Bech.	Humid Forest	Goyder 8329	
Gleicheniaceae	<i>Dicranopteris linearis</i> (Burm.f.) Underw.	Wetland	Goyder 8396	
Thelypteridaceae	<i>Cyclosorus interruptus</i> (Willd.) H.Itô	Wetland	Goyder 8317	Moxico
	<i>Thelypteris confluens</i> (Thunb.) Morton	Wetland	Barker et al. 139	

Family	Species	Habitat	Vouchers	New Records
ANGIOSPERMAE: MAGNOLIIDS				
Annonaceae	<i>Annona stenophylla</i> Engl. & Diels subsp. <i>nana</i> (Exell) N.Robson	Grassland	Goyder & Maiato 8759; Goyder & Maiato 8843	
	<i>Artobotrys antunesii</i> Engl. & Diels	Woodland	Goyder 8436	Moxico
	<i>Uvaria angolensis</i> Welw. ex Oliv.	Woodland	Goyder 8034; Goyder 8414; Goyder 8438	
	<i>Xylopia odoratissima</i> Welw. ex Oliv.	Woodland	Frisby 3067; Goyder & Maiato 8806	
	<i>Xylopia tomentosa</i> Exell	Woodland	Barker et al. 50; Frisby 3057; Goyder 8027; Goyder 8048; Goyder 8096; Goyder 8288; Goyder 8918	Bié
Cabombaceae	<i>Brasenia schreberi</i> J.F.Gmel.	Wetland	Goyder 8295	Moxico
Lauraceae	<i>Casytha pondoensis</i> Engl. var. <i>pondoensis</i>	Woodland	Goyder 8104	
Nymphaeaceae	<i>Nymphaea heudelotii</i> Planch.	Wetland	Barker et al. 44; Goyder 8259	
	<i>Nymphaea nouchali</i> Burm.f. var. <i>caerulea</i> (Savigny) Verdc.	Wetland	Frisby 4013; Goyder 8296; Goyder 8376	
	<i>Nymphaea sulphurea</i> Gilg.	Wetland	Baum 657; Frisby 3050; Frisby 3064; Frisby 3072; Goyder 8097; Goyder 8297; Goyder 8393	
ANGIOSPERMAE: MONOCOTS				
Alismataceae	<i>Limnophyton angolense</i> Buchenau	Wetland	Frisby 3093; Goyder 8375; sight record 15	Moxico
Amaryllidaceae	<i>Boophone disticha</i> (L.f.) Herb.	Grassland	Goyder & Maiato 8829	
	<i>Crinum binghamii</i> Nordal & Kwembeya	Wetland	sight record 42	
	<i>Cryptostephanus densiflorus</i> Welw. ex Baker	Woodland	Goyder 8258	Moxico
	<i>Cyrtanthus welwitschii</i> Hiern ex Baker	Wetland	Frisby 4023	Cuando Cubango
Asparagaceae	<i>Asparagus africanus</i> Lam. var. <i>puberulus</i> (Baker) Sebsebe	Grassland	Goyder 8439	
	<i>Chlorophytum colubrinum</i> (Baker) Engl.	Grassland	Baum 611	
	<i>Chlorophytum fasciculatum</i> (Baker) Kativu	Grassland	Baum 683; Goyder 9495	
	<i>Chlorophytum sphacelatum</i> (Baker) Kativu	Grassland	Goyder 9495a	
	<i>Chlorophytum</i> sp.	Grassland	Goyder 8263	
	<i>Dipcadi viride</i> (L.) Moench	Wetland	Goyder & Maiato 8801	
	<i>Sansevieria aubrytiana</i> Carrière	Woodland	Goyder & Maiato 8838	Moxico
	<i>Schizocarphus nervosus</i> (Burch.) Van der Merwe	Grassland	Goyder & Maiato 8779	Moxico
Asphodelaceae	<i>Aloe nuttii</i> Baker	Grassland	Baum 698	
	<i>Aloe zebrina</i> Baker	Woodland	Goyder 8255	
	<i>Trachyandra arvensis</i> (Schinz) Oberm.	Grassland	Frisby 3062; Goyder 8494; Goyder & Maiato 8816; Goyder & Maiato 8820	
Colchicaceae	<i>Gloriosa sessiliflora</i> Nordal & M.G.Bingham	Wetland	Frisby 4035; Goyder & Maiato 8822	Cuando Cubango; Moxico
	<i>Gloriosa simplex</i> L.	Woodland	Goyder 8425	Moxico
Commelinaceae	<i>Aneilema plagiocapsa</i> K. Schum.	Woodland	Barker et al. 82; Baum 716; Goyder 8244	Moxico
	<i>Commelina africana</i> L. var. <i>lancispatha</i> C.B.Clarke	Woodland	Goyder 8245	
	<i>Commelina sphaerorrhiza</i> Faden & Layton	Woodland	Goyder 8243	Moxico

Family	Species	Habitat	Vouchers	New Records
Commelinaceae	<i>Commelina welwitschii</i> C.B.Clarke	Grassland	Baum 814	
	<i>Cyanotis longifolia</i> Benth.	Grassland	Goyder & Maiato 8832	Moxico
Costaceae	<i>Costus spectabilis</i> (Fenzl) K.Schum.	Grassland	Goyder 8947	
Cyperaceae	<i>Abildgaardia ovata</i> (Burm.f.) Kral	Wetland	Frisby 3041	
	<i>Bulbostylis laniceps</i> C.B.Clarke ex T.Durand & Schinz	Grassland	Goyder 8290	Moxico
	<i>Cyperus chrysoccephalus</i> (K.Schum.) Kük.	Wetland	Frisby 3071	
	<i>Cyperus denudatus</i> L.f.	Wetland	Goyder 8931	
	<i>Cyperus erinaceus</i> (Ridl.) Kük.	Grassland	Goyder 8334	
	<i>Cyperus hensii</i> T.Durand & Schinz	Wetland	Frisby 3081	
	<i>Cyperus kipasensis</i> Cherm.	Wetland	Goyder 8939	
	<i>Cyperus margaritaceus</i> Vahl	Grassland	Goyder 8335; Goyder & Maiato 8831; Goyder 8925	
	<i>Cyperus pectinatus</i> Vahl	Wetland	Goyder 8294	
	<i>Cyperus proteus</i> (Welw.) Bauters	Wetland	Barker et al. 63; Baum 627; Baum 628; Frisby 3009; Goyder 8005; Goyder 8365	
	<i>Cyperus proteus</i> (Welw.) Bauters var. <i>bellidiflora</i> Welw.	Wetland	Goyder 8936	
	<i>Cyperus rhynchosporoides</i> Kuk.	Grassland	Goyder & Maiato 8830	
	<i>Cyperus subtrigonos</i> (C.B.Clarke) Kük.	Wetland	Goyder 8940	
	<i>Cyperus unioloides</i> R.Br.	Wetland	Goyder 8941	Angola
Eriocaulaceae	<i>Cyperus</i> sp. not matched	Grassland	Barker et al. 71; Barker et al. 111	
	<i>Eleocharis acutangula</i> (Roxb.) Schult. subsp. <i>acutangula</i>	Wetland	Goyder 8945	
	<i>Fimbristylis dichotoma</i> (L.) Vahl var. <i>dichotoma</i>	Wetland	Goyder 8942	
	<i>Fuirena umbellata</i> Rottb.	Grassland	Barker et al. 136; Goyder 8924	
	<i>Lipocarpha chinensis</i> (Osbeck) J.Kern.	Wetland	Goyder 8938	
	<i>Rhynchospora candida</i> (Nees) Boeck.	Wetland	Barker et al. 62; Goyder 8302; Goyder 8368	
	<i>Rhynchospora rugosa</i> (Vahl) Gale subsp. <i>brownii</i> (Roem. & Schult.) T.Koyama	Grassland	Barker et al. 65	
	<i>Scleria erythrorrhiza</i> Ridl.	Wetland	Barker et al. 57; Goyder 8933	
	<i>Scleria griegiiifolia</i> (Ridl.) C.B.Clarke	Wetland	Goyder 8239; Goyder 8360; sight record 41	
	<i>Eriocaulon lanatum</i> H.E.Hess	Wetland	Goyder 8202; Goyder 8369	Moxico
Gramineae	<i>Eriocaulon teucsii</i> Engl. & Ruhland	Wetland	Goyder 8099; Goyder 8364	Moxico
	<i>Mesanthemum glabrum</i> Kimpouni	Wetland	Baum 645; Frisby 3065; Goyder 8004; Goyder 8201; Goyder 8238; Goyder 8358	Moxico
	<i>Mesanthemum reductum</i> H.E.Hess	Wetland	Barker et al. 115	
	<i>Syngonanthus angolensis</i> H.E.Hess	Wetland	Goyder 8237; Goyder 8359	Moxico
	<i>Syngonanthus wahlbergii</i> (Wikstr. ex Körn.) Ruhland	Wetland	Goyder 8100	
	<i>Aristida nemorivaga</i> Henrard	Woodland	Barker et al. 108	
	<i>Brachiaria dura</i> Stapf	Grassland	Barker et al. 59; Goyder 8289	Cuando Cubango
	<i>Ctenium newtonii</i> Hack.	Grassland	Barker et al. 54	Cuando Cubango
	<i>Digitaria milanjiana</i> (Rendle) Stapf	Woodland	Goyder 8306	

Family	Species	Habitat	Vouchers	New Records
Gramineae	<i>Diheteropogon amplexens</i> (Nees) Clayton var. <i>amplectens</i>	Grassland; Woodland	Goyder 8274; Goyder 8285	Moxico
	<i>Diheteropogon filifolius</i> (Nees) Clayton	Grassland	Barker et al. 60; Goyder 8407	Cuando Cubango
	<i>Eleusine coracana</i> (L.) Gaertn.	Wetland	Baum 693	
	<i>Elymandra grallata</i> (Stapf) Clayton	Grassland	Barker et al. 98; Barker et al. 105	Cuando Cubango
	<i>Eragrostis brainii</i> (Stent) Launert	Woodland	Goyder 8268; Goyder 8282	
	<i>Eragrostis thollonii</i> Franch.	Woodland	Goyder 8284	Moxico
	<i>Heteropogon contortus</i> (L.) P.Beauv.	Grassland	Goyder 8272; Goyder 8404	
	<i>Hyparrhenia newtonii</i> (Hack.) Stapf	Grassland	Goyder 8042; Goyder 8923	Bié
	<i>Leersia hexandra</i> Sw.	Wetland	Goyder 8930	
	<i>Loudetia angolensis</i> C.E.Hubb.	Wetland	Goyder 8264	Moxico
	<i>Loudetia densispica</i> (Rendle) C.E.Hubb.	Grassland	Barker et al. 109; Goyder 8273; Goyder 8442	Moxico
	<i>Loudetia lanata</i> (Stent & J.M.Rattray) C.E.Hubb.	Woodland	Goyder 8281	Moxico
	<i>Loudetia simplex</i> (Nees) C.E.Hubb.	Grassland	Goyder 8228; Goyder 8269; Goyder 8403	Bié; Moxico
	<i>Loudetia</i> sp. nov. aff. <i>L. densispica</i>	Grassland	Barker et al. 55	
	<i>Misanthus junceus</i> (Stapf) Pilg.	Wetland	Goyder 8299	Moxico
	<i>Monocymbium ceresiiforme</i> (Nees) Stapf	Grassland	Goyder 8275; Goyder 8405	Moxico
	<i>Panicum natalense</i> Hochst.	Grassland	Goyder 8271; Goyder 8409	Moxico
	<i>Pennisetum polystachion</i> (L.) Schult.	Ruderal	Goyder 8043	Bié
	<i>Phragmites mauritianus</i> Kunth	Wetland	Goyder 8935	
Hydrocharitaceae	<i>Pogonarthria squarrosa</i> (Roem. & Schult.) Pilg.	Grassland	Barker et al. 97	
	<i>Rhytachne robusta</i> Stapf	Woodland	Goyder 8283	
	<i>Schizachyrium claudopus</i> (Chiov.) Chiov.	Grassland	Barker et al. 58	
	<i>Sporobolus welwitschii</i> Rendle	Grassland	Goyder 8291	
	<i>Trachypogon spicatus</i> (L.f.) Kuntze	Grassland	Goyder 8913	
	<i>Tristachya hubbardiana</i> Conert	Grassland	Goyder 8408	Moxico
	<i>Tristachya nodiglumis</i> K.Schum.	Grassland	Barker et al. 72	
	<i>Tristachya rehmannii</i> Hack.	Grassland	Goyder 8270; Goyder 8406	
	<i>Blyxa radicans</i> Ridl.	Wetland	Baum 827	
	<i>Ottelia muricata</i> (C.H.Wright) Dandy	Wetland	Barker et al. 118	
Hypoxidaceae	<i>Ottelia ulvifolia</i> (Planch.) Walp.	Wetland	Goyder 8929	
	<i>Hypoxis canaliculata</i> Baker	Grassland	Goyder & Maiato 8790	
Iridaceae	<i>Ferraria welwitschii</i> Baker	Grassland	Frisby 4012; Goyder 8496; Goyder & Maiato 8768	Moxico
	<i>Gladiolus atropurpureus</i> Baker	Grassland	Goyder 8498	Cuando Cubango
	<i>Gladiolus benguillensis</i> Baker	Grassland	Baum 632	
	<i>Gladiolus dalenii</i> Van Geel subsp. <i>dalenii</i>	Wetland	Frisby 3029; Goyder 8461	Moxico
	<i>Gladiolus gregarius</i> Welw. ex Baker	Woodland	Goyder 8401	
	<i>Gladiolus gregarius</i> Welw. ex Baker – anomalous form with filiform leaves and green flowers	Grassland	Goyder 8499	
	<i>Gladiolus laxiflorus</i> Baker	Wetland	Frisby 3010; Frisby 3066; Goyder & Maiato 8793	
	<i>Gladiolus magnificus</i> (Harms) Goldblatt	Grassland	Baum 651; Goyder 8497	

Family	Species	Habitat	Vouchers	New Records
Iridaceae	<i>Gladiolus unguiculatus</i> Baker	Grassland	Frisby 3025; Frisby 3038; Goyder & Maiato 8777; Goyder & Maiato 8778	
Mayacaceae	<i>Mayaca baumii</i> Gürke	Wetland	Barker et al. 117; Baum 811	
Orchidaceae	<i>Brachycorythis congoensis</i> Kraenzl.	Wetland	Frisby 3068	
	<i>Bulbophyllum josephi</i> (Kuntze) Summerhayes	Woodland	Goyder 8419	
	<i>Disa caffra</i> Bolus	Wetland	Goyder & Maiato 8791	
	<i>Disa ochrostachya</i> Rchb. f.	Wetland	Frisby 4005; Goyder & Maiato 8763; Goyder & Maiato 8796	
	<i>Disa bircicornis</i> Rchb.f.	Wetland	Frisby 3075	
	<i>Disa welwitschii</i> Rchb.f.	Wetland	Frisby 3063	
	<i>Eulophia angolensis</i> (Rchb.f.) Summerh.	Wetland	Frisby 3032	
	<i>Eulophia horsfalliae</i> (Bateman) Summerh.	Wetland	Goyder & Maiato 8792	Moxico
	<i>Eulophia longisepala</i> Rendle	Grassland	Goyder & Maiato 8753	Moxico
	<i>Eulophia rolfeana</i> Kraenzl.	Grassland	Frisby 3095; Goyder & Maiato 8755	Moxico
	<i>Eulophia speciosa</i> (R. Br. ex Lindl.) Bolus	Grassland	Goyder & Maiato 8774	Moxico
	<i>Habenaria retinervis</i> Summerh.	Woodland	Goyder 8220	
	<i>Orthochilus aurantiacus</i> (Rolfe) Bytebier	Grassland	Frisby 4002; Goyder & Maiato 8752; Goyder & Maiato 8796	Cuando Cubango
	<i>Phaius occidentalis</i> Schltr.	Wetland	Goyder & Maiato 8761	Moxico
	<i>Polystachya concreta</i> (Jacq.) Garay & H.R.Sweet	Woodland	Goyder 8225	
	<i>Satyrium trinerve</i> Lindl.	Wetland	Frisby 3080; Frisby 4001	
	possibly sp. nov.	Grassland	Goyder 8351	
Smilacaceae	<i>Smilax anceps</i> Willd.	Ruderal	sight record 16	
Xyridaceae	<i>Xyris capensis</i> Thunb.	Wetland	Goyder 8373	
	<i>Xyris congesta</i> Büttner	Wetland	Barker et al. 64; Goyder 8322	
	<i>Xyris foliolata</i> L.A.Nilsson	Wetland	Barker et al. 128	
	<i>Xyris friesii</i> Malme	Wetland	Goyder & Maiato 8800	Moxico
	<i>Xyris imitatrix</i> Malme	Wetland	Goyder 8332	
Zingiberaceae	<i>Aframomum albiviolaceum</i> (Ridl.) K.Schum.	Ruderal	sight record 17	
	<i>Siphonochilus aethiopicus</i> (Schweinf.) B.L.Burtt	Grassland Woodland	Frisby 3089; Goyder & Maiato 8769	
	<i>Siphonochilus puncticulatus</i> (Gagnep.) Lock	Grassland Woodland	Frisby 3076; Goyder & Maiato 8770	
ANGIOSPERMAE: EUDICOTS				
Acanthaceae	<i>Barleria crassa</i> C.B.Clarke subsp. <i>crassa</i>	Woodland	Goyder 8028	
	<i>Barleria</i> sp. nov.	Grassland	Goyder 8343; Goyder 8952	
	<i>Blepharis flava</i> Vollesen	Grassland	Goyder 8277	Moxico
	<i>Blepharis glumacea</i> S.Moore	Grassland	Goyder 8909	
	<i>Justicia subsessilis</i> Oliv.	Grassland	Barker et al. 89	
	<i>Lepidagathis macrochila</i> Lindau	Woodland	Baum 779; Goyder 8040; Goyder 8415	Moxico
	<i>Strobilanthes linifolia</i> (T.Anderson ex C.B.Clarke) Milne-Redh.	Woodland	Barker et al. 107; Goyder 8026	Moxico
	<i>Thunbergia gossweileri</i> S.Moore	Woodland	Goyder 8241	Moxico
Amaranthaceae	<i>Mechowia grandiflora</i> Schinz	Grassland Woodland	Frisby 4010; Goyder 8112; Goyder 8385	Moxico

Family	Species	Habitat	Vouchers	New Records
Anacardiaceae	<i>Lannea gossweileri</i> Exell & Mendonça subsp. <i>gossweileri</i>	Grassland	Goyder & Maiato 8834	
	<i>Ozoroa stenophylla</i> (Engl. & Gilg) R.Fern. & A.Fern.	Grassland	Baum 662; Frisby 3012; Goyder 8310	Moxico
	<i>Ozoroa verticillata</i> (Engl.) R.Fern. & A.Fern.	Grassland	Goyder 8287	Moxico
	<i>Rhus gracilipes</i> Exell	Woodland	Goyder 8254	Moxico
Anisophylleaceae	<i>Anisophyllea boehmii</i> Engl.	Woodland	Goyder 8232	
	<i>Anisophyllea fruticulosa</i> Engl. & Gilg	Grassland	Barker et al. 46; Baum 808†; Gossweiler 2856; Goyder 8106; Goyder & Maiato 8765	
Apocynaceae	<i>Chamaelitandra henriquesiana</i> (Hallier f.) Pichon	Grassland	Barker et al. 81; Goyder & Maiato 8766; Goyder & Maiato 8807	Moxico
	<i>Ceropegia racemosa</i> N.E.Br.	Woodland	Goyder 8402	Moxico
	<i>Cryptolepis oblongifolia</i> (Meisn.) Schltr.	Woodland	Barker et al. 78; Barker et al. 112; Frisby 3037; Goyder 8118; Goyder 8124; Goyder 8300	
	<i>Diplorhynchus condylocarpon</i> (Müll.Arg.) Pichon	Grassland Woodland	Barker et al. 52A; Frisby 3058; Frisby 3061; Goyder 8213; Goyder 8381; Goyder 8445; sight record 1; sight record 8; sight record 36	
	<i>Glossostelma ceciliae</i> (N.E.Br.) Goyder	Grassland	Frisby 4033; Goyder & Maiato 8789	
	<i>Gomphocarpus semiamplectens</i> K.Schum.	Woodland	Barker et al. 121	
	<i>Landolphia camptoloba</i> (K.Schum.) Pichon	Woodland	Barker et al. 49; Barker et al. 122; Baum 669; Frisby 4004; Goyder 8025; Goyder 8400	
	<i>Landolphia cuneifolia</i> Pichon	Woodland	Goyder 8331	
	<i>Landolphia lanceolata</i> (K.Schum.) Pichon	Grassland	Barker et al. 79; Goyder 8019; Goyder 8266; Goyder & Maiato 8803	
	<i>Landolphia thollonii</i> Dewèvre	Grassland	Goyder 8431; Goyder & Maiato 8825 [photographic record]	
	<i>Orthanthera gossweileri</i> C.Norman	Grassland	Frisby 3051; Goyder 8500; Goyder & Maiato 8827	Moxico
	<i>Raphionacme globosa</i> K.Schum.	Grassland	Goyder & Maiato 8797	Moxico
	<i>Raphionacme linearis</i> K.Schum.	Wetland	Frisby 3020; Frisby 3035; Frisby 3039; Frisby 3078; Goyder & Maiato 8776; Goyder & Maiato 8856	
	<i>Raphionacme michelii</i> De Wild.	Grassland	Frisby 3026; Goyder & Maiato 8788; Goyder & Maiato 8809; Goyder & Maiato 8771	Moxico
	<i>Secamone brevipes</i> (Benth.) Pichon	Woodland	Goyder 8330	Moxico
Campanulaceae	<i>Secamone dewevrei</i> De Wild. subsp. <i>elliptica</i> Goyder	Woodland	Goyder 8041; Goyder 8223	
	<i>Strophanthus welwitschii</i> (Baill.) K.Schum.	Woodland	Goyder & Maiato 8837	
	<i>Tabernanthe iboga</i> Baill.	Woodland	Goyder 8226; sight record 18	
	<i>Xysmalobium holubii</i> Scott Elliot	Wetland	Baum 715; Frisby 3034; Goyder & Maiato 8785; Goyder & Maiato 8853	Moxico
	<i>Lobelia</i> sp.	Grassland	Barker et al. 116	
	<i>Wahlenbergia collomoides</i> (A.DC.) Thulin	Grassland	Goyder 8906	
	<i>Wahlenbergia</i> possibly sp. B of Thulin (1975)	Grassland	Barker et al. 94	

Family	Species	Habitat	Vouchers	New Records
Caryophyllaceae	<i>Polycarpea corymbosa</i> (L.) Lam.	Grassland	Barker et al. 132; Baum 818; Goyder 8457	
Celastraceae	<i>Gymnosporia senegalensis</i> (Lam.) Loes.	Wetland	Goyder 8934	
	<i>Salacia bussei</i> Loes.	Grassland	Goyder 8292; Goyder & Maiato 8810	Moxico
Chrysobalanaceae	<i>Parinari capensis</i> Harv.	Grassland	Barker et al. 130; Goyder 8256	
	<i>Parinari curatellifolia</i> Planch. ex Benth.	Woodland	Goyder 8444	
Combretaceae	<i>Combretum dumetorum</i> Exell	Woodland	Goyder 8426	Moxico
	<i>Combretum gossweileri</i> Exell	Woodland	Goyder 8023	
	<i>Combretum platypetalum</i> Welw. ex M.A.Lawson	Grassland	Frisby 3036; Goyder 8121	
	<i>Combretum psidioides</i> Welw. subsp. <i>psidioides</i>	Grassland	Frisby 3053; Goyder 8345	
	<i>Combretum</i> sp. not matched 1	Woodland	Goyder 8307	
	<i>Combretum</i> sp. not matched 2	Grassland	Goyder 8346	
	<i>Pteleopsis anisoptera</i> (Welw. ex M.A.Lawson) Engl. & Diels	Woodland	Goyder 8418	
	<i>Terminalia brachystemma</i> Welw. ex Hiern	Woodland	Frisby 3011; Goyder 8378	
Compositae	<i>Anisopappus chinensis</i> Hook. & Arn.	Grassland	Goyder 8908	
	<i>Bidens crocea</i> Welw. ex O.Hoffm.	Woodland	Goyder 8253	
	<i>Blumea axillaris</i> (Lam.) DC.	Grassland	Barker et al. 134	
	<i>Crassocephalum</i> sp. not matched	Woodland	Goyder 8305	
	<i>Dicoma schinzii</i> O.Hoffm.	Grassland	Barker et al. 85	
	<i>Emilia baumii</i> (O.Hoffm.) S.Moore	Woodland	Baum 707; Goyder 8252; Goyder 8910	Moxico
	<i>Erlangia misera</i> (Oliv. & Hiern) S.Moore	Woodland	Barker et al. 125	
	<i>Hypericophyllum gossweileri</i> S.Moore	Grassland	Goyder 8948	Angola
	<i>Mikania sagittifera</i> B.L.Robb.	Grassland	Barker et al. 104; Baum 679	
	<i>Nidorella resedifolia</i> DC.	Grassland	Barker et al. 126	
	<i>Pascuccardoa baumii</i> O.Hoffm.	Grassland	Frisby 3013; Goyder 8111	
	<i>Pleiotaxis linearifolia</i> O. Hoffm.	Grassland Woodland	Barker et al. 69; Barker et al. 120	
	<i>Pleiotaxis rugosa</i> O.Hoffm.	Woodland	Barker et al. 75	
	<i>Pleiotaxis subcaposa</i> C.Jeffrey	Grassland	Goyder 8279; Goyder 8456	Moxico
	<i>Pseudognaphalium luteoalbum</i> (L.) Hilliard & B.L.Burtt	Grassland	Barker et al. 70; Frisby 3019	
	<i>Senecio strictifolius</i> Hiern	Wetland	Barker et al. 110; Barker et al. 127; Goyder 8915	
	<i>Vernonia</i> sp. nov.	Wetland	Goyder 8357	
	<i>Vernonia</i> sp.	Grassland	Goyder 8459	
	<i>Vernonia gerberiformis</i> Oliv. & Hiern subsp. <i>gerberiformis</i> var. <i>gerberiformis</i>	Grassland	Goyder 8109	
	<i>Vernonia ornata</i> S.Moore	Wetland	Frisby 3091	
	<i>Vernonia poskeana</i> Vatke & Hildebr. subsp. <i>poskeana</i>	Woodland	Barker et al. 84	
	<i>Vernonia subplumosa</i> O.Hoffm.	Woodland	Baum 703; Goyder 8286	Moxico
	<i>Vernonia turbinella</i> S.Moore	Woodland	Goyder 8017	
Convolvulaceae	<i>Ipomoea welwitschii</i> Vatke ex Hallier f.	Grassland	Goyder & Maiato 8828	Moxico
Cucurbitaceae	<i>Acanthosicyos naudinianus</i> (Sond.) C. Jeffrey	Ruderal; Grassland	Barker et al. 119; Goyder 8086	
Dilleniaceae	<i>Tetracera poggei</i> Gilg	Woodland	Goyder 8021; Goyder 8214	Bié; Moxico
Dipterocarpaceae	<i>Monotes dasyanthus</i> Gilg	Woodland	Goyder 8039; sight record 34	

Family	Species	Habitat	Vouchers	New Records
Dipterocarpaceae	<i>Monotes glaber</i> Sprague	Woodland	Goyder 8014; Goyder 8122; sight record 20; sight record 33	
	<i>Monotes gossweileri</i> De Wild.	Grassland	Goyder 8338; Goyder 8951	
Droseraceae	<i>Drosera affinis</i> Welw. ex Oliv.	Wetland	Baum 687; Goyder 8260; Goyder 8356	Moxico
	<i>Drosera burkeana</i> Planch.	Wetland	Goyder & Maiato 8794	
	<i>Drosera madagascariensis</i> DC.	Wetland	Frisby 4011; Goyder 8003; Goyder 8006; Goyder 8372; sight record 40; Goyder & Maiato 8786	
Ebenaceae	<i>Diospyros batocana</i> Hiern	Woodland	Barker et al. 142; Goyder 8029	
	<i>Diospyros chamaethamnus</i> Dinter ex Mildbr.	Grassland	Goyder 8901	
	<i>Diospyros pseudomespilus</i> Mildbr. subsp. <i>brevicalyx</i> F.White	Woodland	Goyder 8032; sight record 32	
	<i>Diospyros virgata</i> (Gürke) Brenan	Woodland	Goyder 8015	
Ericaceae	<i>Erica benguelensis</i> (Welw. ex Engl.) E.G.H.Oliv. var. <i>benguelensis</i>	Grassland	Goyder 8352	
Euphorbiaceae	<i>Acalypha</i> sp. not matched	Grassland	Goyder & Maiato 8802; Goyder & Maiato 8814	
	<i>Maprounea africana</i> Müll.Arg. pyrophytic form	Grassland	Goyder 8312	
	<i>Scleroctron oblongifolius</i> (Müll.Arg.) Kruit & Roebers	Grassland	Goyder 8314; Goyder & Maiato 8844	
Gentianaceae	<i>Faraea salutaris</i> Welw.	Wetland; Grassland	Barker et al. 53; Frisby 4000; Goyder 8216	
	<i>Neurotheca congolana</i> De Wild. & T.Durand	Wetland	Goyder 8234; Goyder 8354	Moxico
	<i>Pycnosphaera buchananii</i> (Baker) N.E.Br.	Wetland	Goyder 8462	
	<i>Schinziella tetragona</i> (Schinz) Gilg	Wetland	Goyder 8333; Goyder 8355	
Gisekiaceae	<i>Gisekia africana</i> (Lour.) Kuntze	Grassland	Barker et al. 124; Goyder 8233; Goyder 8949	Moxico
Hypericaceae	<i>Hypericum oligandrum</i> Milne-Redh.	Wetland	Frisby 4026	
	<i>Psorospermum baumii</i> Engl.	Woodland	Frisby 4003; Goyder 8221	Bié
Ixonanthaceae	<i>Ochthocosmus lemaireanus</i> T.Durand & H.Durand	Woodland	Barker et al. 48; Barker et al. 74; Baum 712; Goyder 8095; Goyder 8311; Goyder 8313; sight record 27	Moxico
Lamiaceae	<i>Alvesia rosmarinifolia</i> Welw.	Woodland	Barker et al. 45; Baum 676; Goyder 8036	
	<i>Clerodendrum baumii</i> Gürke	Grassland	Baum 661; Goyder 8125; Goyder 8367	
	<i>Clerodendrum buchneri</i> Gürke	Grassland	Goyder 8262	
	<i>Clerodendrum formicarium</i> Gürke	Grassland	Goyder & Maiato 8798	
	<i>Endostemon</i> sp. nov.	Grassland	Goyder & Maiato 8762	
	<i>Haumaniastrum katangense</i> (S.Moore) J.Duvign. & Plancke	Grassland	Goyder 8903	
	<i>Haumaniastrum prealtum</i> (Briq.) J.Duvign. & Plancke var. <i>succisifolium</i> (Baker) A.J.Paton	Grassland	Goyder 8341; Goyder 8454	Moxico
	<i>Haumaniastrum sericeum</i> (Briq.) A.J.Paton	Grassland	Barker et al. 87; Goyder 8440	
	<i>Kalaharia uncinata</i> (Schinz) Moldenke	Grassland	Goyder & Maiato 8782	
	<i>Leonotis nepetifolia</i> (L.) R.Br. var. <i>nepetifolia</i>	Ruderal	Baum 822	

Family	Species	Habitat	Vouchers	New Records
Lamiaceae	<i>Ocimum obovatum</i> E.Mey. ex Benth. var. <i>obovatum</i>	Grassland	Goyder & Maiato 8787	
	<i>Plectranthus betonicifolius</i> Baker	Wetland	Goyder 8463	Moxico
	<i>Plectranthus gracillimus</i> (T.C.E.Fr.) Hutch. & Dandy	Grassland	Goyder 8902	
	<i>Plectranthus guerkei</i> Briq.	Grassland	Barker et al. 86	
	<i>Plectranthus mirabilis</i> (Briq.) Launert	Wetland	Barker et al. 140; Baum 794; Goyder 8007; Goyder 8928	
	<i>Pycnostachys gracilis</i> R.D.Good	Woodland	Goyder 8441	
	<i>Tinnea eriocalyx</i> Welw.	Grassland	Goyder 8250	
	<i>Tinnea fusco-luteola</i> Gürke	Grassland	Baum 695	
	<i>Tinnea benguellensis</i> Gürke	Grassland	Baum 697; Goyder 8458	Moxico
	<i>Vitex madiensis</i> Oliv. subsp. <i>milanjiensis</i> (Britten) F.White	Woodland	Frisby 3023; Frisby 3046; Goyder 8044; Goyder 8416; Goyder 8428	
Lecythidaceae	<i>Napoleonaea gossweileri</i> Baker f.	Grassland	Goyder 8107; Goyder & Maiato 8812	Moxico
Leguminosae	<i>Aeschynomene dimidiata</i> Welw. ex Baker	Woodland	Goyder 8392	Moxico
	<i>Aeschynomene glabrescens</i> Welw. ex Baker	Wetland	Goyder & Maiato 8784	
	<i>Albizia adianthifolia</i> (Schumach.) W.Wight	Woodland	Goyder 8212	
	<i>Baphia massaiensis</i> Taub. subsp. <i>obovata</i> (Schinz) Brummitt var. <i>obovata</i>	Woodland	Frisby 3024; Goyder 8092; Goyder 8449; sight record 7; sight record 31; Goyder & Maiato 8780	
	<i>Baphia</i> sp. nov.	Grassland	Goyder & Maiato 8772	
	<i>Bauhinia mendoncae</i> Torre & Hillc.	Woodland	Barker et al. 76; Goyder 8030; Goyder 8391	
	<i>Bauhinia petersiana</i> Bolle subsp. <i>macrantha</i> (Oliv.) Brummitt & J.H. Ross	Woodland	Frisby 4017	
	<i>Bobgunnia madagascariensis</i> (Desv.) J.H. Kirkbr. & Wiersema	Woodland	Goyder 8031; Goyder 8384; Goyder 8429; Goyder 8450	
	<i>Brachystegia bakeriana</i> Hutch. & Burtt Davy	Woodland	Barker et al. 100; Frisby 3014; Goyder 8020; Goyder 8090; Goyder 8116; Goyder 8386; Goyder 8430; Goyder 8432; Goyder 8448; sight record 10	
	<i>Brachystegia longifolia</i> Benth.	Grassland; Woodland	Goyder 8011; Goyder 8328; Goyder 8921	
	<i>Brachystegia spiciformis</i> Benth.	Woodland	Goyder 8038	
	<i>Burkea africana</i> Hook.	Grassland	Goyder 8379; sight record 37; sight record 43	
	<i>Chamaecrista mimosoides</i> (L.) Greene sens. lat.	Woodland	Barker et al. 83	
	<i>Clitoria kaessneri</i> Harms – depauperate form	Grassland	Goyder & Maiato 8758	
	<i>Copaifera baumiana</i> Harms	Grassland; Woodland	Goyder 8018; Goyder 8113; Goyder 8224; Goyder 8388; sight record 3; Goyder & Maiato 8847; Goyder 8919	
	<i>Crotalaria abscondita</i> Welw. ex Baker	Grassland	Goyder 8465	Moxico
	<i>Crotalaria angulicaulis</i> Harms	Grassland	Goyder 8452	Moxico
	<i>Crotalaria annua</i> Milne-Redh.	Grassland	Goyder 8900	
	<i>Crotalaria kambolensis</i> Baker f.	Woodland	Goyder 8424	
	<i>Crotalaria leptoclada</i> Harms	Grassland	Baum 829	
	<i>Crotalaria mendoncae</i> Torre	Woodland	Goyder 8016; Goyder 8103; sight record 26	Cuando Cubango

Family	Species	Habitat	Vouchers	New Records
Leguminosae	<i>Crotalaria stenoptera</i> Welw. ex Baker	Grassland; Wetland; Woodland	Barker et al. 146; Baum 677; Goyder 8093; Goyder 8257	
	<i>Crotalaria youngii</i> Baker f.	Grassland; Woodland	Goyder 8218	Bié
	<i>Crotalaria cf. youngii</i> Baker f.	Grassland; Woodland	Barker et al. 144; Goyder 8944	
	<i>Cryptosepalum exfoliatum</i> De Wild. subsp. <i>pseudotaxus</i> (Baker f.) P.A.Duvign. & Brenan	Woodland	Goyder 8022; Goyder 8323; Goyder 8446; sight record 4; sight record 12; sight record 24	
	<i>Cryptosepalum mimosoides</i> Welw. ex Oliv.	Grassland	Goyder 8337; Goyder & Maiato 8751	Moxico
	<i>Desmodium barbatum</i> (L.) Benth. var. <i>dimorphum</i> (Welw. ex Baker) B.G.Schub.	Grassland	Baum 685; Goyder 8502	
	<i>Dialium englerianum</i> Henriq.	Woodland	Goyder & Maiato 8805	
	<i>Entada arenaria</i> Schinz subsp. <i>arenaria</i>	Grassland; Woodland	Goyder 8390; Goyder & Maiato 8836	Moxico
	<i>Erythrophleum africanum</i> (Welw. ex Benth.) Harms	Woodland	Goyder 8010; Goyder 8380; Goyder 8389; sight record 29; Goyder 8922	
	<i>Eryhrina baumii</i> Harms	Grassland	Frisby 4034; Goyder & Maiato 8767	
	<i>Guibourtia coleosperma</i> (Benth.) J.Léonard	Woodland	Goyder 8035; Goyder 8377; sight record 2; sight record 13; sight record 23; sight record 30; sight record 35	
	<i>Indigofera baumiana</i> Harms	Grassland	Baum 819; Goyder & Maiato 8818	
	<i>Indigofera sutherlandioides</i> Baker	Woodland	Goyder 8046; Goyder 8955	
	<i>Kotschy strobilantha</i> (Welw. ex Baker) Dewit & P.A.Duvign.	Grassland	Barker et al. 56; Goyder 8091; Goyder 8943	
	<i>Julbernardia paniculata</i> (Benth.) Troupin	Woodland	Goyder 8012; Goyder 8089; Goyder 8123; Goyder 8308; Goyder 8443; sight record 11; sight record 19	
	<i>Macrotyloma rupestre</i> (Welw. ex Baker) Verdc.	Woodland	Goyder 8247	Moxico
	<i>Pterocarpus angolensis</i> DC.	Woodland	Barker et al. 52B; Goyder 8009; Goyder 8382; sight record 6; sight record 22; sight record 28	
	<i>Rhynchosia procurrens</i> (Hiern) K.Schum.	Woodland	Barker et al. 77	
	<i>Sphenostylis erecta</i> (Baker f.) Hutch. ex Baker f. subsp. <i>obtusifolia</i> (Harms) Potter & Doyle	Woodland	Goyder 8248; Goyder 8950	Moxico
Lentibulariaceae	<i>Genlisea angolensis</i> R.D.Good	Wetland	Frisby 3073; Goyder 8120; Goyder 8315; Goyder 8371	Moxico
	<i>Utricularia gibba</i> L.	Wetland	Barker et al. 44a; Goyder 8098	
	<i>Utricularia spiralis</i> Sm.	Wetland	Frisby 3094; Goyder 8114	
	<i>Utricularia subulata</i> L.	Wetland	Baum 691; Goyder 8370	
	<i>Utricularia stellaris</i> L.f.	Wetland	Frisby 3088	
Limeaceae	<i>Limeum fenestratum</i> (Fenzl) Heimerl	Grassland	Barker et al. 80; Baum 688	
Linderniaceae	<i>Crepidiorhopalon</i> ?sp. nov.	Grassland	Goyder 8917	
Loranthaceae	<i>Englerina gabonensis</i> (Engl.) Balle	Woodland	Goyder 8413	
	<i>Tapinanthus dependens</i> (Engl.) Danser	Woodland	Barker et al. 137	
Lythraceae	<i>Rotala myriophyloides</i> Welw. ex Hiern	Wetland	Barker et al. 68	
Malvaceae	<i>Grewia falcistipula</i> K.Schum.	Woodland	Frisby 3022	
	<i>Grewia</i> sp.	Ruderal	Goyder & Maiato 8819	
	<i>Triumfetta dekindtiana</i> Engl.	Woodland	Barker et al. 133	

Family	Species	Habitat	Vouchers	New Records
Melastomataceae	<i>Antherotoma debilis</i> (Sond.) Jacq.-Fel.	Wetland	Barker et al. 47; Frisby 4031; Goyder 8094	
	<i>Dissotis brazzae</i> Cogn.	Grassland	Goyder 8927	
	<i>Dissotis rhinanthifolia</i> (Brenan) A.Fern. & R.Fern. var. <i>rhinanthifolia</i>	Wetland	Goyder & Maiato 8823	
	<i>Dissotis welwitschii</i> Cogn.	Wetland	Goyder 8240	Moxico
	<i>Memecylon huillense</i> A.Fern. & R.Fern.	Woodland	Goyder 8399	Moxico
Meliaceae	<i>Trichilia quadrivalvis</i> C.DC.	Woodland	Frisby 3070; Goyder & Maiato 8839	Moxico
Menyanthaceae	<i>Nymphoides forbesiana</i> (Griseb.) Kuntze	Wetland	Goyder & Maiato 8824	Moxico
	<i>Nymphoides indica</i> (L.) Kuntze subsp. <i>occidentalis</i> A.Raynal	Wetland	Barker et al. 113	
Moraceae	<i>Ficus pygmaea</i> Welw. ex Hiern	Wetland	Barker et al. 141	
	<i>Ficus verruculosa</i> Warb.	Wetland	Goyder 8320	
Myricaceae	<i>Morella serrata</i> (Lam.) Killick	Wetland	Goyder 8914	
Myrtaceae	<i>Syzygium cordatum</i> Hochst. ex Krauss subsp. <i>cordatum</i>	Wetland	Barker et al. 145; Goyder 8319	
	<i>Syzygium guineense</i> (Willd.) DC. subsp. <i>huillense</i> (Hiern) F.White	Grassland	Barker et al. 67; Frisby 3045; Goyder 8339; Goyder & Maiato 8835	
	<i>Syzygium owariense</i> (P.Beauv.) Benth.	Humid Forest	Goyder 8326	Moxico
Ochnaceae	<i>Brackenridgea arenaria</i> (De Wild. & T.Durand) N.Robson	Grassland	Frisby 3015; Frisby 3016; Frisby 3060; Goyder & Maiato 8781; Goyder & Maiato 8804	
	<i>Ochna katangensis</i> De Wild.	Grassland	Goyder & Maiato 8754A	
	<i>Ochna manikensis</i> De Wild.	Grassland	Frisby 3031; Goyder 8108; Goyder 8309	Moxico
	<i>Ochna pulchra</i> Hook.	Woodland	Goyder 8013; Goyder 8383; sight record 21	Moxico
	<i>Ochna pygmaea</i> Hiern	Grassland	Frisby 3059; Goyder & Maiato 8754B	Moxico
Olacaceae	<i>Olax gossweileri</i> Exell & Mendonça	Woodland	Goyder & Maiato 8846	Moxico
Oleaceae	<i>Olea capensis</i> L. subsp. <i>macrocarpa</i> (C.H.Wright) I.Verd.	Woodland	Goyder 8437	Moxico
Onagraceae	<i>Ludwigia octovalvis</i> (Jacq.) P.H.Raven	Wetland	Barker et al. 138	
Orobanchaceae	<i>Buchnera prorepens</i> Engl. & Gilg	Grassland	Goyder 8349; Goyder 8451	Moxico
	<i>Buchnera attenuata</i> Skan	Wetland	Frisby 3086; Frisby 4021	
	<i>Buchnera</i> sp. not matched at K	Grassland	Goyder 8276	
	<i>Buchnera welwitschii</i> Engl.	Grassland	Barker et al. 93	
	<i>Cycnium tubulosum</i> (L.f.) Engl. subsp. <i>tubulosum</i>	Grassland	Frisby 4019	
	<i>Gerardiina angolensis</i> Engl.	Wetland	Goyder 8101	
	<i>Gerardiina angolensis</i> Engl. – unusual form with branched inflorescence	Wetland	Goyder 8293	
	<i>Melasma calycinum</i> (Hiern) Hemsl.	Wetland	Frisby 4018; Goyder & Maiato 8760	
	<i>Micrargeriella aphylla</i> R.E.Fr.	Wetland	Goyder & Maiato 8783	Moxico
	<i>Sopubia simplex</i> (Hochst.) Hochst.	Wetland	Frisby 3083; Frisby 4024; Goyder & Maiato 8821	Moxico
Passifloraceae	<i>Striga angolensis</i> K.I.Mohamed & Musselman	Wetland	Goyder 8336	
	<i>Striga bilabiata</i> (Thunb.) Kuntze	Wetland	Frisby 4028; Goyder & Maiato 8795	Moxico
	<i>Basananthe baumii</i> (Harms) W.J. de Wilde var. <i>caerulescens</i> (A.Fern. & R.Fern.) W.J. de Wilde	Grassland	Goyder & Maiato 8826	
Passifloraceae	<i>Paropsia brazzaeana</i> Baill.	Woodland	Barker et al. 101; Goyder 8024; sight record 25; Goyder 8920	
Pedaliaceae	<i>Sesamum calycinum</i> Welw.	Grassland	Frisby 4022	

Family	Species	Habitat	Vouchers	New Records
Peraceae	<i>Clutia benguelensis</i> Müll.Arg.	Grassland	Goyder 8455	Moxico
Phrymaceae	<i>Mimulus gracilis</i> R.Br.	Wetland	Barker et al. 135	
Phyllanthaceae	<i>Bridelia duvigneaudii</i> J.Léonard	Woodland	Goyder 8423	Moxico
	<i>Hymenocardia acida</i> Tul.	Woodland	Goyder 8231	
	<i>Uapaca nitida</i> Müll.Arg.	Woodland	Goyder 8047; Goyder 8427	
	<i>Uapaca nitida</i> Müll.Arg. – pyrophytic form	Grassland Woodland	Goyder 8217	
Picodendraceae	<i>Oldfieldia dactylophylla</i> (Welw. ex Oliv.) J.Léonard	Grassland	Goyder 8267; Goyder 8421	
Plantaginaceae	<i>Limnophylla ceratophylloides</i> (Hiern) Skan	Wetland	Goyder 8318	Moxico
Polygalaceae	<i>Polygala africana</i> Chodat	Wetland	Frisby 4027	
	<i>Polygala arenicola</i> Gürke	Woodland	Barker et al. 123; Goyder 8229	
	<i>Polygala dewevrei</i> Exell	Wetland	Goyder 8361 (blue fls); Goyder 8362 (white fls); Goyder & Maiato 8849; Goyder 8926	Bié
	<i>Polygala gomesiana</i> Welw. ex Oliv.	Wetland	Goyder 8374	Cuando Cubango
	<i>Polygala kalaxariensis</i> Schinz	Grassland	Barker et al. 96	
	<i>Polygala mendoncae</i> E.M.A.Petit	Woodland	Goyder 8037; Goyder 8417	
	<i>Polygala nambalensis</i> Gürke	Grassland	Goyder 8453	Moxico
	<i>Polygala nematophylla</i> Exell	Grassland	Goyder 8366	Moxico
	<i>Polygala paludicola</i> Gürke	Wetland	Barker et al. 92; Frisby 3040; Frisby 3085; Goyder 8119; Goyder 8236	
	<i>Polygala poggei</i> Gürke	Grassland	Goyder 8278	Moxico
	<i>Polygala rivularis</i> Gürke	Grassland	Barker et al. 90	
	<i>Polygala robusta</i> Gürke	Grassland	Baum 704; Frisby 3047; Goyder 8085; Goyder 8280; Goyder 8303; Goyder 8411	Moxico
	<i>Polygala spicata</i> Chodat	Wetland	Frisby 3084; Goyder 8235; Goyder 8363	
	<i>Polygala welwitschii</i> Chodat subsp. <i>pygmaea</i> (Gürke) Paiva	Grassland	Goyder 8350; Goyder 8916	Moxico
Polygonaceae	<i>Securidaca longipedunculata</i> Fresen.	Woodland	sight record 44	
	<i>Oxygonum annum</i> S.Ortíz & Paiva	Grassland	Goyder 8348	Moxico
	<i>Oxygonum fruticosum</i> Dammer ex Milne-Redh.	Woodland	Goyder 8008; Goyder 8105; Goyder 8954	
	<i>Oxygonum pachybasis</i> Milne-Redh.	Grassland	Frisby 3090; Goyder & Maiato 8799	Moxico
Proteaceae	<i>Faurea delevoyi</i> De Wild.	Wetland; Woodland	Goyder 8398	
	<i>Faurea saligna</i> Harv.	Woodland	Barker et al. 102	
	<i>Protea angolensis</i> Welw. var. <i>angolensis</i>	Grassland	Goyder 8410	Moxico
	<i>Protea baumii</i> Engl. & Gilg subsp. <i>baumii</i>	Grassland	Barker et al. 106	
	<i>Protea petiolaris</i> (Hiern) Baker & C.H.Wright subsp. <i>petiolaris</i>	Grassland	Goyder 8412	Moxico
	<i>Protea poggei</i> Engl. subsp. <i>haemantha</i>	Woodland	Baum 709; Goyder 8215; sight record 14; Goyder 8956	Bié; Moxico
	<i>Protea welwitschii</i> Engl.	Grassland	Goyder 8117; Goyder 8353; Goyder 8397; Goyder 8460	Moxico
Ranunculaceae	<i>Clematis villosa</i> DC.	Grassland	Goyder 8912	
Rosaceae	<i>Cliffortia nitidula</i> R.E.Fr. & T.C.E.Fr. var. <i>angolensis</i> (Weim.) Brenan	Grassland	Barker et al. 103; Baum 650; Goyder 8395; Goyder & Maiato 8855; Goyder 8932	
Rubiaceae	<i>Ancylanthos rubiginosus</i> Desf.	Grassland Woodland	Frisby 3056; Goyder 8115; Goyder & Maiato 8775; Goyder & Maiato 8848	

Family	Species	Habitat	Vouchers	New Records
Rubiaceae	<i>Berriera</i> sp.	Humid Forest	Goyder 8325	
	<i>Diodia flavescens</i> Hiern	Grassland	Barker et al. 99	
	<i>Fadogia cienkowskii</i> Schweinf.	Grassland	Frisby 3018; Goyder 8501	
	<i>Fadogia fuchsoides</i> Oliv.	Grassland	Goyder 8340	
	<i>Fadogia gossweileri</i> Robyns	Woodland	Frisby 4032	
	<i>Fadogia tomentosa</i> De Wild. var. <i>flaviflora</i> (Robyns) Verdc.	Woodland	Goyder 8246	
	<i>Ganguelia gossweileri</i> (S.Moore) Robbr.	Grassland	Goyder & Maiato 8815	
	<i>Gardenia imperialis</i> K.Schum.	Wetland	Goyder 8321; Goyder 8394; sight record 39	
	<i>Gardenia resiniflua</i> Hiern subsp. <i>resiniflua</i>	Woodland	Barker et al. 143	
	<i>Gardenia resiniflua</i> Hiern subsp. <i>resiniflua</i> – suffrutescent form	Woodland	Barker et al. 51; Frisby 4007	
	<i>Leptactina benguelensis</i> (Benth. & Hook.f.) R.D.Good	Woodland	Frisby 4029; Goyder & Maiato 8842	
	<i>Morinda angolensis</i> (R.D.Good) F.White	Grassland	Goyder & Maiato 8756; Goyder & Maiato 8851	Moxico
	<i>Pavetta nitidula</i> Hiern	Woodland	Goyder & Maiato 8840	
	<i>Pavetta</i> sp. 1	Woodland	Goyder 8242	
	<i>Pavetta</i> sp. 2	Woodland	Goyder 8249	
	<i>Pavetta</i> sp. 3	Grassland	Goyder 8301	
	<i>Psychotria</i> sp.	Humid Forest	Goyder 8324	
	<i>Pydrax gilletii</i> (De Wild.) Bridson	Woodland	Goyder 8434	
	<i>Pydrax</i> sp.	Woodland	Goyder 8433	
	<i>Pygmaeothamnus zeyheri</i> (Sond.) Robyns	Grassland	Goyder & Maiato 8808; Goyder & Maiato 8811	
Rutaceae	<i>Rothmannia engleriana</i> (K.Schum.) Keay var. <i>engleriana</i>	Woodland	Goyder 8420	
	<i>Rytigynia orbicularis</i> (K.Schum.) Robyns	Woodland	Goyder 8227	
	<i>Tricalysia angolensis</i> A.Rich. ex DC.	Woodland	Barker et al. 73	
	<i>Tricalysia</i> sp.	Woodland	Goyder 8435	
	<i>Vangueria</i> sp. not matched at K	Woodland	Goyder 8265	
	<i>Vangueriopsis</i> cf. <i>lanceiflora</i> (Hiern) Robyns	Woodland	Goyder 8422	
	<i>Zanthoxylum gilletii</i> (De Wild.) P.G.Waterman	Humid Forest	Goyder 8327	Moxico
	<i>Thesium atrum</i> A.W.Hill	Grassland	Goyder 8342; Goyder & Maiato 8813	Moxico
	<i>Thesium subaphyllum</i> Engl.	Grassland	Barker et al. 91; Goyder 8347; Goyder 8937	
	<i>Chrysophyllum bangweolense</i> R.E.Fr.	Woodland	Goyder & Maiato 8841	
Sapotaceae	<i>Englerophytum magalismontanum</i> (Sond.) T.D.Penn.	Woodland	Goyder 8033; Goyder 8387; Goyder 8447; sight record 5	
	<i>Englerophytum magalismontanum</i> (Sond.) T.D.Penn. – pyrophytic form	Grassland Woodland	Goyder & Maiato 8854	
	<i>Hannoia chlorantha</i> Engl. & Gilg	Woodland	Barker et al. 66; Barker et al. 129; Baum 674; Goyder 8946	Moxico
Thymelaeaceae	<i>Craterosiphon quarrei</i> Staner	Woodland	Goyder 8219; Goyder & Maiato 8845	Moxico
	<i>Gnidia gossweileri</i> (S.Moore) B.Peterson subsp. <i>gossweileri</i>	Wetland; Grassland	Barker et al. 88	
	<i>Gnidia kraussiana</i> Meisn.	Grassland	Goyder 8110; Goyder & Maiato 8817	
Umbelliferae	<i>Afrocarum imbricatum</i> (Schinz) Rauschert	Wetland	Goyder 8957	
	<i>Pseudoselinum angolense</i> (C.Norman)	Grassland; Woodland	Goyder 8045; Goyder 8251; Goyder 8953	Bié; Moxico

Acknowledgements

We are extremely grateful to the British de-mining charity The Halo Trust for logistical support and for keeping us safe. John Hilton and Steve Boyes of the Wild Bird Trust, South Africa initiated the Okavango Wilderess Project and ensured all expeditions ran smoothly. We are most grateful to the Collections Unit INBAC of the Angolan Ministry of the Environment in Luanda for facilitating permits and permission.

References

- Albuquerque S (2008) Friedrich Welwitsch. In: Figueiredo E, Smith GF (Eds) *Plantas de Angola* [Plants of Angola]. Strelitzia 22: 2–3.
- African Plant Database (2018) Conservatoire et Jardin botaniques de la Ville de Genève and South African National Biodiversity Institute, Pretoria. <http://www.ville-ge.ch/musinfo/bd/cjb/africa/>
- APG IV (2016) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society* 181(1): 1–20. <https://doi.org/10.1111/boj.12385>
- Barbosa LAG (1970) *Carta Fitogeográfica de Angola*. Instituto de Investigação Científica de Angola, Luanda.
- Cavaco A (1959) Contribution à l'Étude de la Flore de la Lunda d'Après les Récoltes de Gossweiler (1946–1948). *Publicações Culturais da Companhia de Diamantes de Angola* 42: 1–230.
- Cheek M, Feika A, Lebbie A, Goyder D, Tchiengue B, Sene O, Choutou P, van der Burgt X (2017) A synoptic revision of *Inversodicraea* Engl. ex R.E. Fries (Podostemaceae). *Blumea* 62(2): 125–156. <https://doi.org/10.3767/blumea.2017.62.02.07>
- Darbyshire I, Goyder D, Crawford F, Gomes A (2011) Report on the rapid botanical survey of the Lagoa Carumbo Region, Lunda Norte Prov., Angola, April/May 2011, inc. Appendix 2: checklist to the flowering plants, gymnosperms and pteridophytes of Lunda Norte Prov, Angola [537 taxa]. In: Huntley BJ (Ed.) *Biodiversity Rapid Assessment of the Lagoa Carumbo Area, Lunda Norte, Angola*. Annex 3, 59–98. <https://www.scribd.com/document/212410297/Carumbo-Final-Report>
- Darbyshire I, Goyder D, Crawford F, Gomes A (2014) Update to Report on the rapid botanical survey of the Lagoa Carumbo Region, Lunda Norte Prov., Angola for the Angolan Ministry of the Environment, following further field studies in 2013, inc. Appendix 2: checklist to the flowering plants, gymnosperms and pteridophytes of Lunda Norte Prov, Angola [752 taxa]. In: Huntley BJ (Ed.) *Biodiversity Rapid Assessment of the Lagoa Carumbo Area, Lunda Norte, Angola*. Annex 3, 59–98.
- Dargie GC, Lewis SL, Lawson IT, Mitchard ETA, Page SE, Bocko YE, Ifo SA (2017) Age, extent and carbon storage of the central Congo Basin peatland complex. *Nature* 542(7639): 86–90. <https://doi.org/10.1038/nature21048>
- Diniz AC (1973) Característica mesológicas de Angola. *Missão de Inquéritos de Angola, Nova Lisboa*, 482 pp.

- Figueiredo E, Smith GF (Eds) (2008) Plants of Angola/Plantas de Angola. Strelitzia 22: 1–279. <https://www.sanbi.org/sites/default/files/documents/documents/strelitzia-22-2008.pdf>
- Figueiredo E, Soares M, Siebert G, Smith GF, Faden RB (2009) The botany of the Cunene-Zambezi Expedition with notes on Hugo Baum (1867–1950). Bothalia 39(2): 185–211. <https://doi.org/10.4102/abc.v39i2.244>
- Finckh M, Revermann R, Aidar MPM (2016) Climate refugees going underground – a response to Maurin et al. (2014). The New Phytologist 209(3): 904–909. <https://doi.org/10.1111/nph.13567>
- Gonçalves FMP, Goyder DJ (2016) A brief botanical survey into Kumbira forest, an isolated patch of Guineo-Congolian biome. PhytoKeys 65: 1–14. <https://doi.org/10.3897/phytkeys.65.8679>
- Gonçalves FMP, Tchamba J, Goyder DJ (2016) *Schistostephium crataegifolium* (Compositae: Anthemideae), a new generic record for Angola. Bothalia 46(1): a2029. <https://doi.org/10.4102/abc.v46i1.2029>
- Goyder DJ (2016) *Genlisea angolensis* – new data. Plant Carnivora 37: 29–33.
- Goyder DJ, Gonçalves FMP (in press) The Flora of Angola: Collectors, Richness and Endemism. In: Huntley BJ, Russo V, Lages F, Ferrand N (Eds) Biodiversity of Angola. Science and Conservation: A Modern Synthesis. Springer.
- Hind DJN, Goyder DJ (2014) *Stomatianthes tundavalaensis* (Compositae: Eupatorieae: Eupatoriinae), a new species from Huíla Province, Angola, and a synopsis of the African species of Stomatianthes. Kew Bulletin 69(4): 9545. <https://doi.org/10.1007/s12225-014-9545-6>
- Marshall CAM, Wieringa JJ, Hawthorne WD (2016) Bioquality hotspots in the tropical African flora. Current Biology 26(23): 3214–3219. <https://doi.org/10.1016/j.cub.2016.09.045>
- Maurin O, Davies TJ, Burrows JE, Daru BH, Yessoufou K, Muasya AM, van der Bank M, Bond WJ (2014) Savanna fire and the origins of the ‘underground forests’ of Africa. The New Phytologist 204(1): 201–214. <https://doi.org/10.1111/nph.12936>
- Nordal I, Kwembeya EG (2004) *Crinum binghamii* sp. nov. – with a key to *Crinum* species with radially symmetrical flowers in mainland Africa. Kew Bulletin 59(4): 599–603. <https://doi.org/10.2307/4110918>
- Oldeman J, Erb C, Finckh M, Jürgens N (Eds) (2013) Environmental assessments in the Okavango region. Biodiversity & Ecology 5: 1–418. https://doi.org/10.7809/b-e.vol_05
- Reiley J, Page S (2016) Tropical peatland of the world. In: Osaki M, Tsuji N (Eds) Tropical Peatland Ecosystems. Springer, 2–32.
- Roux JP (2009) Synopsis of the Lycopodiophyta and Pteridophyta of Africa, Madagascar and neighbouring islands. Strelitzia 23. South African National Biodiversity Institute, Pretoria. <https://www.sanbi.org/documents/synopsis-of-the-lycopodiophyta-and-pteridophyta-of-africa-madagascar-and-neighbouring-islands-strelitzia-23/>
- Sosef MSM, Dauby G, Blach-Overgaard A, van der Burgt X, Catarino L, Damen T, Deblauwe V, Dessein S, Dransfield J, Droissart V, Duarte MC, Engeldorf H, Fadeur G, Figueira R, Gereau RE, Hardy OJ, Harris DJ, de Heij J, Janssens S, Klomberg Y, Ley AC, Mackinder BA, Meerts P, van de Poel JL, Sonké B, Stévert T, Stoffelen P, Svenning J-C, Sepulchre P, Zaiss R, Wieringa JJ, Couvreur TLP (2017) Exploring the floristic diversity of tropical Africa. BMC Biology 15(1): 15. <https://doi.org/10.1186/s12915-017-0356-8>

- Stropp J, Ladle RJ, Malhado ACM, Hortal J, Gaffuri J, Temperley WH, Skøien JO, Mayaux P (2016) Mapping ignorance: 300 years of collecting flowering plants in Africa. *Global Ecology and Biogeography* 25: 1085–1096. <https://doi.org/10.1111/geb.12468>
- Vorontsova MS, Besnard G, Forest F, Malakasi P, Moat J, Clayton WD, Ficinski P, Savva GM, Nanjarisoa OP, Razanatsara J, Randriatsara FO, Kimeu JM, Luke WRQ, Kayombo C, Linder HP (2016) Madagascar's grasses and grasslands: Anthropogenic or natural? *Proceedings. Biological Sciences* 283(1823): 20152262. <https://doi.org/10.1098/rspb.2015.2262>
- Warburg O (1903) Kunene–Sambesi–Expedition. H. Baum. Kolonial-Wirtschaftliches Komitee, Berlin, 593.
- WCSP (2016) World Checklist of Selected Plant Families. Facilitated by the Royal Botanic Gardens, Kew. <http://apps.kew.org/wcsp/>
- White F (1977) The underground forests of Africa: A preliminary review. *Gardens' Bulletin (Singapore)* 29: 57–71. <https://www.nparks.gov.sg/sbg/research/publications/gardens-bulletin-singapore/-/media/sbg/gardens-bulletin/4-4-29-08-y1977-v29-gbs-pg-57.pdf>
- Zimudzi C, Archer RH, Kwembeya EG, Nordal I (2008) Amaryllidaceae. In: Timberlake JR, Martins ES (Eds) *Flora Zambesiaca* 13(1): 97–134.