

An updated checklist of *Begonia* (Begoniaceae) in Laos, with two new species and five new records

Hong-Bo Ding^{1,2*}, Mya Bhone Maw^{1,3*}, Bin Yang^{1,2},
Somsanith Bouamanivong⁴, Yun-Hong Tan^{1,2}

1 Southeast Asia Biodiversity Research Institute & Center for Integrative Conservation, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Menglun, Mengla, Yunnan 666303, China **2** Center of Conservation Biology, Core Botanical Gardens, Chinese Academy of Sciences, Menglun, Mengla, Yunnan 666303, China **3** University of Chinese Academy of Sciences, Shijingshan District, Beijing 100049, China **4** Ecology Division, Biotechnology and Ecology Institute, Ministry of Science and Technology, P.O. Box: 2279, Vientiane Capital, Lao PDR

Corresponding author: Yun-Hong Tan (tyh@xtbg.org.cn)

Academic editor: Xiao-Hua Jin | Received 11 October 2019 | Accepted 8 November 2019 | Published 10 January 2020

Citation: Ding H-B, Maw MB, Yang B, Bouamanivong S, Tan Y-H (2020) An updated checklist of *Begonia* (Begoniaceae) in Laos, with two new species and five new records. In: Jin X-H, Xia N-H, Tan Y-H (Eds) Plant diversity of Southeast Asia-II. PhytoKeys 138: 187–201. <https://doi.org/10.3897/phytokeys.138.46718>

Abstract

Two new species of *Begonia* L. (Begoniaceae), *B. laotica* (sect. *Parvibegonia*) and *B. hypoleuca* (sect. *Reichenheimia*), from north Laos are described and illustrated. *Begonia augustinei*, *B. dryadis*, *B. lancangensis*, *B. sizemoreae* and *B. silentensis* subsp. *mengyangensis* were newly recorded taxa in Laos. Furthermore, an updated checklist of *Begonia* of Laos is also compiled.

Keywords

sect. *Parvibegonia*, sect. *Reichenheimia*, taxonomy, new records

Introduction

Begonia Linnaeus (1753: 1056) is the sixth largest genera of angiosperms and the number of accepted species of *Begonia* currently stands at 1947 (Hughes et al. 2015) and is likely to rise to well over 2000 (Moonlight et al. 2018; Tian et al. 2018). According to

* Contributed equally as the first authors.

the annotated checklist of south east Asia *Begonia* by Hughes (2008) and recent taxonomic publications (Hughes 2007; Newman et al. 2007; De Wilde et al. 2011; Averyanov and Nguyen 2012; Souvannakhoummane et al. 2016, 2018; Yang et al. 2018; Averyanov et al. 2019), 19 species of *Begonia* have been described from Laos, so far.

As part of the botanical inventory of China-Laos transboundary biodiversity conservation, we carried out floristic surveys in the Phou Hin Phee National Biodiversity Conservation Area, Oudomxay Province, the Nam Ha National Biodiversity Conservation Area in Luang Namtha Province and the Phou Dean Din National Bio-Diversity Park, Phongsaly Province of northern Laos. During the fieldwork from 2018 to 2019, some interesting species of *Begonia* have been collected. After reviewing the literature and herbarium specimens, we described and illustrated two species as new to science and five species as new records for the flora of Laos. We provided an updated total of 26 species for the *Begonia* flora of Laos.

Begonia laotica is classified under *B.* sect. *Parvibegonia* A. de Candolle (1859: 136) according to its small and tuberous habitat, axile placentation, 2-locular ovary, 2 styles and fruit with 3 unequal wings (Doorenbos et al. 1998; Moonlight et al. 2018). *Begonia hypoleuca* belongs to *B.* sect. *Reichenheimia* (Klotzsch, 1855: 174) A. de Candolle (1864: 385), based on the morphological characteristics, including 2 tepals in male flowers, axile and entire placentation, 3-locular ovary, 3 styles and fruit with 3 wings (Doorenbos et al. 1998; Moonlight et al. 2018).

Material and methods

Measurements and morphological character assessments of the new species have been examined, based on fresh materials and dried specimens. They have been compared with morphologically similar species by affinities inferred using descriptions (Ku et al. 2007; Camfield and Hughes 2018) and type specimens in herbaria (BM, E, K, B, KUN, P, HITBC, YNU, LE and FOF). Protogues and images of type specimens were gathered from JSTOR Global Plants (<http://plants.jstor.org>).

Taxonomic treatments

Begonia laotica Y.H.Tan & H.B.Ding, sp. nov.

urn:lsid:ipni.org:names:77204211-1

Figure 1

sect. *Parvibegonia* A. de Candolle

Diagnosis. The new species is mostly similar to *B. josephii* A. de Candolle (1859: 126) in its peltate, stemless, tuberous habit, but significantly differs by fleshy reddish bristles at the petiole apex (vs. without), ovary 2-locular (vs. 3-locular) and styles 2 (vs. 3).




Figure 1. *Begonia laotica* Y.H.Tan & H.B.Ding, sp. nov. **A** habitat **B** leaf blade adaxially **C** leaf blade abaxially (showing fleshy reddish bristles at the petiole apex) **D** flowers (front view) **E** flowers (lateral view) **F** inflorescence **G** mature fruits **H** habit **I** staminate flower **J** pistillate flower (young) **K** stamens (lateral view) **L** stamens (front view) **M** cross section of ovary. Photographed by H.B. Ding.

Type. Laos. Oudomxay Province, Maung Xai, Phou Hin Phee National Biodiversity Conservation Area, 20°43'18.69"N, 102°08'47.97"E. 1378 m elev., 25 October 2018, flowering, Y.H. Tan & H.B. Ding L0827 (holotype: HITBC!; isotypes: HNL!).

Description. Monoecious herb, stemless, tuberous, 18–25 cm high, tuber small, subglobose, 12–15 mm in diameter. **Stipule:** broadly lanceolate, 2–3 mm long, membranous, sparsely glandular hairy, margin fringed by hairs, deciduous. **Leaf:** **petiole** 6–15 cm long, glabrous or sparsely puberulous; blade ovate, nearly symmetric, 9.2–13.5 × 5.8–11 cm, peltate, base rounded, apex acuminate, margin slightly denticulate, with sparse short hairs, upper surface green, glabrous, lower surface paler, venation palmate-pinnate, tomentose on the veins, fleshy reddish bristles at the petiole apex, thinly succulent in life, thinly papery when dried. **Inflorescence:** axillary, terminal cyme; peduncle puberulent, 7.2–10.9 cm long, branching 2–3 times; bracts narrowly ovate, ca. 4.0 × 2.5 mm, membranous, reddish line along the middle, slightly glandular hairy, margin fringed by glandular hairs. **Staminate flower:** pedicel 1.1–1.8 cm, membranous, covered with glandular hairs, tepals 4, pink, unequal, outer 2, ovate, 10–13 × 7–17 mm, glandular hairs on the outer surface, margin entire, inner 2, elliptic, 7–10 × 2–4 mm, glabrous, margin entire; stamens 15–20, anther obovate, ca. 1 mm. **Pistillate flower** (young): pedicel 0.9–1.2 cm long, membranous, glandular hairy, tepals 5, pink, unequal, outer 2, ovate, 4–6 × 3–4 mm, inner 2, elliptic, 3–5 × 1–3 mm, glandular hairs on the outer surface, especially at the base; **ovary** 2-locular, placentae axial, placentae 2 per locule, **styles** 2, stigmas bifid with twisted bands, highly convolute, golden yellow. **Capsule** nodding, obovate, pinkish tomentose, unequally 3-winged, abaxial wing oblong, ca. 13 mm long, lateral wings shorter, ca. 3 mm long.

Phenology. Flowering in October–November and fruiting in December.

Distribution. Endemic to the type locality, Oudomxay Province, Maung Xai, Phou Hin Phee National Biodiversity Conservation Area, Laos.

Ecology. Restricted to the shaded base of limestone cliff at ca. 1378 m elevation.

Etymology. The specific epithet '*laotica*' refers to the type locality in Laos.

Conservation status. Data Deficient (DD) (IUCN 2017). The new species was only collected from the limestone cliff of Hin Phee National Biodiversity Conservation Area. Due to its remote habitat, we suggested that the species may not suffer from strong human disturbance. According to the available information, we proposed the preliminary conservation status of this new species as Data Deficient (DD) (IUCN 2017).

Notes. *Begonia laotica* is phenotypically most similar to *B. josephii* in its tuberous habit, ovate and peltate leaves, but it can easily be distinguished by its glabrous upper leaf surface (vs. hispid all over), fleshy reddish bristles at the petiole apex (vs. without), bracts narrowly ovate and fringed by glandular hairs (vs. lanceolate and glabrous), male flowers pedicels with glandular hair (vs. glabrous), outer tepals ovate (vs. broadly obovate), inner tepals elliptic (vs. spathulate), anther obovate (vs. ellipsoid), female flowers pedicels with sparsely glandular hair (vs. glabrous), glandular hair on the outer surface (vs. glabrous), ovary 2-locular (vs. 3-locular) and styles 2 (vs. 3). This new species also shares similar characteristics with *B. subperfoliata* Parish ex Kurz (1873: 81) in tuberous habit and peltate leaves, but it can easily be distinguished by its ovary 2-locular (vs. 3-locular) and styles 2 (vs. 3).

***Begonia hypoleuca* Y.H.Tan & H.B.Ding, sp. nov.**

urn:lsid:ipni.org:names:77204213-1

Figure 2

sect. *Reichenheimia* (Klotzsch) A. de Candolle

Diagnosis. The new species is mostly similar to *B. henryi* Hemsley (1887: 322), but differs by its glabrous leaves, petiole with linear red dots and pubescent, slightly denticulate leaf margin and fewer stamens.

Type. Laos. Luang Namtha Province, Nam Ha National Biodiversity Conservation Area, Nam O. Village, 20°04'48.16"N, 101°06'58.46"E. 656 m elev., 24 October 2018, Y.H. Tan, B. Yang, H.B. Ding & X.D. Zeng L0792 (holotype: HITBC!).

Description. Monoecious herb, stemless, tuber small, globose with fibrous roots, 10–13 mm in diameter. **Stipule:** caducous. **Leaf:** petiole 7–10 cm long, succulent, with reddish linear dots, sparsely pubescent or subglabrous. **Blade** slightly triangular ovate or broadly ovate, 6–12 × 5–8.5 cm, base cordate, apex acuminate, margin slightly denticulate, upper surface green, lower surface paler or silver, glabrous, venation palmate-pinnate, midrib pale green, lateral veins 4–6 pairs. **Inflorescence** axillary, cymose, peduncle ca. 6 cm long, reddish, glabrous, branching 2–4 times, 2–6 male flowers, ca. 2 pistillate flowers, bracts caducous, ca. 2 × 1 mm. **Staminate flower:** pedicel 1.0–1.2 cm long, membranous, reddish at the basal part and pink, glabrous, tepals 2, pink, margin entire, glabrous on both surfaces, broadly ovate, 12 × 10 mm, apex acute, base rounded; stamens 10–15, filaments 1.5–2 mm long, anthers 1–1.5 mm, ellipsoid, yellow. **Pistillate flower:** pedicel ca. 1.8 cm long, reddish to pink, tepals unknown, **ovary** 3-locular, placentae axial, entire, **styles** 3. **Capsule** pale green, nodding, ellipsoid, dorsal wing enlarged, 7–10 mm long, two lateral wings, smaller, 2–3 mm long.

Phenology. Flowering and fruiting in October–November.

Distribution. The species is known only from the type locality, Luang Namtha Province, Nam Ha National Biodiversity Conservation Area, Laos.

Ecology. The species was collected on shaded moist limestone rock surfaces of stone forest at 656 m elevation.

Etymology. The specific epithet ‘*hypoleuca*’ refers to the lower surface of its paler or silver leaf.

Conservation status. Data Deficient (DD). *Begonia hypoleuca* has been collected on only one occasion from the type locality. However, the type locality is located in the Nam Ha National Biodiversity Conservation Area and we did not discover any strong pressure on the species. Thus, the species preliminarily has been assigned in the Data Deficient (DD) category as being appropriate according to the guidelines for using the IUCN Red List Categories and Criteria (IUCN 2017).

Notes. *Begonia hypoleuca* is most similar in morphological characteristics to *B. henryi* Hemsl. under the section *Reichenheimia*. It can, however, be distinguished by the following characteristics, including reddish linear red dots on petiole (vs. without), sparsely pubescent on petiole (vs. brown villous), slightly denticulate leaf margin (vs. crenate), glabrous leaves (vs. pubescent), leaf upper surface green (vs. green with dark green patches), lower surface




Figure 2. *Begonia hypoleuca* Y.H.Tan & H.B.Ding, sp. nov. **A** habitat **B** flowers (front view) **C** flowers (lateral view) **D** habit **E** inflorescence **F** leaves **G** tepals of staminate flower (back view) **H** stamens with peduncle **I** tepals of staminate flower (front view) **J** mature fruit with peduncle **K** mature fruit **L–N** serial cross sections of ovary. Photographed by H.B. Ding.

pale green (vs. pale green with reddish patches), male flower tepals broadly ovate (vs. oblate-orbicular), apex acute (vs. rounded), stamens number fewer 10–15 (vs. up to 30).

New records for Laos

Begonia augustinei Hemsley

Figure 3 A–E

Begonia augustinei Hemsley, Gard. Chron 3 (28): 286. 1900; T.C. Ku in T.L. Wu (ed.), Fl. Reipubl. Popularis Sin. 52(1): 256. 1999; S.H. Huang & Y.M. Shui in C.Y. Wu (ed.), Fl. Yunnan. 12: 213. 2006; T.C. Ku et al. in C.Y. Wu & P.H. Raven (eds), Fl. China 13: 163. 2007. Type: China, Yunnan Province, Simao, A. Henry 12333A (holotype: B100365120!).

Specimens examined. LAOS. Luang Namtha Province, Nam Ha National Biodiversity Conservation Area, Near Nalan Neua Village, 20°49'54.92"N, 101°19'24.85"E. 646 m elev., 13 October 2018, Y.H. Tan, B. Yang, H.B. Ding & X.D. Zeng L0687 (HITBC; HNL); Oudomxay Province, Maung Xai, Phou Hin Phee National Biodiversity Conservation Area, 20°43'30.86"N, 102°08'42.39"E. 1288 m elev., 25 October 2018, Y.H. Tan & H.B. Ding L0819 (HITBC; HNL); Oudomxay Province, Maung Xai, Phou Hin Phee National Biodiversity Conservation Area, 20°43'23.74"N, 102°08'42.84"E. 1224 m elev., 25 October 2018, Y.H. Tan & H.B. Ding L0822 (HITBC).

Distribution. China, Laos.

Note. This species is characterised by elongate rhizomes, extremely asymmetric leaves, base cordate to deeply cordate, margin irregularly serrulate (Hemsley 1900; Ku 1999; Huang and Shui 2006; Ku et al. 2007).

Begonia dryadis Irmscher

Figure 3 F–K

Begonia dryadis Irmscher, Not. Bot. Gard. Edinb. 21(1): 41. 1951; T.C. Ku in T.L. Wu (ed.), Fl. Reipubl. Popularis Sin. 52(1): 246. 1999; S.H. Huang & Y.M. Shui in C.Y. Wu (ed.), Fl. Yunnan. 12: 215. 2006; T.C. Ku et al. in C.Y. Wu & P.H. Raven (eds), Fl. China 13: 171. 2007. Type: China, Yunnan Province, Kuen-ger, Che-li Hsien, 1100 m elev., October 1936, C.W. Wang 79349 (holotype: KUN0370809).

Specimens examined. LAOS. Oudomxay Province, Maung Xai, Phou Hin Phee National Biodiversity Conservation Area, 20°43'18.52"N, 102°08'47.16"E. 1368 m elev.,




Figure 3. **A–E** *Begonia augustinei* Hemsley: **A** habitat **B** staminate flowers (front view) **C** leaf blade (adaxially and abaxially) **D** pistillate flowers **E** cross section of ovary **F–K** *Begonia dryadis* Irmscher: **F** habitat **G** staminate flowers (front view) **H** pistillate flower (front view) **I** pistillate flower (lateral view) **J** fruit **K** cross section of ovary. Photographed by H.B. Ding.

30 March 2018, Y.H. Tan, B. Yang, H.B. Ding & X.D. Zeng L0356 (HITBC; HNL); Oudomxay Province, Maung Xai, Phou Hin Phee National Biodiversity Conservation Area, 20°43'18.74"N, 102°08'47.84"E. 1364 m elev., 25 October 2018, Y.H. Tan & H.B. Ding L0815 (HITBC).

Distribution. Chian, Laos.

Note. This species was first discovered and reported in the forest understorey, by streams in a valley of South Yunnan, China. The discovery of this species in Laos shows the geographical linkage between two type localities. The species is differentiated from the allied species by its short, stout rhizome and the presence of caudine leaves. (Irmscher 1951; Ku 1999; Huang and Shui 2006; Ku et al. 2007).

Begonia lancangensis S.H. Huang

Figure 4 A–C

Begonia lancangensis S.H. Huang, Acta Bot. Yunnan. 21(1): 13. 1999; S.H. Huang & Y.M. Shui in C.Y. Wu (ed.), Fl. Yunnan. 12: 230. 2006; T.C. Ku et al. in C.Y. Wu & P.H. Raven (eds), Fl. China 13: 181. 2007. Type: China, Yunnan Province, Lancang Xian, Fazhan He, 1600 m elev., 1995, Huang Suhua 95001 (holotype: YNUU).

Specimens examined. Laos. Luang Namtha Province, Nam Ha National Biodiversity Conservation Area, Near Na Lun Village, 20°50'39.62"N, 101°19'41.46"E. 687 m elev., 23 March 2018, Y.H. Tan, B. Yang, H.B. Ding & X.D. Zeng L0055 (HITBC).

Distribution. China, Laos.

Note. The species is characterised by its dioecious and erect stems with ovate or ovate-oblong leaves and fleshy berry-like fruits (Shui and Huang 1999, Ku et al. 2007). Additionally, *B. handelii* Irmscher (Irmscher 1921: 24) and *B. acetosella* Craib (Craib 1912: 153) have been discovered in the same locality. We suspect that this species may be a natural hybrid species of them.

Begonia sizemoreae Kiew

Figure 4 D–F

Begonia sizemoreae Kiew, in Gard. Bull. Singapore 56: 95. 2004; Kiew R. in Adansonia sér. 3, 29(2): 234. 2007. Type: Vietnam, Ba Vi National Park, Ha Tay Province, ca. 80 km west of Hanoi, Accession No. 20020399 cult. in Singapore Botanic Gardens ex Palm Hammock Orchid Estate, Miami, U.S.A. R. Kiew 5304 (holotype: SING, isotype: HN).

Specimens examined. Laos. Phongsaly Province, Phou Dean Din National Bio-Diversity Park, 21°46'02.39"N, 102°31'28.58"E. 1111 m elev., 8 April 2018, Y.H. Tan, B. Yang, H.B. Ding & X.D. Zeng L0559 (HITBC).



Figure 4. **A–C** *Begonia lancangensis* S.H. Huang: **A** habitat **B** staminate flower (front view) **C** staminate flower (back view) **D–F** *Begonia sizemoreae* Kiew: **D** habitat **E** staminate flower **F** pistillate flower **G–I** *Begonia silletensis* (A. DC.) C.B. Clarke subsp. *mengyangensis* M.C. Tebbitt & K.-Y. Guan: **G** habitat **H** inflorescence **I** staminate flower (front view). Photographed by H.B. Ding and Y.H. Tan.

Distribution. Vietnam, Laos.

Note. *Begonia sizemoreae* was originally described in north Vietnam. Now, its distribution in Laos has been confirmed. This species is very similar to *B. rex* Putz. (Putzey 1857: 141) from Assam, India, in its leaf shape, fine variegated leaves and flower structure, but it is distinct in its leaf margin with very hairy upper surface of the lamina with hairs 5–10 mm long and its conspicuous deep crimson tertiary venation in the centre and outer part of the lower leaf surface (Kiew 2004, 2007).

***Begonia silletensis* subsp. *mengyangensis* M.C. Tebbitt & K.Y. Guan**

Figure 4 G–I

Begonia silletensis (A. DC.) C.B. Clarke subsp. *mengyangensis* M.C. Tebbitt & K.-Y. Guan, Novon 12(1): 134. 2002; T.C. Ku et al. in C.Y. Wu & P.H. Raven (eds), Fl. China 13: 198. 2007. Type: China, Yunnan Province, Xishuangbanna, on the way from Puwen to Mengyang, at the bottom of valley in a wet area on the slope facing N, dense forest, 21 April 1957, *Sino-Soviet Union expedition* 9633 (holotype KUN0533687!).

Specimens examined. Laos. Luang Namtha Province, Nam Ha National Biodiversity Conservation Area, Near Na Lun Village, 20°50'34.97"N, 101°20'05.69"E. 632 m elev., 23 March 2018, Y.H. Tan, B. Yang, H.B. Ding & X.D. Zeng L0095 (HITBC; HNL).

Distribution. India, China, Bangladesh, Myanmar and Laos.

Note. The species is characterised by its dioecious and prostrate stems with large broad leaves and fleshy berry-like fruits (Tebbitt and Guan 2002; Ku et al. 2007).

An updated checklist of *Begonia* species in Laos

Begonia sect. *Diploclinium*

Begonia adscendens C.B. Clarke

Laos: Champasak: J.K. Munzinger 250 (L, P).

Begonia binnamnoensis Souvann & Lanors

Laos: Khammouan: Lamxay et al. HNN227 (holotype HNL, isotypes FOF, E, SING).

Begonia khammouanensis Souvann. & Lamxay

Laos: Khammouan: Lamxay et al. HNN138 (holotype HNL, isotypes FOF, E, SING).

Begonia modestiflora Kurz

Laos: Bassac: Maxwell 97-1081 (L); Me-Kong: C. Threl 2239 (BM, P).

Begonia tatianae Aver.

Laos: Khammouan: *L. Averyanov*, AL 820b.1 (holotype LE01049623), *L. Averyanov et al.* AL820b (paratype LE01049002).

Begonia viscosa Aver. et H.Q. Nguyen

Laos: Vientiane: *L. Averyanov*, P.V. The, CPC2438 (holotype LE, isotype CPC).

Begonia sect. *Jackia*

Begonia cladotricha M.Hughes

Laos: Khammouan: *M.F. Newman et al.* LAO985 (holotype E).

Begonia sect. *Parvibegonia*

Begonia integrifolia Dalzell

Laos: Champasak: *J.F. Maxwell* 97-1150 (CMU, L), *C. Thorel* 2226 (P[2]); *C. Thorel* 2226A(P[2]).

Begonia laotica Y.H. Tan & H.B. Ding

Laos: Oudomxay: *Y.H. Tan, H.B. Ding* L0827 (holotype HITBC!, isotypes HNL!).

Begonia namkadingensis C.-J. Yang, Souladeth & Tagane

Laos: Bolikhhamxay: *Tagane S. et al.* L1202 (holotype FOF, isotypes HAST, HNL, KYO), *Tagane S. et al.* L960 (FOF, KYO, TAI).

Begonia procridifolia Wall. ex A.DC.

Laos: *A.F.G. Kerr* 21188 (K, P).

Begonia sect. *Petermannia*

Begonia lamxayiana Souvann

Laos: Bolikhhamxay: *V. Lamxay et al.* VL2198 (holotype HNL).

Begonia sect. *Platycentrum*

Begonia acetosella Craib

Laos: *E. Poilane* 20721 (P); Luang Namtha: *Tan et al.* L0050 (HITBC!, HNL!), *Tan et al.* L0083 (HITBC!); Oudomxay: *Tan et al.* L0202 (HITBC!, HNL!).

Begonia augustinei Hemsl

Laos: Luang Namtha: *Tan et al.* L0687 (HITBC; HNL); Oudomxay: *Tan et al.* L0819 (HITBC; HNL), *Tan et al.* L0822 (HITBC).

Begonia dryadis Irmsch

Laos: Oudomxay: *Tan et al.* L0356 (HITBC; HNL), *Tan et al.* L0815 (HITBC).

Begonia handelii var. ***handelii*** Irmsch.

Laos: Luang Namtha: *L.J. Ahnby* 129 (E), *Tan et al.* L0061 (HITBC; HNL); Phongsaly: *Tan et al.* L0502 (HITBC).

Begonia handelii var. ***prostrata*** (Irmsch.) Tebbitt

Laos: Xiangkhoang: *A.F.G. Kerr* 21772 (K).

Begonia lancangensis S.H. Huang

Laos: Luang Namtha: *Tan et al.* L0055 (HITBC).

Begonia palmata D. Don

Laos: Houaphan: *E. Poilane* 1938 (P), *E. Poilane* 2002 (P); Khammouan: *M.F. Newman* LAO1429 (E); Xiangkhoang: *J. Delacour* (P).

Begonia quadripetiolata Aver. et H. Q. Nguyen

Laos: Vientiane: *L. Averyanov et al.* LA-VNI510/1 (holotype LE01049395, isotype LE01049479).

Begonia sizemoreae Kiew

Laos: Phongsaly: *Tan et al.* L0559 (HITBC).

Begonia siamensis Gagnep.

Laos: Attapu: *F.J. Harmand* 1387 (P); Phongsaly: *Tan et al.* L0476 (HITBC; HNL), *Tan et al.* L0416 (HITBC).

Begonia silleensis subsp. ***mengyangensis*** M.C. Tebbitt & K.Y. Guan

Laos: Luang Namtha: *Tan et al.* L0095 (HITBC; HNL).

Begonia sect. *Reichengeimia****Begonia hymenophylla*** Gagnep.

Laos: Champasak: *C. Thorel* 2358 (P), *C. Thorel* 2958 (P).

Begonia hypoleuca Y.H. Tan & H.B. Ding

Laos: Luang Namtha: *Tan et al.* L0792 (holotype HITBC!).

Begonis sect. *Tetraphila****Begonia afromigrata*** J.J. de Wilde

Laos: Muang Awn: *Kerr* 20938 (BM, K, L); Vientiane: *Rodda*, *Simonsson* MR106 (TO), *Rodda*, *Simonsson* MR107 (FI).

Acknowledgements

The authors are grateful to the Biotechnology and Ecology Institute (BEI) of Lao PDR for permission to conduct this study in Northern Lao PDR and for their support and collaboration. We are grateful to Dr. Liang Song, Dr. Hua-Zheng Lu, Dr. Su Li, Mr. Ren Li and Mr. Xiao-Dong Zeng for their kind help in the field. This work was financially supported by a project of the Lancang-Mekong Cooperation (LMC) Special Fund (Biodiversity Monitoring and Network Construction along Lancang-Mekong River Basin project) and the CAS 135 programme (No. 2017XTBG-F03), CAS “the Belt and Road” Master Fellowship Program and CAS President’s International Fellowship Initiative (PIFI) and the project of the Southeast Asia biodiversity research institute, Chinese Academy of Sciences (Y4ZK111B01).

References

- Averyanov LV, Nguyen QH (2012) Eleven new species of *Begonia* L. (Begoniaceae) from Laos and Vietnam. *Turczaninowia* 15(2): 5–32.
- Averyanov LV, Nguyen KS, Maisak TV, Nguyen HT, Nguyen HQ (2019) Two new species of *Begonia* L. (Begoniaceae) from central Laos. *Turczaninowia* 22(2): 134–141. <https://doi.org/10.14258/turczaninowia.22.2.9>
- Camfield R, Hughes M (2018) A revision and one new species of *Begonia* L. (Begoniaceae, Cucurbitales) in Northeast India. *European Journal of Taxonomy* 396: 1–116. <https://doi.org/10.5852/ejt.2018.396>
- Craig WG (1912) Contributions to the Flora of Siam. *Bulletin of Miscellaneous Information, Kew* 1912(3): 144–154. <https://doi.org/10.2307/4104569>

- De Candolle A (1859) Mémoire sur la famille des Bégoniacées. Annales des Science Naturelles. Botanique 4(11): 92–149.
- De Candolle A (1864) Prodromus systematis Naturalis Regni Vegetabilis (Vol. 15(1)). Sumptibus Victoris Masson et filii, Parisiis [Paris]. <https://doi.org/10.5962/bhl.title.286>
- De Wilde JJFE, Hughes M, Rodda M, Thomas DC (2011) Pliocene intercontinental dispersal from Africa to Southeast Asia highlighted by the new species *Begonia afromigrata* (Begoniaceae). Taxon 60(6): 1685–1692. <https://doi.org/10.1002/tax.606013>
- Doorenbos J, Sosef MSM, De Wilde JJFE (1998) The sections of *Begonia* including descriptions, keys, and species lists (Studies in Begoniaceae VI). Wageningen Agricultural University Papers 98: 1–266.
- Hemsley WB (1887) Begoniaceae. In: Forbes FB, Hemsley WB (Eds) An enumeration of all the plants known from China proper, Formosa, Hainan, Corea, the Luchu Archipelago, and the island of Hongkong, together with their distribution and synonymy. The Journal of the Linnean Society, Botany 23: 321–323. <https://doi.org/10.1111/j.1095-8339.1887.tb00532.x>
- Hemsley A (1900) New or noteworthy plants. The Gardeners' Chronicle, London 2: 1–286.
- Huang SH, Shui YM (2006) Begoniaceae. In: Wu CY (Ed.) Flora of Yunnanica (Vol. 12). Science Press, Beijing, 143–237.
- Hughes M (2007) *Begonia cladotricha* (Begoniaceae): A new species from Laos. Edinburgh Journal of Botany 64(1): 101–105. <https://doi.org/10.1017/S0960428607000777>
- Hughes M (2008) An Annotated Checklist of Southeast Asian *Begonia*. Royal Botanic Garden, Edinburgh, 176 pp.
- Hughes M, Moonlight P, Jara A, Tebbitt M, Wilson H, Pullan M (2015–Present) Begonia Resource Centre. <http://padme.rbge.org.uk/begonia/> [accessed 26 July 2019]
- Irmscher E (1921) Plantae novae sinenses, diagnosibus brevibus descriptae a Dr. Henry Han-del-Mazzetti. Anzeiger der Akademie der Wissenschaften in Wien. Mathematische-naturwissenschaftliche Klasse 58: 24–27.
- Irmscher E (1951) Some new Chinese species of *Begonia*. Notes from the Royal Botanic Garden Edinburgh 21(1): 35–45.
- IUCN Petitions Subcommittee (2017) Guidelines for Using The IUCN Red List Categories and Criteria. version 13. Prepared by the Standards and Petitions Subcommittee. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Kiew R (2004) *Begonia sizemoreae* Kiew (Begoniaceae), a handsome new *Begonia* from Vietnam. Gardens' Bulletin (Singapore) 56: 95–100.
- Kiew R (2007) Notes on Vietnamese Begonia (Begoniaceae), including three new species. Adansonia sér. 3 29(2): 229–238.
- Klotzsch JF (1855) Begoniaceen – Gattungen und Arten. Abh. Königl. Akad. Wiss. Berlin 1854: 121–255.
- Ku TC (1999) Begoniaceae. In: Ku TC (Ed.) Flora Reipublicae Popularis Sinica (Vol. 52(1)). Science Press, Beijing, 126–269.
- Ku TC, Peng CI, Turland NJ (2007) Begoniaceae. In: Wu CY, Raven PH (Eds) Flora of China (Vol. 13). Science Press & Missouri Botanical Garden, Beijing & St. Louis, Missouri, 153–207.
- Kurz S (1873) New Burmese Plants. Journal of the Asiatic Society of Bengal, part 2, Natural History, Calcutta, 59–110.

- Linnaeus C (1753) Species Plantarum 2. Laurentius Salvius, Stockholm, 1056 pp.
- Moonlight PW, Ardi WH, Padilla LA, Chung KF, Fuller D, Girmansyah D, Hollands R, Jara-Munoz A, Kiew R, Leong WC, Liu Y, Mahardika A, Marasinghe LDK, O'Connor M, Peng C-I, Perez AJ, Phutthai T, Pullan M, Rajbhandary S, Reynel C, Rubite RR, Sang J, Scherberich D, Shui YM, Tebbitt MC, Thomas DC, Wilson HP, Zaini NH, Hughes M (2018) Dividing and conquering the fastest-growing genus: Towards a natural sectional classification of the mega-diverse genus *Begonia* (Begoniaceae). *Taxon* 67(2): 267–323. <https://doi.org/10.12705/672.3>
- Newman MF, Ketphanh S, Svengsuksa B, Thomas P, Sengdala K, Lamxay V, Armstrong K (2007) A Checklist of the Vascular Plants of Lao PDR. Royal Botanic Garden, Edinburgh, 73–74.
- Putzey JAAH (1857) *Begonia rex*. In: Van Houtte L. Flores des Serres et des Jardins de l'Europe 12: 141–146.
- Shui YM, Huang SH (1999) Notes on the genus *Begonia* from Yunnan. *Yunnan Zhi Wu Yan Jiu* 21(1): 11–23.
- Souvannakhounmane K, Hughes M, Lanorsavanh S (2016) *Begonia lamxayiana* Souvann. (Begoniaceae): A new species from Lao PDR. *Thai Journal of Botany* 8(1): 1–5.
- Souvannakhounmane K, Lanorsavanh S, Lamxay V (2018) Two new species of *Begonia* L. (Begoniaceae) from central Laos. *Taiwania* 63(3): 188–194. <https://doi.org/10.6165/tai.2018.63.188>
- Tebbitt MC, Guan K-Y (2002) Emended circumscription of *Begonia silletensis* (Begoniaceae) and description of a new subspecies from Yunnan, China. *Novon* 12(1): 133–136. <https://doi.org/10.2307/3393252>
- Tian DK, Xiao Y, Tong Y, Fu NF, Liu QQ, Li C (2018) Diversity and conservation of Chinese wild begonias. *Plant Diversity* 40(3): 75–90. <https://doi.org/10.1016/j.pld.2018.06.002>
- Yang CJ, Tagane S, Souladeth P, Okabe N, Hu JM, Yahara T (2018) Flora of Nam Kading National Protected Area III: *Begonia namkadingensis* (Begoniaceae), a new species in limestone area. *Phytotaxa* 334(2): 195–199. <https://doi.org/10.11646/phytotaxa.334.2.8>