

A revision of infrageneric classification in *Astelia* Banks & Sol. ex R.Br. (Asteliaceae)

Joanne L. Birch^{1,2}

1 Department of Botany, University of Hawaii at Manoa, Honolulu, HI, 96822, USA **2** Present address: Royal Botanic Gardens Victoria, Birdwood Avenue, Melbourne, Victoria 3004, Australia

Corresponding author: Joanne L. Birch (joanne.birch@rbg.vic.gov.au)

Academic editor: L. Peruzzi | Received 27 February 2015 | Accepted 28 May 2015 | Published 13 July 2015

Citation: Birch JL (2015) A revision of infrageneric classification in *Astelia* Banks & Sol. ex R.Br. (Asteliaceae). *PhytoKeys* 52: 105–132. doi: 10.3897/phytokeys.52.4768

Abstract

Systematic investigations and phylogenetic analyses have indicated that *Astelia*, as currently circumscribed, is paraphyletic, with *Colospermum* nested within it. Further, *A.* subgenus *Astelia* is polyphyletic, and *A.* subgenera *Asteliopsis* and *Tricella* are paraphyletic, as currently circumscribed. Revision of the subgeneric classification of *Astelia* is warranted to ensure classification accurately reflects the evolutionary history of these taxa. *Colospermum* is relegated to synonymy within *Astelia*. *Astelia* is dioecious or polygamodioecious, with a superior ovary, anthers dorsi- or basifix, pistillodes or pistils that have a single short or poorly defined style, a 3 lobed stigma, and fleshy uni- or trilocular fruit with funicular hairs that are poorly to well developed. *Astelia* subgenus *Colospermum* (Skottsb.) Birch is described. A key to *Astelia* sections is provided. *Astelia hastata* Colenso, *A. montana* Seem., and *A. microsperma* Colenso *pro parte* are resurrected and the new combination *A. samoense* (Skottsb.) Birch, **comb. nov.** is made.

Keywords

Asparagales, Asteliaceae, Austral, Australia, dioecy, New Zealand, Pacific

Introduction

Astelia Banks & Sol. ex R.Br. is the largest genus in Asteliaceae Dumort., containing twenty-six species and three non-nominotypical varieties with an Austral-Pacific distribution. *Astelia* species exhibit a range of growth forms including low, cushion-forming and tall, clustered habits. *Astelia* species grow in a diverse range of habitats including

coastal, lowland wetlands, tropical and temperate lowland forests, tropical montane cloud forests, sub-alpine heath, alpine fellfields and grasslands, and bog habitats. Many *Astelia* species are facultatively epiphytic and three species are primarily epiphytic. *Collospermum* Skottsb. includes four species that occur in lowland forests and in lowland and tropical montane cloud forests in New Zealand, the Independent State of Samoa and the Republics of Fiji and Vanuatu. All species exhibit a tall, clustered habit and are primarily epiphytic, although plants that fall to the ground can persist for long periods.

Astelia and *Collospermum* share many morphological characters (Bayer et al. 1998, Rudall et al. 1998). All Asteliaceae genera have branched hairs that are otherwise uncommon in the Lilianae (Bayer et al. 1998) and those of *Astelia* and *Collospermum* are dense, at least in the young leaves of most species. The tomentum of *Astelia* and *Collospermum* consist of a multi-celled stalk, frayed unicellular filaments that form a basal wool adjacent to the stalk, and linear or ovate scales that are a single cell thick (McCarthy 1928, Skottsberg 1934, L. B. Moore 1980, Rudall et al. 1998). In *Astelia* the scales are linear, may be short or long, and the stalk is attached at the base of the scale. In *Collospermum*, the scales are ovate, short, and the stalk is peltate. Scales may be present on the adaxial and/or abaxial leaf surfaces, inflorescences, and flowers. On the leaves of *Astelia* species the scales may become fused to form a membranous pellicle over the leaf epidermis. *Astelia* and *Collospermum* have superior ovaries that are uni- or trilocular and trilocular, respectively. Published chromosome numbers for *Astelia* taxa range from $2n = 60$ to $2n = 210$ (Wheeler 1966, Dawson and Beuzenberg 2000, de Lange et al. 2004). Darlington and Wylie (1955) proposed a basic chromosome number for *Astelia* of $x = 8$, but Wheeler (1966) considered the basic chromosome number to be $x = 5, 7$, or 35 . *Collospermum hastatum* and *C. microspermum* are the only *Collospermum* taxa for which chromosome numbers are known and both have a chromosome number of $n = 35$.

Skottsberg (1934) segregated *Collospermum* from *Astelia* based on the presence of simple lateral racemes, dimorphism of staminate and pistillate plants, basifixated anthers, long style papillae, and mucilaginous seed hairs of the former, which differ from the predominantly paniculate lateral racemes, versatile anthers, and poorly developed mucilaginous seed hairs of the latter (Birch unpublished PhD thesis 2011, Birch et al. 2012). However, morphological (Bayer et al. 1998, Birch unpublished PhD thesis 2011), cytological (Wheeler 1966, Moore 1980), and molecular (Birch unpublished PhD thesis 2011, Birch et al. 2012) data suggested a close evolutionary relationship between these genera. Moore (1980) considered that a re-evaluation of the circumscriptions of *Astelia* and *Collospermum* was warranted due to the production of viable progeny from intergeneric crosses. *Collospermum*, while monophyletic, was nested within *Astelia* in phylogenetic reconstructions based on combined chloroplast and nuclear sequence data applying Bayesian inference, maximum likelihood, and maximum parsimony criteria (Birch unpublished PhD thesis 2011, Birch et al. 2012). As a result, those authors recognized a broadly circumscribed *Astelia* s.l., including *Collospermum*.

Within *Astelia*, Skottsberg (1934) recognized three subgenera (*A.* subg. *Astelia* Skottsb., *A.* subg. *Asteliopsis* Skottsb., and *A.* subg. *Tricella* Skottsb.) based on open or cushion-forming growth form, degree of tepal fusion, ovary division, and seed shape.

Within these subgenera, he recognized seven sections (*A.* sect. *Astelia* Skottsb., *A.* sect. *Desmoneuron* Skottsb., *A.* sect. *Isoneuron* Skottsb., *A.* sect. *Micrastelia* Skottsb., *A.* sect. *Palaeastelia* Skottsb., *A.* sect. *Periastelia* Skottsb., and *A.* sect. *Tricella* Skottsb.) based on leaf venation, pistillode size, seed surface features, and extent of funicle development. In phylogenetic analyses (Birch unpublished PhD thesis 2011, Birch et al. 2012), each of Skottsberg's (1934) sections, except *A.* sect. *Tricella*, were monophyletic, *A.* subg. *Astelia* and *A.* subg. *Asteliopsis* were polyphyletic, and *A.* subg. *Tricella* was paraphyletic. A revised circumscription of *Astelia* subgenera is proposed that accurately reflects the evolutionary relationships within the genus. *Collospermum* is relegated to synonymy under *Astelia*. Skottsberg's sections are retained as they are monophyletic and accurately capture the extensive morphological diversity that is present within the subgenera.

A revision based on recognition of monophyletic taxa is proposed here. Multiple characters support the proposed circumscription of *Astelia*. All taxa are dioecious or polygamodioecious, with a superior ovary, dorsi- or basifixed anthers, pistillodes or pistils that have a single short or poorly defined style, a 3 lobed stigma, and fleshy uni- or trilocular fruit with funicular hairs that are poorly to well developed.

Methods

Taxonomic sampling

All *Astelia* taxa, (twenty-six species and three non-nominotypical varieties) and all *Collospermum* (four species) were included in this study. Herbarium specimens were examined from the following herbaria: Auckland War Memorial Museum (AK), Herbarium Pacificum (BISH), Allan Herbarium (CHR), Harvard University (GH), Kew Royal Botanic Gardens (K), National Herbarium of Victoria (MEL), Missouri Botanical Garden (MO), Herbier National de Paris (P), National Tropical Botanical Garden (PTBG), United States National Herbarium (US), and Museum of New Zealand Te Papa Tongarewa (WELT). Type specimens were examined from AK, BISH, MEL, P, WELT and digital images of type specimens were examined from CHR and K (Herbarium abbreviations follow Index Herbariorum (Thiers continuously updated)).

Morphological data and analyses

Morphological data were obtained for 410 herbarium specimens (Appendix 1). Data were obtained for ten specimens per species, including five staminate and five pistillate specimens for species with unisexual flowers. Measurements and scores were averaged across all specimens to give a mean value for each taxon. Flower and fruit color data were obtained from multiple sources including field observations, specimen label data, and taxon descriptions in national floras (Drake del Castillo 1893, L. B. Moore and Edgar 1976, Coode 1978, D. M. Moore 1983, Williams 1987, Wagner et al. 1999).




Figure 1. Scanning electron micrographs of *Astelia* seeds showing size, shape, and funicle characters. **A** *Astelia* (sect. *Astelia*) *linearis* var. *linearis*; ovoid, funicle long **B** *Astelia* (sect. *Tricella*) *petriei*; ovoid, funicle short **C** *Astelia* (sect. *Tricella*) *chathamica*; ovoid, funicle short **D** *Astelia* (sect. *Isoneuron*) *banksii*; polygonal-turbinate, funicle ribbed **E** *Astelia* (sect. *Micrastelia*) *pumila*; ovoid, funicle short **F** *Astelia* (sect. *Desmoneuron*) *solandri*; obovoid-reniform, funicle ribbed **G** *Astelia* (sect. *Isoneuron*) *neocaledonica*; obovoid, funicle ribbed **H** *Astelia* (sect. *Tricella*) *menziesiana*; fusiform, funicle short. Scale bars = 0.5 mm. SEM images created by J.L. Birch.

Morphological characters that varied at or below the genus rank were measured or scored for all *Astelia* and *Collospermum* taxa in the field and/or herbarium. Herbarium specimens were studied under a dissecting microscope and measurements obtained using digital calipers. Pollen and seed characters were examined directly from material obtained from herbarium specimens after coating with gold-palladium using a Hitachi S-4800 field emission scanning electron microscope (SEM) at the Biological Electron Microscope Facility, Pacific Biosciences Research Center of the University of Hawai‘i at Mānoa. Images were digitally processed and the final plates were prepared in Photoshop 10.0.

Taxonomic treatment

Astelia Banks & Sol. ex R.Br.

Astelia Banks & Sol. ex R.Br., Prodr. 291. 1810. *nomen conservandum* (International Botanical Congress and JH Wiersema 2015). Type: *Astelia alpina* R.Br.

Funckia Willd., Mag. Neuesten Entdeck. Gesammten Naturk. Ges. Naturf. Freunde Berlin 2: 19. 1808, *nomen rejiciendum*. Type: *Funckia magellanica* Willdenow, *nomen illegitimum* (*Melanthium pumilum* G. Forster)

Hamelinia A.Rich., Voy. Astrolabe 1: 158. 1832. Type: *Hamelinia veratroides* A.Rich.

Note. Herbaceous perennials, terrestrial or epiphytic, often growing in clusters with three ramets in trigonal arrangement, some species turf-forming, rhizomatous, dioecious or polygamodioecious. Leaves: 3-ranked, linear, ensiform, or subulate; leaves usually keeled, margins erect or revolute; leaf sheath closed, with surface obscured by dense long white hairs; parallel venation, variously incrassate; tomentum composed of scales and lanate wool at base of scale stalk, scales with basal stalk or peltate. Inflorescence: a terminal panicle, sometimes reduced to a few flowers; lateral branches racemes or sub-panicles, subtended by foliaceous or membranous, linear or lanceolate spathes; peduncle tomentum composed of distinct, narrow scales with dense basal wool. Flowers: pedicillate; bracts membranous, linear or spatulate; perianth membranous or fleshy, 6 tepals in 2 series; connate at base into tube of variable length; outer tepals triangular to lanceolate, with three veins, scales present over entire surface; inner tepals linear with one midvein, scales present along midvein only. Staminate flowers: lobes recurved; stamens 6; filaments filamentous, adnate to tepals at base of tepal lobes; anthers elliptic or linear-hastate, dorsifixed and versatile or basifixed and immobile, latrorse; pistillode present, style undifferentiated or distinct; stigma not formed. Pistillate flowers: 6 reduced staminodes present, adnate to base of tepal lobes, filament filamentous, anthers flattened, sterile; ovary superior, uni- or trilocular, placentation parietal from three placentas or axile, with subapical placentas, ovules few to many; style distinct or undifferentiated, stigmas 3. Fruit: berry, stigma typically persistent. Seeds: black, ovoid, ellipsoid, fusiform, or polygonal; testa smooth or sculptured; funicle with mucilaginous funicular hairs poorly or well developed, funicle hairs surrounding the seeds and either adhering to the testa or not.

Astelia* Banks & Sol. ex R.Br. subg. *Astelia

Note. Flowers with a very short perianth tube (0.1–0.8 mm); anthers dorsifixed, versatile; ovary unilocular; seeds ovoid; funicle long, curved, with mucilaginous funicular hairs poorly developed that surround but do not adhere to the seed.

Remarks. Molecular (Birch et al. 2012) and morphology-based phylogenetic analyses (Birch unpublished PhD thesis 2011) indicate that *A. subg. Astelia*, as circumscribed by Skottsberg (1934), is polyphyletic. *Astelia* subg. *Astelia* is revised to include *A. sect. Palaeastelia* and *A. sect. Astelia*, which form a clade. *Astelia* sect. *Desmoneuron* is placed in a different clade and it is excluded from *A. subg. Astelia*.

Astelia* sect. *Astelia

Note. Low, compact, growth form (including cushion or turf forming taxa); leaves, linear, ensiform, or subulate; reduced inflorescences bearing few flowers; staminate flowers with short filaments (0.5–1.6 mm); pistillate flowers with long outer tepals (4.5–7.0 mm); ovary unilocular, long (4.3–8.3 mm); fruit ovoid or oblong; few or many seeds per fruit (< 25), seeds short and narrow (1.1–2.0 × 0.5–1.3 mm).

Included taxa and distribution. Australia *A. alpina* R.Br. var. *alpina*, *A. alpina* var. *novae-hollandiae* Skottsb. Indonesia (Papua Province), Papua New Guinea *A. papuana* Skottsb.. New Zealand *A. linearis* Hook.f. var. *linearis*, *A. linearis* var. *novae-zelandiae* Skottsb., *A. subulata* Cheeseman

Habitat. Lowland (low latitudes) to sub-alpine (mid and higher latitudes) herb-fields particularly on wet substrates (seeps, swamps etc.).

***Astelia* sect. *Palaeastelia* Skottsb.**

Astelia sect. *Palaeastelia* Skottsb., Kongl. Svenska Vetenskapsakad. Handl. ser. 3, 14(2): 24. 1934.

Note. Open growth form; leaves linear; large inflorescences bearing many flowers; staminate flowers with intermediate length filaments (1.7–4.6 mm); pistillate flowers with short outer tepals (2.3–4.4 mm); ovary unilocular, intermediate length (3.2–4.2 mm); fruit ovoid, many seeds per fruit (< 20); seed intermediate length and narrow (2.4–2.9 × 1.2–1.3 mm). Type: *A. hemichrysa* (Lam.) Kunth.

Included species and distribution. Mascarene Islands, Réunion Island. *A. hemichrysa* (Lam.) Kunth

Habitat. Tropical forest.

***Astelia* subg. *Asteliopsis* Skottsb.**

Astelia subg. *Asteliopsis* Skottsb., Kongl. Svenska Vetenskapsakad. Handl. ser. 3, 14(2): 46. 1934.

Note. Flowers with a short perianth tube (0.2–1.1 mm); anthers dorsifixed, versatile; ovary uni- (*A. sect. Desmoneuron*) or trilocular (*A. sect. Isoneuron*); seeds obovoid, obovoid-reniform, or turbinate-polygonal; funicle ribbed, mucilaginous funicular hairs well developed that surround but do not adhere to the seed; Type: *A. trinervia* Kirk, designated here.

Remarks. Phylogenetic analyses indicate that *A. subg. Asteliopsis*, as circumscribed by Skottsberg (1934), is polyphyletic (Birch unpublished PhD thesis 2011, Birch et al. 2012). *Astelia* subg. *Asteliopsis* is revised to include *A. sect. Isoneuron* and *A. sect. Desmoneuron*, which form a clade. But, as *A. sect. Periastelia*, is placed in a different clade, it is excluded from *A. subg. Asteliopsis*.

***Astelia* sect. *Desmoneuron* Skottsb.**

Astelia sect. *Desmoneuron* Skottsb., Kongl. Svenska Vetenskapsakad. Handl. ser. 3, 14(2): 34. 1934.

Note. Open growth form; leaves linear with a group of three subequal lateral nerves conspicuous in lower half; inflorescences bearing many flowers; staminate flowers with short filaments (0.6–2.1 mm); pistillate flowers with short outer tepals (1.9–4.2 × 0.7–1.8 mm); ovary unilocular, short (1.3–2.8 mm); fruit ampulliform, many seeds per fruit (8–32); seeds short and narrow (1.4–1.8 × 0.5–1.0 mm). Type: *A. trinervia* Kirk, designated here.

Included species and distribution. New Zealand: *A. solandri* A.Cunn., *A. trinervia* Kirk. Society Islands, Tahiti: *A. nadeaudii* Drake & F.Br.

Habitat. Temperate forest (*A. solandri* and *A. trinervia*) and tropical montane cloud forest (*A. nadeaudii*).

***Astelia* sect. *Isoneuron* Skottsb.**

Astelia sect. *Isoneuron* Skottsb., Kongl. Svenska Vetenskapsakad. Handl. ser. 3, 14(2): 51. 1934.

Note. Open growth form; leaves linear; inflorescences bearing many flowers; staminate flowers with short filaments (0.6–1.9 mm); pistillate flowers with short outer tepals (2.7–4.0 × 0.7–2.0 mm); ovary trilocular, intermediate length (2.2–4.4 mm); fruit ovoid, many seeds per fruit (11–18); seeds small and narrow (1.3–2.2 × 0.8–1.4 mm). Type: *A. banksii* A.Cunn., designated here.

Included species and distribution. New Caledonia: *A. banksii* A.Cunn., *A. neocaldonica* Schltr.

Habitat. Lowland coastal cliffs (*A. banksii*) and lowland tropical forest (*A. neocaldonica*).

***Astelia* subg. *Collospermum* (Skottsb.) Birch, stat. nov.**

urn:lsid:ipni.org:names:77148154-1

Basionym: *Collospermum* Skottsb., Kongl. Svenska Vetensk. Acad. Handl. Ser. 3, 14(2): 72. 1934. Type: *Collospermum hastatum* (Colenso) Skottsb. [Lectotypified by Moore and Edgar (1976)].

Note. Open growth form; leaves linear or ensiform with dark coloration at base and peltate branched hairs; inflorescences bearing many flowers; flowers with a long perianth tube (1.3–3.1 mm); staminate flowers with long filaments (3.8–8.6 mm), anthers basifixated, immobile; pistillate flowers with long outer tepals (1.5–6.2 × 0.9–1.6 mm); ovary trilocular, intermediate length (2.1–4.8 mm); fruit globose or obpyriform, with few to many seeds per fruit (1–22); seeds ellipsoid or ovoid, small and narrow (1.3–2.1 × 0.1–1.1 mm); funicle short, truncate, with well-developed mucilaginous funicular hairs that adhere to the seed.

Remarks. Species published under *Collospermum* must be transferred as *Astelia* has nomenclatural priority. Synapomorphies recognized for the genus *Collospermum* (Skottsberg 1934) remain valid for *A. subg. Collospermum*.

Included species and distribution. New Zealand: *A. hastata* Colenso, *A. microsperma* Colenso *pro parte*. Republic of Fiji, Viti Levu, Vanua Levu, Kandavu; Republic of Vanuatu, Espiritu Santo, Tanna, Aneityum; *A. montana* Seem. Independent State of Samoa, Savai'i, Upolu: *A. samoense* (Skottsb.) Birch.

Habitat. Lowland temperate forest (*A. hastata* and *A. microsperma*) and montane tropical cloud forest (*A. montana* and *A. samoense*).

***Astelia hastata* Colenso**

Astelia hastata Colenso, Trans. & Proc. New Zealand Inst. 19. 265. 1887.

Funckia hastata Kuntze, Revis. Gen. Pl. 2: 711. 1891, *nomen rejiciendum*.

Astelia furfuracea Banks et Solander MSS, *fide* C. Skottsberg, Kongl. Svenska Vetensk. Acad. Handl. Ser. 3, 14(2): 77. 1934.

Collospermum hastatum (Colenso) Skottsb., Kongl. Svenska Vetensk. Acad. Handl. Ser. 3, 14(2): 77. 1934; based on *A. hastata* Colenso.

Type. NEW ZEALAND. North Island. Hilly country north of Napier, County of Wairoa. January 1886, *A. Hamilton s.n.* (Lectotype: K [000524883, digital image!], staminate, designated by Skottsberg, 1934, 79; Isolectotypes: K [000524884, digital image!], staminate, pistillate; AK [3191!], staminate, pistillate).

***Astelia microsperma* Colenso pro parte**

Astelia microsperma Colenso pro parte, Trans. & Proc. New Zealand Inst. 17: 251. 1885 (description of fruit only). Type: NEW ZEALAND. North Island. Seventy-mile Bush, between Norsewood and Danniverke, County of Waipawa. 1884, W. Colenso s.n. (Lectotype: K [000524879, digital image!], fruiting material in packet, designated by Skottsberg, 1934, 81).

Collospermum microspermum Skottsb., Kongl. Svenska Vetensk. Acad. Handl. Ser. 3, 14(2): 82. 1934; based on *A. microsperma* Colenso.

Funckia microsperma Kuntze, Revis. Gen. Pl. 2: 711. 1891, *nomen rejiciendum*.

Astelia albicans Colenso, Trans. & Proc. New Zealand Inst. 17: 252. 1885.

Type: NEW ZEALAND. North Island. East slopes of Ruahine mountain range, county of Waipawa. Jan. 1884, *A. Hamilton* s.n. (Syntype: K [000524880, digital image!], staminate, pistillate). Skottsberg (1934, 88) identified a single specimen at K as the type material and the specimen in his plate 20 (K000524880) is consistent with the specimen described as the type. As *Astelia* is dioecious, the inclusion of one staminate and one pistillate inflorescence on the specimen means that it represents two distinct collections and each is rendered a syntype.

Funckia albicans Kuntze, Revis. Gen. Pl. 2: 711. 1891, *nomen rejiciendum*.

Astelia graminifolia Colenso, Trans. & Proc. New Zealand Inst. 19: 267. 1887.

Type: NEW ZEALAND. North Island. Woods, hilly country north of Napier, County of Wairoa, 1886, *A. Hamilton* s.n. (Lectotype: K [000524881, digital image!], pistillate, designated by Skottsberg, 1934, 85).

Funckia graminifolia Kuntze, Revis. Gen. Pl. 2: 711. 1891, *nomen rejiciendum*.

Astelia planifolia Colenso Trans. & Proc. New Zealand Inst. 20: 209-210. 1888.

Type: NEW ZEALAND. North Island, Pohue, hilly country west of Napier, Hawke's Bay. 1884. *A. Hamilton* s.n.; no specimens located.

Remarks. *Astelia microsperma* was described by Colenso based on a specimen at Kew that contained material from two species (Skottsberg 1934). The species description “referred to both, but mainly to the pistillate raceme in the envelope” (Skottsberg 1934, 82). Skottsberg (1934) lectotypified the fruiting material in the packet (K000524879) as the type material of *A. microsperma* (as syn. *C. microspermum*) and determined the remaining material on the specimen (K000524882) as *A. hastata* (as syn. *C. hastatum*).

Excluded species. *Astelia spicata* Colenso, Trans. & Proc. New Zealand Inst. 14: 335. 1882, *nomen illegitimum*. Type: NEW ZEALAND. North Island. In the forests about Kopua and Norsewood, Colenso. (Lectotype: K [000524878, digital image!], pistillate, designated by Skottsberg, 1934, 81). Moore (1966) regarded the type and other specimens examined to represent very small individuals of either *A. hastata* or *A. microsperma*.

Collospermum spicatum (Colenso) Skottsb., Kongl. Svenska Vetensk. Acad. Handl. Ser. 3, 14(2): 80. 1934; based on *A. spicata* Colenso *nomen illegitimum*.

Astelia nana Carse, Trans. & Proc. New Zealand Inst. 57: 91. 1926, *nomen illegitimum*.
Type: NEW ZEALAND. North Island. Kaiaka (Mangonui County), Maungatapere (Whangarei County), Mauku (Franklin County), *H. Carse s.n.*; synonym of *A. spicata* Colenso *nomen illegitimum*. This is regarded by Skottsberg (1934) as a synonym of *Astelia spicata* Colenso *nomen illegitimum*. (Syntypes: CHR [328212, digital image!], staminate, pistillate; [328213, digital image!], pistillate; AK [3227!], pistillate; [3228!], pistillate; [222913!], staminate; [303282!], pistillate).

Funckia spicata Kuntze, Revis. Gen. Pl. 2: 711. 1891, *nomen rejiciendum*.

Astelia montana Seem.

Astelia montana Seem., Fl. Vit. [Seemann] 313, figs 1–6, pl. 95, 1865, non Reinecke 1898, nec Rechinger 1908. Type: FIJI ISLANDS. Kadavu, summit of Mbuke Levu mountain. Seemann 641. (Holotype: K [000524876, digital image!], pistillate; Isotypes: K [000524875, digital image!], vegetative; GH [00029835, digital image!], pistillate; BM [000990536, digital image!], pistillate).

Collospermum montanum (Seem.) Skottsb., Kongl. Svenska Vetensk. Acad. Handl. Ser. 3, 14(2): 73. 1934; based on *Astelia montana* Seem.

Funckia montana Kuntze, Revis. Gen. Pl. 2: 711. 1891, *nomen rejiciendum*.

Remarks. Seemann stated in the preface of *Flora Vitiensis* (1965, iv) that “the first set of specimens collected by me were deposited at the Royal Herbarium, Kew and from these the plates accompanying this work have chiefly been taken”. The type material of *A. montana*, which was effectively published in *Flora Vitiensis*, can reasonably be expected to have been accessioned at Kew. Two sheets containing *Astelia montana* specimens collected by Seemann are accessioned at K. One sheet (K000524875) includes a vegetative plant collected at Mt. Mbuke Levu and a second sheet (K000524876) includes a leaf fragment annotated as collected at “Vuna, June 1860” and a pistillate inflorescence labeled as collected by Seemann (n. 641) in 1860. Sheet K000524876 includes the illustrations of the pistillate flower, staminode, berry and seed that appear in the plate that accompanies the protologue of *A. montana* (Seemann 1965).

The pistillate inflorescence labeled as Seemann’s collection n. 641 on sheet K000524876 represents the holotype. According to Seemann’s (1962, 1965) accounts of his field collections, he successfully ascended Mt. Mbuke Levu only once, on 6 September 1860. The vegetative specimen on sheet K000524875 was also collected by Seemann on “Buke Levu” [*sic*]. A type specimen can be mounted on multiple sheets “as long as the parts are clearly labeled as being part of that same specimen” (International Botanical Congress and JH Wiersema 2015); Article 8.3). Although it is likely that sheets K000524875 and K00524876 represent a single specimen that was mounted on separate sheets (Birch pers. comm., Smith 1979), they were not clearly labeled as such. Therefore, K000524875 is considered a duplicate. The leaf fragment

on sheet K000524876 annotated as collected in “Vuna June 1860” may represent a fragment of an earlier collection from Vuna on the island of Taveuni (Smith 1979) where Seemann spent time during June 1860 (Seemann 1962).

***Astelia samoense* (Skottsb.) Birch, comb. nov.**

urn:lsid:ipni.org:names:77148140-1

Basionym: *Collospermum samoënsē* Skottsb., Kongl. Svenska Vetenskapsakad. Handl.

ser. 3, 14(2): 75. 1934. Syntypes: SAMOA ISLANDS. Upolo, 7 July, 1905, *F.*

Vaupel n. 356 (staminate); Aug. 1905, *K. & L. Rechinger n. 4334* (pistillate).

Astelia montana ex Reinecke, Die Flora der Samoa-Inseln. in Bot. Jahrb. Syst. vol. 25. 595. 1898; based on *A. montana* Seem.

Astelia montana ex Rechinger, Vegetationsbilder, series 6, issue 1. tafel 6. 1908; based on *A. montana* Seem.

Remarks. Two specimens were identified by Skottsberg (1934) as types in the list of specimens examined. This included a staminate specimen (Vaupel 356 noted by Skottsberg (1934) as accessioned at HBG) and a pistillate specimen (K. and L. Rechinger n. 4334 noted by Skottsberg (1934) as accessioned at W). The pistillate specimen of Rechinger is not extant at W (pers. comm. A. Löckher, Department of Botany at Naturhistorisches Museum Wien). Efforts are underway to locate type material of *A. samoense* held in herbaria globally.

Diacritical signs, which should be suppressed in species names (International Botanical Congress and JH Wiersema 2015; Article 60.6) have not been transferred from *Collospermum samoënsē*.

***Astelia* subg. *Tricella* Skottsb.**

Astelia subg. *Tricella* Skottsb., Kongl. Svenska Vetenskapsakad. Handl. ser. 3, 14(2): 58. 1934.

Note. Flowers with a short (*A.* sect. *Periastelia*) or long (*A.* sect. *Tricella*) perianth tube (0.3–3.5 mm); anthers dorsifixed, versatile; seeds ellipsoid, fusiform, or ovoid; funicle short, truncate, mucilaginous funicular hairs poorly developed; Type: *A. nervosa* Banks & Sol. ex Hook.f., designated here.

Remarks. *Astelia* subg. *Tricella* is revised to include both *A.* sect. *Tricella* and *A.* sect. *Periastelia*, which form a well supported clade in phylogenetic analyses (Birch unpublished PhD thesis 2011, Birch et al. 2012). *Astelia* sect. *Periastelia* is monophyletic; however *A.* sect. *Tricella* is present as a grade and relationships within each of these sections remain equivocal.

***Astelia* sect. *Tricella* Skottsb.**

Astelia sect. *Tricella* Skottsb., Kongl. Svenska Vetenskapsakad. Handl. ser. 3, 14(2): 58. 1934.

Note. Compact to open growth form; leaves linear or ensiform; inflorescences bearing many flowers; staminate flowers with short or long filaments (0.8–3.5 mm); pistillate flowers with long outer tepals (2.3–5.3 × 1.2–2.6 mm); ovary trilocular, long (2.9–6.8 mm); fruit globose with few seeds per fruit (4–12); seeds long and wide (2.2–3.7 × 1.3–2.3 mm). Type: *A. nervosa* Banks & Sol. ex Hook.f., designated here.

Included taxa and distribution. Australia: *A. australiana* (J.H.Willis) L.B.Moore, *A. psychrocharis* F.Muell. New Zealand: *A. chathamica* (Skottsb.) L.B.Moore, *A. fragrans* Colenso, *A. graminea* L.B.Moore, *A. grandis* Hook.f. ex Kirk, *A. nivicola* Cockayne ex Cheeseman var. *nivicola*, *A. nivicola* var. *moriceae* L.B.Moore, *A. nervosa* Banks & Sol. ex Hook.f., *A. petriei* Cockayne, *A. skottsbergii* L.B.Moore.

Habitat. Lowland scrub (*A. chathamica*), temperate rainforest (*A. australiana*, *A. fragrans*, *A. grandis*, *A. nivicola* var. *moriceae*), sub-alpine and alpine herb-fields (*A. graminea*, *A. nervosa*, *A. nivicola* var. *nivicola*, *A. petriei*, *A. psychrocharis*, *A. skottsbergii*).

***Astelia* sect. *Periastelia* Skottsb.**

Astelia sect. *Periastelia* Skottsb., Kongl. Svenska Vetenskapsakad. Handl. ser. 3, 14(2): 46. 1934.

Note. Open growth form; leaves linear or ensiform; inflorescences bearing many flowers; staminate flowers with short or long filaments (0.9–2.8 mm); pistillate flowers with long outer tepals (2.3–5.3 × 1.2–2.6 mm); ovary trilocular, intermediate length (1.8–4.6 mm); fruit globose, few seeds per fruit (4–12); seeds long and narrow (1.8–3.6 × 0.7–1.6 mm). Type: *A. argyrocoma* A.Heller & Skottsb., designated here.

Included species and distribution. Austral Islands, Rapa: *A. rapensis* Skottsb. Marquesas Islands, Ua Pou, Nuku Hiva: *A. tovii* F.Br. USA, Hawaii: *A. argyrocoma* A.Heller & Skottsb., *A. menziesiana* Sm., *A. waialealae* Wawra.

Habitat. Lowland mesic forest (*A. menziesiana*, *A. rapensis*), tropical montane cloud forest (*A. argyrocoma* and *A. menziesiana*) and alpine swamps (*A. menziesiana* and *A. waialealae*).

Incertae sedis***Astelia* sect. *Micrastelia* Skottsb.**

Astelia sect. *Micrastelia* Skottsb., Kongl. Svenska Vetenskapsakad. Handl. ser. 3, 14(2): 56. 1934.

Note. Low, compact, turf forming growth form; leaves caniculate; inflorescence bearing few flowers; staminate flowers with short filaments (0.6–1.9 mm); pistillate flowers with short outer tepals (2.6–3.8 × 0.8–1.3 mm); ovary trilocular, intermediate length (2.6–3.8 mm); fruit ellipsoid, many seeds per fruit (17–24); seeds short and narrow (1.1–1.7 × 0.5–0.8 mm). Type: *A. pumila* (Forst.) Gaudich.

Remarks. The relationships of *A.* sect. *Micrastelia* are poorly resolved, with alternate relationships with *A.* subg. *Asteliopsis* and the clade containing *A.* subg. *Tricella* and *Collospermum* (Birch unpublished PhD thesis 2011, Birch et al. 2012). *Astelia* sect. *Microastelia* contains a single species, *Astelia pumila*, which is a compact, turf-forming plant and dominant component of *Astelia* moorland in Chile, the Falkland Islands, and Tierra del Fuego. As a cushion-forming species, it differs morphologically from *A.* subg. *Asteliopsis*, which contains species with open, spreading growth form that are epiphytic or terrestrial and primarily found of the understory in lowland to montane forests. *Astelia pumila* does share morphological features with *A.* subg. *Asteliopsis* (e.g. short pistillode or pistil) and, alternatively, with *A.* subg. *Collospermum* and *A.* subg. *Tricella* (e.g. seeds with a short, truncate funicle). The subgeneric placement of *A.* sect. *Micrastelia* remains equivocal and the section is unplaced (*incertae sedis*).

Included species and distribution. Chile, Falkland Islands, Tierra del Fuego: *A. pumila* (Forst.) Gaudich.

Habitat. Lowland (low latitudes) to sub-alpine (mid and higher latitudes) herb-fields particularly on wet substrates (seeps, swamps etc.).

Synoptic key

- 1 Leaf tomentum with scales with stalk attached at base; anthers versatile; pollen densely echinate; ovary/pistillode with weakly to moderately-developed mucilaginous hairs that do not adhere to seed surface on drying..... 2
- Leaf tomentum with peltate scales; anthers immobile; pollen sparsely spinulous; ovary/pistillode with well-developed mucilaginous hairs that adhere to seed surface on drying..... *A. subg. Collospermum*
- 2 Ovary or pistillode unilocular, placentation parietal..... 3
- Ovary or pistillode trilocular, placentation axile 5
- 3 Plants <40 cm tall; leaves generally less than 30 cm long; compact growth form including cushion and turf forming plants; inflorescence panicle (very reduced in *A. subulata*) < 7 cm long, 1–few flowered *A. sect. Astelia*
- Plants > 40 cm tall; leaves generally greater than 30 cm long; open growth form; inflorescence panicle > 9 cm long, many flowered 4
- 4 Leaves with acute apex; outer staminate tepals lanceolate, 5.0–6.4 mm long; pistillode bottle-shaped (lageniform); ovary ovoid; funicle long and curved .. *A. sect. Palaeastelia*

- Leaves with long acuminate apex; outer staminate tepals ovate, 2.5–5.0 mm long; pistillode ovoid-conical; ovary ampulliform or obpyriform; funicle short and ribbed *A. sect. Desmoneuron*
- 5 Leaves generally < 10 cm long; compact turf forming growth form; inflorescence panicle < 7 cm long, 1-few flowered *A. sect. Micrastelia*
- Leaves generally > 10 cm long; compact or open growth form, but not turf forming; inflorescence panicle > 9 cm long, many flowered 6
- 6 Pistillode ampulliform, < 1.2 mm long; pistillate flowers with short outer tepals (2.7–4.0 × 0.7–2.0 mm); fruit ovoid, white, pink or maroon; seeds 1.3–2.2 mm long, 11–18 seeds per fruit; funicle ribbed.... *A. sect. Isoneuron*
- Pistillode ovoid, >1.0 mm, long; pistillate flowers with long outer tepals (2.3–5.3 × 1.2–2.6 mm); fruit globose, orange; seeds 2.0–3.6 mm long, 4–12 seeds per fruit; funicle not ribbed..... 7
- 7 Perianth tube 0.8–4.0 mm long; ovary 1.6–4.6 mm; seeds fusiform and narrow (0.7–1.6 cm)..... *A. sect. Tricella*
- Perianth tube 0.1–0.7 mm long, ovary 2.9–6.8 mm; seeds ovate and wide (1.3–2.3 cm)..... *A. sect. Periastelia*

Acknowledgements

I wish to thank the following individuals and organizations for their generous support of this research. Herbarium specimens were examined from the following herbaria: Auckland War Memorial Museum (AK), Herbarium Pacificum (BISH), Allan Herbarium (CHR), Harvard University (GH), Kew Royal Botanic Gardens (K), National Herbarium of Victoria (MEL), National Tropical Botanical Garden (PTBG), Missouri Botanical Garden (MO), Herbier National de Paris (P), United States National Herbarium (US), and Museum of New Zealand Te Papa Tongarewa (WELT). Staff at BISH graciously provided access to collections, extensive research support and shared their expertise during this research. Tom Ranker, Michael Thomas, and student volunteers at Joseph F. Rock Herbarium (HAW) provided curation support. Tina Carvalho at the Biological Electron Microscope Facility, Pacific Biosciences Research Center of the University of Hawai‘i at Mānoa provided training in and assistance with scanning electron microscopy. I thank Neville Walsh and two reviewers who provided guidance with nomenclature and helpful comments on the manuscript. Funding for field work and sequencing was provided by the American Society of Plant Taxonomists, Explorers Club, National Science Foundation DDIG (DEB-0910402), Sigma-Xi, University of Hawai‘i Foundation and multiple affiliates of the University of Hawai‘i at Mānoa including the Botany Department, Ecology, Evolution, and Conservation Biology Program, Graduate Division, Graduate Student Organization, and the Office of Community and Alumni Relations.

References

- Bayer C, Appel O, Rudall PJ (1998) Asteliaceae. In: Kubitzki K, Huber H (Eds) Flowering Plants, Monocotyledons: Liliaceae (except Orchidaceae) Vol. 3. Springer-Verlag, Berlin, 141–145. doi: 10.1007/978-3-662-03533-7_17
- Birch JL, Keeley SC, Morden CW (2012) Molecular phylogeny and dating of Asteliaceae (Asparagales): *Astelia* s.l. evolution provides insight into the Oligocene history of New Zealand. Molecular Phylogenetics and Evolution 65: 102–115. doi: 10.1016/j.ympev.2012.05.031
- Brown R (1810) Prodromus Florae Novae Hollandiae et Insulae van-Diemen. Hafner Publishing Co., New York.
- Carse H (1926) Botanical notes with descriptions of new species. Transactions and Proceedings of the New Zealand Institute, Wellington 57: 89–92.
- Colenso W (1885) A description of a few new plants from our New Zealand forests. Transactions and Proceedings of the New Zealand Institute, Wellington 14: 329–341.
- Colenso W (1885) A description of some newly-discovered and rare indigenous plants, being a further contribution towards the making known the botany of New Zealand. Transactions and Proceedings of the New Zealand Institute, Wellington 17: 237–265.
- Colenso W (1887) A description of some newly-discovered and rare indigenous Phaenogamic plants, being a further contribution towards making known the botany of New Zealand. Transactions and Proceedings of the New Zealand Institute, Wellington 19: 259–271.
- Colenso W (1888) On new Phaenogamic plants of New Zealand. Transactions and Proceedings of the New Zealand Institute, Wellington 20: 188–211.
- Coode MJE (1978) *Astelia*. In: Bosser J (Ed.) Flore des Mascareignes: La Réunion, Maurice, Rodrigues. 177. Iridacées à 188. Joncacées, Sugar Industry Research Institute, Mauritius, 25–27.
- Darlington C, Wylie AP (1955) Chromosome atlas of flowering plants. Allan & Unwin, London.
- Dawson MI, Beuzenberg EJ (2000) Contributions to a chromosome atlas of the New Zealand flora - 36. Miscellaneous families. New Zealand Journal of Botany 38: 1–23. doi: 10.1080/0028825X.2000.9512671
- de Lange PJ, Murray BG, Datson PM (2004) Contributions to a chromosome atlas of the New Zealand flora - 38. Counts for 50 families. New Zealand Journal of Botany 42: 873–904. doi: 10.1080/0028825X.2004.9512936
- Drake del Castillo E (1893) Flore de la Polynésie Française. Kessinger Publishing, Paris.
- International Botanical Congress and JH Wiersema (2015) International code of nomenclature for algae, fungi and plants (Melbourne Code); adopted by the Eighteenth International Botanical Congress, Melbourne, Australia, July 2011. Appendices II–VIII. Regnum Vegetabile 157: Appendix 3, 163. Koeltz Scientific Books, Königstein, Germany.
- Kuntze O (1891) Revisio Generum Plantarum: vascularium omnium atque cellularium multarum secundum leges nomenclaturae internationales cum enumeratione plantarum exoticarum in itinere mundi collectarum. Volume 2, A. Felix, Leipzig, Germany. doi: 10.5962/bhl.title.327
- McCarthy EM (1928) The structure and development of *Astelia nervosa* var. *sylvestris*. Transactions and Proceedings of the New Zealand Institute 59: 343–360.
- Moore LB (1966) Australasian asteliads (Liliaceae). New Zealand Journal of Botany 4(2): 201–240. doi: 10.1080/0028825X.1966.10429041

- Moore DM (1983) Flora of Tierra del Fuego. Anthony Nelson/Missouri Botanical Garden, Oswestry, England/St. Louis, Missouri.
- Moore LB (1980) Hybrid asteliads (Liliaceae). New Zealand Journal of Botany 18: 37–42. doi: 10.1080/0028825X.1980.10427229
- Moore LB, Edgar E (1976) Flora of New Zealand. Indigenous Tracheophyta: Monocotyledones except Gramineae (Vol. 2). Botany Division, Department of Scientific and Industrial Research, Wellington.
- Rechinger K (1908) Samoa. In: Karsten G, Schenck H (Eds) Vegetationsbilder Series 6(1): tafel 6.
- Reinecke F (1898) Die Flora der Samoa-Inseln. In: Engler A (Ed.) Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie 25: 579–708.
- Richard A (1832) Essai d'une Flore de La Nouvelle-Zélande. In: d'Urville JD, Lesson A, Richard A. Voyage de Découvertes de l'Astrolabe, exécuté par ordre du roi, pendant les années 1826–1827–1828–1829. J. Tastu, Paris.
- Rudall PJ, Chase MW, Cutler DF, Rusby J, de Brujin AY (1998) Anatomical and molecular systematics of Asteliaceae and Hypoxidaceae. Botanical Journal of the Linnean Society 127: 1–42. doi: 10.1111/j.1095-8339.1998.tb02086.x
- Smith AC (1979) Collospermum. In: Flora Vitiensis nova: A new flora of Fiji (Spermatophytes only), Volume 1. National Tropical Botanical Garden, Lawai, Hawaii. doi: 10.5962/bhl.title.44033
- Seemann B (1862) Viti: an account of a government mission to the Vitian or Fijian Islands in the years 1860–61. MacMillion & Co., Cambridge.
- Seemann B (1865) Flora Vitiensis: a description of the plants of the Viti or Fiji islands, with an account of their history, uses, and properties. L. Reeve & Co., London.
- Skottsberg C (1934) Studies in the genus *Astelia* Banks et Solander. Kungliga Svenska Vetensaps-Akademiens Handlingar 14(2): 1–128.
- Thiers B (continuously updated) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/>
- Wagner WL, Herbst DR, Sohmer SH (1999) Manual of the Flowering Plants of Hawai'i (Vol. 2). University of Hawai'i Press/Bishop Museum Press, Honolulu, Hawai'i.
- Wheeler JM (1966) Cytotaxonomy of the large asteliads (Liliaceae) of the North Island of New Zealand. New Zealand Journal of Botany 4: 95–113. doi: 10.1080/0028825X.1966.10443957
- Willdenow CL (1808) Nähtere Bestimmung einiger Liliengewächse, besonders aber derjenigen, die zur sechsten Klasse und dritten Ordnung des Linnéschen Systems gehören. Magazin für die neuesten Entdeckungen in der gesammten Naturkunde, Gesellschaft Naturforschender Freunde zu Berlin, 1: 14–30.
- Williams JB (1987) *Astelia*. In: George AS (Ed.) Flora of Australia (Vol. 45). Australian Government Publishing Service, Canberra, 165–169.

Appendix I

Specimens examined for generation of morphological data. Herbarium abbreviations follow Index Herbariorum (Thiers [continuously updated]). For each specimen, the following data are provided: sampled taxa, voucher specimen information [collection location, date, collector, collection number, herbarium, and herbarium accession number]

Astelia argyrocoma A.Heller & Skottsb. - HAWAIIAN ISLANDS. Kauai: 10–16 Sept 1895, *A. A. Heller* 2752 (BISH120976); Awaawaphui Trail, Honopu, 29 Dec 1956, *H. St. John* 36004 (BISH413786); Kawaihau District, Lihue-Koloa Forest Reserve, Powerline trail from Wailua to Princeville, 27 Feb 1987, *T. Flynn* 2063, *R. Read, B. Read, D. Harder, and S. Weiss* (PTBG018401); Kawaihau District, north facing cliffs and forested slopes below Kekoiki, 9 Feb 1993, *K. R. Wood* 2366, *T. Flynn, D. Lorence, and S. Perlman* (PTBG014508); Na Pali-Kona Forest Reserve, Awaawapuhi Trail, 6 Dec 2007, *J. L. Birch* 137, *D. Lorence, K. Wood, and R. Aqurainja* (BISH751095); Na Pali-Kona Forest Reserve, Pihea Trail, 6 Dec 2007, *J. L. Birch* 141, *D. Lorence, K. Wood, and R. Aqurainja* (BISH751094); Koloa District, Lihue-Koloa Forest Reserve, Wahiawa Bog, 10 Apr 1987, *D. Lorence* 5171, *T. Flynn, R. DeLappe, W. H. Wagner, Jr. and F. Wagner* (PTBG018403); Koloa District, Lihue-Koloa Forest Reserve, Wahiawa Mountains, jeep road to microwave towers on Mt. Kahili, 2 Mar 1987, *D. Lorence* 5105 and *T. Flynn* (PTBG018404); Na Pali-Kona Forest Reserve, N. W. end of Alakai Swamp, 27 Dec 1930, *H. St. John* 10768 (BISH120985); Pihea trail, on the flats above Lehua maka noe, 30 Jul 1963, *W. Takeuchi* 140a (BISH507424); Wahiawa Mts., Aug 1909, *C.N. Forbes* 280K and *J. M. Lydgate* (BISH120978); Waimea Drainage Basin, west side, 3–18 Jul 1917, *C.N. Forbes* 849K (BISH120969).

***A. alpina* R.Br. var. *alpina*-AUSTRALIA.** Tasmania: 1844, *M. Verreaux*s.n. (P00614764); Table Mountain, *R. Brown* 5652 (K000524938); *R. Brown* s.n. (P00614768); Table Mountain, *R. Brown* s.n. (MEL727772); *R. Brown* s.n. (P00614766); *F. v. Mueller* s.n. (K000524934); Mt. Field National Park, Mt Field East, 1876, *F. v. Mueller* s.n. (K00524936); 24 Jan 1983, *S. J. Forbes* 1277 (MEL1523734); 14 Feb 1989, *N. G. Walsh* 2300 (MEL1577414); Mt. Field National Park, Mt. Rodway tow, 8 Jan 2009, *J. L. Birch* 369 (MEL); Mt. Field National Park, Mt. Rodway tow, 8 Jan 2009, *J. L. Birch* 370 (MEL); Mt. Field National Park, Uraquat Trail, 8 Jan 2009, *J. L. Birch* 372 (SP094921); Mt. Field National Park, Uraquat Trail, 8 Jan 2009, *J. L. Birch* 373 (CHR, BISH751060); Cradle Mountain-Lake St. Clair National Park, Marian's Lookout trail, 11 Jan 2009, *J. L. Birch* 379 (MEL); Southwest National Park, Mt. Eliza Plateau, 13 Jan 2009, *J. L. Birch* 387 and *A. Buchanan* (BISH751062); Tasmania: Southwest National Park, Mt. Eliza Plateau, 13 Jan 2009, *J. L. Birch* 388A and *A. Buchanan* (MEL); Southwest National Park, Mt. Eliza Plateau, 13 Jan 2009, *J. L. Birch* 389 and *A. M. Buchanan* (MEL); Southwest National Park, Mt. Eliza Plateau, 13 Jan 2009, *J. L. Birch* 390 and *A. M. Buchanan* (MEL).

A. alpina var. novae-hollandiae Skottsb. - AUSTRALIA. Victoria: Mt. Baw Baw, 1863, *B. F. v. Mueller s.n.* (P00614770); Mt. Baw Baw Alpine Resort, between summit t-bar and Painted Run t-bar, 18 Dec 2008, *J. L. Birch* 343 (MEL); Mt. Baw Baw Alpine Resort, between summit t-bar and Painted Run t-bar, 18 Dec 2008, *J. L. Birch* 344 (MEL); Mt. Baw Baw Alpine Resort, between summit t-bar and Painted Run t-bar, 18 Dec 2008, *J. L. Birch* 345 (MEL); Mt. Baw Baw Alpine Resort, Village Trail, 18 Dec 2008, *J. L. Birch* 346 (MEL); Mt. Buffalo National Park, Lyrebird Plains, 19 Dec 2008, *J. L. Birch* 347 (MEL); Mt. Buffalo National Park, Bogong Plains, 19 Dec 2008, *J. L. Birch* 348 (MEL); Mt. Buffalo National Park, Bogong Plains, 19 Dec 2008, *J. L. Birch* 349 (MEL); Alpine National Park, Big River Fire trail, 20 Dec 2008, *J. L. Birch* 350 (MEL); Alpine National Park, Big River Fire trail, 20 Dec 2008, *J. L. Birch* 351 (BISH751063). New South Wales: Kosciuszko National Park, Kosciuszko summit trail, 30 Dec 2008, *J. L. Birch* 357 and *A. Beehag* (MEL).

A. australiana (J.H.Willis) L.B.Moore - AUSTRALIA. Victoria: Beenak, 20 Dec 1970, *A. C. Beaglehole and B. A. Fuhrer ACB38542* (MEL534348); Beenak, 10 May 1984, *N. G. Walsh 1244* (MEL662242); Eastern Highlands, 11 Jul 1981, *N. G. Walsh s.n.* (MEL599643–44); Eastern Highlands, 18 Mar 1967, *J. H. Williss.s.n.* (MEL224481–3, MEL235432); Great Otway National Park, Brown Town Track, 22 Dec 2008, *J. L. Birch* 352 and *S. McDougall* (MEL); Otways, c. 150 m along Brown Town track from the Lavers Hill-Beech Forest road (just into State Forest on right hand side of Brown Town track), 6 Apr 1990, *R. Robinson s.n.* (MEL1581443–44); Otway Range, Browntown Road, Weeaproinah (head of a tributary of Youngs Creek), 22 Nov 1990, *G. Beilby s.n.* (MEL227997–8); Pioneer Creek, +/- 4 miles S.E. of Powelltown, 22 Nov 1969, *J. H. Willis s.n. and A. Morrison* (MEL224479); Yarra State Forest, 16 Dec 2008, *J. L. Birch* 342 and J. Downe (BISH751064).

A. banksii A.Cunn. - NEW ZEALAND. *E. Cosson s.n.* (P00614762-3). Little Barrier Island: South of Parihakoakoa Stream, base of cliffs above shore, *R. Melville 6573 and E. H. Godley* (CHR129107). North Island: Auckland, cliffs at Takapuna Beach, Apr 1887, *D. Petrie s.n.* (SP084324); Auckland, *T. Kirk s.n.* (MEL5315); Auckland Ecological Region, Awhitu Ecological District, Matakawau, end of Hatton Road, sandstone cliffs bordering Manakau Harbour, 20 Apr 2001, *P. A. Aspin s.n.* (AK253724); Omaha Islets, Eastern Northland and Islands Ecological Region and District, south end Oakura Bay, top E. side of W. islet, 26 Apr 2003, *E. K. Cameron 11668* (AK280946, AK280988); Opotiki, Hukutaia Domain, 15 Jun 1962, *N. Potts s.n.* (CHR132070); Waitakere Ranges, Piha Beach, low cliff adjacent to trailhead to Laird Thomson trail, 16 Dec 2007, *J. L. Birch* 228 (BISH751076); Waitemata, *T. Kirk s.n.* (SP084485). Poor Knights Islands: Tawhiti Rahi Island, Whangarei County, near campsite in Landing Bay, 22 Mar 1984, *A. E. Wright 6337* (AK166389, AK166423). Rakitu (Arid) Island: Hauraki Gulf, west of Te Akau Point, 6 Jan 1981, *E. K. Cameron A183* (AK268645).

A. chathamica (Skottsb.) L.B.Moore - NEW ZEALAND. Chatham Islands: *J. D. Hector M.D. s.n.* (SP030867); Oct, *Travers* 48 (MEL5298); F. v. Mueller (MEL5297); Chatham Island, *H. Travers s.n.* (SP034015A-B); *H. N. Travers s.n.* (SP033784); Chatham Islands source (cultivated at 17 Holmwood Road, Christchurch), 8 Apr 1966, L. B. Moore s.n. (CHR150989A-C); Chatham Islands source (cultivated at 17 Holmwood Road, Christchurch), 7 Nov 1967, L. B. Moore s.n. (CHR181613A-D); *W. M. Martin s.n.* (SP080105A-B); Chatham Islands source (G. W. Ramsay) (cultivated in Botany division garden), 29 Feb 1972, *L. B. Moore s.n.* (CHR233293A-B). Chatham Island: 8 Sept 1966 (SP030867C,G,H); Tuku Valley, Timihunga, 19 Oct 1980, *A. M. Ringer s.n.* (AK170715). Pitt Island: Canister Cove, 5 Jan 1970, *N. C. Simpson s.n.* (SP0 42759).

A. fragrans Colenso - NEW ZEALAND. North Island: Bay of Islands County, Puketi Forest, side ridge south of Bramley's Ridge, north of Waikape Stream, 12 Oct 1984, *P. J. Bellingham s.n.* (AK170933); Kahikatea bush, Elsthorpe Scenic Reserve, Hawkes Bay (near Otane), 28 Oct 1963, *I. M. Morice* 41 (CHR146207A-B); Egmont National Park, Curtis Falls Track, 11 Dec 2007, *J. L. Birch* 205 (AK334003); Smith's Creek, Kaitoke, 19 Oct 1964, *L. B. Moore s.n.* (CHR141163); South Auckland Land District, Ikawhenua Ranges, Galatea, near Hikurangi Trig, 28 Dec 1993, *K. A. Ford* 6/94 (CHR507187A-B). South Island: Haast side of Grassy Creek (ca. 2 miles), 3 Dec 1962, *I. M. Morice s.n.* (CHR133152A-C); Kahurangi National Park, Heaphy Track, 3 Jan 2008, *J. L. Birch* 243 and *K. E. Brown* (HAW); Karamea bluffs, Plot 2, 9 Jan 1985, *P. Wardle s.n.* (CHR419194A-C); Kohaihai River (1 mile south of river); Murchison, 12 Oct 1965, *L. B. Moore and J. Clarke s.n.* (CHR159055); Pelorus River Bridge, Marlborough, 12 Oct 1965, *J. B. Irwin s.n.* (CHR159046A-B), Resolution Bay, Marlborough Sounds, 18 Oct 1965, *J. B. Irwin s.n.* (CHR159046A-B); Taumarunui Co., Scenic Reserve at莫erangi, 21 Nov 1981, *R. O. Gardner* 3165 (AK158745); W. Nelson, 1 Jan 1964, *I. M. Morice* 81 (CHR146986A-E). Stewart Island: Northeastern Long Island, off Stewart Island, 8 Nov 1968, *J. Dugdale s.n.* (CHR188040).

A. graminea L.B.Moore - NEW ZEALAND. South Island: E Nelson, Bryant Range, 5 km. S. W. of Mt. Duppia, Nov 1982, *A. P. Druce* (CHR393919); Mt. Richmond Forest Park, Dun Saddle, Maungatapu Track, 18 Jan 2008, *J. L. Birch* 271 (BISH751067); Mt. Richmond Forest Park, Dun Saddle, Maungatapu Track, 18 Jan 2008, *J. L. Birch* 272 (SP094949). NW Nelson, Boulder Lake Basin, near Darby Pond and Owen Creek, 10 Apr 1966, *I. M. Morice* 327 (CHR170909); NW Nelson, Douglas Range, Jan 1966, *K. H. Marshall s.n.* (CHR141454); NW Nelson, Gouland Downs, (grown in shade house at Lincoln), 26 Apr 1968, *J. Clarke s.n.* (CHR182561); NW Nelson, Mt. Arthur summit trail, 15 Jan 2008, *J. L. Birch* 257, *S. Courtney, and R. Gaskill* (BISH751070); NW Nelson, Mt. Arthur summit trail, New Zealand, 16 Jan 2008, *J. L. Birch* 259, *S. Courtney, and R. Gaskill* (CHR); Nelson, Granity Pass, Mt. Owen, 11 Jan 1983, *K. H. Platt s.n.*

(CHR520447); Nelson, Travers Range, Third Basin, 23 Apr 1962, *L. B. Moore s.n.* (CHR125634B); Nelson, Travers Range, Third Basin, 24 Apr 1962, *L. B. Moore s.n.* (CHR129010A-B).

Astelia grandis Hook.f. ex Kirk - NEW ZEALAND. North Island: Dairy flat, Waitemata Co. 16 Dec 1963, *C. Earwaker s.n.* (CHR146924A-B); Junction of Puketaha Road/Gordontown Road, Hamilton, 14 Dec 2007, *J. L. Birch 223* (BISH751075); Egmont National Park, Potaema Track, 11 Dec 2007, *J. L. Birch 201A-B* (AK334001); North Cape, North Cape Scientific Reserve, Ngaroku Stream, 29 Dec 1986, *R. E. Beever 87003* (AK176615); Northland, Karikari, Tokerau Beach, Hector Busby property, 17 Oct 2006, *M. E. Young and L. J. Forster s.n.* (AK297667-70); South Auckland province, Whanganui, 15 Oct 1895 (SP084426); W Hokio, west of Levin, Oct 1967, *A. P. Druce s.n.* (CHR179566); Waitomo Co., Waitanguru, 15 Feb 1967, *R. Bell and N. Potts s.n.* (CHR174847A-B). South Island: Lake Ianthe (cultivated at Lincoln), 10 Nov 1966, *J. Clarke s.n.* (cultivated) (CHR566424A-B); Moeraki Valley, South Westland, 3 Apr 1978, *P. Wardle s.n.* (CHR321183A-C); Te Kinga near Lake Brunner, 14 Nov 1965, *N. Lambrechtsen s.n.* (CHR168155A-C).

A. *bastata* Colenso - NEW ZEALAND. 1875, *M. Filhol s.n.* (P00614792). North Island: Atuanui State Forest, 6 Apr 1969, *R. E and J. Beever 236* (CHR 195414A-B); Auckland Ecological Region, Awhitu Ecological District, Pollok, S off Barthow Road, Craig's Bush, lower part eastern end, 9 Jun 2005, *P. A. Aspin s.n.* (AK294016-7); Auckland Ecological Region, Hunua Ecological District, Hunua Range, Mangatangi Kauri area, 4 Feb 1971, *I. L. Barton* (AK208895); Egmont National Park, Stony River, May 1961, *A. P. Druce s.n.* (CHR129729); Rotorua County, Lake Rotoehu, 29 Jan 1963, *R. Mason 100059* (CHR140253A-C); South Manukau, Mauku, Jun 1901 (AK222913); Tainui Ecological Region, Kawhia Ecological District, Mount Pirongia, Ruapane, 13 Feb 1999, *P. J. de Lange 3801* (AK237952); Takaka, Rameka Gorge, 22 Feb 1964, *V. M. Scott s.n.* (CHR148397); Waikanae, Wellington, 5 Feb 1966, *I. M. Morice 312* (CHR150828A-D); Wairoa, Hawkes Bay, Jan 1886, *A. Hamilton s.n.* (AK3191); Whangarei, Maungatapere, Aug 1989, *H. Carse s.n.* (AK3228). Poor Knights Islands: Motukapiti Island, (summit) Nature Reserve, 20 Nov 1984, *A. Penniket s.n.* (CHR418304A-B).

A. *hemichrysa* (Lam.) Kunth - MASCARENE ISLANDS. Reunion Island: Forêt de Bébour, 26 Mar 1973, *D. H. Lorence s.n.* (MO2232208); Forêt de Bébour, 23 Feb 1979, *D. H. Lorence 2422* and *T. Cadet* (MO 2715190); Plateau de Belouve, 1956, *J. Bosser 9462* (P00636603); Morne du Patates à Durand, 18 Mar 1974, *J. Bosser 21,643* (P00636604); *I. M. Morice P619* (CHR182114). Mauritius: Pétrin Nature Reserve, 3 Feb 1973, *D. Lorence M53* (MO222048); Pétrin Nature Reserve, 8 Oct 1978, *D. Lorence 2100* (MO2715777).

A. linearis Hook.f. var. **linearis** - NEW ZEALAND. South Island: NW Nelson, Lake Sylvester, 28 Mar 1964, *L. B. Moore s.n.* (CHR148475); Lake Sylvester above Cobb Dam, 9 Jan 1961, *P. Hynes 70325* (AK70325); Boulder Lake, Jan 1957, *A. E. Esler s.n.* (AK216795); Southland, Longwood Range, Bald Hill, 22 Jan 2008, *J. L. Birch 279 and P. Michel* (HAW); Southland, Longwood Range, Bald Hill, 22 Jan 2008, *J. L. Birch 281 and P. Michel* (CHR); Southland, Longwood Range, Bald Hill, 22 Jan 2008, *J. L. Birch 276 and P. Michel* (BISH751078); Westland, 3 Nov 1985, *P. N. Johnson 469 and P. Wardle* (CHR420097); Denniston, near Westport, 5 Aug 1942, *I. A. McNeur 25407* (CHR25407); Port Pegasus, Stewart Island, *G. M. Thomson s.n.* (SP084464).

A. linearis var. **novae-zelandiae** Skottsb. - NEW ZEALAND. North Island: *W. Collenso 1587* (K000524926); Wellington Land District, Southern Ruahine Range, Mar 1971, *A. P. Druce s.n.* (CHR245761). South Island: Southland, Longwood Range, Bald Hill, 21 Jan 2008, *J. L. Birch 280 and P. Michel* (HAW); Fiordland National Park, Borland Saddle/Mt. Burns track, 24 Jan 2008, *J. L. Birch 290 and P. Michel* (MEL); Fiordland National Park, Borland Saddle/Mt. Burns track, 24 Jan 2008, *J. L. Birch 289 and P. Michel* (AK334015); Westland Land District, S. W. of Lewis Pass, 27 Dec 1962, *D. R. Given s.n.* (CHR142821); Wilberg Range, 27 Apr 1993, *P. Wardle s.n.* (CHR499828); Mt. Davy. *W. R. B. Oliver s.n.* (SP010664). Stewart Island: Port Pegasus, Bald Cove, Apr 1981, *I. M. Ritchie s.n.* (CHR372824); Rakiura National Park, Mt. Anglem, 26 Jan 2008, *J. L. Birch s.n.* (HAW); Rakiura National Park, Mt. Anglem summit, 26 Jan 2008, *J. L. Birch 300 and J. Blythal* (P02141677).

A. menziesiana Sm. - HAWAIIAN ISLANDS. *J. F. Rock 8415* (BISH121100). Hawai‘i Island: ca. 0.5 miles east of National Park Road junction, 04 Aug 1964, *M. R. Crosby and W. R. Anderson 1972* (BISH121099); Hawaii Volcanoes National Park, Kilauea Research Center, 13 Aug 2009, *J. L. Birch 391* (HAW); Kau Forest Reserve, near boundary of Kapapala Forest Reserve, 1 Jul 1981, *J. Davis 560* (BISH657788); Kealakekua Ranch, S. Kona, 2 Apr 1980, *J. Davis 274* (BISH656028); Saddle Road, Kipuka No. 9. 27 July 1983, *T. Flynn 494* (PTBG019408); Olaa Forest Reserve, road from Hilo to Kulani Prison, 31 Oct 1950, *W. H. Hatheway 441* (BISH121085); South Kohala District, Umipoho Gulch area above Koia sanctuary, 16 Oct 1995, *K.R. Wood 4676* (PTBG033654); Waiakea mauka, July 1986, *W. Takeuchi and K. Shimabukuro 2699* (BISH503251); East Maui: Haleakala, upper Keanae valley, 19 Jul 1927, *O. Degener 4057* (BISH121037); Hana District, Hana Forest Reserve, above N rim of Kipahulu, 1974, *B. Harrison 395* (BISH429287); Hana District, E Haleakala, 2 mi N. E. of Paliku Cabin, N facing slope above Wai Anapanapa, *J. Henrickson and R. Vogl 3556* (BISH35823); Rainforest SW of Kaunuohua, 20 May 1982, *J. Davis 759* (BISH657637); Ukulele, 1919, *C. N. Forbes s.n.* (BISH121031); West Maui: 20 Jul 1964, *M. R. Crosby and W. R. Anderson 1864* (BISH121024); Puu

Kukui Watershed Preserve, along boardwalk to summit at mile marker 2300, 31 Oct 2007, *J. L. Birch 154, R. Bartlett, D. Cole, L. Dunn, and D. Tanaka* (HAW); Puu Kukui Watershed Preserve, forest below bog, 31 Oct 2007, *J.L. Birch 158, R. Bartlett, D. Cole, L. Dunn, and D. Tanaka* (MEL); Puu Kukui Watershed Preserve, forest below bog, 31 Oct 2007, *J.L. Birch 159, R. Bartlett, D. Cole, L. Dunn, and D. Tanaka* (HAW); Molokai: Waikolu Valley, head of valley, 15 Mar 1952, *O. Degener 22160 and C. Tousley* (BISH10425); between Waikolu Valley and N. Puu Alii, 10 Apr 1928, *O. Degener 4065* (BISH121013); Pepeopae Bog, 25 m after end of boardwalk through bog, 22 Jun 2007, *J. L. Birch 149 and C. W. Morden* (BISH751101); ridge E of Mapulehu Valley, 29 Dec 1932, *H. St. John 12840* and F. Fosberg (BISH121017); Kawele, ridge to Pelekunu Pali, 17 Mar 1910, *J. F. Rock 6095* (BISH121053); Oahu. Waianae Range: Mt. Kaala, 14 Sept 2007, *J. L. Birch 179* (HAW); Mt. Kaala summit trail (Waianae access), 26 Sept 2009, *J. L. Birch 393* (HAW); Mt. Kaala summit, 24 Nov 1929, *H. St. John 10071* (BISH121055); Kaukawahua gulch, N. fork, 15 May 1909, *J. F. Rock 4005* (BISH121110); Pukaloa gulch, above Schofield, off ridge trail to Puu Kala-na, near Kumakalii, 16 Apr 1987, *S. Perlman 5640 and B. Hill* (BISH617222); Puu Hapapa, northeast ridge, 7 May 1939, *O. Degener 12382* (AK71043); Puu Hapapa, 16 Mar 1930, *E. Christopherson 1288B* (BISH121111); Puu Hapapa, 16 Mar 1930, *E. Christopherson 1288A* (BISH121113); Ridge between Puu Kanehoa and Puu Kaua, 23 Jun 1940, *O. Degener 12966* (BISH121146). Koolau Range: Kulouou Valley, summit, 23 Jun 1935, *O. Degener 10471*, K. Park, and D. Topping (BISH121114); N Kaaawa, 12 Apr 1931, *E. P. Hume 189* (BISH121171).

- A. *microsperma*** Colenso - NEW ZEALAND. North Island: Auckland Ecological Region, Rodney Ecological District, Mt. Tamahunga, 28 Feb 1993, *M. E. Young s.n.* (AK212022); Blue Mountains, near Pinehaven, Hutt Valley, 1 Jan 1963, *I. M. Morice s.n.* (CHR132056A-C); Kaimai Mamaku Forest Park, Mt. Te Aroha, 13 Dec 2007, *J. L. Birch 248, C. Gemmill, E. Grove, and N. Wakefield* (SP094950); Mangonui County, Maungataniwha Range, Feb 1908, *H. Carse 516/1* (CHR328214 A-B); Mt. Egmont, near Punihou Hut, 20 Jan 1963, *I. M. Morice s.n.* (CHR132058); Ohakune, Mt. Road, 13–14 Dec 1962, *J. M. Wheeler s.n.* (CHR141652); Originally from Ruahine Range, west Tamaki River (cultivated in shade house at Lincoln), 2 Jan 1968, *J. Clarke s.n.* (CHR182141); Pureora State Forest Park, West Taupo, 26 Jan 1982, *J. E. Braggins 97* (AK270463); Tararua Forest Park, trail to Mt. Holdsworth, 12 Jan 2008, *J. L. Birch 244 and S. J. Birch* (BISH751080); Tongariro National Park, road between Ohakune and Turoa, 17 Mar 1999, *L. Perrie and L. Sheppard s.n.* (SP083500).

- A. *montana*** Seem. - VANUATU ISLANDS. Aneityum: crête du Nezwon Ne-tounemla, 21 Jul 1971, *J. Raynal and M. Schmid RSNH 16129* (CHR 298987; P00636616); Mt. Inréro, peak south of mountain, 21 Jul 1971, *P. S. Green in RSNH 1149* (P00636643). Espiritu Santo: Mt. Tabwemasana, 17 Aug 1985, *P.*

Cabalion 2816 (P000636609); Mt Tabwemasana, 2–8 Sept 1971, *C. Wee-Lek RSNH* 249 (P00636644). Tanna: Mt. Toukosméréu, 29 Jul 1971, *P. S. Green* 1244 (P00636642). FIJIAN ISLANDS. Viti Levu: Water reserve near Suva, Jul 1965, *J. W. Dawson s.n.* (CHR170581, SP85966A–B). Vanua Levu: Thakaundrove, Natawa Peninsula, Uluingala, 15 Jun 1934, *A. C. Smith* 1978 (P00636610). Kandavu: Mt. Mbuke Levu summit, 6 Sep 1860, *B. Seemann* 641 (GH00029835).

A. nadeaudii Drake & F.Br. - SOCIETY ISLANDS. Tahiti: Mt. Aorai, 27 Apr 1858, *J. Nadeaud* 250 (P0636625–28); Mt. Aorai, trail to summit from Fare Rau Ape, above Papeete on ridge east of Fautaua Valley, slopes of Rocher du Diable, 30 Mar 1973, *F. R. Fosberg* 54,701 and *M.-H. Sachet* (US2680616, US2680617); Mt. Aorai, 14 Dec 2006, *J. Meyers s.n.* (BISH751087); Mt. Aorai, sentier de l'Aorai, entre Fare Ata et le sommet, 15 Feb 1983, *J. Florence* 4545 (P00636605); Mt. Marau, sentier du Pic. Vert., 28 Jan 1982, *J. Florence* 2307 (P00636617, US3186926); Mt. Orohena, crête ouest de la Papenoo, sentier de l'Orohena, 20 Oct 1982, *J. Florence* 3992 (P00636618); Tahiti, 1847, *M. J. Lépine s.n.* (P00636635); Tahiti, *J. Nadeaud* 250 (P00636622–24). Raiatea: Mt. Toomaru, Tevaitoa, crête sommitale N. du Mt. Toomaru, 27 Nov 1987, *J. Florence* 8936 (P00636640).

A. neocaldonica Schltr. - NEW CALEDONIA. 1868–1870, *M. Balansa* 950 (P00636652); 19 Apr 1910, *M and Mme Le Rat* 2874 (P00636651); *I. Franc* 3125 (MO977220, MEL600656). Grande Terre: Forêt de Sailles, 5 Dec 2001, *J. Muzinger* 1244, *B. Suprin* and *F. Carriconde* (MO5839966); Mt. Koghi, Mar 1929, *M. Franc* 2325 (P00636650); Mt. Mandjélia, 24 May 1980, *J. W. Dawson s.n.* (SP085969); Mt. Mandjélia, below radio tower ca. 5 air-km of Pouébo, 11 Apr 1980, *G. McPherson* 2531 (PTBG023027–8); Mt. Mandjélia, below radio tower ca. 5 air-km of Pouébo, 11 Apr 1980, *G. McPherson* 2532 (MO2922540); Mt. Mau, crête sommitale, 21 Aug 1940, *M. R. Virot* 284 (P0636648); Province Nord, Mont Görö Até, 19 Nov 2002, *F. Tronchet* 469 and *J. Munzinger*, *D. and I. Létocart*, *J.-P. Butin*, *A. Oddi*, and *A. Obry* (MO4781294); Province du Nord, upper Amoa River Valley, trail from Alain Obry property to Görö Até, 23 Apr 2002, *I. Létocart* 5644, *P. Lowry II*, *G. D. McPherson*, *F. Carriconde*, and *D. Létocart* (MO5666374).

A. nervosa Banks & Sol. ex Hook.f. - NEW ZEALAND. North Island: Coromandel County, Moehau summit, 2 Nov 1980, *R. O. Gardner* 2763 (AK153040); Kokianga Co., Waima Forest, ridge between summit of Mount Misery and “highest point in Northland”, 16 Jan 1990, *A. E. Wright* 9684 (CHR192727); Egmont National Park, Ski field Road, 11 Dec 2007, *J. L. Birch* 207 (CHR); Kaimai Mamaku Forest Park, Mt. Te Aroha, Waikato, 13 Dec 2007, *J. L. Birch* 220, *C. Gemmill*, *E. Grove*, and *N. Wakefield* (SP094947); Otorohanga County, Ranginui summit, Rangitoto Range, 16 Dec 1981, *R. O. Gardner* 3209 (AK158716); Table Mountain, Kauaeranga Valley, Coromandel Range, 10 Apr 1971, *I. M. Morice*

484B (CHR208245); Thames County, Lookout rocks, inland from Taran, 10 Dec 1986, *R. O Gardner* 5049 (CHR484212); Tongariro (grown in cultivation at Lincoln), 10 Nov 1966, *J. Clarke s.n.* (CHR566443A–B); Waitemata County, Albany Scenic Reserve, valley lying east of Lonely Track Road, Wright's Road intersection, 6 Oct 1979, *R. O. Gardner s.n.* (AK150794); Wellington District, Tongariro National Park, ca. 1.5 km N. W. of Chateau Tongariro, nr. road to Chateau, Whakapapanui Track, 24 Mar 1970, *P. J. Edwards* 74 (AK129710). South Island: Marlborough Land District, Mt. Stokes, Marlborough Sounds, Mar 1977, *A.P. Druce s.n.* (CHR310138); Mt. Stokes, *T. Kirk s.n.* (SP030865); Mt. Stokes Scenic Reserve, Mt. Stokes summit, Marlborough Sounds, 14 Jan 2008, *J. L. Birch 251 and J. Little* (BISH751065); Mt. Stokes Scenic Reserve, Mt. Stokes summit, Marlborough Sounds, 14 Jan 2008, *J. L. Birch 252 and J. Little* (CHR); Nelson Land District, Nelson, Aniseed Valley Scenic Reserve, 1972, *G. C. and D. Kelly s.n.* (CHR230190). Arthur's Pass, Bealey Track near Margaret's Tarn, 29 Nov 1961, *R. Melville* 5478 (CHR129127, AK156821); Kahurangi National Park, Heaphy Track, 31 Dec 2007, *J. L. Birch 237 and K. E. Brown* (HAW); Kahurangi National Park, Heaphy Track, 2 Jan 2008, *J. L. Birch 240 and K. E. Brown* (SP094946A/B); Kahurangi National Park, Heaphy Track, James Mackay Hutt, 2 Jan 2008, *J. L. Birch 241 and K. E. Brown* (P02141675); Kahurangi National Park, Heaphy Track, trail to Gouland Downs Hut, ca. 1 mile from Percy Hut, 31 Dec 2007, *J. L. Birch 238 and K. E. Brown* (SP094945); NW Nelson, Mt. Arthur, 16 Jan 2008, *J. L. Birch 263* (BISH751073); Mt. Arthur, 16 Jan 2008, *J. L. Birch 267* (CHR); NW Nelson, Boulder Lake, Orater Creek, 11 Apr 1966, *I. M. Morice* 344 (CHR170919); Cobb Valley, Lake Sylvester, just above forest hut, 31 Mar 1964, *L. B. Moore s.n.* (CHR148498A–B); trail to Mt. Arthur, 15 Jan 2008, *J. L. Birch 258, S. Courtney, and R. Gaskill* (SP094941); Mt. Arthur trail to Gordon's Pyramid, 16 Jan 2008, *J. L. Birch 260* (BISH751068); Takaka Hill, on road to Canaan, 11 Jan 1964, *I. M. Morice* 106 (CHR146995); Nelson Lakes National Park, Ridge to Mole Tops, 14 Feb 1964, *M. J. A. Simpson* 4115 (CHR148394); North Canterbury, Upper Clarence Valley, Mt. St. Patrick, 14 Jan 1972 (CHR 228672); Canterbury Land District, Deer Spur Walk, Peel Forest Park, Stop 5, 29 Oct 1970, *B. P. J. Molloy* (CHR212134); West Coast, Paparoa Range, Croesus track, 19 Jan 2008, *J. L. Birch 274* (AK334012); Oparara Arch, 22 Jan 1985, *P. Wardle s.n.* (CHR574281); St. Arnaud Range, 11 Dec 1950, *W. R. B. Oliver s.n.* (SP010631). Fiordland, Dec 1966, *P. K. Dorizac s.n.* (CHR174065); Fiordland, Mt. Gray, 14 Feb 1959, *M. J. A. Simpson* 1154 (CHR111751); Fiordland, Secretary Island, Feb 1967, *P. Wardle* (3) (CHR 566430); Fiordland National Park, Hollyford Valley, 31 Jul 1965, *K. Dorizac s.n.* (CHR141436); Homer, Dec 1943, *J. Salmon s.n. with R. Forster* (SP084347); Milford Sound, Sinbad Gulley, 27 Feb 1975, *P. N. Johnson s.n.* (CHR 261738); Pigeon Saddle, Oct 1957, *A. Esler s.n.* (SP085190); Track from Key Summit to road, 16 Dec 1962, *I. Morice s.n.* (CHR133242); Track from Key Summit to road, 16 Dec 1962, *I. Morice s.n.* (CHR133243).

- A. *nivicola*** Cockayne ex Cheeseman **var. *nivicola*** - NEW ZEALAND. South Island: Fiordland, Hunter Mts., east of summit of unnamed peak north of Green Lake, 9 Jan 1967, *D. Given* 69043 (CHR193928); Fiordland, Key Summit, between upper bog and track upwards, *P. K. Dorizac* (=IMM 474/2) (CHR191766A-B); Fiordland National Park, Borland Saddle/Mt. Burns track, 24 Jan 2008, *J. L. Birch* s.n. and *P. Michel* (HAW); Marlborough, Richmond Range, Mt. Richmond, 23 Nov 1961, *J. I. Townsend* s.n. (CHR366690); Mt. Ollivier, Sealey Range, 12 Feb 1919, *L. Cockayne* 1272 (AK3224); NW Nelson, Mt. Centre, Jan 1977, *A. P. Druce* s.n. (CHR310405, CHR310406); Nelson Land District, NW Nelson, Mt. Centre, Jan 1977, *A. P. Druce* s.n. (CHR310406); Southland, Longwood Range, Bald Hill, 21 Jan 2008, *J. L. Birch* 277 and *P. Michel* (CHR).
- A. *nivicola* var. *moriceae*** L.B.Moore - NEW ZEALAND. South Island: Collected from Wilmot Pass, (grown in Bot. Div. shadehouse), 5 May 1970, *L. B. Moore* s.n. (CHR199865); Fiordland, near Henry Saddle on George Sound track, 16 Jan 1966, *P. K. Dorizac* s.n. (CHR150752, CHR150753); Nelson, Flora track, head of Pearse River (type locality), 16 July 1964, *L. B. Moore* s.n. (CHR 151212); Nelson, Flora track, halfway between Flora saddle and Flora Hut, 26 Dec 1964, *D. R. Given* 64410 (CHR144190A-B); Paparoa Range, above Roa Mine, 18 Dec 1965, *L. B. Moore*, s.n., *J. Clarke*, and *I. Robins* (CHR168145A-C); On E side 15 minutes from Wilmot Pass, 14 Dec 1962, *I. Morice* s.n. (CHR133164A-C); Track to Boulder Lake, The Pulpit, 6 Jan 1964, *I. M. Morice* 92 (CHR146984A-C); Nelson, track to Mt. Arthur, 13 Apr 1963, *I. M. Morice* 12 (CHR 144249).
- A. *papuana*** Skottsb. - PAPUA NEW GUINEA. Central Highlands, Mt. Wilhelm, 25 Aug 2011, *W. R. and M. N. Philipson* 3493 (CHR198540), Eastern Highlands province, Kainantu sub-province, Mt. Piora, 9 Jan 1975, *M. J. S. Sands* 1604, *G. A. Pattison*, and *J. J. Wood* (US3248428); Goilala sub-district, Mt. Dickson, 11 Feb 1964, *T. G. Hartley* 12985 (US3485933); Mt. Wilhelm, Lake Aunde, 22-26 Sep 1962, *F. W. Went* 223 (MO1806059); South Highland district, Tari sub-district, Mt. Ambua, 29 July 1966, *W. Vink* 17287 (MO2322009); Western Highlands district, Hagen sub-district, Mt. Hagen, 2 July 1967, *J. M. Wheeler* ANU 6385 (US3321035); West Highlands, Mt. Kegum, 8 Apr 1977, *J. F. Veldkam* 7598 and *A. Vinas* (MO2682556); West Sepik district, Telefomin sub-district, Star Mountains, Mt. Scorpion, 25 May 1975, *J.R. Croft* and *G.S. Hope* LAE 68018 (US2895090); West Sepik district, Telefomin sub-district, Mt. Capella north of east summit, 18 Apr 1975, *W. R. Barker* LAE 67438 and *T. Umbo* (MO3270313).
- A. *petriei*** Cockayne - NEW ZEALAND. South Island: Arthur's Pass, Temple Basin, 8 Dec 1963, *B. H. MacMillan* 3 (CHR146933); Fiordland National Park, Gertrude Saddle track, 23 Jan 2008, *J. L. Birch* 287 and *P. Michel* (SP094956); Fiordland National Park, Gertrude Saddle track, 23 Jan 2008, *J. L. Birch* 286 and *P. Michel* (CHR); Fiordland National Park, Borland Saddle/Mt. Burns track, 24 Jan

2008, *J. L. Birch* 291 and *P. Michel* (SP094952); Molesworth Ecological Region, Balaclava Ecological District, Island Saddle just north of road summit, 4 Jan 2002, *E. K. Cameron* 10684 (AK255636); N. Canterbury, Upper Clarence Valley, Mt. St. Patrick, 14 Jan 1972 (CHR228674); NW Nelson, Mt. Arthur, 15 Jan 2008, *J. L. Birch* 254, *S. Courtney*, and *R. Gaskill* (BISH751072); NW Nelson, Mt. Arthur, 15 Jan 2008, *J. L. Birch* 255, *S. Courtney*, and *R. Gaskill* (MEL); Otago Province, Lake Harris Saddle (cultivated in Dunedin), 1923, *W. A. Thomson s.n.* (SP085977); Westland National Park, Fox Glacier, Chancellor Hut, 2 Feb 1973, *P. Wardle s.n.* (CHR218793A–B).

A. *psychrocharis* F.Muell. - AUSTRALIA. New South Wales: Mt. Kosciuszko, 19 Feb 1990, *M. G. Corrick* 10668 (MEL1578958); Mt. Kosciuszko, June 1901, *C. H. Grove s.n.* (MEL 2213620); Kosciuszko National Park, Mt. Kosciuszko, 30 Dec 2008, *J. L. Birch* 359 and *A. Beehag* (MEL); Kosciuszko National Park, Mt. Kosciuszko, 30 Dec 2008, *J. L. Birch* 360 and *A. Beehag* (MEL); Kosciuszko National Park, Mt. Kosciuszko, 31 Dec 2008, *J. L. Birch* 361 and *A. Beehag* (MEL); Kosciuszko National Park, Mt. Kosciuszko, 31 Dec 2008, *J. L. Birch* 362 and *A. Beehag* (BISH751056); Kosciuszko National Park, Southern Tablelands, near end of Mt. Blue Cow, “Snowtube”, 31 Jan 1990, *J. H. Willis s.n.* (MEL 2117076).

A. *pumila* (Forst.) Gaudich. - ARGENTINA. Tierra del Fuego: Isla de los Estados, Bahia San Antonio, Puerto Hoppner, 8 Nov 1971, *T. R. Dudley*, *R. N. P. Goodall*, and *G. Crow* 1596 (MO2300439); Northwest side of Bahia Thetis, 14 Nov 1969, *R. N. P. Goodall* 2248 (US2626023). CHILE. Chiloé Island: River Toigoi, Chepu, 26 Oct 1958, *E. J. Godley* 349a (CHR 547077B); River Toigoi, Chepu, 26 Oct 1958, *E. J. Godley* 348 (CHR 55303). Magallenes y de la Antártica Chilena: Churucua, 30 Jan 1879, *Lud Savatier* 189 (P00614743); Détroit de Magellan-Port Famine. *M. le Guiillou* 188 (P00614753). Riesco Island: Peninsula Cordova, Pto. Condor, 29 Aug 1970, *E. Pisano V.* 2639 (MO2384477). FALKLAND ISLANDS. [Iles Maclovian] 1825, *Gaudichaud s.n.* (P00614740); [Iles Maclovian] 1825, *Gaudichaud s.n.* (P00614741).

A. *rapensis* Skottsb. - AUSTRAL ISLANDS. Rapa: Kaimaru, south ridge of Mt. Perahu, 13 Jul 1934, *H. St. John* and *J. Maireau* 15,513 (BISH509590, P00636647).

A. *samoense* (Skottsb.) Birch - SAMOAN ISLANDS. Savai‘i: Cloud forest near Mt. Silisili, 17 June 1992, *A. Whistler* 8861 (housed at HAW); Mata-ole-afi cinder cone, 29 May 1975, *A. Whistler* W2492 (housed at HAW); Matavanu Crater, 24 July 1931, *E. Christophersen* and *E. Hume* 2130 (MO1631444); Mt. Silisili, 30 May 1975, *A. Whistler* W2518 (housed at HAW). Upolu: 22 Nov 1973, *A. Whistler* AW1172 (housed at HAW); between Mt. Lepu‘e and Mt. Fito, 15 Oct 2008, *A. Whistler* 12035 (housed at HAW); Lake Lano, 6 Aug 1971, *A. Whistler* W270 (housed at HAW); Laka Lanoto‘o, 15 Aug 1970, *A. Whistler* W2177 (housed at

HAW); Mt. Siga'ele, 19 July 1973, *A. Whistler W358* (housed at HAW); Mt. Lepu'e east of Tiavi, 7 Dec 1973, *A. Whistler W1251* (housed at HAW).

- A. *skottsbergii*** L.B.Moore - NEW ZEALAND. South Island: Douglas Range, 15 May 1963, *K. H. Marshall s.n.* (CHR146875A-D); NW Nelson, Boulder Lake, Brown Cow Saddle, 12 Apr 1966, *I. M. Morice 350* (CHR170925); NW Nelson, Douglas Range, 1 Jun 1964, *K. Marshall s.n.* (CHR151057); NW Nelson, Mt. Domett, Jan 1977, *A. P. Druce s.n.* (CHR311624); NW Nelson, Mt. Goul, 19 Jan 1973, *M. J. A. Simpson 7215* (CHR278243); Mt. Arthur Plateau, Gordon's Pyramid, 26 Feb 1965, *R. Macfarlane s.n.* (CHR141321); Kahurangi National Park, Mt. Arthur, 16 Jan 2008, *J. L. Birch 261* (BISH751069); Kahurangi National Park, Mt. Arthur, 16 Jan 2008, *J. L. Birch 262* (CHR); Waingaro Peak, where Douglas and Lockett Ranges meet, on North facing slope, 16 Jan 1970, *I. M. Morice s.n.* (CHR199791); Wangapeka, Mt. Luna, 23 Jan 1971, *I. M. Moore s.n.* (CHR215917A-B).
- A. *solandri*** A.Cunn. - NEW ZEALAND. Little Barrier Island: North Auckland Land District, 24 Oct 1965, *R. Melville 6672* (CHR129112). North Island: Anatoki Valley, 9 Jan 1964, *I. M. Morice 100* (CHR147598); Auckland, 1875, *M. Filhol s.n.* (P00614785); Auckland Ecological Region, Awhitu Ecological District, Matakawau Reserve, south of Matakawau Road, 23 Apr 2001, *P. A. Aspin s.n.* (AK253912, AK254486); Coromandel, Dec 1905, *D. Petrie s.n.* (SP85958); Egmont Ecological Region and District, Mt. Egmont, Aug 1958, *A. E. Esler s.n.* (AK219600); Hunua, Dec 1868, *T. Kirk s.n.* (SP84518); Hunua, Dec 1868, *T. Kirk s.n.* (SP084519); Kaimai Mamaku Forest Park, Mt. Te Aroha summit trail, 13 Dec 2007, *J. L. Birch 219, C. Gemmill, E. Grove, and N. Wakefield* (SP094962); Mauku, Franklin County, 5 May 1901, *H. Carse s.n.* (CHR328230); Egmont National Park, Potaema Track, 11 Dec 2007, *J. L. Birch 202* (SP094922); Egmont National Park, road below Dawson Falls, 20 Jan 1964, *I. M. Morice 119* (CHR148342); Mt. Wellington lava fields near Auckland, Apr 1897, *D. Petrie s.n.* (SP084507); Mt. Wellington, Auckland, 4 Apr 1922, *H. Carse s.n.* (CHR292211); North Tararua Range, Makotukutuku St. near Mt. Kaiparoro, Jan 1977, *C. Ogle s.n.* (CHR286974); Orotere, south of Kaeo, 26 Aug 1965, *R. Cooper s.n.* (AK118465); Tararua Forest Park, Mt. Holdsworth, 12 Jan 2008, *J. L. Birch 250 and S. J. Birch* (BISH751081). South Island: Nelson (SP08453). Poor Knights Islands: Whangarei County, Aorangi Island, 31 Aug 1984, *A. E. Wright 6564* (AK169414). Rangitoto Island: North Auckland Land District, 13 Feb 1965, *I. M. Morice s.n.* (CHR 566383).

- A. *subulata*** Cheeseman - NEW ZEALAND. Campbell Island: Homestead Ridge, 26 Dec 1998, *C. D. Meurk s.n.* (CHR537469); Tucker Cove, 14 Jan 1947, *W. B. Brockie s.n.* (CHR223725). South Island: Paparoa Range, Westland, 14 Jan 1967, *L. B. Moore s.n. and J. Clarke* (CHR174718). Stewart Island: Table Hill, 12 Jan

1940, *L. B. Moore* s.n. (CHR24184); Rakiura National Park, Mt. Anglem, 26 Jan 2008, *J. L. Birch* s.n. and *J. Blythal*, (HAW); Rakiura National Park, Mt. Anglem, 26 Jan 2008, *J. L. Birch* 299 and *J. Blythal*, (CHR); Rakiura National Park, Mt. Anglem, 26 Jan 2008, *J. L. Birch* 301 and *J. Blythal* (BISH751079).

- A. *tovii*** F.Br. - MARQUESAS ISLANDS. Nuku Hiva: Summit of the ridge south of Tekao, between the new airport road at peak #1227 M. and Tekao, the main ridge above Toovii, 25 Sept 1995, *S. Perlman* 15,055 (BISH660310, MO5601362, PTBG021995); Mt. Tapuaooa, about 2 km. from the Tapuaooa shelter and 3 km. W of Mt. Ooumu, 10 Jul 1970, *G. W. Gillett* 2166 (US 3485393, P00636646); Toovii, épaulement SE du Mt. Tekao, 28 May 1984, *J. Florence* 6818 (P00636614); Tovii, ridge above L'Economie Rurale to Ooumu, 17 Jul 1988, *W. L. Wagner*, *D. Lorence*, *J. Florence*, and *S. Perlman* 6119 (MO4330721, PTBG008791, P00636608, US3206556); Toovii Plateau, trail behind L'Économie Rurale toward Ooumu peak, near summit crest, 16 Jul 1988, *S. Perlman* 10,127 (PTBG008286).
- A. *trinervia*** Kirk - NEW ZEALAND. North Island: Auckland Ecological Region, Awhitu Ecological District, Matakawau Reserve, south of Matakawau Road, 23 Apr 2001, *P. A. Aspin* s.n. (AK253912); Eastern Northland Ecological Region, Eastern Northland & Islands Ecological District, Orotore, south of Kaeo, 26 Aug 1965, *R. C. Cooper* s.n. (AK118465); North Auckland Land District, Franklin Co., Mauku, 5 May 1901, *H. Carse* s.n. (CHR328230); North Auckland, Hunua, Wairoa [Hunua] Falls, Dec 1868, *T. Kirk* s.n. (SP084519); North Auckland Land District, Mt. Wellington, Auckland, lava field, 4 Apr 1922, *H. Carse* s.n. (CHR292211); North Auckland, Auckland City, Mt. Wellington lavafield, Apr 1897, *D. Petrie* s.n. (SP084507). Little Barrier Island: North Auckland Land District, summit, 9 Mar 1962, *R. Melville* 6672 (CHR129112). Rangitoto Island: North Auckland Land District, 13 Feb 1965, *I. M. Morice* s.n. (CHR 566383).
- A. *waialealae*** Wawra - HAWAIIAN ISLANDS. Kaua'i: Alakai Swamp, transect across Sincock Bog, Bog 1, 13 Feb 1989, *S. Perlman* 10,641, *T. Pratt*, and *J. Lepson* (PTBG011706); Alakai Swamp, 1 Sep 1977, *P. van Royen* 11716 and *J. Davis*, *S. Perlman* (BISH419748A); Waialealae, 23 Sep 1909, *J. F. Rock* 5041 (BISH121147); Waialealae, 20 Oct 1911, *J. F. Rock* 8878 (BISH121150); Waimea District, Sincock Bog, Halehaha area, *S. Perlman* and *K. Wood* 3395 (PTBG019596).