

A new species of *Besleria* (Gesneriaceae) from the Serranía El Pinche (Cauca), southwestern Colombia

Jhon A. Sánchez-Taborda¹, Alejandro Zuluaga^{2,3}, Laura Clavijo^{3,4}

1 Fundación Ecohabitats, Popayán, Colombia **2** Departamento de Biología, Facultad de Ciencias, Universidad del Valle, Cali, Colombia **3** Grupo de Investigación Ecología y Diversidad Vegetal, Universidad del Valle, Cali, Colombia **4** Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Apartado 7495, Bogotá, Colombia

Corresponding author: Laura Clavijo (lvclavijor@unal.edu.co)

Academic editor: Ricardo Kriebel | Received 27 June 2020 | Accepted 7 September 2020 | Published 7 October 2020

Citation: Sánchez-Taborda JA, Zuluaga A, Clavijo L (2020) A new species of *Besleria* (Gesneriaceae) from the Serranía El Pinche (Cauca), southwestern Colombia. *PhytoKeys* 162: 71–80. <https://doi.org/10.3897/phytokeys.162.55891>

Abstract

A new species of the genus *Besleria* (Gesneriaceae), endemic to the department of Cauca, Colombia, is described and illustrated here. The new species, *Besleria santaclarensis* Clavijo & Sánchez-Taborda, was discovered in the Regional Protective Forest Reserve “Serranía El Pinche”, Cordillera Occidental of the Colombian Andes. *B. santaclarensis* is distinguished by the epedunculate inflorescences, usually in the leafless axils near the base, with up to eight orange flowers, and by the magenta calyx that covers 2/3 of the corolla.

Resumen

Una nueva especie del género *Besleria* (Gesneriaceae), endémica del departamento del Cauca, Colombia, se describe e ilustra aquí. La nueva especie, *Besleria santaclarensis* Clavijo & Sánchez-Taborda, se descubrió en la Reserva Forestal Protectora Regional “Serranía El Pinche”, Cordillera Occidental de los Andes colombianos. *B. santaclarensis* se distingue por las inflorescencias epedunculadas, usualmente en las axilas de nudos sin hojas ubicados hacia la base, con hasta ocho flores anaranjadas por inflorescencia y cáliz magenta que cubre 2/3 de la corola.

Keywords

Andes, Cauca, Choco biogeographic, Protective Forest Reserve, San Juan de Micay, taxonomy

Introduction

Besleria Plum. ex L. includes 160 species (Clark et al. 2020) of terrestrial herbs, shrubs and small trees that grow in the rainforest understory. It is strongly supported as mono-

phyletic in the tribe Beslerieae (Smith 2000; Roalson and Clark 2006; Clark et al. 2010). Biogeographic analyses of the family suggest that it diversified in the Andes 15 Mya (Perret et al. 2013; Roalson and Roberts 2016). *Besleria* is one of the largest and least known genera among the New World Gesneriaceae; it occurs in most Neotropical rainforests with the highest diversity in the tropical Andes (>100 species), followed by Central America (20 species) (Skog and Boggan 2007; Ferreira et al. 2016). Colombia is the country with the highest diversity of *Besleria*, with more than 80 species (Cortés 2013), followed by Peru with 35 (Kvist et al. 2005), Ecuador with about 20 (Skog and Kvist 1997) and Panama with 15 (Skog 1978). In Colombia, most of the species are found in the humid forests of the Andes and the Choco Biogeographic region, whereas few species grow in the Amazon (Cortés et al. 2017). In the Andes, the largest number of species is found in the Cordillera Occidental of Colombia. For example, *Besleria* is the third largest Gesneriaceae genus with 10 species out of the 96 recorded in northern Valle del Cauca (Clavijo et al. 2014). Likewise, Pedraza and Betancur (2015) recorded five species of *Besleria* out of the 70 Gesneriaceae species collected in the National Natural Park Orquídeas (Antioquia), north of the Cordillera Occidental.

The Serranía El Pinche in Argelia (Cauca) is located in the southern portion of the Cordillera Occidental of the Colombian Andes, and is part of the Munchique-Pinche corridor, a region known for high levels of biodiversity (Paz-B et al. 2018). Biogeographically, the Serranía El Pinche is more similar to the Nudo de los Pastos and the Colombian Massif than to the northern portions of the Cordillera, mainly due to the lower elevation Paramos on the Pacific slopes (Becking 1995). For the first settlers of the Serranía, the conservation of native ecosystems and sustainable rural development were paramount in this biodiverse region. Early settlers inspired in their children the importance of preserving their natural resources by creating alliances that would allow the proper management and establishment of a formal protected area. As a result of this initiative, the Regional Protective Forest Reserve “Serranía El Pinche” was created in 2008. The Reserve is part of the San Juan de Micay river basin that ranges in elevation from 1040 to 3744 meters above sea level; it has 7,256 ha, however, after its expansion it will have in total 11,930 ha. 90.4% of the reserve corresponds to primary forests and 9.6% to perturbed forests, crops and pastures. The Reserve “Serranía El Pinche” and the buffer zone include 913 households (Paz-B et al. 2018).

The upper San Juan de Micay river basin is a unique biodiverse area in the Pacific slopes of the Andes that still has large extensions of primary tropical rainforests (Becking 1995), despite high deforestation rates caused by agriculture and the presence of illicit crops. The preservation of protected areas led by local communities in regions suffering rapid deforestation is essential to conserve species and entire ecosystems. In fact, private and community-owned protected areas have been successful in preserving natural ecosystems in the northern Andes, where biological diversity is high (Joppa et al. 2008; Armenteras et al. 2009; Rodríguez et al. 2013). Therefore, following the legacy of the first naturalists and ecologists that explored La Serranía El Pinche, we aim to contribute to the floristic knowledge of this underexplored rainforest in south-western Colombia. The plant inventory and discovery of new taxa in this region will

provide important information to support the future expansion of the Reserve toward the coast, in the municipalities of Guapi and Timbiquí, achieving a broad elevation coverage from the Pacific coast to the paramos of the Andean highlands.

Methods

During a rapid ecological evaluation carried out in September of 2017 to characterize the vegetation and establish the baseline for expanding the Regional Protective Forest Reserve “Serranía El Pinche”, we discovered a new species of the genus *Besleria* which is described and illustrated here.

The collections of the new species were processed at the herbarium of the Universidad del Cauca (CAUP) and were deposited at the Colombian National Herbarium (COL), the herbarium of the Universidad del Valle (CUVC) and the Botanical Garden of Medellin (JAUM). Specimens of *Besleria* from COL, CUVC, and JAUM were studied to confirm the identity of the species. The photographs were taken with a Nikon camera model D 5300. For the general botanical terminology we followed Beentje (2010) and Moreno (1984).

Taxonomic treatment

Besleria santaclarensis Clavijo & Sánchez-Taborda, sp. nov.

urn:lsid:ipni.org:names:77211931-1

Figs 1, 2

Type. COLOMBIA, Cauca: Municipio Argelia, corregimiento Santa Clara, vereda Santa Clara, Reserva Forestal Protectora Regional “Serranía El Pinche”, flanco oriental de la Serranía que se desprende de la vertiente pacífica de la Cordillera Occidental, camino por el borde de la quebrada La Planada, 2°23.938'N, 77°18.863'W, 1620 m. 25 Sep 2017 (fl), Jhon Alexander Sánchez-Taborda, Álex Cortés, Andrea Borrero, Fernando Joaquín, Andrés Pérez, Erminson Buitrago, Julian Uetochambo 2552 (holotype: COL!; isotype: CUVC!).

Besleria santaclarensis is distinguished by epedunculate inflorescences, usually in the leafless axils near the base of the stem, with up to 8 glabrous orange flowers, and magenta glabrous calyx that covers 2/3 of the corolla.

Terrestrial subshrub, 1–1.5 m tall. Stem erect, sometimes scandent, branched, terete in cross section, 1.9–4 mm diam., subwoody, green, surface smooth to longitudinally striated, strigose toward the apex, trichomes < 1 mm long, unbranched, white; internodes 1.7–5.8 cm long. Leaves opposite, equal in a pair, sometimes subequal, older leaves usually caducous; petiole 1.6–4.9 cm long, slightly winged in cross-section, glabrate in basal leaves, strigose in apical leaves, trichomes < 1 mm long, white; blades elliptic, coriaceous, papery when dry, 6.4–17.6 × 2.8–6.8 cm, green and glossy adaxially, olive green




Figure 1. Drawing of *Besleria santaclarensis* Clavijo & Sánchez-Taborda, sp. nov. **A** habit **B** leaf detail, adaxial view **C** calyx, lateral view **D** flower, frontal view **E** corolla, lateral view **F** interior view of corolla showing stamens and staminode **G** gynoecium with anular nectary gland **H** fruit, lateral view. Illustration by Adriana Sanín, based on the holotype Sánchez-Taborda et al. 2552.

abaxially, apex acuminate, base cuneate, sometimes oblique, margin entire, glabrate on both surfaces, 7–10 pairs of secondary veins, obscure adaxially and raised abaxially with scarce and whitish indument, higher order of venation only evident abaxially. Inflorescence a pair-flowered cyme, axillary, usually in the leafless axils near the base of the stem, up to 8 flowers per inflorescence; peduncle and bracts absent. Pedicel oblique to perpendicular relative to the stem, 8.6–17.1 mm long, maroon, glabrous. Calyx magenta, mem-




Figure 2. *Besleria santaclarensis* Clavijo & Sánchez-Taborda, sp. nov. **A** branch showing leaf and inflorescence **B** inflorescence showing flower in lateral view **C** inflorescence showing an immature fruit **D** frontal view of the corolla. (Photographs by Jhon A. Sánchez-Taborda from the paratype).

branous, persistent in fruit, venation evident, reticulated; calyx lobes 5, 4 nearly equal, free, apex acute, base truncate to cordate, margin entire, glabrous on both surfaces, ventral and lateral lobes $9\text{--}13 \times 6.2\text{--}10.6$ mm, ovate, dorsal lobe $10.1\text{--}11.9 \times 4.3\text{--}5.3$ mm, oblong; Corolla zygomorphic, protandrous, slightly gibbous, thick, $14.4\text{--}15.4$ mm long, orange, glabrous; corolla tube slightly constricted above the base and then slightly ventricose ventrally, oblique relative to calyx, $12.3\text{--}13.5$ mm long, $5.7\text{--}6.2$ mm at its widest part, constriction above the base $3.7\text{--}4$ mm diam., base $5.4\text{--}6.8$ mm wide; throat $2.3\text{--}2.8$ mm diam., inner surface with glandular trichomes; corolla lobes 5, subequal, orange, spreading, ovate, apex rounded, margin entire, glabrous on both surfaces, ventral lobe $1.4\text{--}2.6 \times 2.4\text{--}3.1$ mm, lateral lobes $2.5\text{--}3.4 \times 2.4\text{--}3.9$ mm, dorsal lobes $0.8\text{--}1.1 \times 1.8\text{--}2.3$ mm. Androecium of 4 stamens, didynamous, included; filaments $7.1\text{--}10.1$ mm long, adnate to the corolla tube for $1.9\text{--}2.6$ mm, forming a sheath, glabrous, coiling after anthesis, staminode $4.9\text{--}5.2$ mm long; anthers reniform, $1.2\text{--}1.4 \times 1.1\text{--}1.7$ mm, coher-

ent by the apex and lateral walls, dehiscence by longitudinal slits. Gynoecium with an annular nectary gland, 0.5–0.7 mm tall, glabrous; ovary superior, 3–3.1 × 2.5–2.7 mm wide, rounded, glabrous; style included, 6.4–6.8 mm long, glabrous; stigma bilobed. Fruit a berry olive green; seeds numerous.

[Measurements from flowers during the mature gynoecium phase.]

Distribution and ecology. *Besleria santaclarensis* is endemic to Colombia and known only from the type locality in the municipality of Argelia (Cauca) in the Regional Protective Forest Reserve “Serranía El Pinche” and surrounding areas (Fig. 3). This species grows in the lower montane rainforest (Holdridge 1967) between 1300 and 1600 m above sea level, on the Pacific slope of the Cordillera Occidental of the Andes. It is frequent in open areas and in forest clearings, usually near to crops, pastures and remnants of secondary and riparian forests, whose canopies reach up to 35 m and 25 m, respectively. These forests have typical Andean floristic elements, represented by the genera *Saurauia* Willd (Actinidaceae), *Schefflera* J.R. Forst. and G. Forst. (Araliaceae), *Axinaea* Ruiz and Pav. (Melastomataceae), *Ladenbergia* Klotzsch (Rubiaceae), *Wettinia* Poepp, *Socratea* H. Karst and *Iriarteia* Ruiz and Pav. (Arecaceae), *Cyathea* Sm and *Alsophila* R. Br. (Cyatheaceae).

Phenology. *Besleria santaclarensis* has been found in flower in February and September and in fruit in February.




Figure 3. Distribution map of *Besleria santaclarensis* and *Besleria membranacea*.

Etymology. *Besleria santaclarensis* is named after the type locality, the Santa Clara village, in the municipality of Argelia, Cauca. The territory currently known as Argelia and areas surrounding El Pinche mountain range were initially inhabited by the *Gua-pios* indigenous people, until the arrival of the colonists who named it the Agua Clara path, due to the streams of crystalline waters present in the area. Afterwards, during evangelization, a Franciscan priest changed the name to Santa Clara because of the fertility of the land, which he called holy (Alveiro Bolaños, personal communication).

Preliminary conservation status. *Besleria santaclarensis* is only known from the Regional Protective Forest Reserve “Serranía El Pinche” and surrounding areas. Although the Reserve is a protected area, deforestation and soil degradation caused by agriculture and livestock systems, as well as the presence of illicit crops in the surrounding areas, may have a significant impact on the integrity of this narrow endemic species. *B. santaclarensis* is present in mature forests, but it is also frequent in open areas in remnants of secondary and riparian forests, suggesting it is resilient and capable of thriving in disturbed areas. A preliminary designation of Endangered (EN) category is provided, according to the criterium B2ab (IUCN 2012, 2017), based on the small population of the species, with an estimated area of occupancy of less than 20 km², and the continued habitat loss, due to the high concentration of illicit crops in the area (UNODC 2019).

Discussion

Besleria santaclarensis can be readily distinguished from its congeners by the epedunculate inflorescences, usually in the leafless axils near the base of the stem, with up to 8 glabrous orange flowers, and magenta glabrous calyx that covers 2/3 of the corolla. *B. santaclarensis* is similar to *B. membranacea* C.V. Morton in their foliage, their flowers with membranous and glabrous calyx, and their slightly gibbous and glabrous corolla tube. *B. membranacea* is endemic to Peru, particularly to the Amazon and the eastern foothills of the Andes, in the Departments of Loreto and San Martín, mainly in the basin of the Huallaga and Aguaytía rivers, between 260 and 880 m of elevation (Salinas and León 2006). *B. santaclarensis* differs from *B. membranacea* by shorter petioles (1.6–4.9 cm vs. (3.2–)6.5–11.6 cm), inflorescences with up to 8 flowers (vs. up to 4), shorter pedicels (8.6–17.1 mm vs. 13.1–20 mm), calyx base truncate to cordate (vs. never cordate), larger calyx lobes (9.0–13.0 × 4.3–10.6 mm vs. 5.8–9.3 × 1.6–4.7 mm) that cover up to 2/3 of the corolla (vs. covering up to 1/2 of the corolla), and corolla tubes shorter (14.4–15.4 mm vs. 14.9–18 mm long) and orange (vs. yellow and white). Additionally, *B. santaclarensis* is similar to an undescribed species collected in the Province of Zamora-Chinchipe, Ecuador, by Dr. John Clark (*Clark et al. 10815*). The two species share the inflorescences with several orange flowers with magenta ovate calyx lobes, but differ in that *B. santaclarensis* has glabrous calyx (vs. pilose) and slightly gibbous and glabrous corolla tube (vs. gibbous and pilose).

With the discovery and description of *B. santaclarensis* we aim to contribute to the floristic knowledge of this underexplored rainforest in southwestern Colombia, and to

provide new information to support the future expansion of the Reserve that will warrant the conservation of this and many other species.

Additional specimens examined (paratypes). COLOMBIA. Cauca: municipio Argelia, corregimiento Santa Clara, vereda El Pinche, zona aledaña a la Reserva Forestal Protectora Regional “Serranía El Pinche”, flanco oriental del Cerro Pinche, camino entre el Plateado y Guapi, en zona cercana a cultivos y pasturas con algunos relictos de bosques secundarios y riparios de la quebrada El Pinche, 2°28.809'N, 77°18.014'W, 1475 m. 11 Feb 2018 (fl, fr). *Jhon Alexander Sánchez-Taborda, Fernando Joaquín and Andrés Pérez* 2938 (JAUM).

Acknowledgments

We thank Liliana Patricia Paz-Betancourt and Luis Alfonso Ortega for their dedication and commitment to the conservation of the forests in the department of Cauca (Colombia), and for securing the resources to carry out the field expeditions; the Fund for Partnerships for Critical Ecosystems (CEPF) for financing the study; and Marnix Leonard Becking for managing the funds to make the first explorations to the high parts of the mountain range possible. We also want to thank the Santa Clara Serranía El Pinche Agro-Environmental Association, especially Alveiro Bolaños, community leader, for his dedication to the Reserve; to the field guides Fernando Joaquín and Andrés Perez for their company and teachings during the extensive field days; to the people of Argelia for allowing us to explore part of their territory and for taking the initiative to expand the Reserve; to the Cauca Regional Corporation (CRC) for providing logistical support during the expeditions; to the directors of the herbaria CAUP, COL, CUV, HUA and JAUM for granting us access to their collections; and to John L. Clark and Ricardo Kriebel for providing helpful reviews and information about collections of species similar to *Besleria santaclarensis*. Finally, we are grateful to Professor Alvaro Jose Negret (q.e.p.d) for opening the paths that would lead us to explore and conserve these wonderful mountains.

References

- Armenteras D, Rodríguez N, Retana J (2009) Are conservation strategies effective in avoiding the deforestation of the Colombian Guyana Shield. *Biological Conservation* 142(7): 1411–1419. <https://doi.org/10.1016/j.biocon.2009.02.002>
- Becking ML (1995) Sinecología, Diversidad y Conservación de la Vegetación Altoandina en el sur de la Cordillera Occidental, Argelia, Cauca, Colombia. Fundación Universitaria de Popayán, 400 pp.
- Beentje H (2010) The Kew Plant Glossary: An Illustrated Dictionary of Plant Identification Terms. Bibliovault OAI Repository, University of Chicago Press, 160 pp.
- Clark JL, Neill DA, Weber A, Gruhn JA, Katan T (2010) *Shuarua* (Gesneriaceae), an arborescent new genus from the Cordillera del Cóndor and Amazonian Ecuador. *Systematic Botany* 35(3): 662–674. <https://doi.org/10.1600/036364410792495917>

- Clark JL, Skog LE, Boggan JK, Ginzburg S (2020) Index to names of New World members of the Gesneriaceae (Subfamilies Sanangoideae and Gesnerioideae). *Rheede* 30: 190–256. <https://doi.org/10.22244/rheede.2020.30.01.14>
- Clavijo L, Marín OH, Zuluaga A (2014) Reservas Naturales Cerro El Inglés y Galápagos (Valle del Cauca): Gesneriaceae. Guide #606. Rapid Color Guides, Environmental and Conservation Programs, The Field Museum, Chicago. <https://fieldguides.fieldmuseum.org/guides/guide/606>
- Cortés O (2013) Sistemática y fitogeografía del complejo *Besleria aggregata* (Mart.) Hanst. (Gesneriaceae) en Colombia. Msc Tesis, Universidad del Valle, Colombia.
- Cortés O, Giraldo A, Torres A (2017) Two new species of *Besleria* (Gesneriaceae) from the department of Valle del Cauca, Colombia. *Phytotaxa* 332(2): 181–188. <https://doi.org/10.11646/phytotaxa.332.2.5>
- Ferreira G, Costa I, Araujo A, Hopkins M, Chautems A (2016) Three new species of *Besleria* (Gesneriaceae) from the southeastern Brazilian Atlantic rainforest. *Phytotaxa* 263(3): 270–278. <https://doi.org/10.11646/phytotaxa.263.3.7>
- Holdridge LR (1967) Life Zone Ecology. San José, Costa Rica, Tropical Science Center, 206 pp.
- IUCN (2012) IUCN Red List categories and criteria: Version 3.1, ed. 2. IUCN Species Survival Commission, Gland, Switzerland and Cambridge, UK.
- IUCN (2017) Guidelines for using the IUCN red list categories and criteria. Version 13. Prepared by the Standards and Petitions Subcommittee. <https://www.iucnredlist.org/resources/redlistguidelines> [Accessed May 25, 2020]
- Joppa LN, Loarie SR, Pimm SL (2008) On the protection of “protected areas”. *Proceedings of the National Academy of Sciences of the United States of America* 105(18): 6673–6678. <https://doi.org/10.1073/pnas.0802471105>
- Kvist L, Skog L, Amaya M, Salinas I (2005) Las Gesneriáceas de Perú. *Arnaldia* 12(1–2): 16–40.
- Moreno NP (1984) Glosario Botánico Ilustrado. Instituto Nacional de Investigaciones sobre Recursos Bióticos. Xalapa, Veracruz, 300 pp. <http://reservaeleden.org/plantasloc/>
- Paz-B LP, Ortega LA, Bolaños JA, Sánchez-Taborda JA (2018) Plan de manejo de la Reserva Forestal Protectora Serranía El Pinche 2019–2023. Acuerdo de Donación Conservación Internacional-Fundación Ecohabitats-Fondo de Alianzas para Ecosistemas Críticos (CEPF).
- Pedraza P, Betancur J (2015) Flora of Las Orquideas National Park: vascular plants of the Colombian Andes and Chocó. The New York Botanical Garden, Bronx, New York. <http://sweetgum.nybg.org/science/projects/orquideas/>
- Perret M, Chautems A, Araujo AO, Salamin N (2013) Temporal and spatial origin of Gesneriaceae in the New World inferred from plastid DNA sequences. *Botanical Journal of the Linnean Society* 171(1): 61–79. <https://doi.org/10.1111/j.1095-8339.2012.01303.x>
- Roalson EH, Clark JL (2006) Phylogenetic patterns of diversification in the Beslerieae (Gesneriaceae). In: Sharma AK, Sharma A (Eds) *Plant Genome: Biodiversity and Evolution*, Phanerogams 1C. Science Publishers, Inc., Enfield, New Hampshire, 251–268.
- Roalson EH, Roberts WR (2016) Distinct Processes Drive Diversification in Different Clades of Gesneriaceae. *Systematic Biology* 65(4): 662–684. <https://doi.org/10.1093/sysbio/syw012>
- Rodríguez N, Armenteras D, Retana J (2013) Effectiveness of protected areas in the Colombian Andes: Deforestation, fire and land-use changes. *Regional Environmental Change* 13(2): 423–435. <https://doi.org/10.1007/s10113-012-0356-8>

- Salinas I, León B (2006) Gesneriaceae endémicas del Perú. *Revista Peruana de Biología* 13(2): 359–365. <https://doi.org/10.15381/rpb.v13i2.1862>
- Skog LE (1978) Gesneriaceae. *Flora of Panama*. Family 175. Gesneriaceae. *Annals of the Missouri Botanical Garden* 65: 1–998. <https://doi.org/10.2307/2395351>
- Skog LE, Boggan JK (2007) World checklist of Gesneriaceae. Department of Botany, Smithsonian Institution, Washington. <http://botany.si.edu/Gesneriaceae/Checklist>
- Skog LE, Kvist LP (1997) Las Gesneriaceae de Ecuador. In: Valencia R, Balslev H (Eds) *Estudios Sobre Diversidad y Ecología de Plantas, Memorias del II Congreso Ecuatoriano de Botánica*. Pontificia Universidad Católica del Ecuador, Quito, 13–23.
- Smith JF (2000) A phylogenetic analysis of tribes Beslerieae and Napeantheae (Gesneriaceae) and evolution of fruit types: Parsimony and maximum likelihood analyses of *ndhF* sequences. *Systematic Botany* 25(1): 72–81. <https://doi.org/10.2307/2666674>
- UNODC (2019) Monitoreo de territorios afectados por cultivos ilícitos 2018. Oficina de las Naciones Unidas contra la Droga y el Delito, Gobierno de Colombia, 1–116. https://www.unodc.org/documents/colombia/2019/Agosto/Informe_de_Monitoreo_de_Territorios_Afectador_por_Cultivos_Illicitos_en_Colombia_2018_.pdf [accessed 06.09.2020]