

A new species of *Thinouia* (Paullinieae, Sapindaceae) from the Amazon and its phylogenetic placement

Herison Medeiros^{1,3}, Jenifer de Carvalho Lopes¹,
Pedro Acevedo-Rodríguez², Rafaela Campostrini Forzza³

1 Universidade de São Paulo, Instituto de Biociências, Departamento de Botânica, Rua do Matão, 277, 05508-090 São Paulo, SP, Brazil **2** Department of Botany, MRC-166 Smithsonian Institution, P.O. Box 37012, Washington D.C. 20013-7012, USA **3** Jardim Botânico do Rio de Janeiro, Pacheco Leão, 915, 22460-030, Rio de Janeiro, Rio de Janeiro, Brazil

Corresponding author: Herison Medeiros (medeiros.herison@gmail.com)

Academic editor: Marc Appelhans | Received 7 August 2020 | Accepted 2 October 2020 | Published 28 October 2020

Citation: Medeiros H, de Carvalho Lopes J, Acevedo-Rodríguez P, Forzza RC (2020) A new species of *Thinouia* (Paullinieae, Sapindaceae) from the Amazon and its phylogenetic placement. *PhytoKeys* 165: 115–126. <https://doi.org/10.3897/phytokeys.165.57341>

Abstract

Thinouia is a Neotropical genus of lianas with approximately 12 species and is the only genus in tribe Paullinieae with actinomorphic flowers. During a taxonomic revision of the genus and fieldwork in southwestern Amazonia, we found a new species that appears similar to *Thinouia trifoliata* (ex *Allosanthus*) because of its racemiform inflorescence. However, before describing the new species, we had to confirm that *Allosanthus* was congeneric with *Thinouia* so we could place the new species in the correct genus. The results of the phylogenetic analysis, based on molecular data (*trnL* intron and ITS sequences), show that *Allosanthus* should be included in *Thinouia*. Thus, the new taxon is described here as *Thinouia cazumbensis* sp. nov. The new species is described, illustrated and phylogenetic trees showing relationships within supertribe Paulliniodeae and *Thinouia* and the congeneric *Allosanthus* are given.

Keywords

Allosanthus, Amazonia, Brazil, lianas, neotropical biodiversity, Paulliniodeae, Paullinieae, Sapindales, Sapindaceae, taxonomy, *Thinouia*

Introduction

Thinouia is a neotropical genus of lianas that includes around 12 species, of which *T. myriantha* Planch. & Triana is widely distributed, including records from Mexico, Central America and northern South America (Ferrucci and Somner 2008; Acevedo-Rodríguez et al. 2011). The remaining species are distributed in Brazil, Bolivia, Paraguay and Peru, except for *T. tomentosa* Standl. which is restricted to Mexico, Belize and Guatemala. Most *Thinouia* species occur in rainforest; a few species occur in savannah (BFG 2015).

Thinouia was proposed by Triana and Planchon (1862). It is characterised by the presence of umbelliform and racemiform thyrses, actinomorphic flowers with marginal or bifid petal appendages, an annular disc and schizocarpic fruits that split into three mericarps, each with a distal wing (Ferrucci and Somner 2008; Acevedo-Rodríguez et al. 2017).

Molecular phylogenetic studies show that *Thinouia* is a monophyletic group in Sapindaceae. In the most recent phylogenetic study, tribe Paullinieae (i.e. *Cardiospermum*, *Lophostigma*, *Paullinia*, *Serjania*, *Thinouia* and *Urvillea*) is a well-supported clade with *Thinouia* sister to the remaining genera (Acevedo-Rodríguez et al. 2017). In the same work, the monospecific genus *Allosanthus* (*A. trifoliatus* Radlk.) was maintained as a synonym of *Thinouia*, based on morphological characters. The only differentiating character (i.e. a racemiform inflorescence) was not considered worthy of generic recognition (Acevedo-Rodríguez et al. 2011, 2017).

During a taxonomic revision of the genus and fieldwork in south-western Amazonia, we found a new species of *Thinouia* that is similar to *Thinouia trifoliata* (Radlk.) Acev.-Rodr. & Ferrucci because of its racemiform inflorescence. Since we now have high-quality DNA material for the taxa previously assigned to *Allosanthus*, we re-analysed the placement of *Allosanthus* within *Thinouia* and further tested the monophyly of *Thinouia s.l.*, which revealed the correct position of the new species.

Material and methods

Plant material

We collected the new species in Reserva Extrativista do Cazumbá-Iracema in Sena Madureira, Acre, Brazil. The collection was pressed and dried for vouchers, leaves were collected in silica gel for DNA extraction and reproductive structures were fixed in 70% alcohol for morphological analyses, which were performed using a stereomicroscope. The morphological structures were described using the terminology in Radford et al. (1974) and Weberling (1989). The herbarium abbreviations cited in the text follow Thiers (2020, cont. upd.).

Phylogenetic analysis

The phylogenetic analysis included the same taxa and molecular markers of Acevedo-Rodríguez et al. (2017), 93 taxa, plastid marker trnL intron and nuclear ribosomal internal transcribed spacer, ITS. Six samples (*Allosanthus* sp., *Allosanthus trifoliatus*, *Thinouia mucronata*, *T. myriantha*, *T. obliqua* and *Thinouia* sp.), including the new species, were added to the analysis, using the same molecular markers. For these additional taxa, approximately 60 mg of leaf tissue were pulverised with Tissuelyzer (Qia- gen, Duesseldorf, Germany) for 3 min at 60 hz. The DNA extraction used the DNA NucleoSpin Plant II kit (Machery-Nagel, GmbH & Co. KG, Dueren, Germany) following the manufacturer's protocol. Primers and the PCR amplification were used, as described in Acevedo-Rodríguez et al. (2017). Products were purified and sequenced by Macrogen (Seoul, South Korea). All sequences, vouchers and GenBank accession numbers are summarised in Appendix I.

The alignments were performed using MAFFT (Katoh et al. 2002) using the default parameters implemented in Geneious 2020.0.5 (Kearse et al. 2012). Poorly-aligned regions were removed and adjusted manually. We used jModelTest 2.0 (Guindon et al. 2010; Darriba et al. 2012) and the Akaike Information Criterion (AIC) to select the best-fit model of nucleotide substitution for each dataset. The GTR+I+G was selected as the best model for the ITS dataset, whereas the GTR+G was selected as the best model for the trnL dataset. Bayesian Inference (BI) analyses were conducted using MrBayes 3.2.2 (Ronquist et al. 2012) in the online CIPRES Science Gateway interface (Miller et al. 2015) with four Markov Chain Monte Carlo (MCMC) runs using a random starting tree and 10 million generations, with a sampling frequency of one every 1000 generations. We used Tracer 1.7 (Rambaut et al. 2018) to check for convergence of the MCMC and to check for stationarity. We discarded 25% of the trees as burn-in.

Phylogenetic trees were plotted and built inside the R environment (R Core Team 2020), version 3.6.2, using the packages ggplot2 (Wickham et al. 2020), ggtree (Yu et al. 2017; Yu and Lam 2020) and cowplot (Wilke 2019).

Results

Phylogenetic results

The ITS dataset included 99 terminals and 876 bp, the trnL dataset included 99 terminals and 727 bp and the combined dataset included 99 terminals and 1604 bp. Phylogenetic trees from the analyses of the combined dataset showed high posterior probability values (PP > 0.8). Only the topology from the combined analysis is described here (Fig. 1). Separate analyses of each locus did not reveal any strong groupings that would indicate incongruences.




Figure 1. **A** bayesian 50% majority-rule consensus tree from a Bayesian analysis of the combined, two-marker dataset for Paulliniodae and outgroups **B** relationships of *Thinouia* and the congeneric *Allosanthus* [= *Thinouia trifoliata* (Radlk.) Acev.-Rodr. & Ferrucci], including the newly-described *Thinouia cazumbensis* sp. nov. Bayesian posterior probability values are indicated above the branches.

Supertribe Paulliniodae is strongly supported as monophyletic (Fig. 1 A, PP = 1.0). The tribe Paullinieae is also strongly supported as monophyletic (PP = 1.0) and the genus *Thinouia* (including *Allosanthus*) is recovered as the clade, sister to the remaining genera of the tribe Paullinieae (PP = 1.0). *Thinouia* species are grouped in two main clades that are in a polytomy with the new species *Thinouia cazumbensis*. The first clade (PP = 1.0) includes *Thinouia obliqua*, *T. mucronata*, *T. restingae* and *T. cf. mucronata* species. The second one (PP = 0.8) includes *Thinouia* sp., *T. myriantha* and *T. trifoliata* (= *Allosanthus trifoliatus* Radlk.) (Fig. 1).

Taxonomic treatment

Thinouia cazumbensis Medeiros, sp. nov.

urn:lsid:ipni.org:names:77212573-1

Figure 2

Diagnosis. The new species differs from *Thinouia trifoliata* by the 5-lobed floral disc, fruits with trichomes and basal petal appendages smaller than the petals, versus annular disc, glabrous fruits and marginal petal appendages larger than the petals.

Type. Brazil. Acre. Sena Madureira. Reserva Extrativista do Cazumbá-Iracema, Núcleo Cazumbá, castanhal coletivo, floresta ombrófila aberta com bambu, 9°8'30"S, 68°56'23"W, 20 Jul 2018, H. Medeiros, M. Silveira & E.M. Soares 3401, (holotype RB!; isotypes: INPA!, SPF!, UFACPZ!, US!).

Description. Tendrilled liana 6–8 m long; stem puberulent, with yellowish to whitish indumentum, lenticellate; cross section simple, cylindrical. Leaves trifoliol-




Figure 2. *Thinouia cazumbensis* **A** fruiting branch **B** detail of leaf, abaxial view **C** racemiform inflorescence with a pair of basal tendrils **D** detail of inflorescence (cincinnus) **E** flower with removed petals showing a 5-lobed nectary disc **F** detail of fruit **G** infructescence (**A–G**) from H. Medeiros 3401 (RB). Photos by H. Medeiros.

ate; stipules ca. 2 mm long, hirsute-tomentose, linear triangular to lanceolate; petiole 2–8.5 cm long, canaliculate; petiolules of lateral leaflets 0.2–0.8 cm long; leaflets 7–14 × 3–9 cm, oblong to ovate- rhomboidal, apex acute, mucronate, margins entire to dentate-serrate, with 2–4 teeth reduced to inconspicuous glands, ciliate, base trunc-




Figure 3. Geographic distribution of *Thinouia cazumbensis*.

cate, rounded to obtuse, sometimes cuneate on the distal leaflet, glabrous on both surfaces, domatia sometimes in the axils of abaxial secondary veins. Thyrses axillary, racemiform, ca. 8.5–16 cm long, peduncle 1.1–2.8 cm long, rachis of 7.5–16 cm long; numerous cincinnus, sessile. Flowers ca. 2 mm long, pedicel ca. 0.5 mm long; sepals 5, ca. 1 mm long, fused at the base, lobes ovate, acute, glabrous and with prominent veins on the internal surface, external surface villous; petals 5, ca. 1.5 mm long, obovate, obtuse, clawed, villous on the central part and margins, the rest glabrous; petal appendages rudimentary, bifid, smaller than the petals, basally adnate, villous; nectary disc glabrous, 5-lobed, lobes ca. 1 mm long; staminate flower: stamens 8, 1.5 mm long, filaments villous for more than half of their length, anthers glabrous, pistillode ca. 1.5 mm long; pistillate flower: staminodes ca. 1 mm long, pistil ca. 1.5 mm long, style 0.5 mm long, with 3 stigmas, ovary ca. 1 mm long. Fruits ovate, chartaceous, 5–5.5 × 2–2.3 cm; cocci slightly inflated, 1.2–1.4 × 1.1–1.4 cm, including the ca. 2–3 mm long stipe constricted at junction with wing; epicarp densely strigose (simple trichomes of same length) on cocci, strigose on wings; endocarp glabrous. Seeds trigonous ovoid, ca. 6 × 4 mm, basally attached, glabrous, mature embryo not observed.

Thinouia cazumbensis is differentiated from most species of *Thinouia* by the thyrses axillary, racemiform (Fig. 2A, C) and the 5-lobed nectary disc, a character that is unique and for the first time recorded in the genus (Fig. 2E). The lobed nectary disc

within *Thinouia* should be further investigated through morpho-anatomical studies to understand how nectaries evolved within the genus.

Distribution and ecology. *Thinouia cazumbensis* is known only from the Reserva Extrativista do Cazumbá-Iracema (Fig. 3) where it is an infrequent liana that reaches the canopy of the open rainforest with abundant bamboo (*Guadua* spp.) (Silveira 2005).

Phenology. Collected in flower and fruit during July.

Etymology. The epithet *cazumbensis* refers to Reserva Extrativista do Cazumbá-Iracema, where the species was collected. In the 1980s, local rubber tappers and extrativists fought against the area becoming a rural settlement and on 19 September 2002 succeeded in getting the area designated as a conservation unit (ICMBio 2007). Situated in the State of Acre between the municipalities of Sena Madureira and Manoel Urbano, the Reserva Extrativista do Cazumbá-Iracema covers an area of 750,794.70 hectares of the Western Amazon Corridor, one of the seven major ecological corridors proposed for Brazil (Ricardo and Lima 2004).

Conservation status. The species is only known from one locality in Acre and is categorised as Data Deficient (DD) according to IUCN (2019). Further field studies are needed to evaluate its conservation status more accurately.

Discussion

The broader relationships that we recovered within supertribe Paulliniodae largely agree with those in Acevedo-Rodríguez et al. (2017). Additionally, with the inclusion of new sequences of *Thinouia* in this study merged with sequence data from Acevedo-Rodríguez et al. (2017), our results recovered the same clades in tribe Paullinieae, where *Thinouia* forms a clade that is the earliest diverging lineage. Therefore, our phylogenetic results reinforce including *Allosanthus* in *Thinouia* as proposed by Acevedo-Rodriguez et al. (2011), based on morphological characters. The only differentiating morphological character (i.e. the racemiform inflorescence) was not considered worthy of generic recognition by Acevedo-Rodriguez et al. (2011, 2017) and the molecular data in the present study corroborate this conclusion. The position of the new species as a member of *Thinouia* is strongly supported albeit its relationship to other species is not fully resolved, perhaps because of our limited sampling of *Thinouia* or because only two markers have been sequenced.

Conclusion

Thinouia cazumbensis is supported as a distinct taxon, based on morphological and molecular sequence data. Its position within the genus is still undetermined, highlighting the need for in-depth taxonomic studies on this genus. Ongoing systematics studies, based on molecular and morphological analyses of *Thinouia*, should provide additional insights into the evolution and biogeographic history of this neotropical genus (H. Medeiros et al. in prep.).

Acknowledgements

This paper was possible through the financial support from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) to HM, by Fundação de Amparo à Pesquisa de São Paulo (FAPESP; 2018/11272-5) to JCL and by research grants from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq; 303420/2016-2) and Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ; E-26/202.778/2018) to RCF. The authors thank Jéssica Francisco and Ricardo Perdiz for assistance with the preparation of the phylogeny and comments on an earlier version of the manuscript, Genise Somner, Maria S. Ferruci, Marc Appelhans and the anonymous reviewer for their numerous suggestions that greatly improved this manuscript and Mark T. Strong (US) for proofreading the manuscript. We also thank ICMBio in Sena Madureira for support during field work.

References

- Acevedo-Rodríguez P, van Welzen PC, Adema F, van der Ham RWJM (2011) *Sapindaceae*. In: Kubitzki K (Ed.) The families and genera of vascular plants. Eudicots: Sapindales, Cucurbitales, Myrtaceae. Springer, Berlin, 357–407. https://doi.org/10.1007/978-3-642-14397-7_17
- Acevedo-Rodríguez P, Wurdack KJ, Ferrucci MS, Johnson G, Dias P, Coelho RG, Somner GV, Steinmann VW, Zimmer EA, Strong MT (2017) Generic relationships and classification of tribe Paulliniaeae (Sapindaceae) with a new concept of supertribe Paulliniodeae. Systematic Botany 42(1): 96–114. <https://doi.org/10.1600/036364417X694926>
- BFG (2015) Growing knowledge: An overview of seed plant diversity in Brazil. Rodriguésia 66: 1085–1113. <https://doi.org/10.1590/2175-7860201566411>
- Darriba D, Taboada GL, Doallo R, Posada D (2012) jModelTest 2: More models, new heuristics and parallel computing. Nature Methods 9(8): 1–772. <https://doi.org/10.1038/nmeth.2109>
- Ferrucci MS, Somner GV (2008) *Thinouia restingae* (Sapindaceae: Paulliniaeae), una nueva especie de Brasil. Brittonia 60(4): 371–376. <https://doi.org/10.1007/s12228-008-9041-0>
- Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New algorithms and methods to estimate maximum likelihood phylogenies: Assessing the performance of PhyML 3.0. Systematic Biology 59(3): 307–321. <https://doi.org/10.1093/sysbio/syq010>
- ICMBio [Instituto Chico Mendes de Conservação da Biodiversidade] (2007) Plano de Manejo da Reserva Extrativista do Cazumbá-Iracema. Sena Madureira. <https://www.icmbio.gov.br/portal/resex-do-cazumba-iracema>
- IUCN (2019) Guidelines for using the IUCN Red List Categories and Criteria. Version 14. Standards and Petitions Committee. <http://www.iucnredlist.org/documents/RedList-Guidelines.pdf> [accessed 14.09.2020]
- Katoh K, Misawa K, Kuma KI, Miyata T (2002) MAFFT: A novel method for rapid multiple sequence alignment based on fast Fourier transform. Nucleic Acids Research 30(14): 3059–3066. <https://doi.org/10.1093/nar/gkf436>

- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thaler T, Ashton B, Meintjes P, Drummond A (2012) Geneious basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* (Oxford, England) 28(12): 1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>
- Miller MA, Schwartz T, Pickett BE, He S, Klem EB, Scheuermann RH, Passarotti M, Kaufman S, O'Leary MA (2015) A RESTful API for access to phylogenetic tools via the CIPRES Science Gateway. *Evolutionary Bioinformatics Online* 11: 43–48. <https://doi.org/10.4137/EBO.S21501>
- R Core Team (2020) R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>
- Radford AE, Dickison WC, Massey JR, Bell CR (1974) Vascular Plant Systematics. Harper & Row Publishers, New York.
- Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA (2018) Posterior summarisation in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* 67(5): 901–904. <https://doi.org/10.1093/sysbio/syy032>
- Ricardo MM, Lima RX (2004) Corredores ecológicos e suas políticas de implementação. Congresso Brasileiro de Unidades de Conservação, 4., Anais (Vol. 2). Seminários. FBPN e Rede Nacional Pró-Unidades de Conservação, Curitiba.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Silveira M (2005) A floresta aberta com bambu no sudoeste da Amazônia: padrões e processos em múltiplas escalas. EDUFAC, Rio Branco, 157 pp.
- Thiers B (2020) [continuously updated] Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih/> [accessed 14.09.2020]
- Triana MM, Planchon JE (1862) Prodromus Florae Novo-Granatensis. Annales des sciences naturelles Ser. 4. Botanique 18: 258–381.
- Weberling F (1989) Morphology of Flowers and Inflorescences. Press Syndicate of the University of Cambridge, Cambridge.
- Wickham H, Chang W, Henry L, Pedersen TL, Takahashi K, Wilke C, Woo K, Yutani H, Dunnington D (2020) Ggplot2: Create elegant data visualizations using the grammar of graphics. <https://CRAN.R-project.org/package=ggplot2>
- Wilke CO (2019) Cowplot: Streamlined plot theme and plot annotations for 'Ggplot2'. <https://CRAN.R-project.org/package=cowplot>
- Yu G, Lam TT (2020) Ggtree: An R package for visualization of tree and annotation data. <https://yulab-smu.github.io/treedata-book/>
- Yu G, Smith D, Zhu H, Guan Y, Lam TT (2017) Ggtree: An R package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods in Ecology and Evolution* 8(1): 28–36. <https://doi.org/10.1111/2041-210X.12628>

Appendix I

Voucher and GenBank information for the taxa included in the phylogenetic analyses. Listed as: taxon, collection, herbarium, place of origin and GenBank accession numbers (ITS, trnL intron). Herbarium acronyms follow Index Herbariorum (Thiers, continuously updated).

Allophylastrum frutescens Acev.-Rodr., Lima 812 (K), Brazil, KX584885, KX584982. *Allophylus abyssinicus* (Hochst.) Radlk., Desissa & Binggeli DD-318 (MO), Ethiopia, KX584886, KX584983. *Allophylus africanus* P. Beauv., Balkwill 4206 (MO), South Africa, KX584887, KX584984. *Allophylus arboreus* Choux, Wohlhauser & Stieffle 60072 (MO), Madagascar, KX584888, KX584985. *Allophylus bicruris* Radlk., Barthelat 828 (MO), Mayotte, KX584889, KX584986. *Allophylus bojerianus* (Cambess.) Blume, Ratovoson 961 (MO), Madagascar, KX584890, KX584987. *Allophylus chaunostachys* Gilg, Mwangoko 729 (MO), Tanzania, KX584891, KX584988. *Allophylus chirindensis* Baker f., Hizza 26 (MO), Tanzania, KX584892, KX584989. *Allophylus cominia* Sw., Acevedo-Rodríguez 12216 (US), Mexico, KX584893, KX584990. *Allophylus crassinervis* Radlk., Acevedo-Rodríguez s.n. (no voucher), Puerto Rico, KX584894, KX584991. *Allophylus decipiens* (E. Mey.) Radlk., Phillipson 4194 (MO), South Africa, KX584895, KX584992. *Allophylus gardineri* Summerh., Pignal 1834 (MO), Mayotte, KX584897, KX584994. *Allophylus hirtellus* (Hook. f.) Radlk., Cheek 5059 (?), KX584898, KX584995. *Allophylus pervillei* Blume, Hoffmann 399 (MO), Mayotte, KX584899, KX584996. *Allophylus poungouensis* Pellegr., McPherson 16109 (MO), Gabon, KX584900, KX584997. *Allophylus puberulus* (Cambess.) Radlk., Somner 1069 (US), Brazil, KX584901, KX584998. *Allophylus racemosus* Sw., Acevedo-Rodríguez 12180 (US), Mexico, KX584902, KX584999. *Allophylus rubifolius* (A. Rich.) Engl., Kuchar 23357 (MO), Tanzania, KX584903, KX585000. *Allophylus* sp., Acevedo-Rodríguez 14847 (NY), Brazil, KX584904, KX585001. *Athyana weinmanniifolia* (Griseb.) Radlk., Acevedo-Rodríguez 11166 (US), Bolivia, KX584906, KX585003. *Balsas guerrerensis* Cruz Durán & K. Vega, Vega Flores 1318 (US), Mexico, KX584908, KX585005. *Bridgesia incisifolia* Cambess., Landrum 9824 (NY), Chile, KX584909, KX585006. *Cardiospermum corindum* L., Harder & Bringham 3495 (MO), Zambia, KX584912, KX585007. *Cardiospermum cuchujaquense* Ferrucci & Acev.-Rodr., Van Devender 92-1012 (ARIZ), Mexico, KX584914, KX585008. *Cardiospermum grandiflorum* Sw., ATBP 603 (MO), Uganda, KX584915, KX585009. *Cardiospermum grandiflorum* Sw., Gildenhuys H1 (?), Hawaii, KM062277, KM062362. *Cardiospermum heringeri* Ferrucci, Urdampilleta 437 (US), Brazil, KX584917, KX585010. *Cardiospermum urvilleoides* (Radlk.) Ferrucci, Urdampilleta 425 (US), Brazil, KX584922, KX585013. *Chimborazoa lachnocarpa* (Radlk.) H.T. Beck, Wiggins 11060 (US), Ecuador, KX584923, KX585014. *Diatenopteryx sorbifolia* Radlk., Zardini 43371 (MO), Paraguay, EU720534, EU721303. *Dictyoneura obtusa* Blume, Edwards KE142 (JCT), Australia, EU720428, EU721187. *Diploglottis campbellii* Cheel, Chase 2048 (K), Australia, EU720457, EU721224. *Guindilia dissecta* (Covas & Burkart) Hunz., Ferrucci 2928 (CTES), Argentina, KX584926, KX585017. *Guioa villosa* Radlk., McPherson

18040 (MO), New Caledonia, EU720544, EU721314. *Haplocoelum inoploeum* Radlk., Lap 117 (?), FJ514259, FJ514265. *Houssyanthus binternatus* (Weath) Rzed. & Calderón, Catalán & Terán 837 (MO), Mexico, KX584927, KX585018. *Houssyanthus incanus* (Radlk.) Ferrucci, Ferrucci 2710 (CTES), Argentina, KX584928, KX585019. *Jagera javanica* (Blume) Kalkman, Chase 2130 (K), Bogor, EU721236, EU720468. *Lepisanthes senegalensis* (Poir.) Leenhardt, Callmander 627 (MO), Madagascar, EU720492, U72126. *Lophostigma plumosum* Radlk., Acevedo-Rodríguez 6554 (US), Bolivia, KX584929, KX585020. *Macphersonia gracilis* O. Hoffm., Rabenantoandro 1081 (MO), Madagascar, EU720550, EU721320. *Matayba guianensis* Aubl., Acevedo-Rodríguez 12342 (US), French Guiana, EU720527, EU721294. *Melicoccus lepidopetalus* Radlk., Acevedo-Rodríguez 11128 (US), Bolivia, EU720443, EU721206. *Paullinia clathrata* Radlk., Acevedo-Rodríguez 14305 (US), Peru, KX584930, KX585021. *Paullinia coriacea* Casar., Somner 1070 (RBR), Brazil, KX584931, KX585022. *Paullinia cuneata* Radlk., Acevedo-Rodríguez 14255 (US), Peru, KX584932, KX585023. *Paullinia elegans* Cambess., Acevedo-Rodríguez 14976 (US), Brazil, KX584933, KX585024. *Paullinia hystrix* Radlk., Acevedo-Rodríguez 14417 (US), Peru, KX584934, KX585025. *Paullinia imberbis* Radlk., Schunke Vigo 14928 (US), Peru, KX584935, KX585026. *Paullinia olivacea* Radlk., Schunke Vigo 16002 (US), Peru, KX584936, KX585027. *Paullinia pinnata* L., Acevedo-Rodríguez 11088 (US), French Guiana, KX584937, KX585028. *Paullinia prevostiana* Acev.-Rodr., Acevedo-Rodríguez 11113 (US), French Guiana, KX584938, KX585029. *Paullinia rubiginosa* Cambess., Thomas 12995 (US), Brazil, KX584939, KX585030. *Paullinia spicata* Benth., Acevedo-Rodríguez 12344 (US), French Guiana, KX584941, KX585032. *Paullinia stellata* Radlk., Acevedo-Rodríguez 14958 (US), Brazil, KX584942, KX585033. *Paullinia xestophylla* Radlk., Hoffman 5955 (US), Suriname, KX584943, KX585034. *Plagioscyphus unijugatus* Capuron, Buerki 145 (NEU), Madagascar, EU720475, EU721245. *Sapindus oligophyllus* Merr. & Chun, How 70627 (US), China, KX584944, KX585035. *Serjania altissima* (Poepp.) Radlk., Acevedo-Rodríguez 14953 (US), Brazil, KX584945, KX585036. *Serjania ampelopsis* Planch. & Lind., Acevedo-Rodríguez 11181 (US), Bolivia, KX584946, KX585037. *Serjania caracasana* (Jacq.) Willd., Acevedo-Rodríguez 15107 (US), Mexico, KX584947, KX585038. *Serjania cf. caracasana* (Jacq.) Willd., Acevedo-Rodríguez 3483 (US), Guyana, KX584948, KX585039. *Serjania clematidifolia* Cambess., Somner 1078 (RBR), Brazil, KX584949, KX585040. *Serjania communis* Cambess., Somner 1334 (US), Brazil, KX584950, KX585041. *Serjania cuspidata* Cambess., Somner 1400 (US), Brazil, KX584951, KX585042. *Serjania emarginata* Kunth, Acevedo-Rodríguez 15135 (US), Mexico, KX584954, KX585043. *Serjania erythrocaulis* Acev.-Rodr. & Somner, Acevedo-Rodríguez 3729 (US), Brazil, KX584955, KX585044. *Serjania eucardia* Radlk., Somner 1072 (RBR), Brazil, KX584956, KX585045. *Serjania fuscifolia* Radlk., Somner 1455 (RBR), Brazil, KX584957, KX585046. *Serjania ichthyoctona* Radlk., Somner 1081 (RBR), Brazil, KX584960, KX585048. *Serjania lethalis* St. Hil., Roque 1860 (ALCB), Brazil, KX584961, KX585049. *Serjania lethalis* St. Hil., Somner 1381 (RBR), Brazil, KX584962, KX585050. *Serjania marginata* Casar., Acevedo-Rodríguez 11131 (US), Bolivia, KX584963, KX585051. *Serjania mexicana* (L.) Willd., Acevedo-

Rodríguez 12014 (US), Jamaica, KX584965, KX585052. *Serjania mexicana* (L.) Willd., Acevedo-Rodríguez 15080 (US), Mexico, KX584966, KX585053. *Serjania paniculata* Kunth, Acevedo-Rodríguez 15143 (US), Mexico, KX584967, KX585054. *Serjania perulacea* Radlk., Acevedo-Rodríguez 11134 (US), Bolivia, KX584968, KX585055. *Serjania unguiculata* Radlk., Acevedo-Rodríguez 15081 (US), Mexico, KX584969, KX585056. *Serjania yucatanensis* Standl., Acevedo-Rodríguez 12183 (US), Mexico, KX584970, KX585057. *Talisia nervosa* Radlk., Pennington 628 (MO), ~, EU720474, EU721244. *Talisia obovata* A.C. Sm., Lombello 13 (MO), Brazil, EU720485, EU721255. *Thinouia cazumbensis* sp. nov., Medeiros 3401 (RB) Brazil, MT853074, MT847016. *Thinouia mucronata* Radlk., Keller 6919 (US), Argentina, KX584971, KX585058. *Thinouia cf. mucronata* Radlk., Medeiros 3800 (RB) Brazil, MT853076, MT847018. *Thinouia myriantha* Radlk., Torke 2024 (HSTM), Brazil, MT853071, MT847013. *Thinouia obliqua* Radlk., Medeiros 3793 (RB) Brazil, MT853075, MT847017. *Thinouia* sp., Medeiros 2193 (RB), Brazil, MT853072, MT847014. *Thinouia restingae* Ferrucci & Somner, Somner 1074 (RBR), Brazil, KX584972, KX585060. *Thinouia trifoliata* (Radlk.) Acev.-Rodr. & Ferrucci, Medeiros 3331 (RB), Brazil, MT853073, MT847015. *Thouinia acuminata* S. Watson, Liston 633-2, —, EU720478, EU721249. *Thouinia villosa* DC., Hall 825 (US), Mexico, KX584975, KX585062. *Tristiropsis acutangula* Radlk., Chase 1358 (K), Bogor, EU720453, EU721220. *Urvillea chacoensis* Hunz., Acevedo-Rodríguez 11133 (US), Bolivia, KX584976, KX585063. *Urvillea chacoensis* Hunz., Keller 6834 (US), Argentina, KX584977, KX585064. *Urvillea pterocarpa* (Radlk.) Acev.-Rodr. & Ferrucci, Urdampilleta 321 (US), Brazil, KX585012, KX584921. *Urvillea rufescens* Cambess., Somner 1073 (RBR), Brazil, KX584978, KX585065. *Urvillea ulmacea* Kunth, Acevedo-Rodríguez 15145 (US), Mexico, KX584979, KX585066. *Urvillea ulmacea* Kunth, Reyes-García 5585 (MO), Mexico, KX584980, KX585067. *Vouarana guianensis* Aubl., Acevedo-Rodríguez 5031 (US), French Guiana, KX584981, KX585068.