

An updated checklist and key to the open-panicled species of *Poa* L. (Poaceae) in Peru including three new species, *Poa ramoniana*, *Poa tayacajaensis*, and *Poa urubambensis*

Steven P. Sylvester^{1,2,*}, Robert J. Soreng^{3,*}, Paul M. Peterson³,
Mitsy D.P.V. Sylvester^{1,4}

1 Institute of Systematic and Evolutionary Botany, University of Zurich, Zollikerstrasse 107, 8008 Zürich, Switzerland **2** Department of Geography, Philipps-Universität Marburg, Deutschhausstraße 10, D-35032 Marburg, Germany **3** Department of Botany, National Museum of Natural History, Smithsonian Institution Washington, DC, 20013-7012, USA **4** Universidad Nacional de San Antonio Abad del Cusco, Avenida de la Cultura 733, Cusco, Perú

Corresponding author: Steven P. Sylvester (steven_sylvester@hotmail.com)

Academic editor: R.P. Oliveira | Received 11 November 2015 | Accepted 3 June 2016 | Published 30 June 2016

Citation: Sylvester SP, Soreng RJ, Peterson PM, Mitsy Sylvester MDPV (2016) An updated checklist and key to the open-panicled species of *Poa* L. (Poaceae) in Peru including three new species, *Poa ramoniana*, *Poa tayacajaensis*, and *Poa urubambensis*. *PhytoKeys* 65: 57–90. doi: 10.3897/phytokeys.65.7024

Abstract

We provide an updated checklist and key to the 30 *Poa* species with open panicles from Peru which includes previously circumscribed *Dissanthelium* and *Aphanelytrum* species, new taxon records, and three undescribed species. *Poa compressa*, *P. grisebachii*, and *P. leioclada* are recorded from Peru for the first time. A number of species are placed in synonymy: *Poa carazensis*, *P. ferreyrae* and *P. tovarii* are synonymized under the name *P. fibrifera*; *Poa adusta* (tentatively) and *P. pilgeri* are synonymized under *P. candamoana*; *Poa superata* is synonymized under *P. grisebachii*; and *Poa paramoensis* is synonymized under *P. huancavelicae*. Included within this treatment are three new species, *Poa ramoniana*, *P. tayacajaensis* and *P. urubambensis*, which are described and illustrated. *Poa ramoniana*, found growing near lakes in high elevation Puna grasslands of Junín, is similar to a small form of *P. glaberrima*, but differs in having rhizomes and growing to only 5 cm tall. *Poa tayacajaensis*, found from shrublands on Andean slopes of Huancavelica and Huánuco, bears similarities to *Poa aequatoriensis* but differs in having shorter lemmas which are pubescent between the veins, densely scabrous sheaths with smooth, glabrous throats, and shorter ligules. *Poa urubambensis*, a common element of the undisturbed *Polylepis* forest understory of the Cordillera

* Authors contributed equally to this work

Urubamba, Cusco, is distinct from all other members of open-panicled *Poa*s by having glabrous lemmas with a smooth and glabrous callus, and notably small anthers. The type material for the name *Poa adusta* is discussed and a lectotype is selected.

Resúmen

Aportamos una lista actualizada y una clave para las 30 especies de *Poa* con panículas abiertas de Perú que incluye las especies de *Dissanthelium* y *Aphanelytrum* anteriormente circunscritas, nuevos registros de taxones y tres especies no descritas. *Poa compressa*, *P. grisebachii*, y *P. leioclada* se registran para Perú por primera vez. Un número de especies son sinonimizadas: *Poa carazensis*, *P. ferreyrae* y *P. tovarii* son sinonimizadas bajo el nombre de *P. fibrifera*; *Poa adusta* (tentativamente) y *P. pilgeri* son sinonimizadas bajo el nombre de *P. candamoana*; *Poa superata* es sinonimizada bajo *P. grisebachii*; y *Poa paramoensis* es sinonimizada bajo *P. huancavelicae*. Dentro de este tratamiento se incluyen dos especies nuevas, *Poa ramoniana*, *P. tayacajaensis* y *P. urubambensis* que a continuación se describen e ilustran. *Poa ramoniana*, que se encuentra creciendo en pastizales de alta elevación cercanos a lagos en la Puna de Junín, es similar a la forma pequeña de *P. glaberrima*, pero se diferencia por tener rizomas y crecer hasta sólo 5 cm de altura. *Poa tayacajaensis*, que se encuentra en matorrales de las laderas andinas de Huancavelica y Huánuco, tiene un parecido a *Poa aequatoriensis*, pero se diferencia por tener lemas cortas que son pubescentes entre las venas, vainas densamente escabrosas con suaves gargantas glabras, y lígulas cortas. *Poa urubambensis*, un elemento común de sotobosque no perturbado de los bosques de *Polylepis* de la Cordillera Urubamba, Cusco, se distingue de todos los otros miembros de *Poa* con panícula-abierta por tener lemas glabras con un callo suave y glabro, y sus notables anteras pequeñas. El material tipo del nombre *Poa adusta* es discutido y un lectotipo es seleccionado.

Keywords

Checklist, Gramineae, new species, Peru, *Poa*, Pooideae, *Polylepis* forest, Pooideae, Puna grassland, grass taxonomy

Introduction

The genus *Poa* L. is the largest genus of the Poaceae, containing over 500 species with a large distribution across temperate areas of the globe (Soreng et al. 2003, 2010, 2015, 2016). The first taxonomic treatment of *Poa* from Peru comes from Hitchcock's (1927) 'The grasses of Ecuador, Peru and Bolivia' in which he mentions 17 species. Standley (1936), in the Flora of Peru series, produced the first treatment of purely Peruvian grasses in which he added *P. aequatoriensis* to the country records, raising the total number of Peruvian *Poa* to 18. Following this, the renowned Peruvian agrostologist, Óscar Tovar Serpa, began his life-long work on Peruvian grasses producing a number of publications related to *Poa* (Tovar 1965, 1974, 1984, 1986) until his largest cumulatory work in 1993 where he provided a concise treatment of all grasses from Peru (Tovar 1993).

Tovar's (1993) most up-to-date taxonomic treatment, and the Checklist for Peru (Brako and Zarucchi 1993) that was published in the same year, considered the country to have either 40 or 41 species of *Poa*, respectively. This number has since varied due to taxonomic revision placing certain names in synonymy (Soreng et al. 2003, 2016), and discoveries of undescribed species (Negritto and Antón 2006; Davidse et al. 2010; Soreng

and Peterson 2010; Peterson and Soreng 2016). Additionally, DNA studies (Gillespie et al. 2007, 2008; Refulio-Rodríguez et al. 2012; Peterson and Soreng 2016) found four small, mostly closed-panicled, genera to be nested within *Poa* subg. *Poa* supersect. *Homalopoa* (Dumort.) Soreng & L. Gillespie. Species of *Anthochloa* Nees & Meyen, *Tovaro-chloa* T.D. Macfarl. & P. But, and *Dissanthelium* Trin., were thus transferred to *Poa* by Gillespie et al. (2007) and Refulio-Rodríguez et al. (2012), while species of *Aphanelytrum* (Hack.) Hack. were recently transferred to *Poa* by Peterson and Soreng (2016).

To this point, 51 species had been accepted in *Poa* for Peru (Soreng et al. 2016), four of which are considered to be exotic (*P. annua* L., *P. infirma* Kunth, *P. pratensis* L. [subsp. *alpigena* (Lindm.) Hiitonen is indigenous in North America and Patagonia], *P. trivialis* L.). Of these 51 species, 32 have open-panicled inflorescences. Panicle characteristics are good for separating Peruvian *Poa* into two distinct groups. All taxonomic treatments of *Poa* in Peru (Hitchcock 1927; Standley 1936; Tovar 1993), Bolivia (Renvoize 1998) and Ecuador (Hjorth 1991) have provided diagnostic keys that, in the first couplet, separate species into those with a congested spike-like panicle, with panicle branches appressed, and those with an open-panicled inflorescence, the branches spreading. We believe this artificial character to be convenient and reliable in separating Peruvian *Poa*.

Our objective is to provide an up-to-date summary of the open-panicled species of *Poa* in Peru including locality information using verified specimens and discussion of nomenclatural and taxonomic attributes, with the new species, *Poa ramoniana*, *P. tayacajaensis* and *P. urubambensis*, being described and illustrated. Two keys are provided to aid with identification of the open-panicled *Poa*. The main key first uses anther length to separate taxa while the Suppl. material 1 first uses lemma indumentum. We hope users of our keys will have a better chance of accurately identifying Peruvian specimens of *Poa*.

Materials and methods

In this treatment, glabrous means without pubescence (in the sense of slender, relatively soft hairs). Smooth indicates no prickle-hairs with broad bases and/or hooked or pointed apices (i.e., pubescence can occur on a smooth surface, and a rough or scabrous surface can be glabrous). Specimen localities in the checklist are cited by political region (also historically called 'departamento') (capital letters) and then province. Only herbaria where specimens have been checked and verified by the authors have been cited (acronyms following Thiers, continuously updated): Mainly MO (material on loan to US, c. 240 collections of *Poa* from Peru [*Peterson* duplicates excepted], and many more from across South America), and US, but USM specimens and types were examined in-situ by RJS (in 2006 and 2007), and CUZ and Z specimens were examined by SPS. Almost all *P.M. Peterson Poa* collections (first set at US, c. 460 collections from Peru) are duplicated at USM, although the USM duplicates have not been re-checked for this paper. Excluded species are discussed at the end of the checklist.

Results

Of the 32 species of *Poa* with open-panicles previously recognized in Peru (Soreng et al. 2016), we consider three of these species records to be erroneous and have placed five species names in synonymy. Following the discovery of three undescribed species and three new country records, we now recognize 30 species of *Poa* with open-panicles in Peru.

The new species, *Poa urubambensis*, was found in remote areas of the Cordillera Urubamba, southern Peru, during recent fieldwork by the first author. While reviewing Peruvian specimens of open-panicled *Poa* from collections in the United States National Herbarium and Missouri Botanical Garden, a further two undescribed species, *Poa ramoniana* and *Poa tayacajaensis*, were discovered. *Poa ramoniana* was discovered from collections by eminent Peruvian botanist, Ramón Alejandro Ferreyra, from Junín. *Poa tayacajaensis* was discovered from collections by the renowned Peruvian agrostologist, Óscar Tovar Serpa, from the province of Tayacaja, Huancavelica.

We report new species records for: *Poa* cf. *leioclada* Hack., previously considered endemic to Ecuador (León Yáñez et al. 2011), and *P. grisebachii* R.E. Fr., previously considered endemic to Argentina (Giussani et al. 2012). *Poa compressa* L., originating from Europe, is also reported for the first time. Three species: *Poa carazensis* Pilg., *P. ferreyrae* Tovar, and *P. tovarii* Soreng, appear to be morphologically indistinct from *P. fibrifera* Pilg. and have been placed under that name. *Poa adusta* J. Presl, known only from the type collection, and the recently described *P. pilgeri* Negritto and Antón (2006: 87) are synonymized under *P. candamoana* Pilg. Specimens of *Poa superata* Hack., previously known only from Argentina and Chile, have been collected from Peru and were determined to be morphologically indistinct from *P. grisebachii* R.E. Fr. *Poa paramoensis* Læggaard, previously considered endemic to Ecuador (Læggaard 1998), has been found to be morphologically indistinct from *P. huancavelicae* Tovar. Reports of *Poa lilloi* Hack. and *P. supina* Schrad. for Peru are considered erroneous, and *P. bromoides* Vahl (accepted by Brako and Zarucchi 1993) is currently accepted as *Eragrostis bromoides* (Vahl) Steud. by Soreng et al. (2016). These species were removed from the checklist.

Key to the open-panicled species of *Poa* in Peru

- 1 Lemmas with bifid apexes that are mucronate to short-awned; spikelets glabrous, with long rachillas 1.2–4.2 mm long; glumes short, less than ½ the length of the florets, sometimes absent **2**
- Lemmas with obtuse to acute apexes, never bifid, mucronate or short awned; spikelets usually with some form of indumentum, less often glabrous, with short rachillas rarely longer than 1.2 mm; glumes less than half to equaling the length of the spikelet, never absent **3**
- 2 (1) Glumes veinless, 0.1–0.5 (–0.7) mm long, minute or absent; plants straggling and stooling; culms 30–80(–100) cm tall ***P. hitchcockiana***

- Glumes veined, 1–2 mm long, lower glume 1-veined, upper glume 3- or 4-veined; plants caespitose; culms 14–24 cm tall.....***P. sanchez-vegae***
- 3 (1) Glumes exceeding the florets; spikelets 2-flowered; lemmas 3 (rarely 5)-veined, glabrous, smooth or scaberulous; low tufted (sometimes rhizomatous in *P. trollii*) perennial plants mostly less than 10 cm tall; panicles 1–3.3 cm long..... **4**
- Glumes shorter than the proximal floret; spikelets 2–4(–6)-flowered; lemmas 5(or 7)-veined, glabrous or pubescent, smooth or variously scabrous; annual or perennial plants of various habits, ranging mostly from 10–120 cm tall; panicles longer (sometimes short in the annuals)..... **6**
- 4 (3) Anthers 2–2.2 mm long in staminate specimens, vestigial in pistillate specimens; lemmas smooth throughout; plants sometimes rhizomatous...***P. trollii***
- Anthers <1 mm long in lower floret of spikelets, sometimes vestigial in upper floret; lemmas scaberulous, at least on the keels (rarely smooth in *P. calycina*); plants densely tufted **5**
- 5 (4) Leaf blade abaxial surface shiny with veins indistinct ***P. calycina***
- Leaf blade abaxial surface dull with veins apparent..... ***P. swallenii***
- 6 (3) Longest anthers of proximal florets 0.2–1.5 mm long **7**
- Longest anthers of proximal florets (1.2–)1.6–3.2 mm long..... **16**
- 7 (6) Plants annual; palea keels distinctly pubescent in part (very rarely glabrous) always without any hooks; callus glabrous..... **8**
- Plants perennial; palea keels glabrous or pubescent in part, but always scabrous in part; callus glabrous or webbed, i.e. with long silky hairs (sometimes sparse) emerging from below the lemma keel (at least of the lower florets) **9**
- 8 (7) Anthers 0.2–0.5(–0.6) mm long; panicle branches ascending, spikelets usually crowded; foliage light green; plants ephemeral..... ***P. infirma***
- Anthers 0.6–1 mm long; panicle branches ascending to spreading, spikelets loosely arranged; foliage usually darker green; plants infrequently persisting for more than one season ***P. annua***
- 9 (7) Culm nodes strongly compressed, lower culm nodes exposed; culms wiry; plants strongly rhizomatous with isolated shoots; all florets of spikelets hermaphroditic (sometimes anthers aborted late in development)***P. compressa***
- Combination of characters not as above; culm nodes terete or slightly compressed, lower culm nodes usually held within sheaths; culms varying from wiry to robust; plants tufted or rhizomatous; all florets of spikelets hermaphroditic, or upper floret(s) within spikelets sometimes pistillate, with rudimentary stamens (i.e. gynomonocious) **10**
- 10 (9) Callus glabrous ***P. urubambensis***
- Callus webbed, i.e. with long silky hairs (sometimes sparse) emerging from below the lemma keel (at least of the lower florets) **11**
- 11 (10) Leaf blades filiform or slightly broader, involute or subinvolute towards the base, 0.5–2 mm wide when expanded; lower lemma 3–4 (5?) mm long, glabrous; web only (web sometimes v. short and sparse, and present on basal florets only) ***P. pauciflora***

- Leaf blades flat or folded, usually >2 mm wide when expanded; lower lemma 2.3–6 mm long, variously glabrous or pubescent **12**
- 12 (11) Upper ligules 0.9–2(–3) mm long, truncate; plants distinctly rhizomatous; lower sheaths smooth, sometimes lightly pubescent; lower lemma keel and marginal veins distinctly pubescent; spikelets with 2–6 florets; all florets of spikelets hermaphroditic (sometimes anthers aborted late in development)...
..... ***P. pratensis***
- Combination of characters not as above; upper ligules 0.2–10 mm long, acute or rarely truncate; plants tufted (weakly rhizomatous in *P. huancavelicae*); lower sheaths smooth to densely scabrous; lower lemma keel and marginal veins varying in indumentum from glabrous to short pubescent; spikelets with 2–3 florets; upper floret within spikelets sometimes pistillate, with rudimentary stamens **13**
- 13 (12) Leaf blades folded, apex prominently naviculate (prow-tipped); plants weakly rhizomatous; lemma keels and marginal veins smooth or scaberulous, glabrous; callus webbing the only indumentum present in the spikelet ... ***P. huancavelicae***
- Leaf blades flat, apex not, or not prominently, naviculate; plants tufted, occasionally stooling and rooting at nodes; lemma keels short pubescent in the lower ½, sometimes sparingly so on the marginal veins near the base (rarely glabrous in *P. aequatoriensis*); spikelet indumentum consisting of lemma pubescence (as mentioned above) and callus webbing **14**
- 14 (13) Spikelet proximal lemmas pubescent on keel, lateral, and marginal veins; distal lemmas pubescent between the veins; sheaths densely scabrous; ligules 2–3.5 mm long..... ***P. tayacajaensis***
- Combination of characters not as above; spikelet proximal lemmas glabrous or sparingly pubescent on the keel, and sometimes marginal veins; distal lemmas often glabrous throughout; sheaths smooth to densely scabrous; ligules 1–10 mm long..... **15**
- 15 (14) Lower culm sheaths usually puberulent in the throat margins and/or along the collar margins, surfaces smooth to lightly scabrous; upper culm leaf ligules 1–5(–7) mm long, abaxially puberulous or scabrous; lowest floret of spikelets hermaphroditic, upper florets commonly pistillate; spikelets 3.5–5 mm long; lower lemma 3.6–4 mm long, intermediate veins faint to moderately pronounced; palea keels usually finely scabrous to some degree; anthers mostly 0.6–1.5 mm long..... ***P. aequatoriensis***
- Lower culm sheaths glabrous in the throat margins, surfaces nearly smooth to densely scabrous; upper culm leaf ligules 4–10 mm long, abaxially smooth or faintly scabrous; all florets of spikelets hermaphroditic; spikelets 2.3–3.5(–4) mm long; lower lemma 2.3–3(–3.5) mm long, intermediate veins distinctly pronounced; palea keels usually muriculate, sometimes minutely scabrous; anthers (1) 1.3–1.6 (1.8) mm long..... ***P. trivialis***
- 16 (6) Lemmas glabrous, smooth or scabrous (rarely sericeous at the base in *P. ramifera*); callus glabrous..... **17**

- Lemmas, at least of the upper florets, pubescent or villous in their lower half (rarely scabrous-pubescent in *P. kurtzii*), or glabrous but then callus webbed, i.e. with long silky hairs emerging from below the lemma keel; callus glabrous or webbed.....**26**
- 17 (16) Culms erect, aerially branching well up the culm with lateral shoots that persist and flower in subsequent seasons *P. ramifera*
- Culms not branching, or branching only near the base, or from decumbent culms.....**18**
- 18 (17) Summit of sheaths with prominent triangular auricles; spikelets 4–6-flowered ...
.....*P. auriculata*
- Sheaths without auricles; spikelets 2–6-flowered.....**19**
- 19 (18) Lemmas surface completely smooth (sometimes distally obscurely to sparsely scaberulous in *P. ramoniana* but then plants 4–6 cm tall and ligules <1 mm long) .**20**
- Lemmas slightly to strikingly scabrous between and on veins.....**23**
- 20 (19) Plants 4–6 cm tall; rhizomatous; ligules <1 mm long..... *P. ramoniana*
- Plants >10 cm tall; tufted or, if rhizomatous >100 cm tall; ligules 2–7 mm long**21**
- 21 (20) Leaf blades smooth throughout, (2–)4–9 cm long, 1–2(–2.5) mm wide when blade flattened*P. glaberrima*
- Leaf blades densely scabrous throughout, 8–40 cm long, 3–10 mm wide when blade flattened.....**22**
- 22 (21) Leaf blades conspicuously folded; plants 25–35 cm tall, tufted ... *P. gilgiana*
- Leaf blades flat; plants 100–150 cm tall, rhizomatous.....*P. ayacuchensis*
- 23 (19) Leaf blades involute or the margins distinctly involute (rarely simply folded in *P. kurtzii*), densely scabrous (at least abaxially), firm to rigid; plants tufted; ligules (2.5–)5–15 mm long.....**24**
- Leaf blades flat or folded, margins rarely distinctly involute, glabrous or lightly scabrous, lax or firm; plants erect (*P. fibrifera*) or rhizomatous (*P. oscariana*), sometimes tufted; ligules 1–7(–9) mm long**25**
- 24 (23) Ligules 8–15 mm long; panicles narrowly ovate, panicle branches ascending and subappressed, panicles included in the sheaths; lemmas scabrous..... *P. pearsonii*
- Ligules (2.5–)5–8 mm long; panicles amply ovate, panicle branches patent or reflexed, panicles exerted; lemmas scabrous or scabrous-pilose *P. kurtzii*
- 25 (23) Lower leaf sheaths often fibrous; ligules 1.5–5(–6) mm long; blades lax; spikelets 3–5-flowered, 5–9 mm long; anthers 2.4–3.5 mm long; rachilla internodes well exposed.....*P. fibrifera*
- Lower leaf sheaths not fibrous; ligules 6–9 mm long; blades somewhat firm; spikelets 2–3-flowered, 4.5–5 mm long; anthers 1.8–2 mm long; rachilla internodes short (compare with *P. gilgiana*)*P. oscariana*
- 26 (16) Callus glabrous; lemmas (at least the distal ones within a spikelet) softly villous-pubescent in their lower half**27**
- Callus webbed, i.e. with long silky hairs (sometimes sparse) emerging from below the lemma keel (at least of the lower florets); lemmas glabrous or dis-

- tinctly to sparsely villous or serious pubescent along the keel and marginal veins only..... **30**
- 27 (26) Plants (30–)60–150 cm tall; leaf blades flat, sometimes folded towards their apices, usually more than 3 and up to 10 mm wide; inflorescence branches commonly verticillate; plants (sub-)rhizomatous with extravaginal shoots (if blades rather firm and folded but broad as in *P. horridula*, compare with *P. gilgiana*, possibly hybrids) ***P. horridula***
- Plants usually <35 cm tall; leaf blades involute to narrowly convolute and 0.5–2 mm wide, or flat to folded and 1–5 mm wide in *P. grisebachii*; inflorescence branches solitary or paired (often 3 branches in basal nodes of *P. grisebachii*); plants usually with only intravaginal shoots **28**
- 28 (27) Leaf blade abaxial surface densely scabrous; ligules (2.5–)5–8 mm long, acute; lemmas scabrous-pilose; plants of semi-arid habitats ***P. kurtzii***
- Leaf blade abaxial surface glabrous to scaberulous with prickles or hooks usually restricted to the leaf margin; ligules 0.5–3 mm long, truncate; lemmas pilose-villous towards base; plants of mesic or more arid habitats **29**
- 29 (28) Leaf blades usually involute, apex narrowly but abruptly naviculate (prow-tipped); spikelets usually 3-flowered, (2.8–)4.3–5.5 mm long; culm basal sheath bases slightly inflated, shiny, and tough; plants of more mesic Puna, mostly 3700–4500 m ***P. candamoana***
- Leaf blades flat or folded, somewhat lax, apex often tapered to a long slender point; spikelets 3–6-flowered, (5–)6–7.2 mm long; culm basal sheaths not as above; plants of more arid zones between 3000 and 4000 m (appears to hybridize with *P. kurtzii* where the two overlap) ***P. grisebachii***
- 30 (26) Leaf blades filiform or slightly broader, involute or sub-involute towards the base, 0.5–2 mm wide when expanded; lower lemma 3–4 (5?) mm long, glabrous; web only (web sometimes v. short and sparse, and present on basal florets only) ***P. pauciflora***
- Leaf blades flat or folded, usually >2 mm wide when expanded; lower lemma 2.3–6 mm long, variously glabrous or pubescent **31**
- 31 (30) Basal sheaths glabrous and densely scabrous; lemmas smooth (or lightly scabrous near the apex), glabrous; web only ***P. scabrivaginata***
- Basal sheaths glabrous or lightly pubescent, smooth or lightly scabrous, or if densely scabrous then lemmas pubescent at least on the keel; lemmas smooth or scabrous, glabrous or pubescent in part **32**
- 32 (31) Upper ligules 0.9–2(–3) mm long, truncate; plants distinctly rhizomatous; lower sheaths smooth, sometimes lightly pubescent; lower lemma keel and marginal veins distinctly pubescent; spikelets with 2–6 florets; all florets of spikelets hermaphroditic (sometimes anthers aborted late in development) ***P. pratensis***
- Combination of characters not as above; upper ligules 0.2–10 mm long, acute or rarely truncate; plants tufted (usually with short rhizomatous shoots in *P. huancavelicae* and prominent sub-rhizomatous extravaginal shoots present in *P. leioclada*); lower sheaths smooth to densely scabrous; spikelets with

- 2–3(–4) florets; upper floret within spikelets hermaphroditic (i.e. *P. trivialis*) or commonly pistillate, with rudimentary stamens (i.e. *P. huancavelicae*, *P. cf. leioclada*) **33**
- 33 (32) Leaf blades folded, apex prominently naviculate (prow-tipped); plants weakly rhizomatous; lemma keels and marginal veins smooth or scaberulous, glabrous; web only ***P. huancavelicae***
- Leaf blades flat, apex not, or not prominently, naviculate; plants tufted, occasionally stooling and rooting at nodes (or with prominent sub-rhizomatous shoots present at the base in *P. leioclada*); lemma keels short pubescent in the lower ½, sometimes sparingly so on the marginal veins near the base..... **34**
- 34 (33) Spikelets glomerate on branches; culm leaf ligules 1–2 (–2.5) mm long, truncate or obtuse; short sub-rhizomatous shoots usually prominent at the base of the plant, these extravaginal, with brown cataphylls; spikelet proximal lemmas pubescent on keel, lateral, and marginal veins; distal lemmas often pubescent between the veins; spikelets 2–4-flowered; sheaths smooth; leaf blades mostly folded, sometimes flat, firm; lowest floret of spikelets hermaphroditic, upper florets commonly pistillate..... ***P. cf. leioclada***
- Spikelets diffuse throughout the panicle; culm leaf ligules 4–10 mm long, acute, rarely truncate in lower leaves; rhizomatous shoots absent, new shoots obscure, stoloniferous, extravaginal, with green cataphylls; spikelet proximal lemmas glabrous or sparingly pubescent on the keel, and sometimes marginal veins; all lemmas glabrous between the keel and marginal veins; spikelets 2–3-flowered; sheaths nearly smooth to densely scabrous; leaf blades mostly flat, flaccid; all florets of spikelets hermaphroditic..... ***P. trivialis***

Checklist to the open-panicled *Poa* species of Peru

All of the indigenous species covered (including the new ones) are accommodated in *Poa* subg. *Poa* supersect. *Homalopoa* sect. *Homalopoa* Dumort. s.l., except *P. calycina* and *P. wallenii*, which currently reside in *P.* subg. *Poa* supersect. *Homalopoa* sect. *Dissanthelium* (Trin.) Reulio, and *P. hitchcockiana* and *P. sanchez-vegae*, which reside in *Poa* subgen. *Poa* supersect. *Homalopoa* sect. *Dioicopoa* subsect. *Aphanelytrum*. The introduced species are referred to *P.* subg. *Ochlopoa* (Asch. & Graebn.) Hyl. sect. *Micrantherae* Stapf (*P. annua* and *P. infirma*); *P.* subg. *Stenopoa* (Dumort.) Soreng & L.J. Gillespie sect. *Pandemos* Asch. & Graebn. (*P. trivialis*); *P.* subg. *Poa* supersect. *Poa* sect. *Poa* (*P. pratensis*); *P.* subg. *Stenopoa* sect. *Tichopoa* Asch. & Graebn. (*P. compressa*).

Poa aequatoriensis Hack. **Ref:** Standley (1936: 125); Tovar (1965: 45, 1993: 135). **Ill:** Hjorth (1991: fig. 6), Tovar (1965: lam. X, B). **Habitat:** Montane forest, Puna grassland, pathsides and open areas, rocky slopes. 2000–4600 m. **Vouchers:** PERU: AMAZONAS: Bongara, J.J. Wurdack 944 (US). AYACHUCO: Weberbauer 7573 (US fragm. Ex F). CAJAMARCA: Cajamarca, I. Sanchez Vega (MO);

Celendin, I.M. Sánchez V. 2668 (MO); San Miguel, J. Mostacero L. 1282 (MO); Santa Cruz, J. Santisteban C. & J. Guevara B. 169 (F, MO). CUSCO: Urubamba, H.H.C. Ellenberg 48232 (MO). HUÁNUCO: Pachitea, J.F. Macbride 4365 (US). JUNÍN: Concepción, J.F. Macbride 3363 (US). PIURA: Huancabamba, P.M. Peterson 15175 (US). **Discussion:** *Poa aequatoriensis* occurs from northern Peru, Ecuador, and Colombia, although one collection is known from Cusco. Brako and Zarucchi (1993) and Tovar (1993) also state *P. aequatoriensis* to occur in ANCASH, HUANCAVELICA and LIMA. Commonly misidentified as *P. trivialis* and vice-versa. Reports of *P. aequatoriensis* from Bolivia by Hjorth (1991) are, most likely, another taxon. This report was probably based on material called *Poa umbrosa* Trin. by Renvoize (1998; Renvoize & Cope 4071, K, US!), which RJS redetermined as *Poa bradei* Pilger, a species otherwise known only from Brazil, which has spikelets with only perfect flowers and short anthers (0.5–1 mm).

***Poa annua* L.** **Syn:** *Ochlopoa annua* (L.) H. Scholz. **Ref:** Standley (1936: 125); Tovar (1965: 61, 1993: 127). **Ill:** Giussani et al. (2012: sp. 3, p. 294). **Habitat:** Waste and disturbed ground, pathsides, roadsides and fields. 2200–4800 m. **Vouchers:** PERU: ANCASH: Carhuaz, D.N. Smith 9561 (MO); Huari, P.M. Peterson 13872 (MO, US); Huaylas, D.N. Smith 9294 (MO); Pallasca, P.M. Peterson 13947 (MO, US); Recuay, P.M. Peterson 13827 (MO, US). AREQUIPA: Arequipa, P.M. Peterson 18256 (US); Caraveli, P.M. Peterson 16391 (US). AYACUCHO: Huanca Sancos, P.M. Peterson 16258 (US); Lucanas, P.M. Peterson 16309 (US); Parinacochas, P.M. Peterson 16328 (US). CAJAMARCA: Cajamarca, P.M. Peterson 14910 (MO, US); San Ignacio, P.M. Peterson 15129 (MO, US). CUSCO: Calca, S.P. Sylvester 1334 (CUZ, LPB, US, Z); Canchis, J. Farfán 864 (MO); Cusco, P. Núñez V. 7500 (US); Espinar, P. Núñez V. 7619 (MO); La Convención, S.P. Sylvester 1552 (CUZ, US, Z). HUANCAVELICA: Huancavelica, P.M. Peterson 14173 (MO, US). JUNÍN: Junín, P.M. Peterson 14095 (MO, US); Tarma, D.N. Smith 1605 (MO). LA LIBERTAD: Bolívar, R.W. Bussmann 18047 (MO); Otuzco, A. Sagástegui A. 11567 (MO); Trujillo, J. Hudson 1027 (MO). PASCO: Oxapampa, D.N. Smith 5831 (MO). PUNO: Chucuito, P.M. Peterson 14634 (US); El Collao, P.M. Peterson 14593 (MO, US); Sandia, B.C. Bennett 1944 (MO). TACNA: Tarata, P.M. Peterson 14724 (MO, US). **Discussion:** Introduced weed from Europe. This species sometimes survives for more than one growing season and can occur with completely glabrous lemmas, where it is commonly misidentified as *Poa supina* (see ‘Species excluded’ section below). Brako and Zarucchi (1993) and Tovar (1993) also state *P. annua* to occur in HUÁNUCO, LIMA and PIURA.

Poa auriculata Soreng & P.M. Peterson **Ref & Ill:** Peterson and Soreng (2016: 121, fig. 5). **Habitat:** Known only from the type locality growing on a dry cliff face between 3100–3200 m. **Vouchers:** PERU: AMAZONAS: Chachapoyas, J.J. Wurdack 1145 (US-holotype). **Discussion:** Endemic herb. This is the first member of *P.* subg. *Poa* supersect. *Homalopoa* with prominent auricles.

Poa ayacuchensis Tovar **Ref & Ill:** Tovar (1974: 6, 1993: 129, fig. 1, 2, 3). **Habitat:** Wet ground, riversides. c. 3700 m. **Vouchers:** PERU: AYACUCHO: Lucanas, Ó.

Tovar S. & R. Foguel 7007 (MO, US, USM). **Discussion:** Endemic herb, known only from the type collection.

Poa calycina (J. Presl) Kunth **Syn:** *Brizopyrum calycinum* J. Presl; *Deschampsia mathewsii* Ball; *Dissanthelium calycinum* (J. Presl) Hitchc.; *Dissanthelium laxifolium* Swallen & Tovar; *Dissanthelium mathewsii* (Ball) R.C. Foster & L.B. Sm.; *Dissanthelium sclerochloides* Steud. ex E. Fourn.; *Dissanthelium semitectum* Swallen & Tovar; *Dissanthelium supinum* Trin.. **Ref:** Refulio-Rodríguez et al. (2012: 130); Swallen & Tovar (1965: 370–371); Tovar (1993: 154–156). **Ill:** Renvoize (1998: fig. 38 E, F). **Habitat:** Puna grassland. 2500–4500 m. **Vouchers:** PERU: ANCASH: Bolognesi, *P.M. Peterson 17974* (US); Recuay, *P.M. Peterson 17904* (US); Yungay, *P.M. Peterson 21676a* (US). AYACUCHO: Cangallo, *P.M. Peterson 18164* (US); Lucanas, *P.M. Peterson 16466* (US). CUSCO: Calca, *P.M. Peterson 18100* (US); Urubamba, *S.P. Sylvester 1706* (AAU, LPB, US). HUANCVELICA: Huancavelica, *P.M. Peterson 18100* (US). HUÁNUCO: Huamalies, *P.M. Peterson 17923* (US). JUNÍN: Huancayo, *P.M. Peterson 14209* (US); Junín, *P.M. Peterson 14105* (US); Tarma, *P.M. Peterson 14090* (US); Yauli, *P.M. Peterson 18042* (US). LIMA: Canta, *P.M. Peterson 18028* (US). PASCO: Pasco, *P.M. Peterson 18075* (US). **Discussion:** This species has a disjunct distribution. It is found in the high elevation Puna grasslands of Bolivia and Peru and the alpine volcanic slopes of Mexico (Soreng and Peterson 2012). Brako and Zarucchi (1993) report the species to occur in PUNO.

Poa candamoana Pilg. (Nom. Cons. Prop. *In prep.*) **Syn:** *Poa adusta* J. Presl, lectotype here designated *Haenke s.n.* (lectotype: PR-495759!, left-hand specimen; isolecotype: HAL-81886, W-0029634); *Poa pilgeri* Negritto & Antón. **Ref:** Standley (1936: 125); Tovar (1965: 57, 1993: 131); Negritto and Antón (2006: 88). **Ill:** Tovar (1965: lam. XIII, A); Negritto and Antón (2006: fig. 3, as *P. pilgeri*). **Habitat:** Grassland. 3400–4500 m. **Vouchers:** PERU: ANCASH: Paillon, *E. Cerrate 2685* (US); Recuay, *P.M. Peterson 21547* (US); Yungay, *P.M. Peterson 21748* (US). AREQUIPA: Arequipa, *P.M. Peterson 18254* (US). AYACUCHO: Lucanas, *H.H.C. Ellenberg 4945* (MO). CAJAMARCA: Celendin, *P.M. Peterson 21915* (US). CUSCO: Cusco, *A.S. Hitchcock 22469* (US); Espinar, *C. Vargas C. 5629* (MO); Quispicanchis, *P.M. Peterson 20599* (US). HUANCVELICA: Huancavelica, *P.M. Peterson 16422* (US); Churcampa, *O. Tovar S. 805* (US); Huaytara, *P.M. Peterson 20428a* (US); Tayacaja, *O. Tovar S. 2475* (US). HUÁNUCO: Huamalies, *P.M. Peterson 17925* (US). JUNÍN: Junín, *P.M. Peterson 14116* (US); Huancayo, *I.J. Blair 424* (K), *S. Soukup 6103* (US); Yauli, *Hirsch P243* (US). LIMA: Canta, *P.M. Peterson 20286* (US). MOQUEGUA: Gral. Sanchez Cerro, *D.B. Montesinos T. 2494* (MO, USM). PASCO: Pasco, *P.M. Peterson 14099* (US). PUNO: Azángaro, *A. Weberbauer 472* (MO, US); Puno, *O.P. Pearson 52-68* (US). **Discussion:** A common and characteristic element in the high Andean Puna grassland of Peru and Bolivia. Brako and Zarucchi (1993) state *P. candamoana* also occurs in APURIMAC.

There has long been uncertainty regarding the identity of *Poa adusta* (Standley 1936, p. 129; Tovar 1965, p. 61, Soreng et al. 2003), which is known only from the type col-

lection, *T. Haenke s.n.* (PR, HAL, W). A report from Colombia (Giraldo-Cañas 2011) represents a different species. The type description of *P. adusta* can be considered erroneous regarding the lemma indumentum, which was stated to be scabrous while the type specimens examined had pubescence present on the lemma keel marginal and intermediate veins (and at least sparingly between them). After studying the HAL isotype, a solitary flowering shoot with one full leaf, without base, we felt it most likely represented *P. candamoana*. A photo of the W isotype (which has only an inflorescence and a bit of upper culm) is also a match for the HAL specimen. Upon studying the PR isotype sheet of *P. adusta*, it was found to comprise two separate leafy shoots that differ from each other. The PR left-hand plant, has a base (with basal-most sheaths characteristic of *P. candamoana*; ie, the base slightly inflated and lustrous), and it otherwise matches the HAL sample in details. The spikelets of these plants differ from *P. candamoana* in the broad, blunt, somewhat distorted lemmas, and dark coloration (*adustus* means blackened or scorched), and do not match any species we know of. Although the dark coloration might derive from poor preservation (coming as they did from moldy bundles), we have seen similarly distorted and discolored spikelets in a few specimens of *Poa* from the region that we expect resulted from disease (given that in one case other spikelets in the same plant inflorescence were normal in shape, color, and pubescence). The PR right-hand plant, although bearing similar characteristics of lemma indumentum, is obviously quite pubescent between the veins and the lemmas are acute and not so discolored (a few spikelets are discolored and distorted to a lesser degree than in the left hand plant), lacks a base, has blade apices that are not navicular as is normal for *P. candamoana*, and the upper culm blade is far too long (longer than the panicle) and inserted too high on the culm for that species (> 10 cm and exceeding the panicle). The right hand plant is almost certainly *P. horridula*. The left hand plant could be either the result of hybridization between *P. candamoana* and *P. horridula* (see *P. horridula* discussion below), diseased material of *P. candamoana* or another species, or a rare species not yet rediscovered. Our choice is to lectotypify *Poa adusta* on the left hand plant of specimen PR-495759. We will propose conservation of *Poa candamoana* (Pilger 1906) over *P. adusta* (J. Presl 1830), rather than outright rejection, since the former name has been widely used for this commonly collected species, and the identity of the lectotype of the latter is still in doubt.

The origin of the *P. adusta* specimens has been uncertain as they might not have been collected from Peru, even though J. Presl (1830: 271) mentioned 'Peruvia' as place of origin in the protologue. The *P. adusta* specimen at PR was part of a bundle of specimens which were tagged with the note 'Aus verfault P.' (translated as 'from rotten bundle'). The tag, written by Bohemian botanist K.M. Sternberg (sometime between 1821 and 1822), indicates that the specimen came from a damaged bundle of plants, for which the country of origin was not indicated. Haenke's herbarium originally consisted of specimens without labels and, when the Czech National Museum purchased the bundle from the compatriot handling company of Hiecke, Ziencke & Co. in 1821, the origin was indicated only on the top of every bundle of specimens. This is the reason why Haenke's handwriting is generally missing from all of his specimens. 'Peruvia', mentioned in the protologue, is J. Presl's opinion about the origin of these specimens rather than the real place of origin for some portion of them. Hae-

nke's collections (made between 1790 and 1792) may have come from his crossing from Buenos Aires and Rio La Plata to Santiago (where he finally caught up with the Malaspina Expedition), or any (suitable) place along the Pacific Coast of America from Santiago (Chile), Lima (Peru), Acapulco (Mexico), to Monterey (California, then part of Mexico), Nootka Sound (now British Columbia), and Yakutat Bay (now Alaska), where the expedition landed (see Češka 2002). Our determination of the right hand plant at PR as *P. horridula* provides a location for the *Poa adusta* lectotype as the central Andes, where *P. horridula* and *P. candamoana* are common.

Although the set of Haenke's collections that made their way to Prague were purchased by the Czech National Museum, a substantial part of it ended up in the Prague University herbarium (now the herbarium of Charles University in Prague, PRC). J. Presl's brother (who wrote up the Gramineae, Cyperaceae, and Taccaceae), C.B. Presl, was custodian of the PR herbarium (where all or most of Haenke's Poaceae from the expedition now reside), and together, while writing the *Reliquiae Haenkeanae* (C.B. Presl 1825–1835), they offered surplus duplicate specimens for sale to other botanical institutions, which is likely how they arrived at HAL and W (Češka 2002; Otakar Sida [PR], pers. communication).

***Poa compressa* L. Ref:** Soreng and Peterson (2012: 31). **Ill:** Giussani et al. (2012: sp. 11, p. 300). **Habitat:** Cool-temperate, semi-shaded to open habitats on wet ground. **Vouchers:** PERU: JUNÍN: Corpacancha, *G.A. Sullivan* 828 (MO). **Discussion:** This species was possibly introduced to the Neotropics for soil stabilization or as a contaminant in other seed. Originating in Europe, and possibly native, but is introduced and naturalized in North America, and Asia. It is presumed to be introduced in Central and South America. This species usually has a congested inflorescence but infrequently exhibits an open inflorescence.

***Poa fibrifera* Pilg. Syn:** *Poa carazensis* Pilg.; *Poa ferreyrae* Tovar; *Poa geniculata* Tovar; *Poa tovarii* Soreng. **Ref:** Standley (1936: 126); Tovar (1965: 37–44, 1984: 8, 1993: 128–130); Soreng (1998: 200). **Ill:** Tovar (1965: lam. X, A.; VIII, B, as *P. ferreyrae*, IX, B as *P. carazensis*, 1984: fig. 6, 7, 8, as *P. geniculata*). **Habitat:** Montane forest, Puna grassland, rocky slopes. 2500–4500 m. **Vouchers:** PERU: ANCASH: Aquia, *E. Cerrate* 1577 (US); Bolognesi, *E. Cerrate* 2607 (US); Carhuaz, *D.N. Smith* 9493 (MO); Huaraz, *D.N. Smith* 10940 (MO); Huari, *D.N. Smith* 12590 (MO); Huaylas, *D.N. Smith* 9774 (MO); Pallasca, *P.M. Peterson* 13949 (US); Recuay, *P.M. Peterson* 13800 (MO, US); Santa, *A. Weberbauer* 3073 (MO, US); Ticllos, *E. Cerrate* 2650 (US); Yungay, *P.M. Peterson* 21741 (US). AYACUCHO: Lucanas, *P.M. Peterson* 18210 (US). CAJAMARCA: Cajamarca, *I. M. Sánchez V.* 3534 (MO); Celendin, *I. M. Sánchez V.* 3488 (MO); Contumaza, *A. Sagástegui* 9647 (MO); Hualgayoc, *P.M. Peterson* 14941 (MO, US). HUANCVELICA: Huaytara, *P.M. Peterson* 18160 (US). LA LIBERTAD: Contumaza, *E.S. Anderson* 1265 (US); Otuzco, *R. Ferreyra* 7619 (MO, US); Santiago de Chuco, *A. Sagástegui* A. 11809 (MO). LIMA: Canta, *P.M. Peterson* 20262 (US); Huarochiri, *R. Ferreyra* 8252 (US). **Discussion:** Endemic herb (a voucher from Bolivia, *T. Johns* 82-108, was redetermined by RJS as *P. horridula*). *Poa ferreyrae* and *P. geniculata*

(=*P. tovarii*), known only from the type collections at La Libertad-Otuzco, and *P. carazensis*, known from the departamento of Ancash, are seen to be indistinct from *P. fibrifera* and so have been placed as synonyms of this species. Brako and Zarucchi (1993) also state *P. fibrifera* to occur in JUNÍN and HUÁNUCO. Specimens from JUNÍN originally identified as *P. fibrifera* (A.S. Hitchcock 22176; P.M. Peterson 14227) have been redetermined as *P. glaberrima* and *P. kurtzii*, respectively. It is impossible to say if either of these specimens were accepted by Brako and Zarucchi (1993) as only one specimen was cited for the country.

Poa gilgiana Pilg. **Syn:** *Melica expansa* Steud. ex Lechl. **Ref:** Standley (1936: 126); Tovar (1965: 36, 1993: 128). **Ill:** Tovar (1965: lam. VIII, A). **Habitat:** Grassland. 3700–4700 m. **Vouchers:** PERU: AREQUIPA: Arequipa, P.M. Peterson 18255 (US); Caraveli, P.M. Peterson 16394 (US). JUNÍN: Junín, A.S. Hitchcock 22251 (US). LIMA: Canta, P.M. Peterson 18025 (US). PUNO: Azángaro, A. Weberbauer 477 (MO, US); Chucuito, P.M. Peterson 14678 (US). TACNA: Tarata, P.M. Peterson 14830 (MO, US). **Discussion:** Distributed in high elevation Puna grasslands from central Peru to Bolivia. Possibly a variety of *P. horridula* which needs further study. This species is suspected to hybridize with *P. horridula* with intermediate forms being collected from ANCASH-Aquia (*E. Cerrate* 1564, US), HUANCANELICA-Huaytara (P.M. Peterson 20424a, US) and PUNO-Chucuito (P.M. Peterson 14669, MO, US).

Poa glaberrima Tovar **Ref:** Tovar (1965: 40, 1993: 129). **Ill:** Tovar (1965: lam. IX, A). **Habitat:** Puna tussock grassland, humid areas, wet ground. 3300–4700 m. **Vouchers:** PERU: ANCASH: Recuay, P.M. Peterson 13821 (MO, US). APURÍMAC: Ayamaraes, P.M. Peterson 16507 (US). AYACUCHO: Huamanga, P.M. Peterson 20532a (US). CUSCO: Calca, P.M. Peterson 16555 (US). HUANCANELICA: Huancavelica, P.M. Peterson 14168 (MO, US). JUNÍN: Huancayo, *Anonymous* 34 (US); Junín, D.N. Smith 5649 (MO); Tarma, D.N. Smith 2989 (MO); Yauli, P.M. Peterson 14044 (US). LIMA: Canta, P.M. Peterson 20282 (US); Huarochiri, J.R. Swallen 7068 (US). PUNO: Lampa, P.M. Peterson 20744 (US); Melgar, P.M. Peterson 20721 (US); Puno, H.H.C. Ellenberg 249 (US). **Discussion:** Previously considered endemic, but has been found in Ecuador, Bolivia, and Argentina according to Giussani et al. (2012: 304); we have verified specimens from Bolivia (*S.A. Renvoize* 4475, US) and Argentina (P.M. Peterson et al. 19577), but not from Ecuador. This species superficially resembles *P. candamoana* but the latter prefers dry ground and has hairy lemmas and lacks extravaginal shoots. Certain specimens placed under this name from southern Peru (eg. PUNO: Lampa “2 km SW of San Jose on road towards Junin” P.M. Peterson 20744, [US]; Melgar “ca. 7 km WNW of Santa Rosa on Hwy 3 and 1 km W toward Quishuara, along Rio Santa Rosa” P.M. Peterson 20624b [US]), Bolivia and northern Argentina (P.M. Peterson 19577 [US]; Giussani et al. 2012, Figure pg. 304) have odd forms with sparse inflorescences and may actually be a species new to science. They resemble *P. pauciflora*, but have completely glabrous and smooth spikelets. Future study should focus on verifying the status of the southern *P. glaberrima* populations.

Poa grisebachii R.E. Fr. **Syn:** *Poa superata* Hack. **Ref:** Giussani et al. (2012: 305, 336). **Ill:** Negritto and Antón (2000: fig 1.); Giussani et al. (2012: sp. 45, p. 305; sp. 59, p. 366, as *P. superata*). **Habitat:** Dry Puna grasslands and high Andean steppe. 3000–4500 m. **Vouchers:** PERU: AYACUCHO: Parinacochas, *P.M. Peterson 16323* (US). JUNÍN: Yauli, Ó. *Tovar S. 6447* (MO). MOQUEGUA: Carumas, *A. Weberbauer 7310* (US); Mariscal Nieto, *P.M. Peterson 14553* (US). TACNA: Tarata, *P.M. Peterson 14793* (MO, US). **Discussion:** Type collections were verified of *Poa grisebachii* (ARGENTINA: JUJUY: Santa Catalina, *F. Kurtz 11412*, US fragm. ex S!) and *P. superata* (ARGENTINA: TUCUMÁN: Tafí, Cumbres Calchaquies, *T.J.V. Stuckert Herb. ARG. 17738 ex Lillo 5604*, US fragm. ex W!). Other collections were verified from Peru (see vouchers above), Chile (ARICA & PARINACOTA: Zapahuiria, *P.M. Peterson 15700*, US; TARAPACÁ: above Pucapa, *P.M. Peterson 15615*, US) and Argentina (JUJUY: Santa Catalina, *F. Kurtz 11409*, US (fragm. ex S); SALTA: Pampa Grande, *C. Spegazzini 60*, US). Previously, *P. grisebachii* and *P. superata* were both considered endemic to Argentina (Giussani et al. 2012) and this is the first recording of the species from Peru. It is likely that, with further revision of Bolivian *Poa*, this species will also be found to occur in Bolivia. Specimens of *P. superata* were found to be morphologically indistinct from the type of *P. grisebachii* and so have been grouped as a single taxon. All material examined of this taxon exhibited soft pubescence on at least the upper lemmas of the spikelets, with the lower florets being glabrous or softly pubescent. *Poa grisebachii* can be highly variable in terms of lemma pubescence. The lectotype of *P. grisebachii* p.p. has glabrous and smooth lemmas while the isolectotype and other syntypes of *P. grisebachii* p.p. at US frags. ex UPS, and the type specimens of *P. superata* p.p., were all observed with at least sparsely pubescent lemmas. All the Peruvian material of *P. grisebachii* has pubescent lemmas, at least on the distal florets.

The plants resemble *P. fibrifera*, and are sometimes fibrous at their bases, but can usually be distinguished by the distinctly pubescent lemmas. The leaf blades are usually somewhat thin, flat, lax, and scabrous, 1.5–3.5 mm wide, tapering to a very slender apex. Certain specimens were found from AYACUCHO: Lucanas, e.g. *P.M. Peterson 16317* (US) that bore characteristics of *P. horridula*. Another specimen from MOQUEGUA: Mariscal Nieto, *P.M. Peterson 14553* (US), could also not be confidently placed to the species. This species may also hybridize with *P. kurtzii*, with intermediate forms being collected which exhibit narrower, firmer, involute blades and generally smaller and more scabrous spikelets.

Poa hitchcockiana Soreng & P.M. Peterson **Syn:** *Brachyelytrum procumbens* Hack.; *Aphanelytrum decumbens* Hack. ex Sodiro. **Ref & Ill:** Peterson and Soreng (2016: 111, fig. 1). **Habitat:** Paramo or moist Jalca vegetation between 2000–4025 m. **Vouchers:** PERU: CUSCO: Paucartambo, *P.M. Peterson 16571* (US, USM); Calca, *P.M. Peterson 16581* (US, USM); Pillco, *C. Vargas C. 19264* (US); Quispicanche, *P.M. Peterson 20582* (US, USM). MOQUEGUA: El Abra, *C. Vargas C.*

19104 (US). SAN MARTÍN: Huicungo, *A. Cano s.n.* (SI, USM); *B. León* 3797 (USM). **Discussion:** The only wide ranging species of *Poa* subsect. *Aphanelytrum*, this species is found from the Andes of Colombia, Ecuador, Peru, and Bolivia. *Poa hitchcockiana* also has the most unusual spikelet morphology with extremely long rachillas (1.5–4.2 mm long) and very short [0.1–0.5 (–0.7) mm long] to obscure or often absent, unveined glumes.

Poa horridula Pilg. **Syn:** *Melica expansa* Steud ex Lechl.; *Poa dumetorum* Hack.; *Poa piifontii* J. Fernandez Casas, J. Molero & A. Susanna; *Poa pufontii* Fern. Casas, Molero & Susanna; *Poa unduavensis* Hack.. **Ref:** Standley (1936: 127); Tovar (1965: 53, 1993: 131). **Ill:** Hjorth (1991: fig. 4), Tovar (1965: lam. XII, B). **Habitat:** Grasslands, rocky slopes, shrublands. 2900–4700 m. **Vouchers:** PERU: ANCASH: Bolognesi, *P.M. Peterson* 17888 (US); Pallasca, *P.M. Peterson* 21810 (US); Recuay, *P.M. Peterson* 13841 (MO, US); Santa, *A. Weberbauer* 3113 (MO, US); Yungay, *P.M. Peterson* 21631 (US). APURIMAC: Abancay, *P.M. Peterson* 16645 (US); Aymaraes, *P.M. Peterson* 16477 (US). AREQUIPA: Caylloma, *P.M. Peterson* 18298 (US). AYACUCHO: Huamanga, *P.M. Peterson* 20503 (US); Huanca Sancos, *P.M. Peterson* 16283 (US); Lucanas, *P.M. Peterson* 16289 (US); Parinacochas, *P.M. Peterson* 16346 (US). CAJAMARCA: Cajamarca, *I.M. Sánchez V.* 79 (US); Celendin, *I.M. Sánchez V.* 2911 (MO); Hualgayoc, *P.M. Peterson* 14938 (US). CUSCO: Anta, *G.R. Brunel* 283 (MO); Calca, *S.P. Sylvester* 1021 (CUZ, US, Z); Cusco, *A.S. Hitchcock* 22443 (US); La Convención, *S.P. Sylvester* 2028 (LPB, US, Z); Paruro, *P. Núñez V.* 7365 (MO); Quispicanchis, *P.M. Peterson* 20549 (US); Urubamba, *H.H.C. Ellenberg* 4824 (MO). HUANCAMELICA: Huancavelica, *P.M. Peterson* 14175 (MO, US); Huaytara, *P.M. Peterson* 18158 (US); Tayacaja, *O. Tovar S.* 4213 (MO). HUÁNUCO: Huamalies, *P.M. Peterson* 17922 (US); Yarowilca, *J.F. Macbride* 1970 (US). JUNÍN: Huancayo, *Black* 46-711 (US); Jauja, *P.M. Peterson* 14122 (US); Junín, *A.S. Hitchcock* 22213 (US). LA LIBERTAD: Bolivar, *P.M. Peterson* 21944 (US). LIMA: Canta, *P.M. Peterson* 20278 (US); Huarochiri, *P.M. Peterson* 14018 (MO, US); Yauyos, *E. Cerrate* 1083 (US). PASCO: Daniel Alcides Carrión, *A.S. Hitchcock* 22302 (US); Pasco, *A.S. Hitchcock* 22260bis (US). PUNO: Chucuito, *P.M. Peterson* 14672 (US); Puno, *H.H.C. Ellenberg* 342 (US); Sandia, *B.C. Bennett* 2323 (MO). **Discussion:** Found above 3000 m in Puna and Paramo grasslands from Colombia to Bolivia. Brako and Zarucchi (1993) and Tovar (1993) indicate that *P. horridula* also occurs in MOQUEGUA. This species is suspected to hybridize with *P. gilgiana* (see *P. gilgiana* discussion, above) and *P. candamoana*, due to the presence of intermediate forms across the ranges of these species. Specimens *Hirsch* P1399 (US) and *E. Cerrate* 2285 (US) appear to be intermediates between *P. horridula* and *P. candamoana* and could be either a short *P. horridula* or a robust *P. candamoana*.

Poa huancavelicae Tovar **Syn:** *Poa paramoensis* Lægaard. **Ref:** Tovar (1965: 52, 1993: 134); Lægaard (1998: 28). **Ill:** Tovar (1965: lam. XII, A); Lægaard (1998: fig. 2, as *P. paramoensis*). **Habitat:** Puna grassland and *Polylepis* woodland. 4000–4800 m. **Vouchers:** PERU: ANCASH: Bolognesi, *P.M. Peterson* 17953 (US); Huaraz,

D.N. Smith 10839 (MO); Recuay, *P.M. Peterson* 21540 (US); Yungay, *P.M. Peterson* 21766 (US). CUSCO: Calca, *P.M. Peterson* 16604 (US); Carhuaz, *D.N. Smith* 11206 (MO); La Convención, *S.P. Sylvester* 1565 (US, Z); Quispicanchis, *P.M. Peterson* 20589 (US); Urubamba, *H.H.C. Ellenberg* 449 (MO). HUANCVELICA: Castrovirreyna, *Ó. Tovar* S. 28 (MO). **Discussion:** Previously considered endemic to Peru and found in HUANCVELICA and CUSCO (Brako and Zarucchi 1993; Tovar 1993). *Poa paramoensis* was found to be inconsistently morphologically distinct from *P. huancavelicae*; the northern plants branches are usually fairly smooth, but the variation appears to be continuous to more scabrous forms further south in Peru, and so it has been synonymized. Specimens from La Convención, CUSCO, have proximal lemmas much shorter (3–3.5 mm) than in the original species description (4–4.5 mm) and anthers were also much shorter (1–1.3 mm). These plants were also found to be short-rhizomatous/sub-rhizomatous, while *P. paramoensis* p.p. is tufted and the isotype material of *P. huancavelicae* p.p. at MO appears tufted, although Tovar (1993) mentions it to be rhizomatous.

Poa infirma Kunth **Syn:** *Catabrosa thomsonii* Hook. f.; *Colpodium thomsonii* (Hook. f.) Hack.; *Eragrostis infirma* (Kunth) Steud.; *Megastachya infirma* (Kunth) Roem. & Schult.; *Ochlopoa infirma* (Kunth) H. Scholz; *Poa annua* subsp. *exilis* (Tomm. ex Freyn) Asch. & Graebn.; *Poa exilis* (Tomm. ex Freyn) Murb.; *Poa remotiflora* (Hack.) Murb. **Ref:** Müller et al. (1981: 334); Tovar (1986: 56; 1993: 126). **Ill:** Giussani et al. (2012: sp. 27, p. 311). **Habitat:** Dry forest, Puna grassland. c.4400 m. **Vouchers:** PERU: CUSCO: Calca, *P. Núñez V.* 7063 (MO), *S.P. Sylvester* 1390 (US, Z). **Discussion:** Introduced from Europe. Brako and Zarucchi (1993) and Tovar (1993) state *P. infirma* occurs in HUANCVELICA and LIMA, with the specimens from LIMA being found on coastal hills in sandy soil. *Poa infirma* is the maternal diploid parent of *P. annua* (tetraploid), with *P. supina*, another diploid, being the plastid donor (Soreng et al. 2010).

Poa kurtzii R.E. Fr. **Syn:** *Poa altoperuana* R. Lara & Fern. Casas; *Poa asperiflora* Hack.; *Poa munozensis* Hack.; *Poa pflanzii* Pilg. **Ref:** Standley (1936: 125); Tovar (1965: 60, 1993: 133); Giussani et al. (2012: 312). **Ill:** Negritto and Antón (1999: fig. 2); Giussani et al. (2012: sp. 30, p. 313); Tovar (1965: lam. XII, A, as *P. asperiflora*). **Habitat:** High Andean Puna grassland, rocky slopes. 3300–5100 m. **Vouchers:** PERU: ANCASH: Huari, *D.N. Smith* 10138 (MO); Recuay, *P.M. Peterson* 21510 (US). AREQUIPA: Arequipa, *P.M. Peterson* 18260 (US); Caraveli, *P.M. Peterson* 16393 (US). AYACUCHO: Huamanga, *P.M. Peterson* 20531 (US); Huanca Sancos, *P.M. Peterson* 16244 (US); Lucanas, *P.M. Peterson* 16177 (US); Parinacochas, *P.M. Peterson* 16345 (US). HUANCVELICA: Castrovirreyna, *O. Tovar* S. 2838 (US); Huancavelica, *P.M. Peterson* 16416 (US); Huaytara, *P.M. Peterson* 20426 (US). JUNÍN: Huancayo, *P.M. Peterson* 14227 (US). LA LIBERTAD: Trujillo, *H.H.C. Ellenberg* 3779 (MO). MOQUEGUA: Mariscal Nieto, *P.M. Peterson* 14552 (US), *P.M. Peterson* 18312 (US). PUNO: Azángaro, *H.H.C. Ellenberg* 598 (US); Chucuito, *P.M. Peterson* 14677 (US); El Collao, *P.M. Peterson* 14591 (US). TACNA: Tacna, *P.M. Peterson* 14762 (US); Tarata, *P.M. Peterson*

14727 (US). **Discussion:** This species is found from Peru to Argentina. JUNÍN specimen *P.M. Peterson* 14227 is atypical and requires further study. Brako and Zarucchi (1993) state *P. asperifolia* (= *kurtzii*) to also occur in CUSCO and LIMA. The species is highly variable in terms of lemma indumentum, and is found in two leads in both keys (see Appendix 1) to account for this.

Poa* cf. *leioclada Hack. **Ref:** Hitchcock (1927). **Ill:** Hjorth (1991: fig. 8) **Habitat:** Humid Paramo grasslands. 3200–4300 m. **Vouchers:** PERU: ANCASH: Yungay, *D.N. Smith* 9095 (MO); Huaylas, *D.N. Smith* 9320a (MO). PIURA: Huanca-bamba, *P.M. Peterson* 15175 (US). **Discussion:** Specimens are an imperfect match of Ecuadorian material of *Poa leioclada* as they bear characteristics of both *Poa mulalensis* Kunth and *P. leioclada*. Both these species were previously considered Ecuadorian endemics. This taxon needs further study.

Poa oscariana Negritto & Antón **Ref & Ill:** Negritto and Antón (2006: 84 [88], fig. 2). **Habitat:** Puna grassland, rocky slopes. 3600–4100 m. **Discussion:** Endemic herb, only known from CUSCO, Paucartambo. None of the type or paratype specimens have been examined. This may be a variety of *Poa gilgiana*, but further study is needed.

Poa pauciflora Roem. & Schult. **Syn:** *Poa depauperata* Kunth; *Poa pardoana* Pilg. **Ref:** Standley (1936: 128); Tovar (1965: 49, 1993: 133). **Ill:** Hjorth (1991: fig. 11), Tovar (1965: lam. XI, B, as *P. pardoana*). **Habitat:** Puna grassland, rocky slopes. 3200–4900 m. **Vouchers:** PERU: ANCASH: Corongo, *P.M. Peterson* 21777 (US); Huaraz, *D.N. Smith* 10797 (MO); Huari, *P.M. Peterson* 13885 (MO, US); Huaylas, *D.N. Smith* 9930 (MO); Pallasca, *P.M. Peterson* 21842 (US); Recuay, *P.M. Peterson* 13848 (MO, US); Yungay, *P.M. Peterson* 21678 (US). CAJAMARCA: Cajamarca, *P.M. Peterson* 14887 (US), *H.H.C. Ellenberg* 1825 (US); Celendin, *P.M. Peterson* 21905 (US); San Miguel, *P.M. Peterson* 14933 (US); San Pablo, *P.M. Peterson* 14878 (US). JUNÍN: Tarma, *P.M. Peterson* 14065 (US). LA LIBERTAD: Bolivar, *P.M. Peterson* 21936 (US); Sanchez Carrion, *D.N. Smith* 2242 (MO); Santiago de Chuco, *P.M. Peterson* 13953 (US). SAN MARTÍN: Mariscal Caceres, *B. León* 1654 (MO). **Discussion:** Specimens have also been verified from Ecuador, but it is unclear whether this species extends to Colombia and Bolivia.

Poa pearsonii Reeder **Ref:** Tovar (1965: 33, 1993: 127); Giussani et al. (2012: 325). **Ill:** Giussani et al. (2012: sp. 45, p. 325). **Habitat:** Rocky Puna grassland. 4500–4900 m. **Vouchers:** PERU: PUNO: El Collao, *O.P. Pearson* 91 (US). TACNA: Tacna, *P.M. Peterson* 13953 (MO, US). **Discussion:** Found from southern Peru, Bolivia, Argentina, and is here reported for Chile (*P.M. Peterson* 15676). Brako and Zarucchi (1993) state *P. pearsonii* to also occur in LIMA and AREQUIPA.

Poa pratensis L. **Ref:** Standley (1936: 128); Tovar (1993: 135); Giussani et al. (2012: 328). **Ill:** Giussani et al. (2012: sp. 50, p. 329). **Habitat:** Open Andean grasslands. 3500–3900 m. **Vouchers:** PERU: CAJAMARCA: Cajamarca, *I.M. Sánchez V.* 2668 (MO); San Miguel, *P.M. Peterson* 14921 (MO, US). HUANCVELICA: Huancavelica, *P.M. Peterson* 18107 (US). JUNÍN: Huancayo, *P.M. Peterson* 14223 (US); Yauli, *D.N. Smith* 2979 (MO). **Discussion:** Plants in Peru are presumably introduced from Europe, but the species in tropical latitudes tends

to look odd and cannot be confidently placed to subspecies. *Poa pratensis* subsp. *alpigena* (Lindm.) Hiitonon is apparently native in North and South America (in Patagonia). Brako and Zarucchi (1993) state *P. pratensis* to also occur in PUNO.

Poa ramifera Soreng & P.M. Peterson **Ref. & Ill:** Soreng and Peterson (2010: 587, fig. 1, 2, 3). **Habitat:** Shrublands. 2700–3100 m. **Vouchers:** PERU: ANCASH: Corongo, *P.M. Peterson 21804* (MO, US, USM). **Discussion:** Endemic, known only from the type locality.

Poa sanchez-vegae Soreng & P.M. Peterson **Syn:** *Aphanelytrum peruvianum* Sánchez Vega, P.M. Peterson, Soreng & Lægaard. **Ref & Ill:** Peterson and Soreng (2016: 118, fig. 4). **Habitat:** Rocky sites associated with Jalca vegetation (humid alpine grass ecosystems) at 3300 m. **Vouchers:** PERU: CAJAMARCA: Cajamarca, *I. Sánchez-Vega 11781* (CPUN, AAU, F, HAO, HUT, LOJA, MICH, MO, SI, US, USM). **Discussion:** Endemic, known only from the type locality.

Poa scabrivaginata Tovar **Ref:** Tovar (1965: 48, 1993: 134). **Ill:** Tovar (1965: lam. XI, A). **Habitat:** Shrublands. c. 4000 m. **Vouchers:** PERU: HUÁNUCO: Pachitea, Tambo de Vaca, *J.F. Macbride 4354* (MO, US-holotype). **Discussion:** Endemic herb. Known only from the type collection. A paratype from HUÁNUCO, and further specimens sometimes referred to as *P. scabrivaginata* from CAJAMARCA (e.g. Celendin, *I.M. Sánchez V. 2668* [MO]; Santa Cruz, *J. Santistaban C. & J. Guevara B. 169* [MO]) and JUNÍN (e.g. Concepción, *J.F. Macbride 3363* [US]) are better included in *P. aequatoriensis*. In the *P. scabrivaginata* type, lower sheaths are densely and coarsely scabrous, glumes are long (lower 3.5–4 mm, upper 4–4.5 mm), lemmas are glabrous except for the web and smooth except for sparsely scaberulous upper keel and apex, and anthers are 1.5–2 mm long. The other material has shorter glumes, sheaths smooth or lightly scabrous (often scabrous only on the margins), lemmas usually with pubescence on the keel and marginal veins, and anthers usually shorter.

Poa swallenii Refulio **Syn:** *Dissanthelium expansum* Swallen & Tovar. **Ref:** Refulio-Rodríguez et al. (2012: 130); Swallen and Tovar (1965: 374); Tovar 1993: 157). **Habitat:** Puna grasslands. 3600–4600 m. **Vouchers:** PERU: CUSCO: Calca, *P.M. Peterson 16594* (US), *P.M. Peterson 16569* (US), *P.M. Peterson 16612* (US); La Convención, *O.F. Cook 1305* (US), *S.P. Sylvester 1924* (US, USM, Z); Urubamba, *S.P. Sylvester 1071* (CUZ, US, Z). **Discussion:** Endemic herb. Swallen and Tovar (1965) and Tovar (1993) cite a collection of *Macbride & Featherstone 2183* from HUÁNUCO, and *P.C. Hutchison 1215* from LIMA.

Poa trivialis L. subsp. *trivialis* **Ref:** Standley (1936: 129); Tovar (1993: 134); Giussani et al. (2012: 336). **Ill:** Giussani et al. (2012: sp. 60, p. 337). **Habitat:** Andean slopes at middle elevations. **Vouchers:** PERU: JUNÍN: *A.S. Hitchcock 22277* (US), *A.S. Hitchcock 22279* (US), *A.S. Hitchcock 22266a* (US). **Discussion:** Introduced from Europe. Brako and Zarucchi (1993) state *P. trivialis* to also occur in CAJAMARCA. However, the collection cited by Brako and Zarucchi (1993), *J. Mostacero L. 1282* (MO), was redetermined as *P. aequatoriensis*. *Poa trivialis* and *P. aequatoriensis* bear many superficial resemblances and can be easily confused. The Hitchcock specimens had no habitat type or specific locality mentioned.

Poa trollii (Pilg.) Refulio **Syn:** *Dissanthelium trollii* Pilg. **Ref:** Tovar (1986: 51; 1993: 153). **Ill:** Renvoize (1998: fig. 38 G, H). **Habitat:** Dry Puna grassland. 4400–4800 m. **Vouchers:** PERU: PUNO: El Collao, *P.M. Peterson 18303* (US). **Discussion:** Distributed in the high Andean dry Puna grassland in southern Peru and the Potosí, Bolivia. This species normally has congested, spikelike panicles but has been collected with open panicles during anthesis.

Newly described species

***Poa ramoniana* Soreng & S.P. Sylvester, sp. nov.**

urn:lsid:ipni.org:names:77155738-1

Fig. 1

Type. PERU. Región JUNÍN. Prov. Junín: Distr. Carhuamayo, orillas de la laguna de Capillacocha, {est. vicinity: S10.86443°, W75.99256°} entre Carhuamayo y Paucartambo, Puna grassland, 4200–4300 m, 8 Jan 1949, *R. Ferreyra 5200 p.p. a* (holotype: US-2207173!; isotype: USM p.p.)

Plants gynomonoeious. **Perennials;** Rhizomatous, with well developed, slender, lateral tending, rhizomes, small tufted. **Tillers** extravaginal. **Culms** c. 5 cm tall, erect, unbranched, isolated or two together; Culm nodes terete, smooth, included in the sheaths; Culm internodes less than 1 cm long (peduncle c. 3 cm), terete, smooth. **Leaves** equally basal and cauline; Sheaths slightly laterally compressed, keeled, smooth, glabrous; Butt sheaths papery or slightly fibrous in age; Uppermost culm sheath c. 18 mm long, margins fused c. 40% their length, distal sheaths longer than their blades; Collars and throats smooth, glabrous; Ligules 1–1.5 mm long, sometimes with a central dent to 2 mm long, indistinctly decurrent, abaxially moderately densely scabrous, apices obtuse, margin irregular sometimes with a tooth, of sterile shoots similar to those of the culm; Cauline blades to 2 cm long, mostly folded or infrequently flat, with strongly inrolled margins, abaxially, marginally, and adaxially smooth, glabrous, tips distinctly prow shaped; Blades graduated up the culm, the sub-terminal one the longest; Sterile shoot blades to 4.5 cm long. **Panicles** 2–2.5 cm long, open, exerted, c. 1 cm wide, with 13–15 spikelets, proximal internode c. 6 mm long, weakly scabrous angled; Rachis with 1 branch per node; Primary branches spreading to reflexed, the upper ones ascending, distinctly angled, closely scabrous along the angles; Lateral pedicels less than 0.5 mm long, scabrous angled; Longest branches c. 7 mm, with 4–5 spikelets clustered in the distal half. **Spikelets** 3.5–4 mm long, c. 1.5 × longer than wide, broadly ovate, laterally compressed, not bulbiferous, anthocyanic and bronzy; Florets (2–)3, the proximal 1 (or 2 if 3 total) perfect, the distal 1 pistillate; Rachilla internodes 0.4–0.8 mm long, terete, distal internodes terete, smooth, glabrous, mostly hidden; Glumes more or less equal, both broadly lanceolate, or the first lanceolate, sub-lustrous in the scareous-hyaline margins, distinctly keeled, keels smooth or distally smooth or sparsely scaberulous, apices acute and pointed to obtuse and blunt and den-




Figure 1. *Poa ramoniana*. **A** Habit **B** Ligular zone **C** Terminal segment of inflorescence branch with four spikelet cluster **D** Spikelet **E** Lower glume ventral view **F** Proximal floret (perfect) **G** Upper glume dorsal view **H** Palea with immature pistil (pistillate floret), dorsal view **I** Palea with pistil (pistillate floret), staminodes, and lodicules, lateral view. Drawn from type material (*R. Ferreyra 5200 p.p. a*, US-2207173).

ticulate; Lower glumes 2.5–3 mm long, 1–3-veined; Upper glumes 3–3.2 mm long, subequal in width to the lower, 2–3-veined; Calluses glabrous; Lemmas (the lowest) 2.9–3.5, 5-veined, broadly lanceolate to ovate, green proximally, anthocyanic distally with a wide bronzy band apically, strongly laterally compressed, distinctly keeled, thin, keel smooth or obscurely scaberulous distally, smooth or mostly smooth elsewhere, glabrous, intermediate veins distinct, not extending into the scarious apical margin, edges smooth, apices scarious-hyaline bronzy for the distal c. 1 mm, edges smooth or slightly erose to denticulate, obtuse, blunt or slightly pointed; Paleas to 1 mm shorter than the lemma, glabrous, keels smooth or distally sparsely scaberulous. **Flowers** chasmogamous; Lodicules not observed; Anthers c. 2 mm long (vestigial in pistillate flowers). **Caryopses** unknown. $2n$ = unknown.

Distribution. Known only from a single locality in Junín.

Habitat. Puna grassland, 4200–4300 m, in wet margins/shore of lakes, in moss.

Etymology. The species is named in recognition of the eminent Peruvian botanist, Ramón Alejandro Ferreyra (1910–2005) who collected the type and paratype.

Conservation status. Data insufficient.

Additional specimens examined. PERU. Región JUNÍN. Prov. Junín: Distr. Carhuamayo, Capillacochoa cerca a Carhuamayo, Puna grassland, 4200–4300 m, 8 Jan 1949, *R. Ferreyra 5211* (USM!).

Discussion. This new species appears like a small form of *P. glaberrima*, but differs by being extensively rhizomatous and reaching only 5 cm tall, versus densely tufted and 12–45 cm tall for *P. glaberrima*. The US holotype of *P. ramoniana* is a mixed collection, with a second taxon p.p. “b”, which appears to be *Poa gymnantha* Pilg. that is sterile, tightly tufted, with intravaginally branching shoots and involute leaf blades that are adaxially scabrous. The USM isotype also contains two species: the small rhizomatous plant is *P. ramoniana*; the taller plants appear to be *P. glaberrima*. Tovar originally determined the USM type and paratype as *Poa lilloi* (Tovar 1993), which, among other differences, has a dense habit, without rhizomes, and ascending panicles branches, densely scabrous lemmas with narrow white, scarious margins, and sometimes a web on the callus. Tovar also identified the US type as *Poa ovata* Tovar (1965: 17), which RJS considers to be a rare to uncommon sexually reproducing phase of the small form of *P. gymnantha*, a species that is otherwise predominantly pistillate and apomictic (Negritto et al. 2008). Other material determined as *P. lilloi* in Peru has been referred to *P. glaberrima* and *P. candamoana*, or small *P. kurtzii* (see excluded species, below).

***Poa tayacajaensis* Soreng & S.P. Sylvester, sp. nov.**

urn:lsid:ipni.org:names:77155739-1

Figs 2, 3

Type. PERU: Región HUANCAMELICA. Prov. Tayacaja: Distr. Colcabamba, Chuspi-Hda., Tocas, entre Colcabamba y Paucarbamba, monte bajo, 2900 m, [vic. S12.5°, W74.6°], 22 Apr 1954, *Ó. Tovar*. S. 2038 (holotype: US-2181284; isotype: USM p.p.)




Figure 2. *Poa tayajajensis*. Digitized holotype specimen Ó. Tovar. S. 2038 (US-2181284).

Plants gynomonocious. **Perennials**; tufted, without lateral or downward tending shoots. **Tillers** intravaginal. **Culms** 55–65 cm tall, erect or decumbent (when decumbent sometimes extravaginally branching at the lower culm nodes, i.e. *Peterson et al.* 20369); Culm nodes 3–4, terete, smooth, 2–3 nodes exposed at flowering; Culm internodes terete, smooth. **Leaves**; Sheaths slightly laterally compressed, keeled, lower culm and lateral ones densely scaberulous distally; Butt sheaths thin papery, somewhat loosely investing the shoots; Uppermost culm sheaths 10–13 cm long, margins fused 60–75 % their length, slightly shorter than their blades; Collars and throats smooth, glabrous; Ligules 2.0–3.5 mm long, not decurrent, abaxially sparsely to moderately densely scabrous, apices obtuse to subacute, margins densely scabrous, ligules of sterile shoots and lower culm leaves 0.5–1 mm long; Cauline blades 6–15 cm long, 3–5 mm wide, well developed, longer than their sheaths, generally flat, keeled, thin, lax, abaxially, marginally, and adaxially scabrous mainly along the veins, folded near the apex, apex gradually tapered to a slender point; Blades gradually increasing in length up the culm, flag leaf blade 10–15 cm long; Sterile shoot blades like those of the culm, but somewhat shorter and smoother. **Panicles** 18–20 cm long, loose, open, exerted, slightly lax, to 5 cm wide, with more than 100 spikelets, proximal internodes 3.5–4 cm long, smooth; Rachis with (3) 5–6 branches at lower nodes; Primary branches slender, mostly laxly ascending, sometimes spreading, one sometimes reflexed, angled, proximally smooth to moderately scabrous mainly on the angles; Lateral pedicels mostly < 1 mm long, scabrous; Longest branches 6–8 cm long, with 14–22 spikelets in the distal half, slightly overlapping. **Spikelets** 4.5–6 mm long, c. 2 × longer than wide, lanceolate, laterally compressed, not bulbiferous, greyish-green to somewhat anthocyanic at maturity; Florets (3–)4(–5), proximal florets hermaphroditic and distal one pistillate; Rachilla internodes terete, distal internodes 0.7–1 mm long, terete, smooth, glabrous; Glumes unequal, narrow lanceolate to lanceolate, herbaceous and pale green below, sometimes anthocyanic in margins and apex, veins distinct, distinctly keeled, keels sparsely short scabrous distally, surfaces smooth, margins scarious-hyaline, edges entire smooth, apices sharply acute, entire; Lower glumes 1.7–2.5 mm, 2/3–4/5 as long as adjacent lemmas, 1-veined, very narrow, slightly sickle shaped; Upper glume 2.4–3 mm, c. 2 × wider than the lower, 3-veined; Calluses webbed, with a dense, long dorsal tuft of wooly hairs; Lemmas (the lowest) 2.8–3.7 mm long, 5-veined, lanceolate in side view, the proximal one c. 5 × longer than wide at maturity, greyish-green, to strongly anthocyanic at maturity, strongly laterally compressed, distinctly keeled, thin, keel to 3/4 the length and marginal veins and sometimes the intermediate veins to 1/2 the length, loosely sericous to villous, between veins sparsely to moderately densely appressed pubescent or occasionally glabrous on the proximal lemma, keel distally weakly scabrous, intermediate veins distinct, not extending to near the margin, margins inrolling below at maturity, very narrowly hyaline above, edges smooth or with a few hooks, apices acute, briefly hyaline; Paleas shorter than the lemmas by c. 0.5 mm, keels scabrous distally, sometimes weakly so, sparsely puberulent medially or nearly so, glabrous. **Flowers** chasmogamous; Lodicules c. 0.25 mm long, obscurely lobed;



Figure 3. *Poa tayacajaensis*. Photograph of individual floret, anther, and spikelet (from top to bottom) of holotype specimen Ó. Tovar. S. 2038 (US-2181284). (Photograph by R.J. Soreng).

Anthers 1.2–1.4 mm long, vestigial in the upper floret. **Caryopsis** 1.8–2 mm long, strongly laterally compressed, sulcate, honey brown, firm, adherent to the lemma and palea, hilum 0.2 mm long, elliptical. $2n =$ unknown.

Distribution. Endemic to the central Andes of Peru. Known from Huancavelica and Huánuco, although the Huánuco specimen is only tentatively placed.

Habitat. Shrublands on Andean slopes at mid elevations.

Conservation status. Data insufficient.

Additional specimens examined. One other specimen appears to represent this species but is too immature to be certain. The specimen in question has extravaginal shoots branching from lower culm nodes; PERU: Región HUÁNUCO. Prov. Pachitea: Distr. Chagglá, canyon of the Rio Grande, c. 20 km above confluence with Rio Huallaga, E of Huánuco c. 44 air km, 1.7 air km SSW of Estación Huacachay

(Huacachi), 2650 m, S9.86836 W75.83306, 8 Mar 2007, Peterson, Soreng & Romaschenko 20369 (US!).

Discussion. These plants bear similarities to *Poa aequatoriensis* but differ by having lemmas which are generally shorter (2.8–3.7 mm long), pubescent between the veins, and by more densely scabrous sheaths, with more-or-less smooth glabrous throats, ligules generally shorter. Tovar (1993) placed his voucher of this form (2038) in *P. aequatoriensis*. However, among the 20 sheets and the US isotype reviewed of *P. aequatoriensis*, all have lemmas that are smooth and glabrous between the veins (consistent with the description of Ecuadorian material by Hjorth, 1991), and the keel and marginal veins can be glabrous or sparsely puberulent. The species also bears some slight similarity to *Poa myriantha* Hack. and *P. hieronymi* Hack. from the Yungas cloud-forests, Argentina, that differ by size of the anthers and ligule being much smaller (anthers <1 mm long, ligules <1 mm long) and overall habit being larger (culms 60–350 cm long with 10–15 internodes, and panicles 20–36 cm long) and glumes having both antrorse and retrorse scabrocities.

***Poa urubambensis* S.P. Sylvester & Soreng, sp. nov.**

urn:lsid:ipni.org:names:77155740-1

Fig. 4

Type. PERU. Región CUSCO. Prov. Calca: Distr. Calca, top of the prominent tower known by locals as “Kontorqayku”, 5 km NE of Huarán, 4401 m, S13°16'05.9", W72°01'17.2", 27 May 2011, S.P. Sylvester 1317 (holotype: USM!; isotypes: CUZ!, K-000501720!, US!, Z-000099199!)

Plants gynomonocious. **Perennials or (rarely) annuals;** Rhizomatous with well developed, slender, lateral tending, rhizomes, solitary and erect, or rarely tufted, often rooting from nodes. **Tillers** extravaginal. **Culms** (3–)15–30(–42) cm tall, decumbent to ascending, sometimes erect, fairly slender, not branching above the base, leafy; Culm nodes 1–2(–3), terete or slightly compressed, smooth, usually 1 node exposed at flowering; Culm internodes 3–15(–20) cm long, terete, smooth. **Leaves** mostly basal; Sheaths slightly compressed to keeled, smooth or scabrous along the veins; Butt sheaths papery, smooth, glabrous; Uppermost culm sheaths (3.5–)8–16 cm long, margins fused 25–60% the length, 1.8–2.7 × longer than their blades; Collars and throats smooth or scabrous, glabrous, collar margins of sterile shoot leaves sometimes flared; Ligules 1–4 mm long, not decurrent, scarious to hyaline, adaxially glabrous to scabrous, upper margins entire or irregularly dentate, apices acute and entire to irregularly dentate above, sterile shoot ligules generally shorter and more scabrous than those of the culm leaves; Cauline blades 2–15(–22) cm long, (1.5–)2–3 mm wide, flat or folded, margins often becoming involute, thin to moderately thin, soft or (rarely) curved, surfaces abaxially and adaxially lightly to moderately scabrous or rarely smooth, margins scabrous, narrowly to abruptly prow-tipped; Mid-cauline blades the longest, 10–22 cm long, shorter upward, flag leaf blade 3.5–7.7 cm long;



Figure 4. *Poa urubambensis*. **A** Habit **B** Basal portion of plant showing extravaginal shoots **C** Inflorescence **D** Ligular zone **E** Segment of inflorescence **F** Spikelet **G** Lower glume dorsal view **H** Upper glume dorsal view **I** Proximal floret **J** Lemma, lateral view **K** Palea, ventral view **L** Pistil, stamens and lodicules from perfect floret. Drawn from type material (*S.P. Sylvester 1317*, US).

Sterile shoot blades similar to cauline blades, sometimes more involute. **Panicles** (4–)8–13 cm long, erect, loosely contracted to open, ovoid to narrowly pyramidal, sparsely to moderately congested, with 13–40(–80) spikelets, proximal internode 1.4–3 cm long, smooth or scaberulous, usually scabrous towards its apex; Rachis with (1–)2–3(–5) branches per node; Primary branches ascending, fairly flexuous, weakly angled, moderately scabrous; Lateral pedicels mostly $3/4$ to equaling the spikelets, moderately to densely scabrous, prickles moderately coarse; Longest branches 3–5 cm long, with 5–15 spikelets in distal $1/2$, loosely arranged. **Spikelets** 3.7–6.5 mm long, to $3.7 \times$ long as wide, lanceolate, laterally compressed, not bulbiferous, two toned; Florets 2–3(–4), proximal 1 or 2 florets hermaphroditic and distal 1 or 2 pistillate or sterile; Rachilla internodes terete, distal internodes 0.6–1 mm long, terete, smooth, glabrous; Glumes equal to subequal, narrow lanceolate, herbaceous and pale green below, scarious bronzy and sometimes anthocyanic in margins and apex, veins distinct, distinctly keeled, usually scabrous purely on the veins and sometimes between veins, margins scarious-hyaline, edges entire or dentate, smooth, apices acute, entire; Lower glumes 3.1–3.5 mm long, $2/3$ – $4/5$ as long as adjacent lemmas, 1-veined, narrow; Upper glumes 3.4–3.9 mm long, c. $2 \times$ wider than the lower, 3-veined; Calluses glabrous; Lemmas (the lowest) 3.2–3.9 mm long, 5-veined, lanceolate in side-view, the proximal one c. 4 – $8 \times$ longer than wide at maturity, proximally light green and distally bronzy-anthocyanic at maturity, moderately laterally compressed, thin, keeled, keels to $1/3$ – $5/6$ and marginal veins to $3/5$ – $4/5$, proximally smooth, keel and sides distally sparsely to moderately scaberulous, intermediate veins obscure to moderately prominent, not extending to near the margin, margins broadly scarious-hyaline, edges scabrous, apices acute; Paleas to 1.6 mm shorter than the lemma, glabrous, keels distally sparsely to moderately scabrous, between keels narrow (0.3–0.4 mm). **Flowers** chasmogamous; Lodicules c. 0.25 mm long, obscurely to shallowly lobed; Anthers 0.7–1.1(–1.3) mm long, infrequently vestigial in upper florets of spikelets. **Caryopses** c. 1.9 mm long, elliptical in side-view, sulcus broad and shallow, brown, hilum 0.2 mm long, oval, grain free from the palea. $2n =$ unknown.

Distribution. Restricted to undisturbed areas of *Polylepis* woodland in hard to access areas throughout the Cordillera Urubamba, Cusco, Peru, at 4390–4802 m. Known from three localities; 1) Cliff ledges of the prominent SW facing cliff face 1.5 km S (170°) of Cancha Cancha village, Huarán. 2) Ledges of the prominent tower known by locals as “Kontorqayku”, 5 km NE of Huarán. 3) Ridgeline to the W of Laguna Manalloqsa, Área de Conservación Privada (ACP) Mantamay, 10 km up the valley from Yanahuara in the small valley 3 km E of Laguna Ipsaycocha.

Habitat. Relatively dry and exposed sites in montane *Polylepis* forest and forest edges.

Etymology. The name ‘urubambensis’ refers to the Cordillera Urubamba.

Conservation status. This narrow endemic is locally common.

Additional specimens examined. PERU. Región CUSCO. Prov. Calca: Distr. Calca, large ledge situated on the prominent SW facing cliff face 1.5 km S (170°) of Cancha Cancha village, Huarán, 4524 m, S13°14'35.1", W72°01'14.1", 21 March

2011, *S.P. Sylvester 812* (CUZ!, MO!, US!, Z!); Distr. Calca, large ledge situated on the prominent SW facing cliff face 1.5 km S (170°) of Cancha Cancha village, Huarán, 4517 m, S13°14'35.0", W72°01'13.7", 24 March 2011, *S.P. Sylvester 869* (CUZ!, US!, Z!); Distr. Calca, within the SW facing forest at the top of the prominent tower known by locals as "Kontorqayku", 5 km NE of Huarán, 4390 m, S13°16'07.7", W72°01'16.8", 11 June 2012, *S.P. Sylvester 1636* (US!); Distr. Calca, within the SW facing forest at the top of the prominent tower known by locals as "Kontorqayku", 5 km NE of Huarán, 4390 m, S13°16'07.7", W72°01'16.8", 11 June 2012, *S.P. Sylvester 1637* (CUZ!, SI!, US!, Z!); Distr. Calca, top of the prominent tower known by locals as "Kontorqayku", 5 km NE of Huarán, 4401 m, S13°16'05.9", W72°01'17.2", 11 June 2012, *S.P. Sylvester 1695* (CUZ!, US!, Z!); Prov. Urubamba: Distr. Urubamba, ACP Mantamay, 10 km up the valley from Yanahuara in the small valley 3 km E of Laguna Ipsaycocha, ledges on cliff side 250°W of Laguna Manalloqsa, 4676 m, S13°12'01.3", W72°08'47.4", 28 January 2011, *S.P. Sylvester 403* (CUZ!, US!); Distr. Urubamba, ACP Mantamay, 10 km up the valley from Yanahuara in the small valley 3 km E of Laguna Ipsaycocha, topmost of the ridge to the W of Laguna Manalloqsa, 4802 m, S13°12'08.9", W72°08'43.9", 25 June 2012, *S.P. Sylvester 1727* (CUZ!, MO!, US!, Z!).

Discussion. This new species is similar to other members of *Poa* sect. *Homalopoa* s.l. from Peru, Bolivia and Argentina, all of which have open panicles and spikelets with 2–5(–8) florets, the lowermost florets hermaphroditic while the upper florets are pistillate. *Poa urubambensis* is easily recognised in the field by the combination of an open-panicled inflorescence, glabrous lemmas and calluses, and exceptionally small anthers for members of *Poa* supersect. *Homalopoa*. *Poa urubambensis* also bears resemblance to *Poa oscariana*, but is distinguished by a less robust habit, the leaf blades being mainly basal and the anthers being smaller.

This species was found during a large scale ecological study attempting to reconstruct the potential natural vegetation (PNV) and soils of the high-elevation Puna grasslands (see Heitkamp et al. 2014 and Sylvester et al. 2014 for pilot studies). In this research, pristine zonal vegetation, only accessible with mountaineering equipment, was compared with surrounding slopes which have been grazed and burnt consistently over millennia (Thompson et al. 1988; Chepstow-Lusty et al. 1996, 2009; Kuentz et al. 2011). *Poa urubambensis* was a common element in undisturbed Puna vegetation in the Cordillera Urubamba, being found associated with *Polylepis* Ruiz & Pav. forests from three different sites and also found growing alongside other species new to science, e.g. *Bartsia lydiae* S.P. Sylvester (2014: 41). Following indicator species analyses, *Poa urubambensis* has been found as an indicator species for the PNV, due to its frequency and abundance within relict patches of near natural vegetation (Sylvester et al., *unpubl. data*). This species has not been found in accessible, disturbed or secondary, vegetation at similar or lower elevations in the Andes of the Cuzco region, despite a more thorough botanical exploration. This may relate to its susceptibility to disturbance from grazing and burning (Sylvester, *pers. observation*).

Species excluded

Poa androgyna Pilg. **Ref:** Renvoize (1998: 144). **Discussion:** *Poa androgyna* has been described for Chile and Bolivia (Renvoize 1998; Soreng et al. 2003, 2016) with Renvoize (1998) mentioning the species to occur in Peru but not citing specimens and none have been encountered so far. The name is difficult to apply and taxonomists are still unsure of what it is exactly. Past authors (Hitchcock 1927; Standley 1936; Foster 1958; Brako and Zarucchi 1993) have placed this name as a synonym of *P. horridula*, which is quite likely, but we are not certain. Plants identified as *P. androgyna* have narrower leaf blades and more reflexed panicle branches than that typical of *P. horridula*.

Poa bromoides Vahl = ***Eragrostis bromoides*** (Vahl) Steud. **Ref:** Steudel (1854: 276). **Discussion:** Indiscrepancy centers around Brako and Zarucchi's (1993) inclusion of *P. bromoides* for Peru. As the type protologue of *Poa bromoides* states the plant to be cultivated: "Cult. Habitat (L)imae? Ex horto parisino habui" (Vahl 1794: 10), this effectively rules out the specimen being placed as *Poa*. Steudel's (1854) description also includes "spiculis lanceolatis 25-floris glabris" (pp. 276), i.e. spikelets lanceolate, 25-flowered, that places further doubt on the specimen belonging to *Poa*. Grassbase (Clayton et al. 2006 onwards) accepts *Poa bromoides* Vahl as *Eragrostis bromoides* (Vahl) Steud. {appl. uncert. but valid}. Soreng et al. (2016), also accept it as *E. bromoides*.

Poa lilloi Hack. **Ref:** Tovar (1965: 32; 1993: 127); Giussani et al. (2012: 319). **Ill:** Negritto and Antón (2000: fig. 11); Giussani et al. (2012: sp. 36, p. 319). **Discussion:** Specimens of *P. lilloi* cited by Tovar (1993) to occur in Peru, *Ó. Tovar S. 2501* (US), *J.R. Swallen 7060b* (US) and *J.R. Swallen 7068* (US) have been redetermined as stunted versions of *P. glaberrima*. The *J.R. Swallen 7060a* (US) specimen was redetermined as a stunted version of *P. candamoana*. The specimens of *P. lilloi* from JUNÍN cited by Brako and Zarucchi (1993) and Tovar (1993), *R. Ferreyra 5260* (US, USM) and *E. Cerrate 988* (USM[?]; Ill: Tovar 1965, lam. VIV, A.), have not been found. Certain specimens identified as *P. lilloi* may actually be the new species *Poa ramoniana*. The type of *P. ramoniana*, *R. Ferreyra 5200* (US, USM), was previously determined as *P. lilloi*, and the USM isotype appears as a mixed collection of *P. glaberrima* and *P. ramoniana*. It is highly probable that all other specimens previously considered as *P. lilloi* from Peru are stunted versions of either *P. candamoana*, *P. glaberrima* or *P. kurtzii* that were collected from heavily grazed areas. Correctly determined specimens of *P. lilloi* have been found from Bolivia, Chile and Argentina and occur in high elevation Puna grasslands above 3200 m. This species is most easily confused with *P. kurtzii* from which it can be distinguished by being 7–20 cm tall, scabrous across the glume and lemma surfaces, lemmas 3–3.5 mm long, sometimes with a bit of pubescence on the lemma keel, sometimes a tiny web emerging from the dorsal side of the callus. *Poa kurtzii* is 20–70 cm tall, has smooth glumes and densely scabrous lemmas, lemmas 4–4.5 mm long. *Poa kurtzii* is also gynomonocious, while *P. lilloi* is gynodioecious.

Poa supina Schrader **Ref:** Tovar (1993: 126); Soreng 2007: 529. **Ill:** Soreng (2007: p. 529). **Discussion:** This taxon is highly unlikely to occur in South America with all specimens identified as *P. supina*, so far encountered, pertaining to *Poa annua*. These include all specimens encountered from Ecuador and Bolivia (Simon Læggaard *pers. communication*). Tovar (1993) includes *P. supina* in his treatment of Peru separating it from *P. annua* and *P. infirma* by having glabrous lemmas, or only lightly pubescent on the nerves, and swollen culms. However, the main distinction between *P. supina* and *P. annua*, aside from sparser lemma pubescence, is in the length of the anthers with *P. supina* having anthers (1.2–)1.6–2(–2.5) mm while *P. annua* has anthers 0.7–1(–1.2) mm. Specimens examined of *P. supina* from Peru (*J. Espinoza* 2 [US]) were redetermined as *P. annua*, but the voucher collections *Tovar & Rivas Martinez* 7720 and *Tovar* 7855 accepted by Tovar (1993) have not been seen by us. Also see note under *Poa infirma* about the origin of *P. annua*.

Acknowledgements

This research was funded by a studentship grant from The Leverhulme Trust and an SNSF Early Postdoc Mobility Grant (Grant No. P2ZHP3_161988) to SPS and expedition grants to PMP from the Smithsonian Institution's Restricted Endowment Fund, the Scholarly Studies Program, Research Opportunities, Atherton Seidell Foundation, Biodiversity Surveys and Inventories Program, Small Grants, and the National Geographic Society for Research and Exploration (Grant Nos. 8848-10, 8087-06). Thanks go to Uwe Braun (HAL) for loaning the *Poa adusta* isotype. We thank Simon Læggaard (AAU), Diego Giraldo-Cañas (COL), Asunción Cano Echevarría (USM), Otakar Sida (PR) for providing information and checking specimens, and Gerrit Davidse (MO) for loans of *Poa* from Latin America to US. Michael Kessler, ECOAN-Peru and Washington Galiano are thanked for their support, especially with in-country logistics, for SPS, and Asunción Cano, Maria Isabel La Torre and Nancy Refulio Rodríguez (and others at USM) for same to PMP et al. during fieldwork in Peru. We thank Alice R. Tangerini for providing the Illustrations. Tarciso Filgueiras and an anonymous reviewer are thanked for helpful comments on an earlier version of this manuscript.

References

- Brako L, Zarucchi J (1993) Catalogue of the Flowering Plants and Gymnosperms in Perú. Monographs in Systematic Botany from the Missouri Botanical Garden 45: 1–1286.
- Češka A (2002) Tadeá Haenke. Botanical Electronic News BEN#287-288. <http://www.ou.edu/cas/botany-micro/ben/ben288.html> [accessed 24.10.2015]
- Chepstow-Lusty AJ, Bennett KD, Switsur VR, Kendall A (1996) 4000 years of human impact and vegetation change in the central Peruvian Andes – with events paralleling the Maya record? *Antiquity* 70: 824–833. doi: 10.1017/S0003598X0008409X

- Chepstow-Lusty AJ, Frogley MR, Bauer BS, Leng MJ, Boessenkool KP, Carcaillet C, Ali AA, Gioda A (2009) Putting the rise of the Inca Empire within a climatic and land management context. *Climate of the Past* 5: 375–388. doi: 10.5194/cp-5-375-2009
- Clayton WD, Vorontsova MS, Harman KT, Williamson H (2006 [onwards]) GrassBase – The Online World Grass Flora. <http://www.kew.org/data/grasses-db.html> [accessed 10.06.2014]
- Davidse G, Soreng RJ, Peterson PM (2010) *Poa unispiculata*, a new gynodioecious species of cushion grass from Perú with a single spikelet per inflorescence (Poaceae: Pooideae: Poaeae: Poinae). *Journal of the Botanical Research Institute of Texas* 4(1): 37–44.
- Foster RC (1958) A catalogue of the ferns and flowering plants of Bolivia. *Contributions from the Gray Herbarium of Harvard University* 184: 1–223.
- Gillespie LJ, Archambault A, Soreng RJ (2007) Phylogeny of *Poa* (Poaceae) based on *trnT-trnF* sequence data: major clades and basal relationships. *Aliso* 23: 420–434. doi: 10.5642/aliso.20072301.33
- Gillespie LJ, Soreng RJ, Bull R, Jacobs SWL, Refulio-Rodriguez NF (2008) Phylogenetic relationships in subtribe Poinae (Poaceae, Poaeae) based on nuclear ITS and chloroplast *trnT-trnF* sequences. *Botany* 86: 938–967. doi: 10.1139/B08-076
- Giussani LM, Antón AM, Negritto MA, Romanutti AA, Soreng RJ (2012) *Poa* L. In: Zuloaga FO, Rugulo ZE, Antón AM (Eds) *Flora Vascular de la República Argentina: Volúmen 3 Tomo II, Monocotyledoneae: Poaceae: Pooideae*. Graficamente Ediciones, Cordoba, 284–338.
- Giraldo-Cañas D (2011) Catálogo de la familia Poaceae en Colombia. *Darwiniana* 49: 139–247.
- Heitkamp F, Sylvester SP, Kessler M, Sylvester MDPV, Jungkunst HF (2014) Inaccessible Andean sites reveal human induced weathering in grazed soils. *Progress in Physical Geography* 38(5): 576–601. doi: 10.1177/0309133314544918
- Hitchcock AS (1927) The grasses of Ecuador, Perú, and Bolivia. *Contributions from the United States National Herbarium* 24(8): 291–556.
- Hjorth U (1991) Slægt. *Poa*. Unpublished MSc. Thesis, Aarhus University, Århus.
- Kuentz A, Ledru M, Thouret J (2011) Environmental changes in the highlands of the western Andean Cordillera, southern Peru, during the Holocene. *Holocene* 22(11): 1215–1226. doi: 10.1177/0959683611409772
- Læggaard S (1998) New species and names in Ecuadorian grasses (Poaceae). *Novon* 8: 23–30. doi: 10.2307/3391886
- León Yáñez S, Valencia Reyes R, Pitman NCA, Endara L, Ulloa Ulloa C, Navarrete H (2011) *Libro Rojo de Plantas Endémicas de Ecuador* (2 ed.). Herbario QCA, Pontificia Universidad Católica del Ecuador, Quito, 1–957.
- Müller C, Müller GK, Gutte P (1981) Neufunde für die peruanische Flora: Gramineae, Cyperaceae. *Willdenowia* 11(2): 333–341.
- Negritto MA, Antón AM (2000) Revisión de las especies de *Poa* (Poaceae) del noroeste argentino. *Kurtziana* 28(1): 95–136.
- Negritto MA, Romanutti AA, Acosta MC, Moscone EA, Cocucci AE, Antón AM (2008) Morphology, reproduction and karyology in the rare Andean *Poa gymnantha*. *Taxon* 57(1): 171–178.
- Negritto MA, Antón AM (2000) Revisión de las especies de *Poa* (Poaceae) del noroeste argentino. *Kurtziana* 28(1): 95–136.

- Negritto MA, Antón AM (2006) Three new species of *Poa* from the Andes of Colombia and Perú. *Systematic Botany* 31(1): 83–88. doi: 10.1600/036364406775971723
- Peterson PM, Soreng RJ (2016) A revision of *Poa* subsection *Aphanelytrum* (Poaceae, Pooideae, Poaceae, Poinae); and a new species, *Poa auriculata*. *PhytoKeys* 63: 107–125. doi: 10.3897/phytokeys.63.8198
- Pilger RKF (1906) *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 37: 381.
- Presl CB (1825–1835) *Reliquiae Haenkeanae seu descriptiones et icones plantarum, quas in America meridionali et boreali, in insulis Philippinis et Marianis collegit Thaddaeus Haenke, Philosophiae Doctor, Phytographus regis Hispaniae*. Prague.
- Presl J (1830) *Reliquiae Haenkeanae, seu, Descriptiones et icones plantarum quas in America Meridionali et Boreali. Reliquiae Haenkeanae* 1(4–5): 271.
- Refugio-Rodríguez N, Columbus JT, Gillespie LJ, Peterson PM, Soreng RJ (2012) Molecular phylogeny of *Dissanthelium* (Poaceae: Pooideae) and its taxonomic implications. *Systematic Botany* 37(1): 122–133. doi: 10.1600/036364412X616701
- Renvoize SA (1998) *Gramíneas de Bolivia*. Continental Printing, Belgium.
- Soreng RJ (1998) An infrageneric classification for *Poa* in North America, and other notes on sections, species, and subspecies of *Poa*, *Puccinellia*, and *Dissanthelium* (Poaceae). *Novon* 8(3): 187–202. doi: 10.2307/3391995
- Soreng RJ (2007) *Poa* L. In: Barkworth ME, Capels KM, Long S, Anderton LK, Piep MB (Eds) *Flora of North America North of Mexico*, vol. 24. Oxford University Press, New York, 486–601.
- Soreng RJ, Bull RD, Gillespie LJ (2010) Phylogeny and reticulation in *Poa* based on plastid *trnT*-*trnL* and nrITS sequences with attention to diploids. In: Seberg O, Petersen G, Barford AS, Davis JI (Eds) *Diversity, Phylogeny, and Evolution in the Monocotyledons*. Aarhus University Press, Aarhus, 619–644.
- Soreng RJ, Davidse G, Peterson PM, Zuloaga FO, Judziewicz EJ, Filgueiras TS, Morrone O, Romaschenko K (2016) *Catalogue of New World Grasses*. <http://www.tropicos.org/project/cnwg> [accessed 16.6.2016]
- Soreng RJ, Giussani LM, Negritto MA (2003) *Poa* L. In: Soreng RJ, Peterson PM, Davidse G, Judziewicz FO, Zuloaga EJ, Filgueiras TS, Morrone O (Eds) *Catalogue of New World grasses (Poaceae): IV. Subfamily Pooideae*. *Contributions from the US National Herbarium* 48: 505–580.
- Soreng RJ, Peterson PM (2010) *Poa ramifer* (Poaceae: Pooideae: Poaceae: Poinae), a new aerially branching gynomonocous species from Perú. *Journal of the Botanical Research Institute of Texas* 4(2): 587–594.
- Soreng RJ, Peterson PM (2012) Revision of *Poa* L. (Poaceae, Pooideae, Poaceae, Poinae) in Mexico: new records, re-evaluation of *P. ruprechtii*, and two new species, *P. palmeri* and *P. wendtii*. *PhytoKeys* 15: 1–104. doi: 10.3897/phytokeys.15.3084
- Soreng RJ, Peterson PM, Romaschenko K, Davidse G, Zuloaga FO, Judziewicz EJ, Filgueiras TS, Davis JI, Morrone O (2015) A worldwide phylogenetic classification of the Poaceae (Gramineae). *Journal of Systematics and Evolution* 53: 117–137. doi: 10.1111/jse.12150

- Standley PC (1936) Gramineae. In: Macbride JF (Ed) Flora of Perú. Publications of the Field Museum of Natural History, Chicago, Botanical Series 13(1/1): 96–261.
- Steudel EGv (1854) Synopsis Plantarum Glumacearum 1. J.B. Metzler, Stuttgart, 475.
- Swallen JR, Tovar Ó (1965) The grass genus *Dissanthelium*. *Phytologia* 11(6): 361–376. doi: 10.5962/bhl.part.24953
- Sylvester SP (2014) *Bartsia lydiae*, a new species of *Bartsia* sect. *Laxae* (Orobanchaceae) from the southern Peruvian Andes with a revised key to *Bartsia* sect. *Laxae*. *Phytotaxa* 164(1): 041–046.
- Sylvester SP, Sylvester MDPV, Kessler M (2014) Inaccessible ledges as refuges for the natural vegetation of the high Andes. *Journal of Vegetation Science*. doi: 10.1111/jvs.12176
- Thiers B. [continuously updated]. Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/> [accessed 01.02.2014]
- Thompson LG, Davis ME, Mosley-Thompson E, Liu K-b (1988) Pre-Incan agricultural activity recorded in dust layers in two tropical ice cores. *Nature* 336(6201): 763–765. doi: 10.1038/336763a0
- Tovar Ó (1965) Revisión de las especies peruanas del género *Poa*. *Memorias del Museo de Historia Natural "Javier Prado"* 15: 1–65.
- Tovar Ó (1974) Dos especies nuevas de *Poa* (Gramineae) para el Perú. *Boletín de la Sociedad Peruana de Botánica* 7: 6–10.
- Tovar Ó (1984) Seis especies nuevas de Gramineae del Perú. *Publicaciones del Museo de Historia Natural Javier Prado. Serie B. Botánica* 32: 1–12.
- Tovar Ó (1986) Nuevos registros de gramíneas para el Perú. *Revista de Ciencias de la Universidad Nacional Mayor de San Marcos* 74: 48–57.
- Tovar Ó (1993) Las Gramíneas (Poaceae) del Perú. *Ruizia* 13: 1–480.
- Vahl M (1794) *Symbolae Botanicae* 2 [3 in IGS]: 10.

Supplementary material I

Supplementary key to the open-panicled species of *Poa* in Peru

Authors: Steven P. Sylvester, Robert J. Soreng, Paul M. Peterson, Mitsy D.P.V. Sylvester
Data type: Adobe PDF file

Explanation note: This morphological key emphasizes lemma indumentum.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.