

Nicotiana gandarela (Solanaceae), a new species of ‘tobacco’ highly endangered from the Quadrilátero Ferrífero in Brazil

Mariana Augsten¹, Pablo Burkowski Meyer¹, Loreta B. Freitas²,
João A. N. Batista¹, João Renato Stehmann¹

1 Departamento de Botânica, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais – UFMG, Av. Antônio Carlos, 6627, Pampulha, Belo Horizonte, CEP 31270–901, MG, Brazil **2** Departamento de Genética, Universidade Federal do Rio Grande do Sul, 91501–970, Porto Alegre, RS, Brazil

Corresponding author: Mariana Augsten (mariana.augsten@gmail.com)

Academic editor: Sandy Knapp | Received 4 October 2021 | Accepted 17 January 2022 | Published 25 February 2022

Citation: Augsten M, Meyer PB, Freitas LB, Batista JAN, Stehmann JR (2022) *Nicotiana gandarela* (Solanaceae), a new species of ‘tobacco’ highly endangered from the Quadrilátero Ferrífero in Brazil. *PhytoKeys* 190: 113–129. <https://doi.org/10.3897/phytokeys.190.76111>

Abstract

Nicotiana gandarela Augsten & Stehmann (Solanaceae), **sp. nov.**, a small ‘tobacco’ known only from one locality at Serra do Gandarela, in the state of Minas Gerais, Brazil, is described and illustrated. It is morphologically characterized by its rosulate basal leaves, red corolla with a short tube not inflated at the apex, and the peculiar habitat, a shaded site under a rocky outcrop ledge along a forested stream. Phylogenetic analyses based on a combined dataset of nuclear (ITS) and plastid (*ndhF*, *trnLF*, and *trnSG*) DNA sequences revealed that the species belongs to the *Nicotiana* sect. *Alatae* and is sister to the clade with the remaining species in the section. A key for the identification of Brazilian species of the section is given. The unusual habitat, the small population size, and the intense pressure of mining activities in the surroundings made the species assessed as Critically Endangered (CR), needing conservation efforts to avoid its extinction.

Keywords

Endemism, IUCN, Minas Gerais, mining, molecular phylogeny

Introduction

Nicotiana L. is a Solanaceae genus known mostly because of tobacco (*N. tabacum* L.), a crop cultivated worldwide, but its diversity goes further. It is the fifth-largest genus in the family, with 82 species (Knapp 2020). *Nicotiana* occurs in South America and Australia, with one species from Africa (Namibia). The Andean region is recognized as the center of diversity of the group in the Americas, as for many other clades in Solanaceae (Goodspeed 1954; Dupin et al. 2017). Molecular phylogenies are available for the genus (Aoki and Ito 2000; Chase et al. 2003; Clarkson et al. 2004; Clarkson et al. 2017), and sectional revision is provided based on phylogenetic analyses, indicating 13 sections (Knapp et al. 2004).

In Brazil, the known native species belong to the *Nicotiana* sect. *Alatae* Goodsp. (hereafter *Alatae*). This section is morphologically characterized by herbaceous habit, rosulate and sessile leaves, usually viscid pubescent, few leaves on upper stems, and an abrupt dilatation at the throat of the corolla tube (Knapp et al. 2004). The section comprises two cytogenetic groups: one displays chromosome number $2n = 20$ and includes two species, *N. longiflora* Cav. and *N. plumbaginifolia* Viv., distributed from Argentina to the USA; the other presents chromosome number $2n = 18$ and includes *N. alata* Link & Otto, *N. bonariensis* Lehm., *N. forgetiana* Hemsl., *N. langsdorffii* Weinm., *N. mutabilis* Stehmann & Semir and *N. azambujae* L. B. Sm. & Downs, occurring in Brazil, Argentina, Uruguay, and Paraguay (Goodspeed 1954; Clarkson et al. 2004; Knapp et al. 2004). The internal phylogenetic relationships of the section are still not well resolved, with the two groups recovered as sisters (Clarkson et al. 2004) or with the $2n = 18$ group as paraphyletic (Clarkson et al. 2017). According to the plastid or nuclear markers used for the reconstruction, section diversification is proposed between 6.2 Ma and 7.7 Ma, respectively (Clarkson et al. 2017).

Nicotiana can be considered a taxonomically well-studied genus for the Brazilian flora (Vignoli-Silva and Mentz 2005; Vignoli-Silva and Stehmann 2020), in which six native, *N. alata*, *N. bonariensis*, *N. langsdorffii*, *N. forgetiana*, *N. mutabilis*, and *N. azambujae*, and two naturalized species, *N. tabacum*, and *N. glauca* Graham, are recognized. Morphometric analyses based on the corolla size and shape of *N. forgetiana* throughout its geographical distribution showed that the species seems to include two distinct taxonomic groups, one not yet formally described (Teixeira et al. 2021). Among the Brazilian species, only *N. mutabilis* is considered endangered (Vulnerable) because of its small geographic distribution and just a few known populations (Stehmann et al. 2002; CNCFlora [<http://cncflora.jbrj.gov.br/portal/pt-br/profile/Nicotiana%20mutabilis>]). *Nicotiana azambujae* needs further investigation, as it is only known from the type collection.

Within Brazil, the Southern region holds the largest number of *Nicotiana* species, which inhabit ecotonal environments between the grasslands and the *Araucaria* forest (Mixed Ombrophilous Forest) in the Atlantic Forest domain (Teixeira et al. 1986; Overbeck et al. 2007). This is an important biogeographic area for several taxa that have their evolutionary history related to this vegetation mosaic (Iganci et al. 2011;

Fregonezi et al. 2013). Species richness decreases towards the north, with *N. bonariensis* being the species with the northern-most distribution, reaching the grasslands at the high altitude regions of the Mantiqueira Range (Minas Gerais and Rio de Janeiro states) and the Espinhaço Range (Minas Gerais and Bahia states).

During floristic studies in the Quadrilátero Ferrífero, a mountainous area with large mineral reserves in Minas Gerais, with significant biological importance and high anthropic pressures (Drummond et al. 2005; Jacobi et al. 2011), samples of a small *Nicotiana* specimen with red flowers were collected. As the phenotype matched with none of the known *Nicotiana* species from Brazil, we assumed it could be a new species. In this work, we test this hypothesis using morphological and molecular data.

Methods

Taxonomy

Plant samples were collected during field expeditions in April 2018, and voucher specimens were deposited in the BHCB herbarium at the Universidade Federal de Minas Gerais, whereas flowers were preserved in Ethanol 50%. We took measures of vegetative parts from herbarium samples and of entire flowers from fixed material. Descriptive terminology was based on Radford et al. (1976). To compare the morphological characters with congeneric species and build the key, we revised the main taxonomic literature (Goodspeed 1954; Vignoli-Silva and Mentz 2005; Cocucci 2013; Vignoli-Silva and Stehmann 2020), including the protologues, as well as the materials from the following Brazilian herbaria: BHCB, HUEFS, ICN, MBM, OUPR, PAMG, RB, R, SP, SPF, and UEC. We also examined high-resolution images of the types available at Global Plants (<https://plants.jstor.org>).

Sampling for phylogenetic analyses and molecular methods

We included DNA sequences from 56 *Nicotiana* species, mostly generated in previous studies (Chase et al. 2003; Clarkson et al. 2004) and obtained from GenBank (www.ncbi.nlm.nih.gov/Genbank). We selected species from all described sections (Knapp et al. 2004) and used two *Anthocercis* Labill. as outgroups. We generated DNA sequences for three different individuals of the new species, also for *N. mutabilis*, described in 2002 (Stehmann et al. 2002) and for which there were no DNA sequences available in the databases. For all Genbank accessions used in this work, see Suppl. material 1: Table S1. Fresh, young leaves were collected in the field and stored in silica gel. Total DNA was obtained using a modified version of the 2X-CTAB extraction method (Doyle and Doyle 1987) using 50 mg of tissue mass and washing with 70% ethanol. We used nucleotide sequences from the nuclear marker (ITS - rDNA internal transcribed spacers) and three plastid regions (*trnLF*, *trnSG*, and *ndhF*). The selection of markers was based on previous studies (Chase et al. 2003; Clarkson et al. 2004). Amplifications

were performed using the same conditions described in previous works (Chase et al. 2003; Clarkson et al. 2004). Polymerase chain reaction (PCR) products were purified by precipitation with 20% PEG (2.5 M NaCl and 20% polyethylene glycol 8000) and sequenced on demand by a specialized company using the same primers employed for amplification. We obtained bidirectional sequence reads for all the DNA regions, and consensus sequences were obtained using Geneious Prime 2020.0.3 (<https://www.geneious.com>). Sequences were aligned using CLUSTALW (Larkin et al. 2007) followed by manual adjustments in MEGA 11 (Tamura et al. 2021). Sequences were concatenated with MESQUITE (Maddison and Maddison 2019), and we obtained the best substitution model per DNA region using MRMODELTEST2 (Nylander 2004).

Phylogenetic analyses

We analyzed the data using Bayesian inference (BI) and Maximum Likelihood (ML). Initially, nuclear and plastid DNA were separately analyzed and posteriorly combined. We performed BI analyses using MRBAYES 3.2.7a (Ronquist et al. 2012) as implemented in the Cyberinfrastructure for Phylogenetic Research (CIPRES) Portal 2.0 (Miller et al. 2010), treating each DNA region as a separate partition. The “unlink” command was used to estimate model parameters separately for each partition. Each analysis consisted of two independent runs with four chains for 7,500,000 generations, sampling one tree every 1,000 generations and a temperature parameter of 0.2. Convergence between the runs was evaluated using the average standard deviation of split frequencies (< 0.01) and the Potential Scale Reduction Factor – PRSF (= 1.0) and was achieved after 715,000 generations. After discarding the first 2,500 trees (33%) as the burn-in, the remaining trees were used to assess topology and posterior probabilities (PP) in a 50% majority-rule consensus. We considered groups with PP > 0.95 as strongly supported, groups with PP ranging from 0.90–0.95 as moderately supported, and groups with PP < 0.90 as weakly supported. Maximum likelihood (ML) analysis of the concatenated dataset was performed using the online version of RAxML-HPC BlackBox (v.8.2.12) (Stamatakis 2014) through the CIPRES Science Gateway Portal (Miller et al. 2010). Bootstrap values were calculated based on 1000 replicates.

Conservation status

The conservation status was assessed based on the International Union for Conservation of Nature Criteria (IUCN 2019), considering the main threats to the *Quadrilátero Ferrífero* vegetation (Sonder et al. 2014).

Scanning electron microscopy analysis

Seeds were affixed in aluminum stubs using double-sided carbon tape, metalized with 10 nm gold-palladium (Robards 1978). The observations were performed under a JEOL JSM– 6360LV scanning electron microscope, with 5 kV (Jeol Ltd., Tokyo, Japan).

Results

Alignment characterization

The concatenated matrix consisted of 3,483 aligned characters, of which 317 (9.1%) were potentially informative. The ITS region had, proportionately, the highest number of phylogenetically-informative characters (20%). General features of the datasets and a summary of the models implemented for each partition are presented in Table 1.

Table 1. General features of the markers used in the phylogenetic analyses and models implemented for each.

Regions	Terminals	Characters	Variable uninformative characters	Informative characters (%)	Model
ITS	59	683	89	135 (20%)	GTR+I+G
<i>ndhF</i>	60	1056	49	71 (6.7%)	GTR+G
<i>trnLF</i>	60	991	41	58 (5.8%)	GTR+G
<i>trnSG</i>	60	855	38	62 (7.2%)	GTR+G
All plastid regions	60	2799	125	182 (6.5%)	–
All regions	60	3483	214	317 (9.1%)	–

Phylogenetic relationships

In the Bayesian phylogenetic tree based on ITS, *N. gandarela* appears within the section Alatae, the Alatae clade is moderately supported (PP = 0.91), and *N. gandarela* is sister of the 2n = 20 clade (*N. longiflora* and *N. plumbaginifolia*), being weakly supported (PP = 0.76) (Suppl. material 2: Fig. S1). The concatenated plastid regions recovered a Bayesian phylogenetic tree with *N. gandarela* as the sister species of section Noctiflorae, moderately supported (PP = 0.91), and Alatae appears as sister of section Repandae, weakly supported (PP = 0.84) (Suppl. material 3: Fig. S2).

Due to the combined dataset produced higher-supported trees in both ML and BI methods than using nuclear and plastid regions independently, we described the results based on this supermatrix only. As the topology was similar in ML and BI trees, we detailed the species relationships from BI results only. The three individuals of *N. gandarela* formed a well-supported clade in all obtained trees.

The *Nicotiana* species analyzed formed a strongly supported clade (PP = 1.00), and several subclades corresponded to the accepted sections for the genus (Fig. 1). The analysis confirmed the monophyly of the Alatae (PP = 0.99), which is strongly supported as the sister of the *Nicotiana* sect. *Suaveolens* Goodsp. (PP = 1.00). The two newly sequenced species, *N. gandarela* and *N. mutabilis*, were resolved as members of the Alatae. *Nicotiana gandarela* was recovered with high support (PP = 0.99) as sister to the clade with the remaining species in the Alatae, whereas *N. mutabilis* was resolved as closely related to *N. forgetiana* and *N. bonariensis* (PP = 1.00). Similar to previous analyses, low support was observed for the 2n = 18 group and for the relationship of *N. bonariensis* to *N. langsdorffii* (PP = 0.66).




Figure 1. Bayesian 50% majority-rule consensus tree of 56 species of *Nicotiana* L. of the combined nuclear (ITS) and plastid (*ndhF*, *trnLF*, and *trnSG*) datasets. Numbers next to the nodes represent the posterior probabilities (PP) and bootstrap values. Section names (according to Knapp et al. 2004) are indicated on the right. Width of lines defined by PP values.

Taxonomic treatment

Nicotiana gandarela Augsten & Stehmann, sp. nov.

urn:lsid:ipni.org:names:77260722-1

Fig. 2

Type. BRAZIL. Minas Gerais. Santa Bárbara, André do Mato Dentro, 19°59'43"S, 43°38'39"W, 17 Apr 2018 (fl, fr), *M. Augsten & J.R. Stehmann* 1078 (holotype: BHCB acc.#190958 [BHCB190958!]; isotypes: ICN, RB, to be distributed).

Diagnosis. *Nicotiana gandarela* differs from all other species of the *Nicotiana* sect. *Alatae* by its short corolla tube (< 15 mm), vivid red corolla limb, and unusual shaded, cave-mouth habitat.

Description. Annual herb to 0.5 m high, not branched, pubescent with multicellular, glandular trichomes. Leaves simple, rosulate, crowded, 6–21 cm long, (1.7–) 3.5–5.0 cm wide, persistent, spatulate, slightly discoloured, the blade hirsute throughout with simple, predominantly glandular hairs, long-attenuate at the base, margin sinuous, ciliate, midribs, and secondary veins visible at both surfaces, rounded or obtuse, sometimes acute, at the apex. Inflorescence scapose, paniculate, apical or lateral, composed of monochasial cymes; bracts lanceolate, 4.3 mm long, 1.0 mm wide, viscid-hirsute; pedicels 5.8–7.7 mm long. Calyx hirsute-glandular, deeply lobed, the tube 2.5 mm long, lobes 3–5 mm long, lanceolate, unequal, the apex acute. Corolla aestivation conduplicate, with folded basal petals covering the other three; tube 1.2–1.5 cm long, 3.0–3.4 mm diam., infundibuliform, not inflated at the apex, magenta, trichomes rare, opening ca. 3.4 mm; limb zygomorphic, red, cleft into widely-depressed ovate to very widely ovate lobes, patent or slightly reflexed. Stamens 5, didynamous, adnate ca. 5 mm from the base of the corolla tube, four longer, ca. 12.5 mm long, one shorter, ca. 11.4 mm long, all filaments glabrous; anthers 0.9–1.2 mm long, white, ellipsoid, pollen whitish; nectariferous disk present. Ovary slightly conical, glabrous, style ca. 9.4 mm long, stigma bilobed, green. Capsule 2-valved, included in the calyx, 6.0–7.5 mm long, 4.4–5.5 mm in diameter, ovoid, valves with incised apex. Seeds about 0.7–0.8 mm long, 0.6–6.5 mm wide, subglobose to obovoid, testa foveolate, anticlinal walls sinuous. Chromosome number unknown.

Etymology. The specific epithet “gandarela” is a noun in apposition and refers to Serra do Gandarela, the mountain range complex where this species is found.

Distribution and habitat. The only known population of *Nicotiana gandarela* is at the Serra da Gandarela Mountain range, located in the northeast portion of the Quadrilátero Ferrífero in the Minas Gerais Brazilian State (Fig. 3A). The species occurs in a shaded site placed under a rocky outcrop ledge that extends for about 350 m along a stream and surrounded by a forest matrix, making this area a permanent humid environment. We recognized three subpopulations, each up to 150 individuals, including many seedlings (Fig. 2B). This environment resembles an open cave, and the individuals grow in bare powdery red soil originating from rock decomposition, usually with no other species co-occurring. The habitat is unique, and no similar microhabitat exists in this geologically diverse and complex area (Instituto Prístino: Atlas Digital

Geoambiental: Quadrilátero Ferrífero: Geodiversidade [<https://institutopristino.org.br/atlas/quadrilatero-ferrifero/>] Fig. 2A–B.

Phenology. *Nicotiana gandarela* has been collected with flowers and fruits in December and April.

Conservation status. Although the single site for *N. gandarela* would normally suggest the species was Data Deficient (DD), the precarious nature of the region




Figure 2. *Nicotiana gandarela* Augsten & Stehmann **A** peculiar habitat of the species, the shaded sites in the base of the shaded ledge **B** seedlings growing in the site **C** habit highlighting the rosulate leaves and the scapose inflorescence **D**, **E** flowers in lateral and frontal view **F** 2-valvate capsule with many seeds. **G** seed with sinuous anticlinal walls (Scanning Electron Microscopy) **D–G** were obtained from plants of the type population (Augsten and Stehmann 1078, BHCB). Photos by JR Stehmann.

in which it grows leads us to assess it more formally. The species was preliminarily assessed as Critically Endangered – CR (B1, B2a, biii, D), mainly due to its geographic range, as the extent of occurrence (EOO) and the area of occupancy (AOO) are smaller than 100 km² and 10 km², respectively. The only populations of *N. gandarela* are located on private property and not inserted in any integral conservation unit. In addition, the Quadrilátero Ferrífero suffers from constant habitat loss (Salles et al. 2018) as well as strong pressures from the mining sector (Sonter et al. 2014) with concessions required to explore iron and gold (Instituto Prístino: Atlas Digital Geoambiental 2021 [<https://institutopristino.org.br/atlas/quadrilatero-ferrifero/>]) (Fig. 3B).

Additional specimen examined. BRAZIL. Minas Gerais: Santa Bárbara. Andr  do Mato Dentro, trilha para a cachoeira, 31 Dec 2017 (fl., fr.), *D.M.G. Oliveira, P.L. Viana & N.O. Mota 359* (BHCB).

Identification key to the Brazilian species of *Nicotiana* sect. *Alatae*

- 1 Calyx length equal to the corolla tube.... ***N. azambujae* L.B. Sm. & Downs**
- Calyx length shorter than the corolla tube **2**
- 2 Corolla yellow-greenish or light green, actinomorphic, limb almost entire (shortly lobed), pollen blue ***N. langsdorffii* Weinm.**
- Corolla white, pink, magenta, purple or red, zygomorphic, limb clearly lobed; pollen green, white or yellow **3**
- 3 Corolla tube > 30 mm long, limb usually white, filaments adnate to the upper portion of the corolla **4**
- Corolla tube usually < 30 mm long, limb rarely white, filaments adnate to the lower portion of the corolla tube **5**
- 4 Leaves decurrent, limb strongly zygomorphic, stamens inserted in two levels (4 higher and 1 lower) at the corolla tube, filaments all geniculate ***N. alata* Link & Otto**
- Leaves auriculate, not decurrent, limb weakly zygomorphic, stamens inserted in three levels (2 higher, 2 middle, and 1 lower) at the corolla tube, filaments not geniculate ***N. longiflora* Cav.**
- 5 Corolla changing from white to pink or magenta during the anthesis ***N. mutabilis* Stehmann & Semir**
- Corolla keeping the same color during the anthesis **6**
- 6 Leaves evenly distributed and decurrent along the stem; calyx usually > 9 mm; corolla tube usually > 20 mm long, magenta or purple-red ***N. forgetiana* Hemsl.**
- Leaves usually rosulate, not decurrent on the stem; calyx usually < 9 mm long; corolla tube < 20 mm long, limb red or white **7**
- 7 Corolla tubular, distally ventricose; limb white ***N. bonariensis* Lehm.**
- Corolla tube gradually enlarged to the apex, not ventricose, limb red ***N. gandarela* Augsten & Stehmann**




Figure 3. **A** main mountains of the Quadrilátero Ferrífero in Minas Gerais. Star indicates the *Nicotiana gandarela* Augsten & Stehmann occurrence site **B** Google Earth image showing the regional landscape associated with the *N. gandarela* site, the forest matrix impacted by a small village, and open mining. The red lines indicate mining concessions (Instituto Prístino: Atlas Digital Geoambiental [<https://instituto-pristino.org.br/atlas/>]).

Discussion

Overall, the recovered phylogenetic relationships were similar to those previously obtained (Chase et al. 2003; Clarkson et al. 2004; Knapp et al. 2004; Clarkson et al. 2017). Phylogenetic trees generated showed that the samples collected in the Serra da Gandarela in Minas Gerais are genetically distinct from all known species of *Nicotiana* used in our analyses. *Nicotiana gandarela* was recovered as the sister of all other species belonging to the Alatae (both of the groups with distinct cytotypes, $2n = 18$ and $2n = 20$; Clarkson et al. 2004; Knapp et al. 2004). The internal nodes of the group $2n = 18$ showed low support, and the lack of chromosome accounts to the new species does not permit any inferences about the phylogenetic relationships within the section.

The new species is an annual plant, with rosulate leaves, long-attenuate to the base, sessile, viscid-pubescent, and a zygomorphic corolla, all morphological traits associated with the species of the Alatae (Knapp et al. 2004). The decurrent leaf bases on the stem and the abrupt dilatation at the corolla throat, commonly found in the other species of the section (Goodspeed 1954), are lacking in the new species. The habit resembles that of *N. bonariensis* because of its rosulate leaves and scapose inflorescence. However, *N. gandarela* differs from *N. bonariensis* in its ombrophilous and extremely narrowly distributed habitat, its diurnal anthesis with red corollas with tube < 15 mm long, rather than being heliophilous, widespread, with nocturnal anthesis, and white corollas with a tube usually > 15 mm long.

Corolla color is variable in the Alatae, including white, yellow, pink, magenta, and purple-red. Such variation reflects an evolutionary history of radiation to distinct pollinator agents (Ippolito et al. 2004; Kaczorowski et al. 2005; Knapp 2010). Red flowers are usually associated with bird pollination (Faegri and van der Pijl 1979), and hummingbirds were already reported pollinating *N. forgetiana*, a pink to purple-red flowered species distributed in southern Brazil (Ippolito et al. 2004). Empirical data on the effective pollinator of *N. gandarela* are still necessary.

The Quadrilátero Ferrífero in Minas Gerais is located in the southern Espinhaço Range, a mountainous chain where the Cerrado (tropical Savannah) and Brazilian Atlantic Forest, two hotspots of biodiversity, connect (Mittermeier et al. 2004). The vegetation is a mosaic of topologies and vegetation, including grasslands, savannah, and forests (Spósito and Stehmann 2006), whose distributions are influenced by an altitudinal gradient, ranging from about 700 m to 2,080 m (Borges et al. 2011). In the region, besides the new species, a further two species of the Alatae are found, *N. bonariensis* and *N. langsdorffii*. The first usually inhabits the grasslands at higher altitudes known as Campos Rupestres (Silveira et al. 2016), while the second is generally associated with disturbed areas. Both grow mainly in open and sunny sites, as with most species of the Alatae, and the peculiar, shaded habitat of *N. gandarela* seems to be unique in the section. The microhabitat of the new species roughly resembles that of *Petunia exserta* Stehmann, an endemic species from Rio Grande do Sul, in

Southern Brazil, but where the geology and vegetational matrix are totally different (Lorenz-Lemke et al. 2006; Stehmann et al. 2009). In both cases, few or no species share the ground where these two species grow, meaning that they are fragile, with low, competitive capacity, but are presumably adapted to survive in these empty niches. Penumbral plant communities have also been described for the Campos Rupestres in Minas Gerais, where several species were reported as growing in small caves (Alves and Kolbek 1993), but until now, no Solanaceae species have been registered in these areas.

The new species deserves special conservation attention because it inhabits a small and rare habitat, if not unique, in the Quadrilátero Ferrífero, a region that is suffering intensive habitat loss (Salles et al. 2018) and has been the scenario of recurrent environmental disasters (Carmo et al. 2017; Rotta et al. 2019). Floristic inventories looking for new populations should be carried out in the surrounding forested areas, including the Serra do Gandarela National Park, following the drainage lines in the valleys. Also, we suggest the engagement of local people in trying to find other populations because of the impossibility of looking for similar areas since they are hidden by forest in satellite images. In the end, although we emphasize that the species should be primarily preserved in its natural habitat, ex-situ conservation measures might also be necessary (CNCFlora 2016) in order to prevent its extinction.

Acknowledgements

We thank the Coordination for the Improvement of Higher Education Personnel – Brazil (CAPES) –Financing Code 001 and FAPEMIG for the scholarship to MA and CNPQ (APQ 306086/2017–4) for financial support to JRS. Thanks also to the Center for Microscopy at the Universidade Federal de Minas Gerais (<http://www.microscopia.ufmg.br>) for providing the equipment and technical support for experiments involving electron microscopy. We thank the reviewers of the work for their valuable suggestions. And, especially, we thank Miss Gloria Regina Perpétuo for kindly allowing collections on her property.

References

- Alves R, Kolbek J (1993) Penumbral rock communities in campo rupestre sites in Brazil. *Journal of Vegetation Science* 4(3): 357–366. <https://doi.org/10.2307/3235594>
- Aoki S, Ito M (2000) Molecular phylogeny of *Nicotiana* (Solanaceae) based on the nucleotide sequence of the matK gene. *Plant Biology* 2(3): 316–324. <https://doi.org/10.1055/s-2000-3710>
- Borges RAX, Carneiro MAA, Viana PL (2011) Altitudinal distribution and species richness of herbaceous plants in campos rupestres of the Southern Espinhaço Range, Minas Gerais, Brazil. *Rodriguésia* 62(1): 139–152. <https://doi.org/10.1590/2175-7860201162110>
- Carmo FF, Kamino LHY, Tobias R Junior, Campos IC, Carmo FF, Silvino G, Castro KJSX, Mauro ML, Rodrigues NUA, Miranda MPS, Pinto CEF (2017) Fundão tailings dam

- failures: The environment tragedy of the largest technological disaster of Brazilian mining in global context. *Perspectives in Ecology and Conservation* 15(3): 145–151. <https://doi.org/10.1016/j.pecon.2017.06.002>
- Chase MW, Knapp S, Cox AV, Clarkson JJ, Butsko Y, Joseph J, Savolainen V, Parokkonny AS (2003) Molecular systematics, GISH and the origin of hybrid taxa in *Nicotiana* (Solanaceae). *Annals of Botany* 92(1): 107–127. <https://doi.org/10.1093/aob/mcg087>
- Clarkson JJ, Knapp S, Garcia VF, Olmstead RG, Leitch AR, Chase MW (2004) Phylogenetic relationships in *Nicotiana* (Solanaceae) inferred from multiple plastid DNA regions. *Molecular Phylogenetics and Evolution* 33(1): 75–90. <https://doi.org/10.1016/j.ympev.2004.05.002>
- Clarkson J, Dodsworth S, Chase MW (2017) Time-calibrated phylogenetic trees establish a lag phase between polyploidisation and diversification in *Nicotiana* (Solanaceae). *Plant Systematics and Evolution* 303(8): 1001–10212. <https://doi.org/10.1007/s00606-017-1416-9>
- CNCFlora (2016) Estratégia nacional para conservação *ex situ* de espécies ameaçadas da flora brasileira. CNCFlora - Centro Nacional de Conservação da Flora. Instituto de Pesquisas Jardim Botânico do Rio de Janeiro, Rio de Janeiro, 24 pp.
- Cocucci AA (2013) *Nicotiana*. In: Barboza GE (Coord.) Solanaceae. In: Zuloaga FO, Belgrano M, Anton AM (Eds) Flora Argentina: Flora vascular de la República Argentina. Dicotyledoneae. Instituto de Botánica Darwinion, Instituto Multidisciplinario de Biología Vegetal, San Isidro, 75–89.
- Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* 19(1): 11–15. https://webpages.uncc.edu/~jweller2/pages/BINF8350f2011/BINF8350_Readings/Doyle_plantDNAextractCTAB_1987.pdf
- Drummond GM, Martins CS, Machado ABM, Sebaio FA, Antonini Y [Eds] (2005) Biodiversidade em Minas Gerais: um atlas para sua conservação, 2nd edn. Fundação Biodiversitas, Belo Horizonte, 222 pp.
- Dupin J, Matzke NJ, Särkinen T, Knapp S, Olmstead RG, Bohs L, Smith SD (2017) Bayesian estimation of the global biogeographical history of the Solanaceae. *Journal of Biogeography* 44(4): 887–899. <https://doi.org/10.1111/jbi.12898>
- Faegri K, van der Pijl L (1979) The principles of pollination ecology. New York, Pergamon Press, 244 pp. <https://doi.org/10.1016/B978-0-08-023160-0.50020-7>
- Fregonezi JN, Turchetto C, Bonatto SL, Freitas LB (2013) Biogeographical history and diversification of *Petunia* and *Calibrachoa* (Solanaceae) in the Neotropical Pampas grassland. *Botanical Journal of the Linnean Society* 171(1): 140–153. <https://doi.org/10.1111/j.1095-8339.2012.01292.x>
- Goodspeed TH (1954) The genus *Nicotiana*. *Chronica Botanica*, Waltham, 536 pp.
- Iganci JR, Heiden G, Miotto STS, Pennington RT (2011) Campos de Cima da Serra: The Brazilian Subtropical Highland Grasslands show an unexpected level of plant endemism. *Botanical Journal of the Linnean Society* 167(4): 378–393. <https://doi.org/10.1111/j.1095-8339.2011.01182.x>
- Ippolito A, Fernandes GW, Holtsford TP (2004) Pollinator preferences for *Nicotiana alata*, *N. forgetiana*, and their F1 hybrids. *Evolution: International Journal of Organic Evolution* 58(12): 2634–2644. <https://doi.org/10.1111/j.0014-3820.2004.tb01617.x>

- IUCN (2019) Guidelines for using the IUCN red list categories and criteria, version 14. Prepared by the Standards and Petitions Subcommittee of the IUCN Species Survival Commission. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Jacobi CM, do Carmo FF, de Campos IC (2011) Soaring extinction threats to endemic plants in Brazilian metal-rich regions. *Ambio* 40(5): 540–543. <https://doi.org/10.1007/s13280-011-0151-7>
- Kaczorowski RL, Gardener MC, Holtsford TP (2005) Nectar traits in *Nicotiana* section *Alatae* (Solanaceae) in relation to floral traits, pollinators, and mating system. *American Journal of Botany* 92(8): 1270–1283. <https://doi.org/10.3732/ajb.92.8.1270>
- Knapp S (2010) On ‘various contrivances’: Pollination, phylogeny and flower form in the Solanaceae. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 365(1539): 449–460. <https://doi.org/10.1098/rstb.2009.0236>
- Knapp S (2020) Biodiversity (Solanaceae) of *Nicotiana*. *The Tobacco Plant Genome* 21: 21–41. https://doi.org/10.1007/978-3-030-29493-9_2
- Knapp S, Chase MW, Clarkson JJ (2004) Nomenclatural changes and a new sectional classification in *Nicotiana* (Solanaceae). *Taxon* 53(1): 73–82. <https://doi.org/10.2307/4135490>
- Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG (2007) Clustal W and Clustal X version 2.0. *Bioinformatics (Oxford, England)* 23(21): 2947–2948. <https://doi.org/10.1093/bioinformatics/btm404>
- Lorenz-Lemke AP, Mader G, Muschner VC, Stehmann JR, Bonnato SL, Salzano FM, Freitas LB (2006) Diversity and natural hybridization in a highly endemic species of *Petunia* (Solanaceae): A molecular and ecological analysis. *Molecular Ecology* 15(14): 4487–4497. <https://doi.org/10.1111/j.1365-294X.2006.03100.x>
- Maddison WP, Maddison DR (2019) Mesquite: a modular system for evolutionary analysis. Version 3.61. <http://www.mesquiteproject.org>
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: *Proceedings of the Gateway Computing Environments Workshop (GCE)*, 14 Nov. 2010, New Orleans, LA, 1–8. <https://doi.org/10.1109/GCE.2010.5676129>
- Mittermeier RA, Gil PR, Hoffmann M, Pilgrim J, Brooks T, Mittermeier CG, Lamoreux J, Fonseca GAB (2004) *Hotspots Revisited. Earth’s biologically richest and most endangered terrestrial ecoregions.* CEMEX, Mexico, 392 pp.
- Nylander JAA (2004) MrModeltest v2. Program distributed by the author. *Evolutionary Biology Centre, Uppsala University*, Vol. 5 No.18.
- Overbeck GE, Müller SC, Fidelis A, Pfadenhauer J, Pillar VD, Blanco CC, Boldrini II, Both R, Forneck ED (2007) Brazil’s neglected biome: The South Brazilian *Campos*. *Perspectives in Plant Ecology, Evolution and Systematics* 9(2): 101–116. <https://doi.org/10.1016/j.ppees.2007.07.005>
- Radford AE, Dickison WC, Massey JR, Bell CR (1976) *Vascular Plant Systematics* Harper and Row, New York.
- Robards AW (1978) An introduction to techniques for scanning electron microscopy of plant cells. In: Hall JL (Ed.) *Electron microscopy and cytochemistry of plant cells.* Elsevier, New York, 343–403.

- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MRBAYES 3.2: Efficient Bayesian phylogenetic inference and model selection across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Rotta LHS, Alcântara E, Park E, Negri RG, Lin YN, Bernardo N, Mendes TSG, Souza Filho CR (2019) The 2019 Brumadinho tailings dam collapse: Possible cause and impacts of the worst human and environmental disaster in Brazil. *International Journal of Applied Earth Observation and Geoinformation* 90: 102–119. <https://doi.org/10.1016/j.jag.2020.102119>
- Salles DM, do Carmo FF, Jacobi CM (2018) Habitat loss challenges the conservation of endemic plants in mining-targeted Brazilian mountains. *Environmental Conservation* 46(2): 140–146. <https://doi.org/10.1017/S0376892918000401>
- Silveira FAO, Negreiros D, Barbosa NPU, Buisson E, Carmo FF, Carstensen DW, Conceição AA, Cornelissen TG, Echternacht L, Fernandes GW, Garcia QS, Guerra TJ, Jacobi CM, Lemos-Filho JP, Le Stradic S, Morellato LPC, Neves FS, Oliveira RS, Schaefer CE, Viana PL, Lambers H (2016) Ecology and evolution of plant diversity in the endangered campo rupestre: A neglected conservation priority. *Plant and Soil* 403(1–2): 129–152. <https://doi.org/10.1007/s11104-015-2637-8>
- Sonter LJ, Barrett DJ, Soares-Filho BS, Moran CJ (2014) Global demand for steel drives extensive land-use change in Brazil's Iron Quadrangle. *Global Environmental Change* 26: 63–72. <https://doi.org/10.1016/j.gloenvcha.2014.03.014>
- Spósito TCS, Stehmann JR (2006) Heterogeneidade Florística e Estrutural de Remanescentes Florestais da Área de Proteção Ambiental ao Sul da Região Metropolitana de Belo Horizonte (APA Sul-RMBH), Minas Gerais, Brasil. *Acta Botanica Brasílica* 20(2): 347–362. <https://doi.org/10.1590/S0102-33062006000200010>
- Stamatakis A (2014) RAxML Version 8: A tool for Phylogenetic Analysis and Post-Analysis of Large Phylogenies. *Bioinformatics* 30(9): 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Stehmann JR, Semir J, Ippolito A (2002) *Nicotiana mutabilis* (Solanaceae), a new species from southern Brazil. *Kew Bulletin* 57(3): 639–646. <https://doi.org/10.2307/4110993>
- Stehmann JR, Freitas LB, Lorenz-Lemke AP, Semir J (2009) *Petunia*. In: Gerats T, Strommer J (Eds) *Petunia: Evolutionary, developmental and physiological genetics*. Springer Science Business Media, Nijmegen, 1–28. <https://doi.org/10.1093/aob/mcr051>
- Tamura K, Stecher G, Kumar S (2021) MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Molecular Biology and Evolution* 38(7): 3022–3027. <https://doi.org/10.1093/molbev/msab120>
- Teixeira MB, Coura-Neto AB, Pastore U, Rangel Filho ALR (1986) Levantamento de recursos naturais. Vegetação. In: Instituto Brasileiro de Geografia e Estatística (Ed.) Vegetação. Vol. 33. Instituto Brasileiro de Geografia e Estatística, Rio de Janeiro, 541–632. <https://biblioteca.ibge.gov.br/index.php/biblioteca-catalogo?view=detalhes&id=219048> [accessed 15.06.21]
- Teixeira MC, Quintana IV, Segatto ALA, Maestri R, Freitas LB, Augsten M, Stehmann JR, Turchetto C (2021) Changes in floral shape: Insights into the evolution of wild *Nicotiana*

(Solanaceae). Botanical Journal of the Linnean Society. <https://doi.org/10.1093/botlinnean/boab068>

Vignoli-Silva M, Mentz LA (2005) O gênero *Nicotiana* L. (Solanaceae) no Rio Grande do Sul, Brasil. Iheringia. Série Botânica 60(2): 151–173. <https://isb.emnuvens.com.br/iheringia/article/view/193>

Vignoli-Silva M, Stehmann JR (2020) *Nicotiana*. In: Flora do Brasil 2000. <http://servicos.jbrj.gov.br/flora/search/Nicotianahttp://floradobrasil.jbrj.gov.br/reflora/floradobrasil/FB14675> [accessed 14.09.21]

Supplementary material 1

Table S1

Authors: Mariana Augsten, Pablo Burkowski Meyer, Loreta B. Freitas, João A. N. Batista, João Renato Stehmann

Data type: GenBank accessions

Explanation note: Genbank accessions. ITS sequences from Chase et al. 2003, Plastidial sequences (*trnL-F*, *trnS-G* and *ndhF*) from Clarkson et al. 2004. Sequences from this work marked with an * (*Nicotiana gandarela* and *Nicotiana mutabilis*).

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/phytokeys.190.76111.suppl1>

Supplementary material 2

Figure S1

Authors: Mariana Augsten, Pablo Burkowski Meyer, Loreta B. Freitas, João A. N. Batista, João Renato Stehmann

Data type: Phylogenetic

Explanation note: Bayesian 50% majority-rule consensus tree of 56 species of *Nicotiana* L. of the nuclear (ITS) dataset. Numbers next to the nodes represent the posterior probabilities (PP).

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/phytokeys.190.76111.suppl2>

Supplementary material 3

Figure S2

Authors: Mariana Augsten, Pablo Burkowski Meyer, Loreta B. Freitas, João A. N. Batista, João Renato Stehmann

Data type: Phylogenetic

Explanation note: Bayesian 50% majority-rule consensus tree of 56 species of *Nicotiana* L. of the plastid (*trnL-F*, *trnS-G* and *ndhF*) datasets. Numbers next to the nodes represent the posterior probabilities (PP).

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/phytokeys.190.76111.suppl3>