

## Flora of Mongolia: annotated checklist of native vascular plants

Shukherdorj Baasanmunkh<sup>1</sup>, Magsar Urgamal<sup>2</sup>, Batlai Oyunsetseg<sup>3</sup>,  
Alexander P. Sukhorukov<sup>4,5</sup>, Zagarjav Tsegmed<sup>2</sup>, Dong Chan Son<sup>6</sup>,  
Andrey Erst<sup>5,7</sup>, Khurelpurev Oyundelger<sup>8,9</sup>, Alexey A. Kechaykin<sup>10</sup>,  
Joscelyn Norris<sup>11</sup>, Petr Kosachev<sup>10</sup>, Jin-Shuang Ma<sup>12</sup>,  
Kae Sun Chang<sup>13</sup>, Hyeok Jae Choi<sup>1</sup>

**1** Department of Biology and Chemistry, Changwon National University, Changwon 51140, Republic of Korea **2** Laboratory of Plant Systematics and Phylogeny, Botanic Garden and Research Institute, Mongolian Academy of Sciences, Ulaanbaatar 13330, Mongolia **3** Department of Biology, School of Arts and Science, National University of Mongolia, Ulaanbaatar 14201, Mongolia **4** Moscow State University, Leninskiye Gory, 1/12, Moscow, 119234, Russia **5** Tomsk State University, Lenina Pr., 36, Tomsk, 634050, Russia **6** Division of Forest Biodiversity, Korea National Arboretum, Pocheon 11186, Republic of Korea **7** Central Siberian Botanical Garden SB RAS, Zolotodolinskaya St. 101, Novosibirsk, 630090, Russia **8** Technical University Dresden, International Institute (IHI) Zittau, Chair of Biodiversity of Higher Plants, 02763, Zittau, Germany **9** Department of Botany, Senckenberg Museum of Natural History Görlitz, 02826, Görlitz, Germany **10** South Siberian Botanical Garden, Altai State University, Lenina 61, Barnaul, 656049, Russia **11** Rubenstein School of Environment and Natural Resources, University of Vermont, Vermont 05405-0088, USA **12** Institute of Botany, Beijing Botanical Garden, Beijing 100093, China **13** DMZ Forest and Biological Resources Conservation Division, Korea National Arboretum, Yanggu 24564, Republic of Korea

Corresponding author: Hyeok Jae Choi ([hjchoi1975@changwon.ac.kr](mailto:hjchoi1975@changwon.ac.kr))

---

Academic editor: Manuel Luján | Received 23 December 2021 | Accepted 24 February 2022 | Published 14 March 2022

**Citation:** Baasanmunkh S, Urgamal M, Oyunsetseg B, Sukhorukov AP, Tsegmed Z, Son DC, Erst A, Oyundelger K, Kechaykin AA, Norris J, Kosachev P, Ma J-S, Chang KS, Choi HJ (2022) Flora of Mongolia: annotated checklist of native vascular plants. *PhytoKeys* 192: 63–169. <https://doi.org/10.3897/phytokeys.192.79702>

---

### Abstract

In this study, we critically revised and updated the checklist of native vascular plants of Mongolia. The checklist comprises 3,041 native vascular plant taxa (2,835 species and 206 infraspecific species) from 653 genera and 111 families, including 7 lycophytes, 41 ferns, 21 gymnosperms, and 2,972 angiosperms. In the angiosperms, we identified the 14 families with the greatest species richness, ranging from 50 to 456 taxa. Species endemism is also noted here; 102 taxa are endemic to Mongolia, and 275 taxa are sub-endemic that co-occur in adjacent countries. Since 2014, a total of 14 taxa have been described new to

science based on morphological evidences. Moreover, five genera and 74 taxa were newly added to the flora of Mongolia. Based on our critical revisions, names of three families, 21 genera, and 230 species have been changed in comparison to the previous checklist, “Conspectus of the vascular plants of Mongolia” (2014).

### Keywords

Checklist, flora of Mongolia, native taxa, vascular plants

## Introduction

Mongolia is located in the mid-latitude (between 41°35'N–52°09'N and 87°44'E–119°56'E), between Russia and China, covering approximately 1.6 million km<sup>2</sup>, roughly equivalent to the size of western and central Europe. The flora of Mongolia is comprised of native species of different origins including boreal, steppe, desert, and mountainous elements of vegetation (Hilbig 1995; Gunin et al. 1999). The country is divided into sixteen phytogeographical regions which have various vegetation types (Grubov and Yunatov 1952), namely, alpine steppe, forest, meadow steppe, typical steppe, desert steppe, and desert (Gunin et al. 1999; Baasanmunkh et al. 2021a). Mongolia has a significant amount of temperate grasslands and semi-arid desert (Gunin et al. 1999; Wesche et al. 2016), which cover about 80% of the country's area (Pfeiffer et al. 2020). Overall, the species richness of vascular plants in Mongolia is not particularly high compared to other countries in Asia (Du et al. 2020; Li et al. 2020; Wang et al. 2020). However, Mongolia has the world's largest intact grassland with respect to its biodiversity (Batsaikhan et al. 2014; Hurka et al. 2019), which has great importance for the preservation of native vascular plants.

## A brief history of listing of the flora of Mongolia and recent taxonomic revisions

Historically, floristic studies have been very thoroughly conducted in this country, although recent updates are continuously being made. The first checklist of vascular plants included 1,897 species belonging to 555 genera and 97 families (Grubov 1955). Then, Grubov (1982) updated the checklist of vascular plants, which included 2,239 taxa from 599 genera and 103 families, with an identification key and information on their regional distribution and representative habitats. Later, Gubanov (1996) published a checklist with 2,823 higher plant species from 662 genera and 128 families, including notes on their regional distribution. More recently, Urgamal et al. (2014) updated the families of vascular flora according to APG III, with a total of 3,127 taxa that belong to 683 genera and 112 families. Since 2009, nine volumes with selected families have been published by Mongolian botanists, including Cyperaceae (Nyambayar 2009a), Apiaceae to Cornaceae (Urgamal 2009), Huperziaceae to Ephedraceae (Ulziikhutag et al. 2015), Asteraceae (Dariimaa 2014, 2021; Dariimaa and Saruul 2017), Ceratophyllaceae to Zygophyllaceae (Urgamal et al. 2020), Amaranthaceae s.l. (incl. Chenopodiaceae) (Tungalag 2020), Nymphaeaceae to Asphodelaceae (Urgamal et al. 2021).

Additionally, several families and genera have been revised in recent years. For example, a new checklist of the Brassicaceae family, the fifth-largest family in the country, was provided by German (2015). Taxonomic notes and checklists of *Aquilegia* L., *Stipa* L., and *Primula* L. were compiled by Erst et al. (2016), Zhao et al. (2019) and Baasanmunkh et al. (2020a), respectively. Recently, Troshkina (2021) revised and updated Geraniaceae in Mongolia. Baasanmunkh et al. (2021b) compiled a checklist of Orchidaceae, which included notes on their species richness and conservation status. The families Menyanthaceae and Nymphaeaceae were also revised by Baasanmunkh et al. (2022a). Additionally, some thorough regional floristic works were published: for the Khangai (Biazrov et al. 1989) and the Dzungarian Gobi regions (Baasanmunkh et al. 2021a). Moreover, an updated checklist of endemic plant species was recently provided by Baasanmunkh et al. (2021c), which comprises 102 taxa (95 species, 5 subspecies, and 2 varieties) from 43 genera and 19 families in the flora of Mongolia.

### New additions to the flora of Mongolia

Since Urgamal et al. (2014), 13 new species and one infraspecific taxon from Mongolia have been described as new to science (Nobis 2014; Erst et al. 2015, 2016; Kechaykin and Kutsev 2015; Yurtseva et al. 2016; Alexeeva 2018; Gundegmaa and Kechaykin 2018; Ovczinnikova 2019a, 2020; Pyak and Pyak 2019; Zhao et al. 2019; He et al. 2020; Pyak et al. 2020). Many new records of vascular plants have also been reported (Nobis et al. 2014, 2019a; Doronkin et al. 2015; Urgamal et al. 2016, 2019; Baasanmunkh et al. 2019a, b, c, 2020a, b, 2021b, d; Bazarragchaa et al. 2019; Erst et al. 2019; Ovczinnikova 2019b; Knyazev 2020; Shiga et al. 2020; Yano et al. 2021), including five genera new to the country, i.e. *Matthiola* W.T.Aiton, Brassicaceae (German 2015), *Onoclea* L., Onocleaceae (Doronkin et al. 2015), *Aldrovanda* L., Droseraceae (Shiga et al. 2020), *Hydrilla* L., Hydrocharitaceae (Shiga et al. 2020), and *Arctium* L., Asteraceae (Javzandolgor et al. 2021). Additionally, some genera previously listed by Urgamal et al. (2014) were omitted from Mongolian flora based on recent studies. In particular, the genus *Epipactis* Zinn. (Orchidaceae), for example, had two species that have been proven absent in the country due to the inaccurate location written on the herbarium specimens (Baasanmunkh et al. 2021b). On the other hand, some genera were not listed in Urgamal et al. (2014); for example, the genus *Phyllodoce* Salisb. (Ericaceae) was found in northern Mongolia by Oyunmaa and de Priest (2011). Furthermore, representatives of some genera, which are listed in the flora of Mongolia (Gubanov 1996; Urgamal et al. 2014), have been revised in recent studies (Podlech and Zarre 2013; Sukhorukov et al. 2013, 2019; Wang et al. 2014; Global Carex Group 2015; Duan et al. 2016; Drew et al. 2017; Kosachev 2017; Moore and Dillenberger 2017; Nosov et al. 2017; Pimenov 2017; Wiegleb et al. 2017; Boltenkov 2018; Gillespie et al. 2018; Madhani et al. 2018; Sinitsyna et al. 2018; Zhang et al. 2018; Barberá et al. 2019, 2020; Nobis et al. 2019b; Sramkó et al. 2019; Akan et al. 2020; Esput 2020; Friesen et al. 2020; Murakami et al. 2020; Nesom 2020; Ren et al. 2020; Zaika et al. 2020; Al-Shehbaz 2021; Al-Shehbaz et al. 2021; Liu et al. 2021).

## Necessity to update the list of flora of Mongolia


In 2016, the orders and families of flowering plants were updated by the APG IV (2016). Similarly, a new classification of ferns and lycophytes was provided for the first time (PPG I 2016). Given the recent updates to the international plant classification system, as well as a number of recent publications that identify new species and records and their distribution in Mongolia, there is a pressing need to revise and provide an updated list of the floristic diversity. Therefore, our study aims to present a thoroughly revised checklist of Mongolian native vascular flora that comprises the up-to-date names of all species, genera, and families, by conducting comparisons to the latest checklist, *Conspectus of Vascular Plants of Mongolia* by Urgamal et al. (2014) and earlier studies by Grubov (1982) and Gubanov (1996).

## Materials and methods

The systematic order and taxonomic circumscription of the families is based on the following classifications: Ferns and Fern Allies by PPG I (2016), Gymnosperms by Christenhusz et al. (2011), and Angiosperms by APG IV (2016). The names of accepted genera and species mostly follow Govaerts et al. (2021), which is currently maintained by Plants of the World Online (POWO 2021). Additionally, we reference recently published taxonomic revisions of certain families and genera. The authorship of species, genera and families is given after the International Plant Names Index (IPNI 2021); each species is provided with the author and respective publication as a reference. The name changes and most common synonyms, compared to the previous checklist, are also provided. Species endemism is given after Baasanmunkh et al. (2021c). Sub-endemic species are those that have also been found in at least one other country outside Mongolia, such as China, Kazakhstan, or Russia. For each species, we examined representative occurrence records based on the Global Biodiversity Information Facility (GBIF 2021, <https://www.gbif.org/>). We also compiled the phytogeographical regional distribution of all species, because species distribution is important information for species identification. The main herbaria for Mongolian flora (ALTB, LE, MW, MHA, NS, NSK, OSBU, TK, UBA, and UBU; acronyms follow Thiers 2021), Virtual Guide to the Flora of Mongolia (Rilke et al. 2012; Zemrich et al. 2013; <https://floragreif.uni-greifswald.de/floragreif/>), and all literature data for the species' regional distribution, have been checked and studied. The regional distribution of the taxa mostly follows Gubanov (1996), Urgamal et al. (2014), German (2015), and Baasanmunkh et al. (2021a). In addition to these sources, we used a revision of Baasanmunkh et al. (2022b), where numerous species were added or excluded from some phytogeographical regions. Lastly, the checklist comprises only native species, thus non-native taxa are not included and should be addressed in a future publication.

## Results

The current checklist comprises 3,042 native vascular plant taxa (including 2,891 species, 116 subspecies, 29 varieties, and 12 nothospecies), belonging to 653 genera and 111 families (Table 1, Fig. 1). The updated checklist is divided into four major taxonomic groups: lycophytes (2 families and 4 genera), ferns and fern allies (12 families and 17 genera), gymnosperms (3 families and 6 genera), and angiosperms (94 families and 626 genera) (see Table 1 for detailed numbers of taxa). Among these, angiosperms comprise 2,979 taxa, which constitute 97% of Mongolian flora (Table 1). We cross-checked the occurrence of each taxon using GBIF (2021), which includes occurrence data for 2,249 taxa (73% of Mongolian flora).



**Figure 1.** Species richness of genera ( $\geq 24$  taxa) and families ( $\geq 57$  taxa) of vascular flora of Mongolia. The names of only the most species-rich genera and families are shown.

**Table 1.** Number of native Mongolia vascular plant taxa in each taxonomic group.


Major taxonomic groups	Family	Genus	Taxon
Lycophytes	2	4	7
Ferns and fern allies	12	17	41
Gymnosperms	3	6	21
Angiosperms	94	627	2,972
<b>Total</b>	<b>111</b>	<b>653</b>	<b>3,041</b>

**Table 2.** The species richness of the total, endemic, and sub-endemic vascular plants of each phytogeographical region of Mongolia.

Region number	Name of the phytogeographical regions	Taxon	Endemic	Sub-endemic
1	Khuvsgul	1,054	7	63
2	Khentei	1,236	6	48
3	Khangai	1,514	27	87
4	Mongolian Dauria	1,198	6	54
5	Foothills of Great Khyangan	793	3	24
6	Khovd	1,011	10	68
7	Mongolian Altai	over 1400	47	114
8	Middle Khalkh	777	4	49
9	East Mongolia	952	3	50
10	Depression of Great Lakes	882	16	60
11	Valley of Lakes	466	5	39
12	East Gobi	462	2	57
13	Gobi Altai	865	16	76
14	Dzungarian Gobi	913	20	58
15	Transaltai Gobi	356	8	40
16	Alashan Gobi	262	2	43

There are 14 families with a high species richness ( $\geq 9$  genera and  $\geq 57$  taxa): Asteraceae (85 genera and 456 taxa), Fabaceae (24 and 328), Poaceae (58 and 229), Rosaceae (28 and 168), Ranunculaceae (20 and 156), Brassicaceae (51 and 138), Cyperaceae (10 and 130), Lamiaceae (22 and 103), Amaranthaceae (30 and 94), Caryophyllaceae (20 and 97), Boraginaceae (24 and 78), Apiaceae (36 and 66), Polygonaceae (11 and 63), and Orobanchaceae (9 and 57) (Fig. 1). The remaining 97 families comprise a smaller set of taxa. At the genus level, 14 genera represent a high species richness ( $\geq 24$  taxa): *Astragalus* L. (127 taxa), *Artemisia* L. (103), *Carex* L. (99), *Oxytropis* DC. (97), *Potentilla* L. (75), *Saussurea* DC. (55), *Taraxacum* F.H.Wigg. (53), *Allium* L. (50), *Salix* L. (42), *Ranunculus* L. (41), *Pedicularis* L. (36), *Poa* L. (28), *Viola* L. (27), and *Silene* L. (24) which is shown in Fig. 1.

In this study, a total of 275 sub-endemic taxa are provided which account for 9% of the total species of flora of the country. Among these, Fabaceae (74 taxa) show the highest number of sub-endemic taxa along with Asteraceae (60 taxa), Brassicaceae (23 taxa), Poaceae (18 taxa), and Amaranthaceae (9 taxa). The highest number of sub-endemic taxa were found in the Mongolian Altai (114 taxa) followed by Khangai (87 taxa), Gobi-Altai (76 taxa), Khovd (68 taxa), Khuvsgul (63 taxa), and the Depression of Great Lakes (60 taxa). The remaining ten regions have between 24 and 58 sub-endemic taxa (Table 2).



**Figure 2.** The species richness of native vascular plants of each phytogeographical region in Mongolia (1 Khuvgul 2 Khentei 3 Khangai 4 Mongolian Dauria 5 Foothills of Great Khyangan 6 Khovd 7 Mongolian Altai 8 Middle Khalkh 9 East Mongolia 10 Depression of Great Lakes 11 Valley of Lakes 12 East Gobi 13 Gobi Altai 14 Dzungarian Gobi 15 Transaltai Gobi 16 Alashan Gobi).

Based on the species distribution across the 16 phytogeographical regions, six regions with high species richness (over 1,000 taxa) were identified: the Mongolian Altai (more than 1,400 taxa), Khangai (1,514 taxa), Khentei (1,236 taxa), Mongolian Dauria (1,118 taxa), Khuvgul (1,054 taxa), and Khovd (1,011 taxa). The remaining ten regions have between 262 and 952 taxa (Table 2, Fig. 2).

In this study, we primarily excluded non-native species, including archeophytes that were listed in the previous checklist of Mongolia by Urgamal et al. (2014). As a result, 59 plant taxa found in Mongolia are considered non-native, including the families Cannabaceae (*Cannabis sativa* L.) and Portulacaceae (*Portulaca oleracea* L.). Several taxa were found to be archeophytes, which were introduced in “ancient” times and became naturalized as part of the native flora (a full list is given in Table 3).

## Discussion

We revised the checklist of Mongolian vascular plants provided by Urgamal et al. (2014), which included 3,127 taxa, belonging to 683 genera and 112 families. Because the list comprised both native and non-native taxa, we first sorted non-native taxa out to make our study comparable. There were 3,069 native taxa from 682 genera and 110 families listed by Urgamal et al. (2014) and the current checklist has been shortened and comprises 3,041 native vascular plant taxa from 653 genera and 111 families. Since

**Table 3.** List of non-native taxa that were included in the previous checklist of Mongolian flora but are excluded from this checklist.

No	Taxon name	No	Taxon name
	Amaranthaceae		Fabaceae
1	<i>Amaranthus albus</i> L.	30	<i>Lathyrus oleraceus</i> Lam.
2	<i>Amaranthus blitoides</i> S.Watson	31	<i>Lathyrus sativus</i> L.
3	<i>Amaranthus cruentus</i> L.	32	<i>Lotus corniculatus</i> L.
4	<i>Amaranthus retroflexus</i> L.	33	<i>Medicago sativa</i> L.
	Apiaceae	34	<i>Melilotus albus</i> Medik.
5	<i>Anethum graveolens</i> L.	35	<i>Ornithopus perpusillus</i> L.
6	<i>Eryngium planum</i> L.	36	<i>Trigonella caerulea</i> (L.) Ser.
7	<i>Pastinaca sativa</i> L.	37	<i>Vicia angustifolia</i> L.
	Asteraceae	38	<i>Vicia faba</i> L.
8	<i>Cichorium intybus</i> L.	39	<i>Vicia sativa</i> L.
9	<i>Sonchus oleraceus</i> L..	40	<i>Vicia sepium</i> L.
10	<i>Xanthium orientale</i> L.		Malvaceae
11	<i>Xanthium strumarium</i> L.	41	<i>Hibiscus trionum</i> L.
	Brassicaceae		Plantaginaceae
12	<i>Berteroa incana</i> (L.) DC.	42	<i>Veronica peregrina</i> L.
13	<i>Brassica campestris</i> L.		Poaceae
14	<i>Brassica juncea</i> (L.) Czern.	43	<i>Avena fatua</i> L.
15	<i>Bunias orientalis</i> L.	44	<i>Avena sativa</i> L.
16	<i>Camelina caucasica</i> (Sinskaya) Vassilcz.	45	<i>Cenchrus americanus</i> (L.) Morrone
17	<i>Camelina sativa</i> (L.) Crantz	46	<i>Chloris virgata</i> Sw.
18	<i>Eruca sativa</i> Mill.	47	<i>Dactylis glomerata</i> L.
19	<i>Guenthera persica</i> (Boiss. & Hohen.) D.A.German	48	<i>Hordeum aegiceras</i> Royle ex Walp.
20	<i>Neslia paniculata</i> (L.) Desv.	49	<i>Hordeum distichon</i> L.
21	<i>Raphanus raphanistrum</i> L.	50	<i>Hordeum vulgare</i> L.
22	<i>Sinapis arvensis</i> L.	51	<i>Panicum miliaceum</i> L.
23	<i>Sisymbrium altissimum</i> L.	52	<i>Panicum miliaceum</i> subsp. <i>ruderale</i> (Kitag.) Tzvelev
24	<i>Sisymbrium volgense</i> M.Bieb.	53	<i>Secale cereale</i> L.
	Cannabaceae	54	<i>Setaria pumila</i> Roem. & Schult.
25	<i>Cannabis sativa</i> L.	55	<i>Setaria viridis</i> (L.) P.Beauv.
	Caryophyllaceae	56	<i>Triticum aestivum</i> L.
26	<i>Agrostemma githago</i> L.		Polygonaceae
27	<i>Gypsophila vaccaria</i> (L.) Sm.	57	<i>Fagopyrum esculentum</i> Moench
28	<i>Silene banksia</i> (Meerb.) Mabb.	58	<i>Fagopyrum tataricum</i> (L.) Gaertn.
	Convolvulaceae		Portulacaceae
29	<i>Calystegia silvatica</i> (Kit.) Griseb.	59	<i>Portulaca oleracea</i> L.

Grubov (1955) provided the first checklist, more than 1,150 species have been added to the flora of Mongolia. In this study, the family Brassicaceae is based upon the work of German (2015), who recognized 141 species from 59 genera which significantly differs from the 160 species and 61 genera provided by Urgamal et al. (2014); this is because of misidentification of species and records from outside Mongolia being included in the latter publication. Since Urgamal et al. (2014), 14 taxa have been described as new to science based on morphological evidence, the majority of which are from Boraginaceae (Suppl. material 1: Appendix 1). Furthermore, 79 taxa have recently been added to the flora of Mongolia (Suppl. material 1: Appendix 1). On the other hand, many taxa are

synonymized and/or the names and status of numerous taxa were changed based on our critical revisions (list of the synonymized taxa is provided Suppl. material 2: Appendix 2). In particular, accepted names of three families, 21 genera, and 232 taxa have been changed/synonymized. Moreover, 21 taxa listed in the Urgamal et al (2014) were absent from Mongolia based on our extensive research which is given in Suppl. material 2: Appendix 2 (Pimenov 2017; Ovczinnikova 2019a; Nesom 2020; Baasanmunkh et al. 2021b, 2022a). Due to this high number of synonymization, even after dozens of new species have been discovered in Mongolian flora, change in absolute species numbers appeared to be minor.

Previous checklists of vascular plants in Mongolia (Gubanov 1996; Urgamal et al. 2014), listed both native and non-native taxa; however, there were no specific remarks for non-native taxa. Recently, many researchers have published checklists of non-native plants, mainly in Europe and East Asia, for example, concerning China (Xu et al. 2012), Italy (Galasso et al. 2018), South Korea (Korea National Arboretum 2021) as well as the Russian Far East and Siberia (Ebel et al. 2014; Vinogradova et al. 2020). These works aim to increase awareness about invasive species and strengthen their biosecurity regulations. This level of detail is important because invasions have complex and often immense long-term direct and indirect effects on native natural communities (Pagad et al. 2018; van Kleunen et al. 2019; Pyšek et al. 2020). In many Central Asian countries, there is no separate checklist for non-native plants species (Sennikov and Lazkov 2021). The first checklist of alien species was recently published for Kyrgyzstan by Sennikov et al. (2021). In Mongolia, approximately 35 non-native plants taxa are recognized in GBIF (Munkhnast et al. 2020), but this list has not critically revised all non-native plants species of the country. In this study, we primarily excluded non-native species that were listed in the previous checklist of Mongolia, including two families Cannabaceae (*Cannabis sativa* L.) and Portulacaceae (*Portulaca oleracea* L.) and 57 other taxa (Table 3).

Mongolia has a relatively small number of endemic vascular plants, with 102 taxa belonging to 43 genera and 19 families, accounting for only about 3% of the country's total flora (Baasanmunkh et al. 2021c). Notably, the Mongolian Altai and Khangai regions harbor over 70% of the total endemics and sub-endemics of Mongolia, which reflects their high species richness (Fig. 2; Table 2). This could be due to the diverse habitats along the Altai mountains and the large expanse of forest and mountain-steppe in both regions. Regional distribution of new species, new records, and recently revised genera and families, were provided based on literature. It is important to highlight that the study does not entirely revise regional distribution of each taxon. Nonetheless, we have been working on the grid distribution map of vascular plants (see Baasanmunkh et al. 2021b, 2022a) since 2020 based on critical revision of herbarium specimens, literature, and our own field observation data.

Both of the online databases are allowing researchers to collaborate and revise Mongolian taxa more readily and will continue to improve the documentation of Mongolia's flora. To date, a total of 2,249 taxa (ca. 73% of the flora) have been deposited in the database of GBIF (2021). Furthermore, the data for 1,249 species (including herbarium specimens and/or images of living plants) are at least partially available in the database of the Virtual Guide to the Flora of Mongolia (Rilke et al.

2012; Zemmrich et al. 2013; <https://floragreif.uni-greifswald.de/floragreif/>). Moreover, approximately 19,300 images of 1,780 taxa have been observed as part of citizen science contributions to the “Flora of Mongolia” project on the iNaturalist platform (<https://www.inaturalist.org/projects/flora-of-mongolia?tab=observations>), which was established on January 2019.

In this study, we checked more than 70 works published since 2013 that have revised the flora of Mongolia, and provided respective references for each species in our checklist. We reviewed the species status of all vascular flora of Mongolia and made critical changes by adding, synonymizing, and excluding taxa; this work resulted in 265 fewer taxa compared to Urgamal et al. (2014). We believe that our revised checklist serves as an essential background and reference not only for scientists and students, but also for local government administrations and protected areas, for the conservation of Mongolian flora. Having an updated checklist allows researchers and communities to monitor plants as climate and land use changes, and population size and herding pressures increase. We recommend more research be conducted on regional flora, as well as comprehensive revisions of species distribution based on herbarium collections for phytogeographical regions and taxonomic revision of doubtful taxa.

## Annotated checklist of native vascular flora in Mongolia.

The families in the checklist are alphabetically ordered and, within them, the genera, species, and subspecies are alphabetically listed. The currently accepted names are highlighted in bold italics. The most common synonyms (previously used in Urgamal et al. (2014)) and the species’ distribution in phytogeographical regions are provided here. Symbols used in the checklist include endemic [E] and sub-endemic [SE].

### I Lycophtyes

#### 1. Lycopodiaceae P.Beauv. (3 genera and 5 species)

<i>Diphasiastrum alpinum</i> (L.) Holub	[1, 2]
<i>Diphasiastrum complanatum</i> (L.) Holub	[1]
<i>Huperzia selago</i> (L.) Bernh.	[1, 2]
<i>Lycopodium annotinum</i> L.	[1, 2]
<i>Lycopodium clavatum</i> L.	[2]

#### 2. Selaginellaceae Willk. (1 genus and 2 species)

<i>Selaginella borealis</i> (Kaulf.) Spring	[1, 3, 4, 8]
<i>Selaginella sanguinolenta</i> (L.) Spring	[1, 2, 3, 4, 8]

### II Ferns and fern allies

#### 3. Aspleniaceae Newman (1 genus and 5 species)

<i>Asplenium altajense</i> (Kom.) Grubov	[1, 3, 4, 7, 10, 13]
<i>Asplenium ruprechtii</i> Sa.Kurata	[2, 3, 4]

<i>Asplenium ruta-muraria</i> L.	[7, 14]
<i>Asplenium septentrionale</i> (L.) Hoffm.	[7, 8, 10, 14]
<i>Asplenium yunnanense</i> Franch.	[7, 13]
<b>4. Athyriaceae</b> Alston (2 genera and 4 species)	
<i>Athyrium filix-femina</i> (L.) Roth	[1–5]
<i>Athyrium monomachi</i> Kom.	[2, 3, 4, 5]
<i>Athyrium sinense</i> Rupr.	[3, 5, 9]
<i>Diplazium sibiricum</i> (Turcz.) Sa.Kurata	[1–5]
<b>5. Cystopteridaceae</b> Shmakov (2 genera and 4 species)	
<i>Cystopteris fragilis</i> (L.) Bernh.	[1–10, 13, 14, 15]
<i>Cystopteris sudetica</i> A.Braun & Milde	[2]
<i>Gymnocarpium dryopteris</i> Newman	[2, 4]
<i>Gymnocarpium jessoense</i> (Koidz.) Koidz.	[1, 2, 3, 4, 5, 8]
<b>6. Dennstaedtiaceae</b> Losty (1 genus and 1 species)	
<i>Pteridium aquilinum</i> (L.) Kuhn	[2, 3, 4, 5]
<b>7. Dryopteridaceae</b> Herter (1 genus and 3 species)	
<i>Dryopteris dilatata</i> (Hoffm.) A.Gray	[2, 5]
<i>Dryopteris expansa</i> (C.Presl) Fraser-Jenk. & Jermy	[2, 5]
<i>Dryopteris fragrans</i> (L.) Schott	[2, 3, 4, 5, 7, 10]
<b>8. Equisetaceae</b> Michx. (1 genus and 9 species)	
<i>Equisetum arvense</i> L.	[1–10, 14]
<i>Equisetum fluviatile</i> L.	[1–10, 14]
<i>Equisetum hyemale</i> L.	[3, 4, 5]
<i>Equisetum palustre</i> L.	[1–10, 14]
<i>Equisetum pratense</i> Ehrh.	[1–7, 9, 10]
<i>Equisetum ramosissimum</i> Desf.	[14]
<i>Equisetum scirpoideum</i> Michx.	[1, 2, 3, 4, 6]
<i>Equisetum sylvaticum</i> L.	[1–5, 8, 9]
<i>Equisetum variegatum</i> Schleich.	[1, 4]
<b>9. Onocleaceae</b> Pic.Serm. (2 genera and 2 species)	
<i>Matteuccia struthiopteris</i> (L.) Tod.	[2, 4]
<i>Onoclea sensibilis</i> L.	[2]
<b>10. Ophioglossaceae</b> Martinov (1 genus and 2 species)	
<i>Botrychium lanceolatum</i> (Gmel.) Ångstr.	[3]
<i>Botrychium lunaria</i> (L.) Sw.	[1–7, 9]

**11. Polypodiaceae** J.Presl. & C.Presl (2 genera and 2 species)

- Lepisorus clathratus* Ching [13]  
*Polypodium virginianum* L. [1–5, 8]

**12. Pteridaceae** E.D.M.Kirchn. (2 genera and 2 species)

- Cheilanthes argentea* (S.G.Gmel.) Kunze [1–9, 12, 13]  
*Cryptogramma stelleri* (S.G.Gmel.) Prantl [1]

**13. Thelypteridaceae** Ching (1 genus and 1 species)

- Phegopteris connectilis* (Michx.) Watt [2, 4]

**14. Woodsiaceae** Herter (1 genus and 6 species)

- Woodsia calcarea* (Fomin) Shmakov [1, 3, 4]  
*Woodsia glabella* R.Br. [1, 7, 10]  
*Woodsia heterophylla* (Turcz.) Shmakov [1]  
*Woodsia ilvensis* (L.) R.Br. [= *Woodsia acuminata* (Fomin) Sipliv.] [1–9, 13]  
*Woodsia pseudopolystichoides* (Fomin) Kiselev & Shmakov [5]  
*Woodsia subcordata* Turcz. [4, 5, 9]

**III Gymnosperms****15. Cupressaceae** Gray (1 genus and 4 taxa)

- Juniperus communis* L. [1, 2, 3, 4, 6, 7]  
*Juniperus pseudosabina* Fisch. & C.A.Mey. [1–4, 7, 8, 13]  
*Juniperus sabina* var. *davurica* (Pall.) Farjon [= *Juniperus davurica* Pall.] [2]  
*Juniperus sabina* L. var. *sabina* [2–4, 6–8, 10, 11, 13, 14]

**16. Ephedraceae** Dumort. (1 genus and 9 species)

- Ephedra dahurica* Turcz. [= *Ephedra sinica* subsp. *dahurica* (Turcz.) Galanin] [2, 3, 8, 9, 12]  
*Ephedra equisetina* Bunge [3, 6–9, 12–16]  
*Ephedra fedtschenkoi* Paulsen [2, 3, 4, 7]  
*Ephedra glauca* Regel [12, 14, 15]  
*Ephedra intermedia* Schrenk & C.A.Mey. [7, 14, 15]  
*Ephedra lomatolepis* Schrenk [12, 14, 15]  
*Ephedra monosperma* J.G.Gmel. [1–8, 10, 12–14]  
*Ephedra przewalskii* Stapf [6, 7, 10–16]  
*Ephedra sinica* Stapf [2–5, 7–15]

**17. Pinaceae** Spreng. (4 genera and 8 species)

- Abies sibirica* Ledeb. [1, 2]  
*Larix czekanowskii* Szafer [4]  
*Larix gmelinii* (Rupr.) Kuzen. [= *Abies gmelinii* Rupr.] [2, 4]  
*Larix sibirica* Ledeb. [1–4, 6–8, 10, 14]  
*Picea obovata* Ledeb. [1, 2, 3, 6, 7]

<i>Pinus pumila</i> (Pall.) Regel	[2]
<i>Pinus sibirica</i> Du Tour	[1, 2, 3, 6, 7]
<i>Pinus sylvestris</i> L.	[1–5, 8, 9]

**IV Angiosperms****18. Acoraceae** Martinov (1 genus and 1 species)

<i>Acorus calamus</i> L.	[1, 3, 4, 5, 8, 9]
--------------------------	--------------------

**19. Adoxaceae** E.Mey. [including Viburnaceae Raf.] (3 genera and 6 species)

<i>Adoxa moschatellina</i> L.	[1–3, 5–7, 13]
<i>Sambucus sibirica</i> Nakai	[3, 6]
<i>Sambucus williamsii</i> Hance [= <i>Sambucus manshurica</i> Kitag.]	[1, 2, 3, 4, 5, 9]
<i>Viburnum burejaeticum</i> Regel & Herder	[5]
<i>Viburnum mongolicum</i> Rehder [= <i>Lonicera mongolica</i> Pall.]	[4, 5, 8, 9]
<i>Viburnum sargentii</i> Koehne	[5]

**20. Alismataceae** Vent. (2 genera and 4 species)

<i>Alisma gramineum</i> Lej.	[5, 7, 8, 10, 11, 14]
<i>Alisma plantago-aquatica</i> L.	[1–5, 8–10, 14]
<i>Sagittaria natans</i> Pall.	[4, 9, 10, 14]
<i>Sagittaria trifolia</i> L.	[1, 4, 5, 9, 14]

**21. Amaranthaceae** Juss. [including Chenopodiaceae Vent.] (34 genera and 94 taxa)

<i>Agriophyllum pungens</i> (Vahl) Link [= <i>Agriophyllum squarrosum</i> Moq.]	[6–16]
<i>Anabasis aphylla</i> L.	[7, 14]
<i>Anabasis brevifolia</i> C.A.Mey.	[3, 6–8, 10–16]
<i>Anabasis elatior</i> (C.A.Mey.) Schischk.	[14]
<i>Anabasis eriopoda</i> Paulsen	[14]
<i>Anabasis pelliottii</i> Danguy	[14]
<i>Anabasis salsa</i> Paulsen	[14]
<i>Anabasis truncata</i> Bunge	[7, 14]
<i>Atriplex altaica</i> Sukhor.	[7]
<i>Atriplex cana</i> C.A.Mey.	[14]
<i>Atriplex fera</i> (L.) Bunge	[1, 3, 4, 8–10, 12, 13]
<i>Atriplex laevis</i> C.A.Mey.	[3, 4, 8–11, 13–15]
<i>Atriplex sibirica</i> L.	[2, 3, 4, 6–16]
<i>Atriplex tatarica</i> L.	[7, 10, 14]
<i>Axyris amaranthoides</i> L.	[2–5, 8, 9, 13]
<i>Axyris hybrida</i> L.	[2–5, 7–10, 12–14]
<i>Axyris prostrata</i> L.	[1–4, 6–10, 13, 14]
<i>Bassia hyssopifolia</i> (Pall.) Kuntze	[6, 7, 10–14]
<i>Bassia prostrata</i> (L.) Beck [= <i>Kochia prostrata</i> (L.) Schrad.]	[1–15]
<i>Bassia scoparia</i> (L.) A.J.Scott [= <i>Kochia scoparia</i> (L.) Schrad.]	[2, 3, 4, 6–14]
<i>Blitum virgatum</i> L. [= <i>Chenopodium foliosum</i> Asch.]	[3, 4, 6, 7, 12–15]

	<i>Camphorosma monspeliaca</i> subsp. <i>lessingii</i> (Litv.) Aellen	[6, 10, 14]
	<i>Caroxylon gemmascens</i> (Pall.) Tzvelev [= <i>Salsola gemmascens</i> Pall.]	[10]
	<i>Caroxylon passerinum</i> (Bunge) Akhani & Roalson [≡ <i>Salsola passerina</i> Bunge]	[8, 10–16]
	<i>Ceratocarpus arenarius</i> L.	[6, 7, 10, 14]
	<i>Chenopodiastrum hybridum</i> (L.) S.Fuentes, Uotila & Borsch [≡ <i>Chenopodium hybridum</i> L.]	[2, 3, 4, 5, 7–16]
	<i>Chenopodium acuminatum</i> Willd.	[3–14, 16]
	<i>Chenopodium album</i> L.	[1–16]
	<i>Chenopodium ficifolium</i> Sm.	[1, 9, 10, 11, 14]
	<i>Chenopodium frutescens</i> C.A.Mey.	[6, 7, 10]
	<i>Chenopodium iljinii</i> Golosk.	[7, 10]
	<i>Chenopodium karoi</i> Aellen	[1–15]
	<i>Chenopodium novopokrovskyanum</i> (Aellen) Uotila [≡ <i>Chenopodium album</i> subsp. <i>novopokrovskyanum</i> (Aellen) Uotila]	[7]
	<i>Chenopodium strictum</i> Roth	[2, 4, 9, 14]
	<i>Chenopodium vulvaria</i> L.	[3, 6–10, 13, 14]
	<i>Climacoptera affinis</i> (C.A.Mey.) Botsch. [≡ <i>Pyankovia affinis</i> (C.A.Mey.) Mosyakin & Roalson]	[14]
	<i>Climacoptera subcrassa</i> (Popov) Botsch.	[14]
	<i>Corispermum chinganicum</i> Iljin	[1–12]
	<i>Corispermum declinatum</i> Steph. ex Iljin	[3, 4, 8]
	<i>Corispermum elongatum</i> Bunge [≡ <i>Corispermum stauntonii</i> subsp. <i>elongatum</i> (Bunge) Vorosch.]	[10, 13]
SE	<i>Corispermum mongolicum</i> Iljin	[3, 4, 7–16]
SE	<i>Corispermum patelliforme</i> Iljin	[10, 13, 16]
SE	<i>Corispermum tylocardrum</i> Hance [= <i>Corispermum gmelinii</i> Bunge]	[12]
	<i>Dysphania botrys</i> (L.) Mosyakin & Clemans	[7, 14, 15]
	<i>Grubovia dasypylla</i> (Fisch. & C.A.Mey.) Freitag & G.Kadereit [≡ <i>Bassia dasypylla</i> (Fisch. & C.A.Mey.) Kuntze, <i>Kochia dasypylla</i> Fisch. & C.A.Mey.]	[3–16]
	<i>Grubovia krylovii</i> (Litv.) Freitag & G.Kadereit [≡ <i>Kochia krylovii</i> Litv.]	[6, 7, 10–16]
	<i>Grubovia melanoptera</i> (Bunge) Freitag & G.Kadereit [≡ <i>Kochia melanoptera</i> Bunge]	[3, 6, 7, 10–16]
	<i>Halocnemum strobilaceum</i> (Pall.) M.Bieb.	[14]
	<i>Halogeton glomeratus</i> (M.Bieb.) C.A.Mey.	[7, 10, 11, 14, 15]
	<i>Halostachys caspica</i> C.A.Mey.	[14, 15]
	<i>Haloxylon ammodendron</i> (C.A.Mey.) Bunge	[7, 10–16]
	<i>Iljinia regelii</i> (Bunge) Korovin	[14, 15, 16]
	<i>Kalidium caspicum</i> (L.) Ung.-Sternb.	[10, 14]
	<i>Kalidium cuspidatum</i> (Ung.-Sternb.) Grubov	[3, 8, 9, 12–16]
	<i>Kalidium foliatum</i> Moq.	[3, 6–16]

SE	<i>Kalidium gracile</i> Fenzl	[3, 4, 8–16]
	<i>Krascheninnikovia ceratoides</i> (L.) Gueldenst. [= <i>Krascheninnikovia ewersmanniana</i> (Stschegl.) Grubov]	[1, 3, 4, 6–16]
	<i>Micropeplis arachnoidea</i> (Moq.) Bunge [≡ <i>Halogeton arachnoides</i> Moq.]	[4, 6–16]
SE	<i>Nanophyton grubovii</i> U.P.Pratov	[10]
SE	<i>Nanophyton mongolicum</i> U.P.Pratov	[7, 14]
	<i>Oreosalsola abrotanoides</i> (Bunge) Akhani [≡ <i>Salsola abrotanoides</i> Bunge]	[6–13]
	<i>Oxybasis chenopodioides</i> (L.) S.Fuentes, Uotila & Borsch [≡ <i>Chenopodium chenopodioides</i> (L.) Aellen]	[7, 10, 14]
	<i>Oxybasis glauca</i> (L.) S.Fuentes, Uotila & Borsch [= <i>Chenopodium glaucum</i> L.]	[2–16]
	<i>Oxybasis gubanovii</i> (Sukhor.) Sukhor. & Uotila [≡ <i>Chenopodium gubanovii</i> Sukhor.]	[10, 14]
	<i>Oxybasis rubra</i> (L.) S.Fuentes, Uotila & Borsch [≡ <i>Chenopodium rubrum</i> L.]	[3, 7, 10, 14, 15]
	<i>Oxybasis urbica</i> (L.) S.Fuentes, Uotila & Borsch [≡ <i>Chenopodium urbicum</i> L.]	[4, 10]
	<i>Petrosimonia litvinowii</i> Korsh.	[10]
	<i>Petrosimonia sibirica</i> Bunge	[14]
SE	<i>Salicornia altaica</i> Lomon. [= <i>Salicornia perennans</i> subsp. <i>altaica</i> (Lomon.) G.Kadereit & Piirainen]	[7]
	<i>Salsola collina</i> Pall. [= <i>Kali collinum</i> (Pall.) Akhani & Roalson]	[2–15]
SE	<i>Salsola ikonnikovii</i> Iljin [= <i>Kali ikonnikovii</i> (Iljin) Akhani & Roalson]	[7, 11, 12, 13]
	<i>Salsola jacquemontii</i> Moq [= <i>Kali jacquemontii</i> (Moq.) Akhani & Roalson]	[8, 13]
	<i>Salsola laricifolia</i> Litv.	[12, 13, 16]
	<i>Salsola monoptera</i> Bunge [= <i>Kali monopterum</i> (Bunge) Lomon.]	[3, 6–13]
	<i>Salsola paulsenii</i> Litv. [= <i>Kali paulsenii</i> (Litv.) Akhani & Roalson]	[3, 7, 9–11, 14]
	<i>Salsola rosacea</i> L.	[7, 14]
	<i>Salsola tragus</i> L.	[2, 3, 4, 6–16]
	<i>Soda foliosa</i> (L.) Akhani [= <i>Salsola foliosa</i> L. ≡ <i>Neocaspia foliosa</i> (L.) Tzvelev]	[14]
	<i>Suaeda acuminata</i> (C.A.Mey.). Moq.	[6, 10, 14]
	<i>Suaeda corniculata</i> (C.A.Mey.) Bunge subsp. <i>corniculata</i>	[1, 3–16]
	<i>Suaeda corniculata</i> subsp. <i>mongolica</i> Lomon. & Freitag	[3, 4, 7–11]
	<i>Suaeda glauca</i> (C.A.Mey.) Bunge	[4, 9, 16]
	<i>Suaeda heterophylla</i> (Kar. & Kir.) Bunge	[10–15]
	<i>Suaeda kossinskyi</i> Iljin [= <i>Bienertia kossinskyi</i> (Iljin) Tzvelev]	[6, 7, 10, 13, 16]
	<i>Suaeda linifolia</i> Pall.	[10, 14]
	<i>Suaeda prostrata</i> Pall. [= <i>Suaeda maritima</i> auct. non L.]	[6, 9–13, 16]

	<i>Suaeda przewalskii</i> Bunge [ $\equiv$ <i>Bienertia przewalskii</i> (Bunge) G.L.Chu]	[10–13]
	<i>Suaeda salsa</i> (L.) Pall.	[8–13, 15, 16]
	<i>Suaeda sibirica</i> Lomon. & Freitag	[3, 4, 8, 9, 10]
SE	<i>Suaeda tschujensis</i> Lomon. & Freitag	[6, 7]
SE	<i>Suaeda tuvinica</i> Lomon. & Freitag	[3, 6, 10]
	<i>Sympogma regelii</i> Bunge	[7, 10–16]
	<i>Teloxys aristata</i> (L.) Moq. [ $\equiv$ <i>Chenopodium aristatum</i> L.]	[3–13, 15, 16]
	$\equiv$ <i>Dysphania aristata</i> (L.) Mosyakin & Clements	
	<i>Xylosalsola arbuscula</i> (Pall.) Tzvelev [ $\equiv$ <i>Salsola arbuscula</i> Pall.]	[7, 13–16]

## 22. Amaryllidaceae J.St.-Hil. (1 genus and 50 taxa)

Note: According to Sinitsyna et al. (2018), *Allium spirale* is absent in Mongolia and *A. subangulatum* was found in southern Gobi by Friesen et al. (2020).

	<i>Allium altaicum</i> Pall.	[1–3, 6–8, 10, 13, 14]
	<i>Allium amphibolum</i> Ledeb.	[1–4, 6, 7, 10, 13, 14]
	<i>Allium anisopodium</i> Ledeb.	[2–13]
SE	<i>Allium austrosibiricum</i> N.Friesen	[3, 6, 7, 10, 14]
	<i>Allium baicalense</i> Willd. [= <i>Allium senescens</i> subsp. <i>glaucum</i> (Schrader) Dostál]	[1, 3–5, 9, 10]
	<i>Allium bidentatum</i> Fisch.	[1–6, 8–12, 14]
	<i>Allium burjaticum</i> N.Friesen	[3, 4, 8]
	<i>Allium carolinianum</i> Redouté	[14]
	<i>Allium chamarens</i> se M.M.Ivanova	[1, 2, 3]
	<i>Allium clathratum</i> Ledeb.	[3, 6, 7, 10, 11]
	<i>Allium condensatum</i> Turcz.	[5, 9]
	<i>Allium eduardi</i> Stearn	[2, 3, 4, 6–16]
	<i>Allium flavidum</i> Ledeb.	[1–4, 6, 7, 13, 14]
	<i>Allium galanthum</i> Kar. & Kir.	[7, 14]
	<i>Allium hymenorrhizum</i> Ledeb.	[7, 14]
	<i>Allium karelinii</i> Poljakov	[7, 14]
	<i>Allium ledebourianum</i> Schult. & Schult.f.	[7]
	<i>Allium leucocephalum</i> Turcz.	[1–4, 7–14, 16]
	<i>Allium macrostemon</i> Bunge	[8, 9]
	<i>Allium malyschevii</i> N.Friesen	[1, 2, 3]
	<i>Allium maximowiczii</i> Regel	[2, 4, 5, 9]
	<i>Allium microdictyon</i> Prokh.	[1, 2, 3, 4]
	<i>Allium monadelphum</i> Turcz.	[1, 2, 3, 6, 7]
	<i>Allium mongolicum</i> Regel	[3, 4, 6–16]
	<i>Allium neriniflorum</i> G.Don	[4, 5, 9]
	<i>Allium obliquum</i> L.	[7]
	<i>Allium oliganthum</i> Kar. & Kir.	[6, 7, 10, 14]
	<i>Allium pallasii</i> Murray	[14]

	<i>Allium platyspathum</i> Schrenk subsp. <i>platyspathum</i>	[3, 6, 7, 13, 14]
	<i>Allium platyspathum</i> subsp. <i>amblyophyllum</i> (Kar. & Kir.) N.Friesen	[7, 13, 14]
	<i>Allium polyrhizum</i> Turcz.	[1, 2, 3, 4, 7–16]
	<i>Allium prostratum</i> Trev.	[1–13]
SE	<i>Allium pumilum</i> Vved.	[6, 7, 14]
	<i>Allium ramosum</i> L.	[1–13]
	<i>Allium rubens</i> Schrad.	[6, 7, 14]
	<i>Allium schischkinii</i> Sobolevsk.	[3, 6, 7, 10, 11, 13]
	<i>Allium schoenoprasum</i> L.	[1–7, 10]
	<i>Allium schrenkii</i> Regel [= <i>Allium bogdoicola</i> Regel]	[3, 6, 7, 10, 13, 14]
	<i>Allium senescens</i> L.	[1–10, 13]
	<i>Allium splendens</i> Willd.	[1–5, 8, 9]
	<i>Allium spurium</i> G.Don	[1, 2, 4, 5, 9]
	<i>Allium stellerianum</i> Willd.	[1, 2, 3, 4]
	<i>Allium strictum</i> Schrad.	[1–10, 13, 14]
SE	<i>Allium subangulatum</i> Regel	[16]
	<i>Allium subtilissimum</i> Ledeb.	[3, 14]
	<i>Allium tenuissimum</i> L.	[1–5, 7–9, 11–15]
	<i>Allium tuvinicum</i> (N.Friesen) N.Friesen [≡ <i>Allium stellerianum</i> subsp. <i>tuvinicum</i> N.Friesen]	[3, 6, 7, 10]
SE	<i>Allium tytthocephalum</i> Schult.f.	[4, 6, 7, 13]
SE	<i>Allium ubsicola</i> Regel	[6, 10, 14]
	<i>Allium vodopjanovae</i> N.Friesen	[3, 4, 6–8, 10–15]

### 23. Apiaceae Lindl. (36 genera and 66 taxa)

	<i>Aegopodium alpestre</i> Ledeb.	[1–5, 13]
	<i>Angelica czernaevia</i> (Fisch. & C.A.Mey.) Kitag.	[5, 9]
	<i>Angelica dahurica</i> (Hoffm.) Benth. & Hook.f. [≡ <i>Callisace dahurica</i> Hoffm.]	[2, 3, 4, 5, 9]
	<i>Angelica saxatilis</i> Turcz. [= <i>Physolophilum saxatile</i> (Turcz.) Turcz.]	[2]
	<i>Angelica sylvestris</i> L.	[6, 7]
	<i>Anthriscus sylvestris</i> (L.) Hoffm.	[1–10]
	<i>Archangelica decurrens</i> Ledeb. [≡ <i>Angelica archangelica</i> subsp. <i>decurrens</i> (Ledeb.) Kuvaev]	[1–4, 6, 7, 14]
	<i>Aulacospermum anomalum</i> Ledeb.	[6, 7]
	<i>Bupleurum aureum</i> Fisch.	[7]
	<i>Bupleurum bicaule</i> Helm [= <i>Bupleurum pusillum</i> Krylov]	[1–4, 6–13]
	<i>Bupleurum densiflorum</i> Rupr. [= <i>Bupleurum mongolicum</i> V.M.Vinogr.]	[7, 13, 14]
	<i>Bupleurum krylovianum</i> Schischk.	[3, 7]
	<i>Bupleurum multinerve</i> DC. [= <i>Bupleurum longeinvolucratum</i> Krylov]	[1–5, 7, 9, 11]
	<i>Bupleurum scorzonerifolium</i> Willd.	[1–6, 8, 9, 12, 13]
	<i>Bupleurum sibiricum</i> Vest	[2, 4, 8, 9]

	<i>Carum buriaticum</i> Turcz.	[1–6, 8, 9]
	<i>Carum carvi</i> L.	[1–5, 7–10, 14]
	<i>Cenolophium denudatum</i> (Hornem.) Tutin	[3, 7, 10, 14]
	<i>Cicuta virosa</i> L.	[1–15]
	<i>Cnidium dauricum</i> (Jacq.) Turcz. [= <i>Laserpitium dauricum</i> Jacq.]	[2–10]
	<i>Cnidium monnieri</i> Cusson	[4, 9]
SE	<i>Conioselinum longifolium</i> Turcz.	[1, 2, 4, 7, 9, 10]
	<i>Conioselinum tataricum</i> Hoffm. [= <i>Conioselinum vaginatum</i> (Spreng.) Thell.]	[1, 2, 3, 4]
	<i>Elwendia setacea</i> (Schrenk) Pimenov & Kljuykov [= <i>Bunium setaceum</i> (Schrenk) H.Wolff, ≡ <i>Carum setaceum</i> Schrenk]	[6, 7]
SE	<i>Ferula bungeana</i> Kitag.	[5, 8–16]
	<i>Ferula caspica</i> M.Bieb.	[7, 14]
	<i>Ferula dissecta</i> Ledeb.	[3, 6, 7, 10, 14]
	<i>Ferula dshaudshamyr</i> Korovin [= <i>Ferula dubjanskyi</i> Korovin]	[7, 14]
	<i>Ferula feruloides</i> (Steud.) Korovin	[7]
	<i>Ferula potaninii</i> Korovin	[14]
	<i>Ferula soongarica</i> Pall. [= <i>Ferula mongolica</i> (V.M.Vinogr. & Kamelin) V.M.Vinogr. & Kamelin]	[3, 7, 10, 14, 15]
SE	<i>Ferulopsis hystrix</i> (Bunge) Pimenov [= <i>Peucedanum hystrix</i> Bunge]	[2–4, 6–11, 13, 15]
SE	<i>Haloselinum falcaria</i> (Turcz.) Pimenov [= <i>Peucedanum falcaria</i> Turcz.]	[1, 3, 4, 6–8, 10, 11, 13–16]
	<i>Hansenia mongholica</i> Turcz. [= <i>Ligusticum mongholicum</i> (Turcz.) Krylov]	[1, 2]
	<i>Heracleum dissectum</i> Ledeb.	[1–7, 9, 10, 11, 13]
	<i>Heracleum sibiricum</i> L.	[1, 2, 3, 9, 13]
	<i>Kadenia salina</i> (Turcz.) Lavrova & V.N.Tikhom. [= <i>Cnidium salinum</i> Liou]	[2, 3, 4, 8–11, 13]
	<i>Kitagawia baicalensis</i> (Redow.) Pimenov [≡ <i>Peucedanum baicalense</i> (Redow.) Koch]	[1–8, 10]
	<i>Kitagawia terebinthacea</i> (Fisch.) Pimenov [≡ <i>Peucedanum terebinthaceum</i> (Fisch.) Ledeb.]	[2, 4, 5, 9]
SE	<i>Lithosciadium kamelinii</i> (V.M.Vinogr.) Pimenov [≡ <i>Cnidium kamelinii</i> V.M.Vinogr.]	[7]
SE	<i>Lithosciadium multicaule</i> Turcz.	[1, 3, 4, 6, 7, 13]
	<i>Neogaya simplex</i> Meisn. [= <i>Pachypleurum alpinum</i> Ledeb.]	[10]
	<i>Oenanthe aquatica</i> (L.) Poir. [= <i>Peucedanum salinum</i> Pall.]	[1–4, 6–10, 13]
	<i>Ostericum tenuifolium</i> (Pall.) Y.C.Chu [= <i>Pachypleurum alpinum</i> Ledeb.]	[1–4, 6, 7, 13, 14]
	<i>Paraligusticum discolor</i> (Ledeb.) V.N.Tikhom.	[7]
SE	<i>Peucedanum puberulum</i> Turcz.	[2, 3, 6, 8, 13]
	<i>Peucedanum vaginatum</i> Ledeb.	[1–4, 6, 7, 8, 11, 13]

	<i>Phlojodicarpus sibiricus</i> Koso-Pol.	[1–4, 7, 8, 9, 13]
	<i>Phlojodicarpus villosus</i> Turcz.	[1, 2, 3, 6]
	<i>Pimpinella thellungiana</i> H.Wolff	[4, 5, 9]
	<i>Pleurospermum uralense</i> Hoffm.	[1–6, 8, 9]
	<i>Prangos ledebourii</i> Herrnst. & Heyn	[7, 14]
	<i>Sajanella monstrosa</i> (Willd.) Soják	[1, 2]
	<i>Saposhnikovia divaricata</i> (Turcz.) Schischk.	[2–6, 8, 9]
	<i>Schulzia crinita</i> (Pall.) Spreng.	[1, 2, 3, 4, 6, 7]
SE	<i>Seseli abolinii</i> (Korovin) Schischk. [= <i>Libanotis abolinii</i> (Korovin) Korovin]	[7, 10, 11, 13]
	<i>Seseli buchtormense</i> W.D.J.Koch [= <i>Libanotis buchtormensis</i> (Fisch.) DC.]	[7, 14]
	<i>Seseli condensatum</i> Rchb.f. [= <i>Libanotis condensata</i> (L.) Fisch.]	[1–3, 6–8, 10, 14]
	<i>Seseli eriocarpum</i> B.Fedtsch. [= <i>Libanotis eriocarpa</i> Schrenk]	[7, 10, 14]
	<i>Seseli glabratum</i> Willd. [= <i>Libanotis tenuifolia</i> DC.]	[7]
	<i>Seseli grubovii</i> V.M.Vinogr. & Sanchir [= <i>Libanotis grubovii</i> (V.M.Vinogr. & Sanchir) M.L.Sheh & M.F.Watson]	[7, 13, 14, 15]
	<i>Seseli mucronatum</i> (Schrenk) Pimenov & Sdobnina	[14]
	<i>Seseli seseloides</i> (Fisch. & C.A.Mey.) M.Hiroe [= <i>Libanotis seseloides</i> (Fisch. & C.A.Mey.) Turcz.]	[1–7, 9]
	<i>Sium suave</i> Walter	[1–10, 14]
	<i>Sphallerocarpus gracilis</i> Koso-Pol.	[1–4, 6–13]
SE	<i>Stenocoelium athamantoides</i> Ledeb. [= <i>Seseli athamantoides</i> (M.Bieb.) Beck]	[6, 7]

#### 24. Apocynaceae Juss. (3 genera and 10 taxa)

	<i>Apocynum pictum</i> Schrenk [= <i>Apocynum hendersonii</i> Hook.f.]	[7, 11, 14, 15]
	<i>Apocynum venetum</i> L. [= <i>Poacynum venetum</i> (L.) Mavrodiev]	[14]
	<i>Cynanchum acutum</i> subsp. <i>sibiricum</i> (Willd.) Rech.f.	[10–16]
	<i>Cynanchum bungei</i> Decne.	[9]
	<i>Cynanchum chinense</i> R.Br.	[9, 12, 15, 16]
	<i>Cynanchum gobicum</i> Grubov [= <i>Vincetoxicum lanceolatum</i> (Grubov) Grubov]	[12–16]
	<i>Cynanchum mongolicum</i> Hemsl.	[16]
	<i>Cynanchum purpureum</i> K.Schum.	[1, 2, 4, 5, 8, 9, 12, 16]
	<i>Vincetoxicum mukdenense</i> Kitag. [= <i>Cynanchum paniculatum</i> (Bunge) Kitag.]	[4, 5, 9]
	<i>Vincetoxicum sibiricum</i> (L.) Decne. [= <i>Cynanchum thesioides</i> K.Schum.]	[2–16]

#### 25. Araceae Juss. (2 genera and 4 species)

	<i>Lemna minor</i> L.	[1, 3, 4, 7–11, 13]
	<i>Lemna trisulca</i> L.	[1–5, 8–11]
	<i>Lemna turionifera</i> Landolt	[9, 11]

*Spirodela polyrhiza* (L.) Schleid. [5, 9]

**26. Asparagaceae** Juss. (5 genera and 19 species)

	<i>Anemarrhena asphodeloides</i> Bunge	[5, 9]
	<i>Asparagus brachyphyllus</i> Turcz.	[9]
SE	<i>Asparagus burjaticus</i> Peschkova	[4]
	<i>Asparagus dauricus</i> Fisch.	[2–6, 8, 9, 11, 12]
SE	<i>Asparagus gobicus</i> Ivanova	[7–16]
	<i>Asparagus neglectus</i> Kar. & Kir.	[14]
	<i>Asparagus oligoclonos</i> Maxim.	[5]
	<i>Asparagus pallasii</i> Misch.	[7, 10, 11]
	<i>Asparagus schoberioides</i> Kunth	[5]
	<i>Asparagus tamariscinus</i> Ivanova	[10, 14, 15, 16]
	<i>Asparagus trichophyllus</i> Bunge	[10, 12, 14–16]
	<i>Convallaria keiskei</i> Miq.	[2, 5]
	<i>Maianthemum bifolium</i> (L.) F.W.Schmidt	[1, 2, 3, 4, 5]
	<i>Maianthemum dilatatum</i> (Alph.Wood) A.Nelson & J.F.Macbr.	[3, 4, 5, 9]
	<i>Maianthemum × intermedium</i> Vorosch.	[5]
	<i>Maianthemum trifolium</i> (L.) Sloboda	[2]
	<i>Polygonatum humile</i> Fisch.	[4, 5]
	<i>Polygonatum odoratum</i> (Mill.) Druce	[1–5, 8, 9]
	<i>Polygonatum sibiricum</i> Redouté	[1–5, 8, 9, 12]

**27. Asphodelaceae** Juss. [including Xantorrhoeaceae Dumort.] (1 genus and 2 taxa)

	<i>Hemerocallis lilioasphodelus</i> L. var. <i>lilioasphodelus</i>	[4, 5, 9]
	<i>Hemerocallis lilioasphodelus</i> var. <i>minor</i> (Mill.) M.N.Tamura [= <i>Hemerocallis minor</i> Mill.]	[1–5, 8, 9]

**28. Asteraceae** Bercht. & J.Presl (85 genera and 456 taxa)

Note: Some classifications of some genera of Asteraceae have changed after extensive molecular investigations. For example, species of *Scorzonera* L. were split into several genera, and three of them are present in Mongolia: *Lipschitzia* Zaika, Sukhor. & N.Kilian, *Takhtajaniantha* Nazarova, and *Scorzonera* L. s.str. by Zaika et al. (2020). The taxonomic status of *Scorzonera curvata*, *S. grubovii*, and *S. sinensis* is not resolved yet.

	<i>Achillea acuminata</i> Sch.Bip.	[2, 4, 5, 9]
	<i>Achillea alpina</i> L.	[1–6, 8–10]
	<i>Achillea asiatica</i> Serg.	[1–10, 14]
	<i>Achillea impatiens</i> L.	[2, 3, 4]
	<i>Achillea ledebourii</i> Heimerl	[3, 7, 8]
	<i>Achillea millefolium</i> L.	[1, 2, 3, 4, 7]
	<i>Achillea ptarmicoides</i> Maxim.	[2, 4, 8, 9, 10]

	<i>Achillea sergievkiana</i> Shaulo & Shmakov	[7]
SE	<i>Ajania achilleoides</i> Poljakov	[3, 6–8, 10–13, 15, 16]
	<i>Ajania fruticulosa</i> (Ledeb.) Poljakov	[3, 4, 6–16]
E	<i>Ajania grubovii</i> Muldashev [= <i>Chrysanthemum grubovii</i> (Muldashev) H.Ohashi & Yonek.]	[7, 14]
	<i>Ajania trifida</i> (Turcz.) Muldashev [= <i>Hippolytia trifida</i> (Turcz.) Poljakov]	[3, 6–9, 11–13, 16]
	<i>Allardia tridactylites</i> Sch.Bip. [= <i>Waldheimia tridactylites</i> Kar. & Kir. ]	[1, 3, 6, 7, 13]
	<i>Ancathia igniaria</i> DC.	[3, 7, 10, 14, 15]
	<i>Antennaria dioica</i> (L.) Gaertn.	[1, 2, 3, 4, 7]
	<i>Arctium tomentosum</i> Mill.	[4]
	<i>Arctogeron gramineum</i> (L.) DC.	[1–5, 7, 8, 9]
	<i>Arnica angustifolia</i> subsp. <i>iljinii</i> (Maguire) I.K.Ferguson	[7]
	<i>Artemisia adamsii</i> Besser	[2, 3, 4, 6–13]
SE	<i>Artemisia aksaiensis</i> Y.R.Ling	[2, 6, 8, 12–14, 16]
	<i>Artemisia amoena</i> Poljakov	[7, 12]
	<i>Artemisia anethifolia</i> Weber	[2, 3, 4, 7–16]
	<i>Artemisia anethoides</i> Mattf.	[8–16]
	<i>Artemisia annua</i> L.	[2–4, 7–10, 12–16]
	<i>Artemisia argyi</i> H.Lév. & Vaniot	[2, 4, 5, 7–9, 12, 13]
	<i>Artemisia argyrophylla</i> Ledeb.	[1, 3, 6, 7, 13, 15]
E	<i>Artemisia assurgens</i> Filatova [= <i>Seriphidium assurgens</i> (Filatova) K.Bremer & Humphries]	[7, 11, 13–15]
	<i>Artemisia aurata</i> Kom.	[2–5, 8, 9, 13]
	<i>Artemisia bargusinensis</i> Spreng.	[1, 2, 3, 4, 5]
SE	<i>Artemisia blepharolepis</i> Bunge	[11, 12, 13, 16]
	<i>Artemisia borealis</i> Pall.	[1–4, 6, 7, 10, 13]
	<i>Artemisia borotalensis</i> Poljakov [= <i>Seriphidium borotalense</i> (Poljakov) Ling & Y.R.Ling]	[7, 14]
SE	<i>Artemisia brachyloba</i> Franch.	[4, 8, 9]
	<i>Artemisia brachyphylla</i> Kitam.	[5]
	<i>Artemisia caespitosa</i> Ledeb.	[3, 4, 6–16]
	<i>Artemisia capillaris</i> Thunb.	[2, 3, 4, 5, 8, 9]
	<i>Artemisia compacta</i> Fisch.	[3, 6–8, 10–12, 14]
	<i>Artemisia dahurica</i> (Turcz.) Poljakov	[4]
E	<i>Artemisia davazamczii</i> Darijma & Kamelin	[7, 10, 13, 15]
	<i>Artemisia demissa</i> Krasch.	[3, 7–16]
	<i>Artemisia depauperata</i> Krasch.	[1–4, 6–8, 10, 11, 13, 14]
	<i>Artemisia desertorum</i> Spreng. subsp. <i>desertorum</i>	[2, 4, 5, 9, 13]
E	<i>Artemisia desertorum</i> subsp. <i>pseudojaponica</i> Darijma & Kamelin	[5]
SE	<i>Artemisia disjuncta</i> Krasch.	[7, 13]

SE	<i>Artemisia dolosa</i> Krasch.	[1–9, 11, 13]
SE	<i>Artemisia dracunculus</i> var. <i>changaica</i> (Krasch.) Y.R.Ling [= <i>Artemisia changaica</i> Krasch.]	[1, 3, 7, 8, 10, 11, 13]
	<i>Artemisia dracunculus</i> L. var. <i>dracunculus</i>	[1–15]
	<i>Artemisia eriopoda</i> Bunge	[16]
	<i>Artemisia feddei</i> subsp. <i>arschantinica</i> (Darijma) Gubanov & Kamelin [= <i>Artemisia arschantinica</i> Darijma]	[16]
	<i>Artemisia feddei</i> H.Lév. & Vaniot subsp. <i>feddei</i>	[5, 9]
	<i>Artemisia freyniana</i> (Pamp.) Krasch.	[4–6, 8–10, 12, 13]
	<i>Artemisia frigida</i> Willd.	[1–16]
SE	<i>Artemisia giraldii</i> Pamp.	[4]
	<i>Artemisia glauca</i> Pall.	[1–8, 10, 14]
SE	<i>Artemisia globosa</i> Krasch.	[6–8, 10, 12–14]
SE	<i>Artemisia globosoides</i> Ling & Y.R.Ling	[9, 12]
	<i>Artemisia gmelinii</i> Web. var. <i>gmelinii</i>	[2–13]
	<i>Artemisia gmelinii</i> var. <i>messerschmidiana</i> (Besser) Poljakov	[2–5, 8, 9, 12, 13]
	<i>Artemisia gracilescens</i> Krasch. & Iljin	[7, 14, 15]
	<i>Artemisia halodendron</i> Turcz.	[4, 5, 8, 9, 12, 16]
	<i>Artemisia heptapotamica</i> Poljakov [≡ <i>Seriphidium heptapotamicum</i> (Poljakov) Ling & Y.R.Ling]	[7, 14]
	<i>Artemisia implicata</i> T.G.Leonova	[16]
	<i>Artemisia integrifolia</i> L.	[1–5, 8, 9, 13]
	<i>Artemisia klementzae</i> Krasch. [= <i>Artemisia xylorrhiza</i> Krasch.]	[3, 4, 7–13, 16]
	<i>Artemisia laciniata</i> Willd.	[1–5, 7–10, 12, 14]
SE	<i>Artemisia lagocephala</i> Fisch. var. <i>lithophila</i> (Turcz.) Y.R.Ling	[1]
	<i>Artemisia latifolia</i> Ledeb.	[2, 4, 5, 9]
	<i>Artemisia macilenta</i> (Maxim.) Krasch.	[2, 3, 4, 5, 9]
	<i>Artemisia macrantha</i> Ledeb.	[1, 2, 3, 7]
	<i>Artemisia macrocephala</i> Jacquem.	[1–16]
	<i>Artemisia manshurica</i> (Kom.) Kom.	[2, 3, 4, 5, 8, 9]
	<i>Artemisia marschalliana</i> Spreng.	[7]
	<i>Artemisia maximovicziana</i> Krasch.	[4, 5, 9]
	<i>Artemisia medioxima</i> Krasch.	[1, 2, 3, 4, 9]
	<i>Artemisia mongolica</i> (Fisch.) Nakai [= <i>Artemisia vulgaris</i> var. <i>mongolica</i> Fisch.]	[1–15]
	<i>Artemisia mongolorum</i> subsp. <i>gobicum</i> Krasch. [≡ <i>Artemisia gobica</i> (Krasch.) Grubov]	[3, 4, 6–16]
	<i>Artemisia mongolorum</i> Krasch. subsp. <i>mongolorum</i> [≡ <i>Seriphidium mongolorum</i> (Krasch.) Ling & Y.R.Ling]	[3, 4, 6–16]
	<i>Artemisia nitrosa</i> Weber	[3, 4, 8, 9]
	<i>Artemisia obtusiloba</i> subsp. <i>altaicensis</i> (Krasch.) Krasnob. [= <i>Artemisia altaicensis</i> Krasch.]	[3, 6, 7]

	<i>Artemisia obtusiloba</i> Ledeb. subsp. <i>obtusiloba</i>	[3, 6, 7, 10, 13, 14]
	<i>Artemisia obtusiloba</i> var. <i>glabra</i> Ledeb. [= <i>Artemisia glabella</i> Kar. & Kir.]	[3, 6, 10]
SE	<i>Artemisia ordosica</i> Krasch.	[7, 9, 10, 12–16]
SE	<i>Artemisia oxycephala</i> Kitag.	[4, 5, 8, 9]
	<i>Artemisia palustris</i> L.	[1–13]
	<i>Artemisia pamirica</i> C.Winkl.	[3, 6, 7, 10–13]
	<i>Artemisia phaeolepis</i> Krasch.	[1–4, 6–9, 13, 14]
	<i>Artemisia pubescens</i> Ledeb. [= <i>Artemisia commutata</i> Besser]	[1–4, 6–11, 13, 14]
	<i>Artemisia pycnorrhiza</i> Ledeb.	[1–4, 6–8, 10–14]
	<i>Artemisia rubripes</i> Nakai	[2, 3, 4, 5, 8, 9]
	<i>Artemisia rupestris</i> L.	[1–4, 6–8, 10, 14]
	<i>Artemisia rutifolia</i> var. <i>altaica</i> (Krylov) Krasch.	[7]
	<i>Artemisia sacrorum</i> var. <i>messerschmidiana</i> (Besser) Y.R.Ling	[2–5, 8, 9, 12, 13]
	<i>Artemisia saissanica</i> (Krasch.) Filatova	[7, 10, 14]
	<i>Artemisia santolinifolia</i> Turcz.	
	[= <i>Artemisia santolinifolia</i> subsp. <i>stepposa</i> Darijma]	[2, 3, 6–15]
	<i>Artemisia schischkinii</i> Krasch.	[6, 7, 10, 14]
	<i>Artemisia schrenkiana</i> Ledeb.	[3, 6, 10, 14]
	<i>Artemisia scoparia</i> Waldst. & Kit.	[2–12]
	<i>Artemisia selengensis</i> Turcz.	[3, 4, 5, 9]
	<i>Artemisia sericea</i> Weber	[1, 2, 3, 4, 5, 8]
	<i>Artemisia sieversiana</i> Ehrh.	[1–16]
SE	<i>Artemisia sphaerocephala</i> Krasch.	[3, 10–16]
	<i>Artemisia stolonifera</i> (Maxim.) Kom.	[3, 7, 10, 13–16]
SE	<i>Artemisia subchryssolepis</i> Filatova [= <i>Seriphidium subchryssolepis</i> (Filatova) K.Bremer & Humphries]	[7, 14]
	<i>Artemisia subdigitata</i> Mattf. [= <i>Artemisia dubia</i> var. <i>subdigitata</i> (Mattf.) Y.R.Ling]	[3, 4, 7, 10, 12–16]
	<i>Artemisia sublessingiana</i> Krasch. [= <i>Seriphidium gorjaevii</i> (Poljak.) Y.R.Ling]	[14]
	<i>Artemisia subulata</i> Nakai	[1, 5, 9]
	<i>Artemisia succulenta</i> Ledeb.	[7]
	<i>Artemisia sylvatica</i> Maxim.	[4, 5, 9, 10, 15]
	<i>Artemisia tanacetifolia</i> L.	[1–10, 14]
	<i>Artemisia terrae-albae</i> Krasch.	[7, 14, 15]
	<i>Artemisia tomentella</i> Trautv.	[1, 3, 6, 10–12, 14]
	<i>Artemisia tournefortiana</i> Rchb.	[7, 9, 12, 14]
SE	<i>Artemisia transbaicalensis</i> T.G.Leonova	[1, 3]
	<i>Artemisia umbrosa</i> (Besser) Turcz.	[4, 5, 9]
	<i>Artemisia vestita</i> Wall.	[13]

	<i>Artemisia viridis</i> Willd.	[6, 7, 14]
	<i>Artemisia vulgaris</i> subsp. <i>vulgaris</i> L.	[2, 3]
E	<i>Artemisia vulgaris</i> subsp. <i>inundata</i> Darijma [= <i>Artemisia superba</i> Pamp.]	[1–4, 7, 9, 10, 13, 14]
	<i>Artemisia wudanica</i> Liou & W.Wang	[8, 9, 12]
SE	<i>Artemisia xanthochloa</i> Krasch.	[3–16]
	<i>Artemisia xerophytica</i> Krasch.	[6, 7, 8, 10–16]
	<i>Askellia flexuosa</i> (Ledeb.) W.A.Weber	[1, 3, 5–11, 13–16]
	<i>Askellia pygmaea</i> (Ledeb.) Sennikov	[1, 3, 6, 7]
	<i>Aster alpinus</i> L.	[1–10, 13]
	<i>Aster hispidus</i> Thunb.	[2–6, 8–11, 13, 15]
	<i>Aster lingii</i> G.J.Zhang & T.G.Gao [= <i>Rhinactinidium limoniifolia</i> Novopokr.]	[7]
E	<i>Aster maackii</i> Regel	[5]
	<i>Aster sanczirii</i> Kamelin & Cubanov	[5]
	<i>Aster tataricus</i> L.f.	[2, 3, 4, 5, 9]
SE	<i>Astrothamnus alyssoides</i> (Turcz.) Novopokr. [= <i>Aster alyssoides</i> Turcz.]	[8, 12]
SE	<i>Astrothamnus centralasiaticus</i> var. <i>potaninii</i> (Novopokr.) Y.Ling & Y.L.Chen [≡ <i>Astrothamnus potaninii</i> Novopokr.]	[7, 8, 9, 11–16]
SE	<i>Astrothamnus heteropappoides</i> Novopokr.	[6, 7, 10, 14]
	<i>Astrothamnus molliusculus</i> Novopokr.	[12, 15]
	<i>Astrothamnus polifolius</i> Novopokr.	[3, 6, 7, 10, 11, 13–15]
	<i>Bidens cernua</i> L.	[3, 4, 6, 7, 9–11]
	<i>Bidens parviflora</i> Willd.	[3, 4, 6, 8–10, 13]
	<i>Bidens radiata</i> Thuill.	[3, 4, 6–10]
	<i>Bidens tripartita</i> L.	[1–4, 7, 8–10, 14]
SE	<i>Brachanthemum gobicum</i> Krasch.	[12, 13, 16]
SE	<i>Brachanthemum mongolicum</i> Krasch.	[12, 14]
E	<i>Brachanthemum mongolorum</i> Grubov	[9]
	<i>Cancrinia discoidea</i> (Ledeb.) Poljakov	[7, 10–16]
SE	<i>Cancrinia krasnoborovii</i> Khanm.	[10]
	<i>Carduus crispus</i> L.	[2–7, 9]
	<i>Carduus nutans</i> L.	[1, 7]
	<i>Centaurea adpressa</i> Ledeb.	[6]
	<i>Centaurea glastifolia</i> subsp. <i>intermedia</i> (Boiss.) L.Martins [= <i>Centaurea chartolepis</i> Greuter]	[6, 7]
	<i>Centaurea pulchella</i> Ledeb. [= <i>Hyalea pulchella</i> (Ledeb.) K.Koch]	[7, 14]
	<i>Chondrilla lejosperma</i> Kar. & Kir.	[6, 7, 10, 14]
E	<i>Chrysanthemum chalchingolicum</i> Grubov	[5, 9]
	<i>Chrysanthemum mongolicum</i> Ling [= <i>Chrysanthemum zawadzkii</i> var. <i>mongolicum</i> (Ling) Cubanov]	[1, 2, 3]

	<i>Chrysanthemum naktongense</i> Nakai	[9]
SE	<i>Chrysanthemum sinuatum</i> Ledeb. [ <i>Tanacetum sinuatum</i> Sch.Bip.]	[6, 7]
	<i>Chrysanthemum trilobatum</i> (Poljakov) H.Ohashi & Yonek. [≡ <i>Ajania trilobata</i> Poljakov]	[12, 13]
	<i>Chrysanthemum zawadzkii</i> Herbich	[1–5, 8, 9]
	<i>Cicerbita azurea</i> (Ledeb.) Beaverd	[3, 7, 10]
	<i>Cirsium arvense</i> (L.) Scop. [≡ <i>Serratula arvensis</i> L.]	[2–4, 7, 9–11, 13–15]
	<i>Cirsium esculentum</i> C.A.Mey.	[1–4, 6–11, 14]
	<i>Cirsium glabrifolium</i> O.Fedtsch. & B.Fedtsch.	[7]
	<i>Cirsium helenioides</i> (L.) Hill [= <i>Carduus helenioides</i> L.]	[2]
	<i>Cirsium pendulum</i> Fisch.	[2, 4, 5, 9]
	<i>Cirsium serratuloides</i> Hill	[1, 2, 3]
	<i>Cirsium setosum</i> (Willd.) M.Bieb. [≡ <i>Serratula setosa</i> Willd.]	[3, 4, 7–11, 14, 15]
	<i>Cirsium sieversii</i> (Fisch. & C.A.Mey.) Petr. [= <i>Cirsium polyacanthum</i> Kar. & Kir.]	[7]
	<i>Cirsium vlassovianum</i> Fisch.	[2, 5, 9]
	<i>Cousinia affinis</i> Schrenk	[14]
	<i>Crepidiastrum akagii</i> (Kitag.) J.W.Zhang & N.Kilian [= <i>Youngia tenuicaulis</i> (Babc. & Stebbins) Czerep.]	[2, 3, 6–8, 10–15]
	<i>Crepidiastrum sonchifolium</i> (Bunge) Pak & Kawano	[5]
	<i>Crepidiastrum tenuifolium</i> (Willd.) Sennikov [≡ <i>Crepis tenuifolia</i> Willd. ≡ <i>Youngia tenuifolia</i> (Willd.) Babc. & Stebbins]	[1–11, 13, 14]
	<i>Crepis bungei</i> Ledeb.	[1–4, 6–9, 11]
	<i>Crepis chrysantha</i> Froel.	[1, 2, 3, 6, 7, 10]
	<i>Crepis crocea</i> (Lam.) Babc. var. <i>crocea</i>	[2, 3, 4, 6–13]
SE	<i>Crepis crocea</i> var. <i>cuzuensis</i> (Serg.) Tzvelev [≡ <i>Crepis czuensis</i> Serg.]	[6, 7]
E	<i>Crepis lomonosovae</i> Tzvelev	[3, 13]
	<i>Crepis lyrata</i> (L.) Froel.	[1, 7]
	<i>Crepis multicaulis</i> Ledeb.	[3, 7, 10, 13, 14]
	<i>Crepis polysticha</i> Turcz.	[1, 3, 4, 6, 7]
	<i>Crepis praemorsa</i> (L.) Tausch [≡ <i>Hieracium praemorsum</i> L.]	[4, 10]
	<i>Crepis sibirica</i> L.	[2–5, 7, 8, 11]
	<i>Crepis tectorum</i> L.	[2, 4, 7, 10, 14]
	<i>Doronicum altaicum</i> Pall.	[1]
	<i>Doronicum oblongifolium</i> DC.	[7, 14]
	<i>Doronicum turkestanicum</i> Cavill.	[3, 7, 14]
	<i>Echinops davuricus</i> Fisch. [= <i>Echinops latifolius</i> Tausch]	[1–5, 8, 9]
	<i>Echinops gmelinii</i> Turcz.	[3, 7–16]
	<i>Echinops humilis</i> M.Bieb.	[3, 7, 13–15]
	<i>Echinops integrifolius</i> Kar. & Kir.	[6, 7, 14]
	<i>Echinops nanus</i> Bunge	[7, 14]
	<i>Echinops ritro</i> L.	[7, 14]

	<i>Erigeron acris</i> L.	[1–7, 9, 10, 13]
SE	<i>Erigeron altaicus</i> Popov	[7, 14]
	<i>Erigeron baicalensis</i> Botsch.	[1]
	<i>Erigeron eriocalyx</i> (Ledeb.) Vierh.	[1–3, 6, 7, 13]
	<i>Erigeron krylovii</i> Serg.	[3, 7]
	<i>Erigeron lonchophyllus</i> Hook.	[1–7, 9, 10, 13]
	<i>Erigeron oreades</i> Fisch. & C.A.Mey.	[1, 3, 7, 13]
	<i>Erigeron petiolaris</i> Vierh.	[3, 7]
	<i>Erigeron politus</i> Fr.	[1–4, 6, 7, 13]
	<i>Erigeron pseudoeriocephalus</i> Popov	[3]
	<i>Filago arvensis</i> L.	[7, 8, 10, 14]
	<i>Filifolium sibiricum</i> (L.) Kitam. [= <i>Tanacetum sibiricum</i> L.]	[1–5, 8, 9]
	<i>Galatella altaica</i> Tzvelev	[7, 14]
	<i>Galatella angustissima</i> (Tausch) Novopokr.	[1]
	<i>Galatella dahurica</i> DC. [= <i>Galatella macrosciadia</i> Gand. = <i>Galatella songorica</i> Novopokr.]	[1–7, 9, 10]
	<i>Galatella hauptii</i> Lindl.	[7]
	<i>Gnaphalium uliginosum</i> L. [= <i>Gnaphalium baicalense</i> Kirp. & Kuprian.]	[2–4, 7, 9, 10, 14]
SE	<i>Helichrysum arenarium</i> Moench	[7]
SE	<i>Heteropappus altaicus</i> Novopokrov. [= <i>Aster altaicus</i> Willd.]	[1–4, 6–8, 10, 12–16]
	<i>Heteropappus biennis</i> (Ledeb.) Tamamsch.	[1–5, 8, 9]
	<i>Heteropappus medius</i> (Krylov) Tamamsch.	[3, 4, 5, 8, 9]
SE	<i>Hieracium czadanense</i> Tupitz.	[1, 10]
	<i>Hieracium korshinskyi</i> Zahn	[2, 4]
	<i>Hieracium narymense</i> Schischk. & Serg.	[2, 4]
	<i>Hieracium robustum</i> Fr.	[8, 9]
	<i>Hieracium sershukense</i> Üksip	[7]
	<i>Hieracium subramosum</i> Lonnr.	[2, 4]
	<i>Hieracium umbellatum</i> L.	[1–5, 7, 8, 9, 10]
	<i>Hieracium virosum</i> Pall.	[2, 3, 4, 5, 7, 9]
	<i>Holeion maximowiczii</i> Kitam.	[9]
	<i>Hypochaeris maculata</i> L. [= <i>Trommsdorffia maculata</i> (L.) Bernh.]	[4, 8]
	<i>Inula japonica</i> Thunb.	[2]
	<i>Inula linariifolia</i> Turcz.	[5, 8, 10, 11]
	<i>Inula salsoloides</i> Ostenf. [= <i>Limbarda salsoloides</i> Ikonn.]	[8, 11, 12, 13, 15, 16]
	<i>Ixeris chinensis</i> (Thunb.) Kitagawa subsp. <i>chinensis</i> s.l. [= <i>Ixeridium graminifolium</i> (Ledeb.) Tzvelev, = <i>Ixeridium gramineum</i> (Fisch.) Tzvelev]	[2–5, 7–9, 12, 14]
	<i>Jacobaea ambracea</i> (Turcz.) B.Nord. [= <i>Senecio ambraceus</i> Turcz.]	[2, 3, 4, 6–10, 14]
	<i>Jacobaea cannabifolia</i> (Less.) E.Wiebe [= <i>Senecio cannabifolius</i> Less.]	[2, 3, 5, 9]

	<i>Jacobaea erucifolia</i> subsp. <i>argunensis</i> (Turcz.) Veldkamp [≡ <i>Senecio argunensis</i> Turcz.]	[5, 9]
	<i>Jacobaea erucifolia</i> (L.) G.Gaertn., B.Mey. & Scherb. subsp. <i>erucifolia</i> [= <i>Senecio erucifolius</i> L.]	[2–4, 6, 7, 9, 10]
	<i>Jacobaea vulgaris</i> Gaertn. [≡ <i>Senecio jacobaea</i> L.]	[3, 4, 7–10, 14]
	<i>Jurinea chaetocarpa</i> (Ledeb.) Ledeb.	[7, 14]
	<i>Jurinea margalensis</i> Iljin	[7, 14]
SE	<i>Jurinea mongolica</i> Maxim. [= <i>Jurinea potaninii</i> Iljin]	[10–14]
	<i>Jurinea multiflora</i> B.Fedtsch.	[7, 14]
	<i>Karelinia caspia</i> Less.	[14, 15, 16]
	<i>Kaschgaria komarovii</i> (Krasch. & Rubtzov) Poljakov [≡ <i>Tanacetum komarovii</i> Krasch. & Rubtzov]	[7, 14, 15]
	<i>Klasea cardunculus</i> (Pall.) Holub [≡ <i>Serratula cardunculus</i> (Pall.) Schischk.]	[2, 3, 4, 5, 7]
	<i>Klasea centaurooides</i> (L.) Cass. [≡ <i>Serratula centaurooides</i> L.]	[1–5, 7–13]
	<i>Klasea marginata</i> (Tausch) Kitag. [≡ <i>Serratula marginata</i> Tausch]	[1–4, 7–10, 13, 14]
	<i>Klasea sogdiana</i> (Bunge) L.Martins [≡ <i>Serratula sogdiana</i> Bunge, <i>Serratula alatavica</i> C.A.Mey.]	[6]
	<i>Lactuca serriola</i> L. [= <i>Lactuca sativa</i> subsp. <i>serriola</i> (L.) Frietema]	[7, 10, 14, 15]
	<i>Lactuca sibirica</i> Benth.	[2–6, 8, 9, 11]
	<i>Lactuca tatarica</i> C.A.Mey.	[3, 4, 6–16]
	<i>Lactuca undulata</i> Ledeb.	[7, 14]
	<i>Leibnitzia anandria</i> (L.) Turcz.	[2, 3, 4, 5, 9]
	<i>Leontopodium campestre</i> Hand.-Mazz.	[1–3, 6–9, 11, 13, 14]
	<i>Leontopodium conglobatum</i> Hand.-Mazz.	[1–9, 13]
	<i>Leontopodium leontopodioides</i> (Willd.) Beauverd	[1–5, 8, 9, 16]
	<i>Leontopodium nanum</i> (Hook.f. & Thomson) Hand.-Mazz.	[16]
	<i>Leontopodium ochroleucum</i> Beauverd	[1–3, 6, 7, 13]
	<i>Leontopodium palibinianum</i> Beauverd	[2, 4, 5]
	<i>Leuzea carthamoides</i> DC. [= <i>Rhaponticum carthamoides</i> (Willd.) Iljin]	[7]
	<i>Leuzea repens</i> (L.) D.J.N.Hind, [≡ <i>Rhaponticum repens</i> (L.) Hidalgo ≡ <i>Acroptilon repens</i> (L.) DC.]	[6, 7, 10–16]
	<i>Leuzea uniflora</i> (L.) Holub [= <i>Rhaponticum uniflorum</i> (L.) DC.]	[1–5, 8, 9]
	<i>Ligularia altaica</i> DC.	[6, 7]
	<i>Ligularia fischerii</i> (Ledeb.) Turcz.	[2, 3, 4, 5, 9]
	<i>Ligularia glauca</i> (L.) O.Hoffm.	[7]
	<i>Ligularia Hodgsonii</i> Hook.f.	[5, 9]
	<i>Ligularia mongolica</i> DC.	[5, 9]
	<i>Ligularia przewalskii</i> Diels	[9, 12]
	<i>Ligularia sagitta</i> (Maxim.) Mattf. [≡ <i>Senecio sagitta</i> Maxim.]	[4, 5, 9]
	<i>Ligularia sibirica</i> Cass.	[1, 2, 3, 4, 5, 9]

	<i>Lipschitzia divaricata</i> (Turcz.) Zaika, Sukhor. & N.Kilian [≡ <i>Scorzonera divaricata</i> Turcz.]	[6–13, 15, 16]
	<i>Matricaria chamomilla</i> L. [= <i>Matricaria recutita</i> L.]	[2]
	<i>Neopallasia pectinata</i> (Pall.) Poljakov	[1–4, 6–16]
SE	<i>Olgaea leucophylla</i> (Turcz.) Iljin	[8, 9, 11–13]
SE	<i>Olgaea lomonossowii</i> (Trautv.) Iljin	[9]
	<i>Omalotheca supina</i> (L.) DC. [= <i>Gnaphalium supinum</i> L.]	[2, 7]
	<i>Packera cymbalaria</i> (Pursh) W.A.Weber & Á.Löve [≡ <i>Senecio cymbalaria</i> Pursh]	[1, 3, 7]
	<i>Parasenecio hastatus</i> (L.) H.Koyama [≡ <i>Cacalia hastata</i> L.]	[1, 2, 3, 4, 5, 9]
	<i>Pentanema asperum</i> (Poir.) G.V.Boiko & Korniy. [≡ <i>Inula aspera</i> Poir.]	[2, 3, 9]
	<i>Pentanema britannica</i> (L.) D.Gut.Larr. [≡ <i>Inula britannica</i> L.]	[1–11, 13, 14]
	<i>Pentanema salicinum</i> (L.) D.Gut.Larr. [≡ <i>Inula salicina</i> L.]	[2, 3, 4, 5, 9]
	<i>Petasites frigidus</i> (L.) Fr.	[1]
	<i>Petasites radiatus</i> (J.F.Gmel.) Toman	[1]
	<i>Petasites rubellus</i> (J.F.Gmel.) Toman	[1, 3]
	<i>Phalacrachena calva</i> (Ledeb.) Iljin	[10]
	<i>Picris davurica</i> Fisch.	[1, 3, 4, 8, 9]
	<i>Picris hieracioides</i> L.	[2, 3, 4, 5]
	<i>Picris japonica</i> Thunb.	[2, 3, 4, 5, 9]
	<i>Pilosella dublitzkii</i> (B.Fedtsch. & Nevski) Sennikov [≡ <i>Hieracium dublitzkii</i> B.Fedtsch. & Nevski]	[7]
	<i>Pilosella echioioides</i> (L.) F.W.Schultz & Sch.Bip. [≡ <i>Hieracium echioioides</i> L.]	[2, 4]
	<i>Pulicaria vulgaris</i> Gaertn.	[10]
	<i>Rhinactinidia eremophila</i> (Bunge) Novopokr. [= <i>Rhinactinidia eremophila</i> subsp. <i>grubovii</i> Botsch.]	[3, 6, 7, 10, 11, 13, 14]
	<i>Richteria pyrethroides</i> Kar. & Kir. [≡ <i>Pyrethrum pyrethroides</i> (Kar. & Kir.) B.Fedtsch.]	[7]
SE	<i>Saussurea acuminata</i> Turcz.	[2, 4, 5]
SE	<i>Saussurea alaschanica</i> Maxim.	[6, 10]
SE	<i>Saussurea alata</i> DC.	[4, 6, 10]
SE	<i>Saussurea alpina</i> (L.) DC.	[1–3, 6, 7, 13]
SE	<i>Saussurea amara</i> (L.) DC.	[1–5, 7–12, 14]
SE	<i>Saussurea arctecapitulata</i> Lipsch.	[1, 3]
SE	<i>Saussurea baicalensis</i> B.L.Rob.	[1, 2, 3, 7]
SE	<i>Saussurea bogedaensis</i> Yu J.Wang & J.Chen	[14]
SE	<i>Saussurea catharinae</i> Lipsch.	[15]
SE	<i>Saussurea ceterachifolia</i> Lipsch.	[3, 6, 7]
	<i>Saussurea congesta</i> Turcz.	[1]
	<i>Saussurea controversa</i> DC.	[1, 2, 3, 5]
	<i>Saussurea coronata</i> Schrenk [= <i>Saussurea dshungarica</i> Iljin]	[7]

	<i>Saussurea daurica</i> Adams	[3, 6–16]
SE	<i>Saussurea dorogostaiskii</i> Palib.	[1, 2]
	<i>Saussurea elata</i> Ledeb.	[7]
	<i>Saussurea elegans</i> Ledeb. [= <i>Saussurea amoena</i> Kar. & Kir.]	[3, 6, 7]
SE	<i>Saussurea elongata</i> DC.	[1, 2, 4]
	<i>Saussurea foliosa</i> Ledeb.	[6, 7]
	<i>Saussurea glacialis</i> Herder	[1, 3, 6, 7, 13, 14]
SE	<i>Saussurea grubovii</i> Lipsch.	[7, 14, 15]
E	<i>Saussurea gubanovii</i> Kamelin	[15]
	<i>Saussurea involucrata</i> (Kar. & Kit.) Sch.Bip.	[1–3, 6, 7, 13, 14]
	<i>Saussurea japonica</i> (Thunb.) DC.	[9]
	<i>Saussurea klementzii</i> Lipsch.	[7]
SE	<i>Saussurea krasnoborovii</i> S.V.Smirn.	[1]
	<i>Saussurea krylovii</i> Schischk. & Serg.	[7]
	<i>Saussurea laciniata</i> Ledeb.	[3, 4, 6–8, 10, 11, 13–16]
	<i>Saussurea latifolia</i> Ledeb.	[3, 7]
	<i>Saussurea leucophylla</i> Schrenk	[1, 3, 6, 7, 13]
SE	<i>Saussurea lipschitzii</i> Filatova	[7, 13]
	<i>Saussurea mongolica</i> (Franch.) Franch.	[5]
	<i>Saussurea neoserrata</i> Nakai	[2, 5]
	<i>Saussurea odontolepis</i> Sch.Bip.	[5]
E	<i>Saussurea odorata</i> E.Pjak	[7]
	<i>Saussurea orgaadayi</i> Khanm. & Krasnob.	[3, 7]
	<i>Saussurea parviflora</i> (Poir.) DC.	[1–7, 9]
SE	<i>Saussurea popovii</i> Lipsch.	[14]
	<i>Saussurea pricei</i> N.D.Simpson	[3, 6–8, 10, 11, 13, 14]
	<i>Saussurea pseudoalpina</i> N.D.Simpson	[1–3, 6, 7, 13, 14]
	<i>Saussurea pseudosalsa</i> Lipsch.	[15, 16]
	<i>Saussurea pulchella</i> Fisch.	[5, 7, 8, 9]
SE	<i>Saussurea purpurata</i> (Fisch.) Lipsch.	[2, 4]
E	<i>Saussurea ramosa</i> Lipsch.	[3, 10, 11, 15]
	<i>Saussurea recurvata</i> (Maxim.) Lipsch.	[2, 5]
	<i>Saussurea runcinata</i> DC.	[2–4, 7, 8, 10]
E	<i>Saussurea saichanensis</i> Kom.	[1–3, 6, 7, 13, 14]
	<i>Saussurea salicifolia</i> DC.	[2–9]
	<i>Saussurea salsa</i> Spreng.	[3, 5–11, 14, 16]
	<i>Saussurea schanginiana</i> (Wydler) Fisch.	[1–3, 6, 7, 13]
SE	<i>Saussurea squarrosa</i> Turcz.	[1]
	<i>Saussurea stubendorffii</i> Herder	[1, 3]
	<i>Saussurea subacaulis</i> (Ledeb.) Serg.	[1, 3, 6, 7, 13]
SE	<i>Saussurea sukaczewii</i> Lipsch.	[1, 2, 3]
	<i>Saussurea ussuriensis</i> Maxim.	[5]
	<i>Scorzonera albicalvis</i> Bunge	[1, 2, 4, 5, 9]

	<i>Scorzonera curvata</i> (Popl.) Lipsch.	[3, 7, 8, 9, 13]
E	<i>Scorzonera grubovii</i> Lipsch.	[7, 14]
	<i>Scorzonera parviflora</i> Jacq.	[14]
	<i>Scorzonera radiata</i> Fisch.	[1–10, 13, 14]
	<i>Scorzonera sinensis</i> (Lipsch. & Krasch.) Nakai	[9]
	<i>Senecio dubitabilis</i> C.Jeffrey & Y.L.Chen [≡ <i>Senecio dubius</i> Ledeb. nom. illegit. non Beck]	[2, 3, 7, 8, 10–15]
E	<i>Senecio kenteicus</i> Grubov	[2]
	<i>Senecio nemorensis</i> L.	[1, 2, 3]
	<i>Senecio subdentatus</i> Ledeb.	[7, 10, 14, 15]
	<i>Senecio vulgaris</i> L.	[1–4, 7, 8, 10]
	<i>Serratula coronata</i> L.	[5, 9]
	<i>Serratula kirghisorum</i> Iljin	[7]
	<i>Solidago dahurica</i> (Kitag.) Kitag.	[1–5, 7, 9]
	<i>Solidago virgaurea</i> L.	[7]
	<i>Sonchella dentata</i> (Ledeb.) Sennikov [≡ <i>Sonchus dentatus</i> Ledeb.]	[10, 14, 15]
	<i>Sonchella stenoma</i> (Turcz.) Sennikov [≡ <i>Crepis stenoma</i> Turcz.]	[8–15]
	<i>Sonchus arvensis</i> L.	[2–5, 7–11, 13, 14]
	<i>Sonchus brachyotus</i> DC.	[8–10, 13, 14]
	<i>Sonchus uliginosus</i> M.Bieb.	[4, 5, 8, 9, 10]
	<i>Stilpnolepis intricata</i> (Franch.) C.Shih	[3, 4, 7, 9–15]
	<i>Symphyotrichum ciliatum</i> (Ledeb.) G.L.Nesom	[3, 4, 9, 10]
	<i>Synurus deltoides</i> (Aiton) Nakai	[4, 5]
	<i>Takhtajaniantha austriaca</i> (Willd.) Zaika, Sukhor. & N.Kilian [≡ <i>Scorzonera austriaca</i> Willd.]	[2–10, 12–14]
	<i>Takhtajaniantha capito</i> (Maxim.) Zaika, Sukhor. & N.Kilian [≡ <i>Scorzonera capito</i> Maxim.]	[8, 11–16]
	<i>Takhtajaniantha ikonnikovii</i> (Krasch. & Lipsch.) Zaika, Sukhor. & N.Kilian [≡ <i>Scorzonera ikonnikovii</i> Lipsch. & Krasch.]	[3, 6–15]
	<i>Takhtajaniantha mongolica</i> (Maxim.) Zaika, Sukhor. & N.Kilian [≡ <i>Scorzonera mongolica</i> Maxim.]	[10–16]
	<i>Takhtajaniantha pseudodivaricata</i> (Lipsch.) Zaika, Sukhor. & N.Kilian [≡ <i>Scorzonera pseudodivaricata</i> Lipsch.]	[3, 6, 7, 9, 10–16]
	<i>Takhtajaniantha pusilla</i> (Pall.) Nazarova [≡ <i>Scorzonera pusilla</i> Pall.]	[8, 14]
	<i>Takhtajaniantha subacaulis</i> (Regel) Zaika, Sukhor. & N.Kilian [= <i>Scorzonera subacaulis</i> (Regel) Lipsch.]	[6]
	<i>Tanacetum alatavicum</i> Herder [≡ <i>Pyrethrum alatavicum</i> O.Fedtsch. & B.Fedtsch.]	[7]
E	<i>Tanacetum changaicum</i> (Krasch.) K.Bremer & Humphries [≡ <i>Pyrethrum changaicum</i> Krasch.]	[3, 4, 7, 10]
	<i>Tanacetum crassipes</i> (Stschegl.) Tzvelev	[7]
	<i>Tanacetum krylovianum</i> (Krasch.) K.Bremer & Humphries [≡ <i>Pyrethrum krylovianum</i> Krasch.]	[7]

	<i>Tanacetum lanuginosum</i> Sch. [= <i>Pyrethrum lanuginosum</i> (Sch.Bip. & Herder) Tzvelev]	[1, 6, 7, 13]
SE	<i>Tanacetum pulchellum</i> Sch. [= <i>Pyrethrum pulchellum</i> Turcz.]	[7]
	<i>Tanacetum pulchrum</i> Sch. [= <i>Pyrethrum pulchrum</i> Ledeb.]	[3, 6, 7, 13]
	<i>Tanacetum tanacetoides</i> (DC.) Tzvelev	[2, 3, 6, 7]
	<i>Tanacetum vulgare</i> L. [= <i>Tanacetum boreale</i> Fisch. & DC.]	[1–7, 9]
	<i>Taraxacum armeriifolium</i> Soest	[3–5, 7, 9–14, 16]
	<i>Taraxacum asiaticum</i> Dahlst.	[4, 7, 8, 13]
	<i>Taraxacum atrans</i> Schischk.	[7, 13]
	<i>Taraxacum bessarabicum</i> (Hornem.) Hand.-Mazz.	[2–4, 7–10]
	<i>Taraxacum bicorne</i> Dahlst.	[1, 2, 3, 7, 9–12]
SE	<i>Taraxacum bornuurense</i> R.Doll	[3, 4, 6, 7]
	<i>Taraxacum brevirostre</i> Hand.-Mazz.	[3, 7, 13]
	<i>Taraxacum ceratophorum</i> (Ledeb.) DC. [= <i>Taraxacum altaicum</i> Schischk.]	[1–7, 9, 13, 14]
	<i>Taraxacum collinum</i> DC.	[3, 4, 6–10, 14]
	<i>Taraxacum dealbatum</i> Hand.-Mazz.	[1–4, 6–15]
	<i>Taraxacum dissectum</i> Ledeb.	[1–4, 6–10, 12, 13]
	<i>Taraxacum eriopodium</i> DC.	[6, 7, 13, 14]
	<i>Taraxacum erythrospermum</i> Andrz.	[3]
	<i>Taraxacum glabrum</i> DC.	[1–3, 6, 7, 14]
	<i>Taraxacum glaucanthum</i> Nakai	[3, 4, 8]
	<i>Taraxacum goloskokovii</i> Schischk.	[6, 7, 10, 13]
E	<i>Taraxacum inimitabile</i> Kirschner & Štěpánek	[13]
E	<i>Taraxacum junatovii</i> Tzvelev	[3, 7, 13, 14]
	<i>Taraxacum kok-saghyz</i> Rodin	[3, 7, 13]
SE	<i>Taraxacum krasnoborovii</i> Krasnikov	[7]
SE	<i>Taraxacum krylovii</i> Krasnikov & Khanm.	[7]
	<i>Taraxacum leucanthum</i> Ledeb.	[1–4, 6–8, 10–15]
	<i>Taraxacum linczevskyi</i> Schischk.	[7]
SE	<i>Taraxacum longicorne</i> Dahlst.	[1, 2, 5–10]
	<i>Taraxacum luridum</i> G.E.Haglund	[6, 7]
	<i>Taraxacum lyratum</i> (Ledeb.) DC.	[1, 3, 6, 7]
	<i>Taraxacum macilentum</i> Dahlst.	[1, 3, 6, 7]
	<i>Taraxacum microspermum</i> Schischk. [= <i>Taraxacum compactum</i> Schischk.]	[1, 2]
	<i>Taraxacum minutilobum</i> Popov	[7]
	<i>Taraxacum mongolicum</i> Hand.-Mazz.	[1–4, 6, 7, 10, 11, 13]
	<i>Taraxacum mongoliforme</i> R.Doll	[1, 2, 4, 7–11, 13, 15]
	<i>Taraxacum monochlamydeum</i> Hand.-Mazz.	[3, 4, 7, 12–15]
	<i>Taraxacum mujense</i> Petrochenko	[1, 2]
	<i>Taraxacum multisectum</i> Kitag.	[9]
	<i>Taraxacum officinale</i> F.H.Wigg.	[1, 2, 3, 4]

	<i>Taraxacum parvulum</i> DC.	[14]
	<i>Taraxacum pawlodarskum</i> R.Doll [= <i>Taraxacum ustamenum</i> R.Doll]	[7]
	<i>Taraxacum pingue</i> Schischk.	[1, 3, 6, 7]
	<i>Taraxacum pseudoatratum</i> Orazova	[6]
SE	<i>Taraxacum pseudonivale</i> Malyschev	[1]
	<i>Taraxacum puberulum</i> G.E.Haglund	[14]
SE	<i>Taraxacum sangilense</i> Krasnob. & Khanm.	[1, 2, 3, 4, 6, 7]
	<i>Taraxacum scariosum</i> (Tausch) Kirschner & Štěpánek [= <i>Taraxacum stenolobum</i> Stschegl., <i>Taraxacum commixtiforme</i> Soest]	[4, 8–10, 13, 14]
E	<i>Taraxacum selengensis</i> Tzvelev	[3]
	<i>Taraxacum sinicum</i> Kitag. [= <i>Taraxacum borealisinense</i> Kitam.]	[3–16]
SE	<i>Taraxacum smirnovii</i> M.S.Ivanova	[7]
SE	<i>Taraxacum songoricum</i> Schischk.	[6, 7, 13]
	<i>Taraxacum stanjukoviczii</i> Schischk.	[7, 13]
E	<i>Taraxacum submacilentum</i> Tzvelev	[7]
	<i>Taraxacum sumneviczii</i> Schischk.	[1, 7, 13]
	<i>Taraxacum tibetanum</i> Hand.-Mazz.	[13, 14]
	<i>Taraxacum turgaicum</i> Schischk.	[7, 13, 14]
SE	<i>Taraxacum tuvense</i> Krasnob. & Krasnikov	[1]
	<i>Tephroseris flammula</i> (DC.) Holub [= <i>Senecio flammus</i> DC.]	[5]
	<i>Tephroseris integrifolia</i> subsp. <i>atropurpurea</i> (Ledeb.) B.Nord.	[1]
	<i>Tephroseris integrifolia</i> (L.) Holub subsp. <i>integrifolia</i> [= <i>Senecio campestris</i> (Retz.) DC.]	[1–4, 6–9, 13]
	<i>Tephroseris kirilowii</i> (DC.) Holub	[5]
	<i>Tephroseris palustris</i> (L.) Rchb. [= <i>Senecio palustris</i> (L.) Hook.]	[1, 2, 4–10]
SE	<i>Tephroseris porphyrantha</i> (Schischk.) Holub [= <i>Senecio porphyranthus</i> Schischk.]	[1, 7]
	<i>Tephroseris praticola</i> (Sisk. & Serg.) Holub [= <i>Senecio asiatica</i> Schischk. & Serg.]	[1, 2, 3, 7]
	<i>Tephroseris pricei</i> (N.D.Simpson) Holub [= <i>Senecio pricei</i> N.D.Simpson]	[1, 3, 6, 7, 13, 14]
SE	<i>Tephroseris sukaczewii</i> (Schischk.) Holub [= <i>Senecio sukaczewii</i> Schischk.]	[2, 4, 9]
	<i>Tephroseris turczaninovii</i> (DC.) Holub [= <i>Senecio sumneviczii</i> Schischk. & Serg.]	[1, 2, 3, 6, 7]
	<i>Tephroseris vereszczaginii</i> (Schischk. & Serg.) Holub [= <i>Senecio vereszczaginii</i> Schischk. & Serg.]	[7]
	<i>Tibetiodes flaccida</i> (Bunge) G.L.Nesom [= <i>Erigeron flaccidus</i> (Bunge) Botsch.]	[1, 2, 3, 4, 6, 7]
	<i>Tragopogon kasahstanicus</i> S.A.Nikitin	[7]
	<i>Tragopogon orientalis</i> L.	[6, 7]
	<i>Tragopogon ruber</i> S.G.Gmel.	[7, 14]
	<i>Tragopogon songoricus</i> S.A.Nikitin	[6, 7, 13, 14]
SE	<i>Tragopogon trachycarpus</i> S.A.Nikitin	[2–5, 7, 8, 13]

- Tripleurospermum ambiguum* (Ledeb.) Franch. & Sav.  
[≡ *Matricaria ambigua* (Ledeb.) Krylov] [6, 7]
- Tripolium pannonicum* (Jacq.) Dobrocz. [≡ *Tripolium pannonicum* Jacq.] [4, 9–13]
- Trommsdorffia ciliata* (Thunb.) Soják [≡ *Hypochaeris ciliata* (Thunb.) Makino] [5]
- SE *Tugarinovia mongolica* Iljin [11, 12, 13, 16]
- Turczaninovia fastigiata* (Fisch.) DC. [≡ *Aster fastigiatus* Fisch.] [2, 4]
- Vickifunkia songarica* (Fisch.) C.Ren [≡ *Ligularia songarica* (Fisch.) Y.Ling] [14]
- Vickifunkia thomsonii* (C.B.Clarke) C.Ren  
[≡ *Ligularia thomsonii* (C.B.Clarke) Pojark.] [14]
- Vickifunkia thyrsoides* (Ledeb.) C.Ren [≡ *Ligularia thyrsoides* (Ledeb.) DC.] [6, 7, 14]
- 29. Balsaminaceae** A.Rich. (1 genus and 2 species)
- Impatiens noli-tangere* L. [1, 2, 3, 4, 5, 9]
- Impatiens parviflora* DC. [7]
- 30. Berberidaceae** Juss. (1 genus and 2 species)
- Berberis amurensis* Rupr. [5]
- Berberis sibirica* Pall. [1–4, 6, 7, 9, 10, 13]
- 31. Betulaceae** Gray (2 genera and 9 taxa)
- Alnus alnobetula* subsp. *fruticosa* (Rupr.) Raus [1, 2, 4, 9]
- Betula fruticosa* Pall. [1, 2, 3, 4, 5, 6]
- Betula mandshurica* subsp. *tauschii* (Regel) Kamelin [4, 5]
- Betula microphylla* Bunge [1–4, 6–8, 10, 13, 14]
- Betula nana* subsp. *exilis* (Sukachev) Hultén [2, 3]
- Betula nana* subsp. *rotundifolia* (Spach) Malyshev [1, 2, 3, 6, 7]
- Betula ovalifolia* Rupr. [1–5, 7, 8, 13]
- Betula pendula* subsp. *mandshurica* (Regel) Ashburner & McAll. [1–5, 8, 13]
- Betula pendula* Roth subsp. *pendula* [2, 3, 4]
- 32. Biebersteiniaceae** Schnizl. (1 genus and 1 species)
- Biebersteinia odora* Stephan [6, 7]
- 33. Bignoniaceae** Juss. (1 genus and 1 species)
- SE *Incarvillea potaninii* Batalin [13, 15, 16]
- 34. Boraginaceae** Juss. (24 genera and 78 taxa)
- Note: Since Urgamal et al. (2014), several genera and species have been critically revised and updated by Ovczinnikova (2019a). Additionally, six new species of *Craniospermum* Lehm. have been described from Mongolia by Ovczinnikova and

Korolyuk (2016) and Ovchinnikova (2019b, 2020). We follow the treatment of Ovczinnikova (2019a). Furthermore, *Arnebia tibetica* previously known as a synonym of *A. guttata*, differs from *A. guttata* based on floral morphology and plastid genome characteristics discovered by Park et al. (2020).

	<i>Amblynotus rupestris</i> (Pall.) Popov [= <i>Eritrichium rupestre</i> (Georgi) Bunge]	[1–9, 13]
	<i>Anchusa arvensis</i> (L.) M.Bieb.	[7, 10, 11, 14]
SE	<i>Anoplocaryum compressum</i> Ledeb. [= <i>Echinospermum compressum</i> (Ledeb.) Turcz.]	[1, 2, 3, 6, 8]
E	<i>Anoplocaryum tenellum</i> A.L.Ebel & Rudaya [= <i>Microula tenella</i> (A.L.Ebel & Rudaya)]	[7]
SE	<i>Anoplocaryum turczaninovii</i> Krasnob.	[1, 3, 6, 7, 8, 10, 14]
	<i>Arnebia decumbens</i> Coss. & Kralik	[6, 7, 8, 13, 14]
	<i>Arnebia fimbriata</i> Maxim.	[11–13, 15, 16]
	<i>Arnebia guttata</i> Bunge	[3, 7, 10–16]
	<i>Arnebia tibetana</i> Kurz	[7]
	<i>Asperugo procumbens</i> L.	[3, 6, 7, 10, 14]
E	<i>Asperula gobicola</i> Grubov [= <i>Asperula saxicola</i> Grubov]	[13, 16]
SE	<i>Craniospermum canescens</i> DC.	[3, 7, 13, 14]
E	<i>Craniospermum desertorum</i> Ovczinnikova & A.Korolyuk	[7]
E	<i>Craniospermum gubanovii</i> Ovczinnikova	[14]
E	<i>Craniospermum kamelinii</i> Ovczinnikova	[7]
SE	<i>Craniospermum mongolicum</i> I.M.Johnst.	[7, 11–14]
E	<i>Craniospermum montanostepposum</i> Ovczinnikova	[7]
E	<i>Craniospermum pseudotuvicum</i> Ovczinnikova & A.Korolyuk	[10]
SE	<i>Craniospermum tuvinicum</i> Ovczinnikova	[6, 7]
E	<i>Craniospermum volkovae</i> Ovczinnikova	[10]
	<i>Cynoglossum divaricatum</i> Steph.	[3, 4, 8, 9, 13, 14]
SE	<i>Eritrichium alpinum</i> Ovczinnikova	[6]
	<i>Eritrichium pauciflorum</i> DC.	[1–8, 13]
	<i>Eritrichium pectinatum</i> DC.	[3]
SE	<i>Eritrichium pulviniforme</i> Popov	[3, 10, 13]
SE	<i>Eritrichium sajanense</i> (Malysch.) Sipliv.	[1]
	<i>Eritrichium thymifolium</i> (DC.) Y.S.Lian & J.Q.Wang	[3, 4, 6–15]
	<i>Eritrichium tianschanicum</i> Iljin	[6]
	<i>Eritrichium villosum</i> (Ledeb.) Bunge [= <i>Myosotis villosa</i> Ledeb.]	[2, 3, 4, 6, 7, 14]
	<i>Hackelia deflexa</i> (Wahlenb.) Opiz [= <i>Myosotis deflexa</i> Wahlenb.]	[2–5, 7, 9, 10, 13]
	<i>Heliotropium ellipticum</i> Ledeb.	[6, 7, 15]
	<i>Lappula balchaschensis</i> Popov	[7, 13, 14, 15]
	<i>Lappula brachycentroides</i> Popov	[3]
	<i>Lappula consanguinea</i> Gürke	[2–4, 6, 7, 10, 11, 13, 14]
	<i>Lappula coronifera</i> Popov	[3]

	<i>Lappula duplicitarpa</i> Pavlov	[7, 12, 14]
SE	<i>Lappula granulata</i> (Krylov) Popov	[3, 7, 9, 10, 12]
	<i>Lappula heteracantha</i> (Ledeb.) Gürke	[7]
	<i>Lappula intermedia</i> (Ledeb.) Popov	[3, 4, 6, 7, 9, 14, 15]
	<i>Lappula krylovii</i> Ovczinnikova, Pjak & A.L.Ebel	[7]
	<i>Lappula macrantha</i> (Ledeb.) Gürke	[7, 14]
	<i>Lappula microcarpa</i> Gürke	[7, 10]
	<i>Lappula myosotis</i> Wolf	[2–5, 8, 9, 13]
	<i>Lappula patula</i> Asch.	[3, 15]
	<i>Lappula redowskii</i> (Hornem.) Greene	[1–4, 8, 9, 11–13]
	<i>Lappula semiglabra</i> (Ledeb.) Gürke	[7, 11, 14, 15]
	<i>Lappula stricta</i> (Ledeb.) Gürke	[3, 7–12, 14, 15]
	<i>Lappula tadshikorum</i> Popov	[7]
	<i>Lappula tenuis</i> Gürke	[14, 15]
	<i>Lappula tianschanica</i> Popov & Zakirov	[7]
	<i>Lappula tuvinica</i> Ovczinnikova	[6]
	<i>Lindelofia stylosa</i> (Kar. & Kir.) Brand [= <i>Cynoglossum stylosum</i> Kar. & Kir.]	[7, 10, 11, 14]
SE	<i>Mertensia davurica</i> (Sims) G.Don [= <i>Mertensia ochroleuca</i> Ikonn.-Gal.]	[1, 2, 3, 4]
	<i>Mertensia pallasii</i> G.Don	[7]
	<i>Mertensia stylosa</i> DC.	[1, 2, 3]
	<i>Mertensia tarbagataica</i> B.Fedtsch.	[7]
	<i>Microula tibetica</i> var. <i>pratensis</i> (Maxim.) W.T.Wang [= <i>Tretocarya pratensis</i> Maxim.]	[3, 7]
	<i>Myosotis alpestris</i> F.W.Schmidt	[1–4, 6, 7, 9, 14]
	<i>Myosotis austrosibirica</i> O.D.Nikif.	[7, 13]
	<i>Myosotis baltica</i> Sam.	[3, 5]
	<i>Myosotis caespitosa</i> Schultz	[2–5, 9, 10, 14]
	<i>Myosotis krylovii</i> Serg.	[1–4, 6, 7, 13]
	<i>Myosotis scorpioides</i> L.	[2]
	<i>Myosotis stricta</i> Link	[7]
	<i>Nonea caspica</i> G.Don	[7, 10, 11, 14, 15]
	<i>Nonea pulla</i> DC.	[2, 4, 8, 9, 14]
	<i>Nonea rossica</i> Steven	[3]
	<i>Onosma fuyunensis</i> Y.He & Q.R.Liu	[7]
	<i>Onosma gmelinii</i> Ledeb.	[7, 14]
	<i>Onosma setosa</i> Ledeb. subsp. <i>setosa</i>	[7]
	<i>Onosma setosa</i> subsp. <i>transrhytmensis</i> (Klokov) Kamelin	[3, 7, 10]
	<i>Pseudolappula occultata</i> (Popov) Q.R.Liu & D.H.Liu [≡ <i>Lappula occultata</i> Popov]	[14]
	<i>Pulmonaria dacica</i> (Simonk.) Simonk. [= <i>Pulmonaria mollissima</i> A.Kern.]	[2, 4]
	<i>Rindera tetraspis</i> Pall.	[14]

<i>Rochelia bungei</i> Trautv.	[3, 6, 14]
<i>Rochelia leiocarpa</i> Ledeb.	[6, 14]
<i>Stenosolenium saxatile</i> (Pall.) Turcz. [= <i>Anchusa saxatilis</i> Pall.]	[3, 4, 10]
<i>Tournefortia sibirica</i> L. [= <i>Messerschmidia sibirica</i> (L.) L.]	[5, 8–13, 16]

### 35. Brassicaceae Burnett (51 genera and 138 taxa)

Note: The updated checklist and taxonomic notes of Brassicaceae was recently revised by German (2015). In this study, we followed German (2015) where the names of several species and genera changed compared to Urgamal et al (2014). Since 2015, several new records of this family have been found in the flora of Mongolia (Dorofeyev 2019; Dorofeyev and Ekhmaa 2020). For example, *Lepidium gobicum* V.I.Dorof. was newly described from Mongolia and China by Dorofeyev (2019); however, this species should be referred to *Lepidium apetalum* Willd. (German 2020).

<i>Alyssum desertorum</i> Stapf	[3, 6, 7, 8, 10]
<i>Alyssum lenense</i> Adams	[1–5, 7, 8, 9]
<i>Aphragmus involucratus</i> O.E.Schulz	[7, 13]
<i>Arabidopsis thaliana</i> (L.) Heynh.	[6, 7]
<i>Arabis borealis</i> Andrz.	[2, 3, 4, 5, 9]
<i>Barbarea orthoceras</i> Ledeb.	[2, 3, 4, 5, 9]
<i>Barbarea vulgaris</i> W.T.Aiton	[2, 3, 4, 5, 7]
<i>Braya humilis</i> (C.A.Mey.) B.L.Rob. [= <i>Neotorularia grubovii</i> (Botsch.) Botsch., <i>Neotorularia mongolica</i> Botsch. & Gubanov]	[1, 3, 4, 6, 7, 8]
<i>Braya rosea</i> Bunge	[1, 3, 6, 7]
<i>Braya siliquosa</i> Bunge	[1]
<i>Camelina microcarpa</i> Andrz.	[4, 6, 14]
<i>Capsella bursa-pastoris</i> (L.) Medik.	[2–4, 6, 7, 10, 14]
<i>Capsella orientalis</i> Klokov [= <i>Capsella bursa-pastoris</i> subsp. <i>orientalis</i> (Klokov) Tzvelev]	[7, 10]
<i>Cardamine bellidifolia</i> L.	[1, 2, 3, 7]
<i>Cardamine impatiens</i> L.	[6]
<i>Cardamine leucantha</i> (Tausch) O.E.Schulz	[5]
<i>Cardamine macrophylla</i> Willd.	[1, 2, 6, 7]
<i>Cardamine parviflora</i> L.	[2, 3]
<i>Cardamine pratensis</i> L.	[1–5, 7, 9]
<i>Cardamine prorepens</i> Fisch.	[5]
<i>Cardamine trifida</i> (Lam.) B.M.G.Jones	[5]
<i>Catolobus pendulus</i> (L.) Al-Shehbaz [= <i>Arabis pendula</i> L.]	[1–6, 8, 9, 10, 12, 13]
<i>Chorispora bungeana</i> Fisch. & C.A.Mey.	[7]
<i>Chorispora sibirica</i> (L.) DC.	[6, 7, 13, 14]
<i>Chorispora tenella</i> (Pall.) DC.	[7, 14]
<i>Clausia aprica</i> Trotzky	[1–4, 6, 7, 9]

	<i>Clausia trichosepala</i> (Turcz.) F.Dvořák	[4]
	<i>Crucihimalaya mollissima</i> (C.A.Mey.) Al-Shehbaz [= <i>Sisymbrium mollissimum</i> C.A.Mey.]	[6, 7, 9, 13, 14]
SE	<i>Crucihimalaya rupicola</i> (Krylov) A.L.Ebel & D.A.German [= <i>Arabis rupicola</i> Krylov]	[6, 7, 10, 11, 13, 14]
	<i>Dendroarabis fruticulosa</i> (C.A.Mey.) D.A.German & Al-Shehbaz [= <i>Arabis fruticulosa</i> C.A.Mey.]	[1, 7]
	<i>Descurainia sophia</i> (L.) Webb	[1–10, 12–14]
SE	<i>Dontostemon crassifolius</i> (Bunge) Maxim.	[7, 10–16]
	<i>Dontostemon dentatus</i> Ledeb.	[5]
SE	<i>Dontostemon elegans</i> Maxim.	[6, 7, 10, 11, 13–16]
E	<i>Dontostemon gubanovii</i> (D.A.German) D.A.German [= <i>Dontostemon senilis</i> subsp. <i>gubanovii</i> D.A.German]	[6, 7, 10]
	<i>Dontostemon integrifolius</i> (L.) Ledeb.	[1–13, 16]
	<i>Dontostemon micranthus</i> C.A.Mey.	[1–5, 8, 9, 13]
SE	<i>Dontostemon perennis</i> C.A.Mey.	[3, 5–8, 10–13, 15]
	<i>Dontostemon pinnatifidus</i> (Willd.) Al-Shehbaz & H.Ohba [= <i>Cheiranthus pinnatifidus</i> Willd.]	[1, 3, 4, 8, 13]
SE	<i>Dontostemon senilis</i> Maxim.	[6, 7, 8, 10–16]
	<i>Draba alpina</i> L.	[6]
	<i>Draba altaica</i> (C.A.Mey.) Bunge	[6, 7, 13]
SE	<i>Draba baicalensis</i> Tolm.	[3, 6, 7]
	<i>Draba eriopoda</i> Turcz.	[1, 2, 3, 6]
	<i>Draba fladnizensis</i> Wulfen	[1, 2, 3, 6, 7, 13]
	<i>Draba hirta</i> L.	[1, 2, 3, 6, 7, 13]
	<i>Draba kusnetzovii</i> (Turcz.) Hayek	[1, 3, 6, 7, 13]
	<i>Draba lanceolata</i> Royle	[1–4, 6, 7, 13]
	<i>Draba mongolica</i> Turcz.	[1, 3]
	<i>Draba nemorosa</i> L.	[1–10, 13]
	<i>Draba ochroleuca</i> Bunge	[1, 3, 6, 7, 13]
	<i>Draba oreades</i> Schrenk	[1, 3, 6, 7, 13]
SE	<i>Draba pygmaea</i> Turcz.	[1, 3, 6]
	<i>Draba sibirica</i> (Pall.) Thell.	[3, 7]
	<i>Draba stenocarpa</i> Hook.f. & Thomson	[7]
	<i>Draba subamplexicaulis</i> C.A.Mey.	[1–3, 6, 7, 13, 14]
	<i>Draba turczaninowii</i> Pohle	[1, 6, 7, 13]
	<i>Erysimum andrzejowskianum</i> Bess.	[7]
	<i>Erysimum cheiranthoides</i> subsp. <i>altum</i> Ahti	[without indication of regions]
	<i>Erysimum cheiranthoides</i> L. subsp. <i>cheiranthoides</i>	[1–5, 7–11, 13, 14]
	<i>Erysimum cheiranthoides</i> subsp. <i>transiliense</i> (Popov) D.A.German [= <i>Erysimum transiliense</i> Popov]	[7]
	<i>Erysimum flavum</i> (Georgi) Bobrov subsp. <i>flavum</i>	[1–5, 8, 9, 12]
	<i>Erysimum flavum</i> subsp. <i>altaicum</i> (C.A.Mey.) Polozhij	[3, 6, 7, 10]

SE	<i>Erysimum kotuchovii</i> D.A.German	[7]
	<i>Erysimum ledebourii</i> D.A.German	[7]
	<i>Erysimum marschallianum</i> Andrz.	[2–4, 6, 7, 10, 13, 14]
SE	<i>Erysimum mongolicum</i> D.A.German	[7, 14]
	<i>Erysimum sisymbrioides</i> C.A.Mey.	[6, 7, 15]
	<i>Eutrema edwardsii</i> subsp. <i>compactum</i> (O.E.Schulz) A.L.Ebel [= <i>Eutrema compactum</i> O.E.Schulz]	[7]
	<i>Eutrema edwardsii</i> R.Br. subsp. <i>edwardsii</i>	[1–3, 6, 7, 13]
	<i>Eutrema salsugineum</i> (Pall.) Al-Shehbaz & Warwick [≡ <i>Sisymbrium salsugineum</i> Pall.]	[3, 4, 6–10]
E	<i>Galitzkya macrocarpa</i> (Ikonn.-Gal.)Botsch. [= <i>Berteroa macrocarpa</i> Ikonn.-Gal.]	[13, 15]
SE	<i>Galitzkya potaninii</i> (Maxim.)Botsch.	[7, 14, 15]
SE	<i>Goldbachia ikonnikovii</i> Vassilcz.	[6, 7, 8, 10, 11, 13, 14]
	<i>Goldbachia pendula</i> Botsch.	[7, 14]
	<i>Hesperis sibirica</i> L.	[1, 2, 3, 4, 7]
	<i>Hornungia procumbens</i> Hayek	[3, 6, 7, 10, 11, 14]
	<i>Iljinsea planisiliqua</i> (Fisch. & C.A.Mey.) Al-Shehbaz [≡ <i>Conringia planisiliqua</i> Fisch. & C.A.Mey.]	[6, 14]
	<i>Isatis costata</i> C.A.Mey.	[2–4, 6–9, 11–14]
	<i>Isatis gymnocarpa</i> (Fisch.) Al-Shehbaz, Moazzeni & Mumm. [≡ <i>Tauschia gymnocarpa</i> Fisch.]	[14]
	<i>Isatis multicaulis</i> (Kar. & Kir.) Jafri	[14]
	<i>Isatis oblongata</i> DC.	[1, 3, 4, 6–9, 13]
	<i>Leiocarpaea cochlearioides</i> (Murray) D.A.German & Al-Shehbaz [≡ <i>Bunias cochlearioides</i> Murray]	[1]
	<i>Leiospora exscapa</i> (C.A.Mey.) F.Dvořák [= <i>Parrya exscapa</i> C.A.Mey.]	[1, 6, 7]
	<i>Lepidium affine</i> Ledeb. [= <i>Lepidium latifolium</i> subsp. <i>affine</i> (Ledeb.) Kitag.]	[4, 9, 14]
	<i>Lepidium amplexicaule</i> Willd.	[3, 7–11, 14, 15]
	<i>Lepidium apetalum</i> Willd.	[1–5, 7–15]
	<i>Lepidium appelianum</i> Al-Shehbaz	[7, 10, 11, 14–16.]
	<i>Lepidium cartilagineum</i> Thell.	[5–8, 10, 12, 14]
	<i>Lepidium cordatum</i> Willd.	[6–11, 13–16]
	<i>Lepidium lacerum</i> C.A.Mey. [= <i>Lepidium songaricum</i> Schrenk]	[7, 14]
	<i>Lepidium obtusum</i> Basiner	[6, 7, 10, 14, 15]
	<i>Litwinowia tenuissima</i> (Pall.) Woronow	[14]
	<i>Macropodium nivale</i> R.Br.	[1, 7]
	<i>Matthiola superba</i> Conti	[14]
	<i>Megacarpaea megalocarpa</i> Schischk.	[14]
	<i>Meniocus linifolius</i> (Willd.) DC. [= <i>Alyssum linifolium</i> Willd.]	[3, 5, 7]
SE	<i>Microstigma brachycarpum</i> Botsch.	[6, 7, 15, 16]

SE	<i>Microstigma deflexum</i> (Bunge) Juz.	[3, 6, 7, 12, 13, 15, 16]
	<i>Neotorularia brevipes</i> (Kar. & Kir.) Hedge & J.Léonard [= <i>Sisymbrium brevipes</i> F.Muell.]	[7, 14]
	<i>Noccaea ferganensis</i> (N.Busch) Czerep. [= <i>Thlaspi ferganense</i> N.Busch]	[7]
	<i>Noccaea thlaspidioides</i> (Pall.) F.K.Mey. [≡ <i>Lepidium thlaspidioides</i> Pall. = <i>Thlaspi cochleariforme</i> DC.]	[1–9, 13]
	<i>Odontarrhena obovata</i> C.A.Mey. [= <i>Alyssum obovatum</i> (C.A.Mey.) Turcz.]	[1–10]
	<i>Olimarabidopsis pumila</i> (Stephan) Al-Shehbaz [≡ <i>Sisymbrium pumilum</i> Stephan]	[14]
SE	<i>Pachyneurum grandiflorum</i> Bunge	[1, 3, 6, 7, 13]
	<i>Pugionium dolabratum</i> Maxim.	[11, 12, 13, 16]
SE	<i>Pugionium pterocarpum</i> Kom.	[10]
	<i>Rhammatophyllum erysimoides</i> (Kar. & Kir.) Al-Shehbaz & O.Appel [≡ <i>Arabis erysimoides</i> Kar. & Kir.]	[7, 14]
	<i>Rorippa barbareifolia</i> (DC.) Kitag.	[2]
	<i>Rorippa dogadovae</i> Tzvelev	[3, 11]
	<i>Rorippa palustris</i> Besser	[1–11, 13, 14]
	<i>Sisymbrium brassiciforme</i> C.A.Mey.	[7, 9, 14, 15]
	<i>Sisymbrium heteromallum</i> C.A.Mey.	[2–4, 6–8, 10–14]
	<i>Sisymbrium loeselii</i> L.	[3, 4, 14]
	<i>Sisymbrium polymorphum</i> (Murr.) Roth	[3, 4, 6–10, 14]
	<i>Sisymbrium subspinescens</i> Bunge	[14]
	<i>Smelowskia alba</i> (Pall.) B.Fedtsch.	[1, 3, 4, 6, 7, 10, 13, 14]
SE	<i>Smelowskia altaica</i> (Pobed.) Botsch.	[6, 7]
	<i>Smelowskia bifurcata</i> (Ledeb.) Botsch.	[1, 3]
	<i>Smelowskia calycina</i> (Stephan) C.A.Mey. [= <i>Lepidium calycinum</i> Steph.]	[1, 3, 6, 7, 13, 14]
SE	<i>Smelowskia calycina</i> subsp. <i>pectinata</i> (Bunge) D.A.German [= <i>Hutchinsia pectinata</i> Bunge]	[3, 7, 13, 14]
E	<i>Smelowskia mongolica</i> Kom.	[3]
SE	<i>Sterigmostemum violaceum</i> (Botsch.) H.L.Yang [= <i>Oreoloma violaceum</i> Botsch., = <i>Sterigmostemum regeliorum</i> Kamelin & D.German]	[7, 14]
	<i>Stevenia alyssoides</i> Adams & Fisch.	[1, 3]
SE	<i>Stevenia alyssoides</i> subsp. <i>zinaidae</i> (Malyshev) Kamelin [= <i>Stevenia zinaidae</i> Malyshev]	[1, 3]
	<i>Stevenia canescens</i> (DC.) D.A.German [≡ <i>Alyssum canescens</i> DC. ≡ <i>Ptilotrichum canescens</i> (DC.) C.A.Mey.]	[1–4, 6–9, 11–16]
SE	<i>Stevenia cheiranthoides</i> DC. subsp. <i>cheiranthoides</i>	[4, 5, 6, 7, 9]
	<i>Stevenia cheiranthoides</i> subsp. <i>incarnata</i> (Kamelin) D.A.German	[1–4, 6–8, 10]
SE	<i>Stevenia dahurica</i> (Peschkova) D.A.German & Al-Shehbaz [= <i>Alyssum dahuricum</i> (Peschkova) Al-Shehbaz, <i>Ptilotrichum dahuricum</i> Peschkova]	[4, 5, 8, 9]

SE	<i>Stevenia sergievkajae</i> (Krasnob.) Kamelin & Gubanov [ $\equiv$ <i>Alyssum sergievkajae</i> Krasnob.]	[3]
SE	<i>Stevenia tenuifolia</i> (Stephan) D.A.German [ $\equiv$ <i>Alyssum tenuifolium</i> Steph.]	[2–10, 12–15]
	<i>Strigosella africana</i> (L.) Botsch.	[10, 11]
	<i>Strigosella brevipes</i> (Bunge) Botsch.	[14]
	<i>Subularia aquatica</i> L.	[3, 6]
	<i>Tetracme quadricornis</i> (Steph.) Bunge	[7, 14]
	<i>Thlaspi arvense</i> L.	[1–4, 6, 7, 13, 14]
	<i>Thlaspi ceratocarpum</i> (Pall.) Murray [ $\equiv$ <i>Carpoceras ceratocarpum</i> (Pall.) N. Busch]	[6, 10, 14]
	<i>Turritis glabra</i> L.	[7]

**36. Butomaceae** Mirb. (1 genus and 2 species)

	<i>Butomus junceus</i> Turcz.	[1, 8, 9, 10, 14]
	<i>Butomus umbellatus</i> L.	[1–5, 8, 9, 14]

**37. Campanulaceae** Juss. (4 genera and 18 taxa)

E	<i>Adenophora changaica</i> Gubanov & Kamelin	[3]
	<i>Adenophora gmelinii</i> Fisch.	[4, 5, 9]
	<i>Adenophora lamarkii</i> Fisch. [ $\equiv$ <i>Campanula lamarcii</i> D.Dietr.]	[2, 3, 4, 6]
	<i>Adenophora liliifolia</i> (L.) A.DC. [ $\equiv$ <i>Campanula liliifolia</i> L.]	[2, 6, 8]
	<i>Adenophora pereskifolia</i> (Fisch.) G.Don	[4, 5, 9]
	<i>Adenophora stenanthina</i> (Ledeb.) Kitagawa [= <i>Adenophora crispata</i> Turcz.]	[1–5, 8, 9, 13]
	<i>Adenophora tricuspidata</i> A.DC.	[2, 4, 5, 9]
	<i>Adenophora triphylla</i> (Thunb.) A.DC. [ $\equiv$ <i>Campanula triphylla</i> Thunb.]	[2, 4, 9]
	<i>Campanula cervicaria</i> L.	[2]
	<i>Campanula dasyantha</i> M.Bieb.	[1, 2]
	<i>Campanula glomerata</i> L.	[1–7, 9]
	<i>Campanula punctata</i> Lam.	[9]
	<i>Campanula rotundifolia</i> L.	[6]
	<i>Campanula stevenii</i> subsp. <i>altaica</i> (Ledeb.) Fed. [ $\equiv$ <i>Campanula altaica</i> Ledeb.]	[7]
	<i>Campanula stevenii</i> subsp. <i>turczaninovii</i> (Fed.) Victorov [ $\equiv$ <i>Campanula turczaninovii</i> Fed.]	[1, 2, 3, 6, 13]
	<i>Campanula stevenii</i> subsp. <i>wolgensis</i> (P.A.Smirn.) Fed. [ $\equiv$ <i>Campanula wolgensis</i> P.A.Smirn.]	[7]
	<i>Codonopsis clematidea</i> C.B.Clarke	[7]
	<i>Platycodon grandiflorus</i> A.DC.	[5]

**38. Caprifoliaceae** Juss. (5 genera and 24 taxa)

	<i>Linnaea borealis</i> L.	[1, 7, 13, 14]
--	----------------------------	----------------

<i>Lonicera caerulea</i> subsp. <i>altaica</i> (Pall.) Gladkova [= <i>Lonicera altaica</i> Pall.]	
	[1–4, 6, 7, 13, 14]
<i>Lonicera caerulea</i> L. subsp. <i>caerulea</i>	[6]
<i>Lonicera caerulea</i> var. <i>venulosa</i> (Maxim.) Vorosch. [= <i>Lonicera venulosa</i> Maxim.]	[5]
<i>Lonicera chrysanthra</i> Turcz.	[5]
<i>Lonicera hispida</i> Pall.	[3, 6, 7, 13, 14]
<i>Lonicera microphylla</i> Willd. [= <i>Caprifolium microphyllum</i> (Willd.) Kuntze]	
	[3, 6, 7, 9, 10, 13, 14, 16]
<i>Lonicera tatarica</i> L.	[4]
<i>Patrinia heterophylla</i> Bunge	[9]
<i>Patrinia intermedia</i> Roem. & Schult.	[3, 6, 7, 14]
<i>Patrinia rupestris</i> (Pall.) Dufr. [= <i>Valeriana rupestris</i> Pall.]	[1–5, 8, 9]
<i>Patrinia scabiosifolia</i> Fisch.	[4, 5]
<i>Patrinia sibirica</i> (L.) Juss.	[1–7]
<i>Scabiosa comosa</i> Fisch.	[1–5, 8, 9]
<i>Scabiosa ochroleuca</i> L.	[3, 10]
<i>Valeriana altaica</i> Sumnev.	[1, 2]
<i>Valeriana alternifolia</i> Ledeb. [= <i>Valeriana dahurica</i> Sumnev.]	[1–6, 9]
<i>Valeriana capitata</i> Pall.	[1]
<i>Valeriana dubia</i> Bunge	[1, 3, 6, 7, 14]
<i>Valeriana martjanovii</i> Krylov [= <i>Valeriana saichanensis</i> Kom.]	[13]
<i>Valeriana officinalis</i> L.	[1, 2, 3, 4, 8, 9]
<i>Valeriana petrophila</i> Bunge	[1, 3, 6, 7, 13]
SE <i>Valeriana tangutica</i> Batalin	[16]
SE <i>Valeriana transjenisensis</i> Kreyer	[1, 3, 7]

**39. Caryophyllaceae** Juss. (20 genera and 97 taxa)

<i>Acanthophyllum pungens</i> Boiss.	[6, 7, 14]
<i>Arenaria leptoclados</i> Guss.	[7]
<i>Arenaria serpyllifolia</i> L.	[7, 14]
<i>Cerastium alpinum</i> L.	[6]
<i>Cerastium arvense</i> L.	[1–10, 13, 14]
<i>Cerastium cerastoides</i> (L.) Britton [= <i>Dichodon cerastoides</i> (L.) Rchb. ≡ <i>Stellaria cerastoides</i> L.]	[1–7, 10, 14]
<i>Cerastium davuricum</i> Fisch.	[2, 4, 7, 14]
<i>Cerastium falcatum</i> (Gren.) Bunge [= <i>Stellaria falcata</i> Ser.]	[14]
<i>Cerastium holosteoides</i> Fr. [= <i>Cerastium fontanum</i> f. <i>holosteoides</i> (Fr.) M.B.Wyse Jacks.]	[2]
<i>Cerastium lithospermifolium</i> Fisch.	[1, 3, 6, 7, 10, 13]
<i>Cerastium maximum</i> L.	[13]
<i>Cerastium pauciflorum</i> Steven	[1, 2, 3, 6, 7]
<i>Cerastium pusillum</i> Ser.	[1, 2, 3, 6, 7, 14]

	<i>Cherleria arctica</i> (Steven) A.J.Moore & Dillenb.	
	[≡ <i>Minuartia arctica</i> (Steven) Graebn.]	[1, 2, 3, 4, 6, 7]
	<i>Cherleria biflora</i> (L.) A.J.Moore & Dillenb.	
	[≡ <i>Minuartia biflora</i> (L.) Schinz & Thell.]	[1, 2, 3, 6, 7]
	<i>Dianthus chinensis</i> L. [= <i>Dianthus versicolor</i> Fisch.]	[1–11, 13]
	<i>Dianthus crinitus</i> subsp. <i>soongoricus</i> (Schischk.) Kozhevnikov	
	[≡ <i>Dianthus soongoricus</i> Schischk.]	[7, 14]
	<i>Dianthus ramosissimus</i> Pall.	[10]
	<i>Dianthus repens</i> Willd. [= <i>Dianthus chinensis</i> subsp. <i>repens</i> (Willd.) Vorosch.]	
		[6, 7]
	<i>Dianthus superbus</i> L.	[1–10, 13]
	<i>Eremogone androsacea</i> (Grubov) Ikonn. [≡ <i>Arenaria androsacea</i> Grubov]	[13]
SE	<i>Eremogone asiatica</i> (Schischk.) Ikonn. [≡ <i>Arenaria asiatica</i> Schischk.]	[7]
SE	<i>Eremogone capillaris</i> (Poir.) Fenzl [= <i>Arenaria capillaris</i> Poir.]	[1–10, 12, 13]
	<i>Eremogone juncea</i> (M.Bieb.) Fenzl [≡ <i>Arenaria juncea</i> M.Bieb.]	[4, 5, 9]
	<i>Eremogone meyeri</i> (Fenzl) Ikonn. [≡ <i>Arenaria meyeri</i> Fenzl]	
		[2–4, 6, 7, 9, 10, 12, 13]
	<i>Eremogone mongolica</i> (Schischk.) Ikonn. [≡ <i>Arenaria mongolica</i> Schischk.]	[7]
	<i>Gymnocarpos przewalskii</i> Maxim. [= <i>Paronychia przewalskii</i> (Bunge) Rohweder & Urmi-König]	[12, 14, 16]
	<i>Gypsophila altissima</i> L.	[7]
	<i>Gypsophila capituliflora</i> Rupr.	[7, 13, 14, 15]
	<i>Gypsophila cephalotes</i> (Schrenk) F.N.Williams	[6, 7]
	<i>Gypsophila davurica</i> Fenzl	
	[≡ <i>Gypsophila patrinii</i> subsp. <i>davurica</i> (Fenzl) Kozhevnikov.]	[2–5, 8, 9, 13]
	<i>Gypsophila paniculata</i> L.	[3, 4, 7, 10]
	<i>Gypsophila patrinii</i> Ser.	[1, 3, 4, 6–8, 10, 11]
	<i>Gypsophila perfoliata</i> L.	[10]
	<i>Gypsophila sericea</i> (Ser.) Krylov [≡ <i>Arenaria sericea</i> Ser.]	[7]
	<i>Herniaria caucasica</i> Rupr.	[7]
	<i>Herniaria glabra</i> L.	[7]
	<i>Heterochroa desertorum</i> Bunge [≡ <i>Gypsophila desertorum</i> Fenzl]	
		[1–4, 6–13, 16]
	<i>Lepyrodiclis holosteoides</i> (C.A.Mey.) Fenzl [= <i>Gouffeia holosteoides</i> C.A.Mey.]	
		[3, 10]
	<i>Moehringia lateriflora</i> (L.) Fenzl [≡ <i>Arenaria lateriflora</i> L.]	[1–5, 7, 9, 13]
	<i>Moehringia umbrosa</i> (Bunge) Fenzl [≡ <i>Arenaria umbrosa</i> Bunge]	[1, 2, 6, 7]
	<i>Pseudocherleria laricina</i> (L.) Dillenb. & Kadereit	
	[≡ <i>Minuartia laricina</i> Mattf.]	[4, 5]
	<i>Pseudostellaria rupestris</i> (Turcz.) Pax	[1–4, 7, 13]

	<i>Sabulina regeliana</i> (Trautv.) Dillenb. & Kadereit	
	[= <i>Minuartia regeliana</i> (Trautv.) Mattf.]	[3]
	<i>Sabulina stricta</i> (Sw.) Rchb. [= <i>Minuartia stricta</i> (Sw.) Hiern]	[1, 2, 3]
	<i>Sabulina verna</i> Rchb. [= <i>Minuartia verna</i> (L.) Hiern]	[1–3, 6, 7, 14]
	<i>Sagina saginoides</i> (L.) H.Karst.	[7]
	<i>Saponaria floribunda</i> (Kar. & Kir.) Boiss.	
	[≡ <i>Psammophiliella floribunda</i> (Kar. & Kir.) Ilkonn.]	[14]
	<i>Silene alexandrae</i> B.Keller	[14]
	<i>Silene altaica</i> Pers.	[7, 13, 14]
	<i>Silene aprica</i> Turcz. [= <i>Ussuria aprica</i> (Turcz.) Tzvelev]	[1–5, 7–10, 12, 13]
	<i>Silene borysthenica</i> (Gruner) Walters	[3, 10]
	<i>Silene bungei</i> Bocquet	[1, 2, 3, 6]
	<i>Silene chamarensis</i> Turcz. [= <i>Silene tenuis</i> subsp. <i>chamarensis</i> (Turcz.) Kozhevnikov]	[1–3, 6, 7, 9, 10, 12, 13]
	<i>Silene conoidea</i> L.	[7]
	<i>Silene foliosa</i> Maxim.	[4, 12, 13]
	<i>Silene graminifolia</i> Otth [= <i>Silene sobolevskajae</i> Czerep.]	[2, 6, 7, 10, 14]
	<i>Silene gubanovii</i> Lazkov	[6, 7, 13, 14]
SE	<i>Silene intramongolica</i> Lazkov	[7, 14]
	<i>Silene jeniseensis</i> Willd. [= <i>Silene iche-bogdo</i> Grubov]	[1–6, 8, 9, 13]
	<i>Silene latifolia</i> subsp. <i>alba</i> (Mill.) Greuter & Burdet [= <i>Lychnis alba</i> Mill.]	[7]
E	<i>Silene mongolica</i> Maxim.	[10, 13]
	<i>Silene quadriloba</i> Turcz.	[2, 3, 7, 10, 14]
	<i>Silene repens</i> Patrin	[1–10, 12–14]
	<i>Silene samojedorum</i> (Sambuk) Oxelman	
	[≡ <i>Lychnis sibirica</i> subsp. <i>samojedorum</i> Sambuk.]	[1, 2, 3, 4, 5, 9]
	<i>Silene sibirica</i> Pers.	[14]
	<i>Silene songarica</i> (Fisch., C.A.Mey. & Avé-Lall.) Bocquet [= <i>Gastrolychnis brachypetala</i> (Hornem.) Tolm. & Kozhanczikov]	[1–7, 9, 12, 13]
	<i>Silene suaveolens</i> Kar. & Kir. [= <i>Carpophora suaveolens</i> (Kar. & Kir.) Tzvelev ≡ <i>Melandrium suaveolens</i> (Kar. & Kir.) Schischk.]	[7, 10, 14]
	<i>Silene uralensis</i> (Rupr.) Bocquet	[1–3, 6, 7, 10, 13, 14]
	<i>Silene violascens</i> (Tolm.) V.V.Petrovsky & Elven	
	[≡ <i>Gastrolychnis violascens</i> Tolm.]	[7]
	<i>Silene viscosa</i> Schleich.	[3, 7, 10, 14]
	<i>Silene vulgaris</i> (Moench) Garcke [= <i>Behen vulgaris</i> Moench]	[2, 4, 6, 7]
	<i>Spergularia marina</i> (L.) Besser	
	[≡ <i>Arenaria rubra</i> var. <i>marina</i> L.]	[4, 5, 7, 10–15]
	<i>Spergularia segetalis</i> G.Don	[14]
	<i>Stellaria alsinoides</i> Boiss. & Buhse	[7, 14]
	<i>Stellaria amblyosepala</i> Schrenk	[7, 10–16]
	<i>Stellaria brachypetala</i> Bunge [= <i>Stellaria alatavica</i> Popov,	
	<i>Stellaria brachypetala</i> var. <i>alatavica</i> (Popov) Kozhevnikov]	[1, 3–7, 9, 11, 13, 14]

<i>Stellaria bungeana</i> Fenzl [= <i>Hylebia bungeana</i> (Fenzl) Tzvelev]	[2, 3, 6, 7, 9]
<i>Stellaria cherleriae</i> (Fisch.) F.N.Williams [= <i>Arenaria cherleriae</i> Fisch.]	[1–9, 13]
<i>Stellaria crassifolia</i> Ehrh.	[1–11, 14]
<i>Stellaria davurica</i> Willd.	[1–4, 7, 14]
<i>Stellaria depressa</i> Schmid	[7]
<i>Stellaria dichotoma</i> L.	[1–14]
<i>Stellaria dichotoma</i> var. <i>lanceolata</i> Bunge [= <i>Stellaria gypsophilooides</i> Fenzl]	[3, 7–9, 11–13, 15, 16]
<i>Stellaria discolor</i> Turcz.	[4, 5, 9]
<i>Stellaria filicaulis</i> Makino	[2, 3]
<i>Stellaria imbricata</i> Bunge	[6, 7, 14]
<i>Stellaria irrigua</i> Bunge	[1–4, 6, 7, 13]
<i>Stellaria longifolia</i> Muhl.	[1, 2, 3, 4, 5, 9]
<i>Stellaria longipes</i> Goldie [= <i>Stellularia longipes</i> (Goldie) MacMill.]	[1 – 3 , 5–7, 9]
<i>Stellaria martjanovii</i> Krylov [= <i>Mesostemma martjanovii</i> (Krylov) Ikonn.]	[7]
<i>Stellaria media</i> (L.) Vill. [= <i>Alsine media</i> L.]	[2, 3]
<i>Stellaria palustris</i> Ehrh.	[2, 3, 7, 9]
<i>Stellaria petraea</i> Bunge	[1–4, 6, 7, 13]
<i>Stellaria pulvinata</i> Grubov	[6, 7]
<i>Stellaria radians</i> L.	[5, 9]
<i>Stellaria zolotuchinii</i> A.L.Ebel [= <i>Stellaria glandulifera</i> N.Zolot. nom. illegit.]	[3, 10]

**40. Celastraceae** R.Br. (2 genera and 3 species)

<i>Euonymus maackii</i> Rupr.	[5, 9]
<i>Parnassia laxmannii</i> Pall.	[1–4, 6, 10]
<i>Parnassia palustris</i> L.	[1–11, 14]

**41. Ceratophyllaceae** Gray (1 genus and 2 taxa)

<i>Ceratophyllum demersum</i> L.	[1, 4, 8–10, 14]
<i>Ceratophyllum platyacanthum</i> subsp. <i>oryzetorum</i> (Kom.) Les	[10]

**42. Cleomaceae** Bercht. & J.Presl (1 genus and 1 species)

E <i>Cleome gobica</i> Grubov	[15]
-------------------------------	------

**43. Convolvulaceae** Juss. (4 genera and 15 species)

<i>Calystegia hederacea</i> Wall.	[12, 13]
<i>Calystegia pellita</i> G.Don [= <i>Calystegia dahurica</i> Herb.]	[1, 3, 4]
<i>Calystegia sepium</i> (L.) R.Br. [= <i>Convolvulus sepium</i> L.]	[14]
<i>Calystegia subvolubilis</i> G.Don	[2, 3]

<i>Convolvulus ammannii</i> Desr.	[2, 3, 4, 6–14, 16]
<i>Convolvulus arvensis</i> L.	[2, 3, 4, 7–16]
<i>Convolvulus fruticosus</i> Pall.	[7, 10–16]
<i>Convolvulus gortschakovii</i> Schrenk	[7, 8, 10, 11, 13–16]
<i>Convolvulus tragacanthoides</i> Turcz.	[12, 16]
<i>Cuscuta australis</i> R.Br.	[9]
<i>Cuscuta chinensis</i> Lam.	[4, 9, 12, 13, 15, 16]
<i>Cuscuta europaea</i> L.	[3–7, 9, 13, 14]
<i>Cuscuta lupuliformis</i> Krock.	[6, 7, 10, 14]
<i>Cuscuta monogyna</i> Vahl [ $\equiv$ <i>Monogynella monogyna</i> (Vahl) Hadač]	[4, 5, 8, 14]
<i>Merremia sibirica</i> (L.) Hallier f.	[3]

**44. Cornaceae** Bercht. & J.Presl (1 genus and 1 species)

<i>Cornus alba</i> L.	[1, 2, 3, 4, 5, 9]
-----------------------	--------------------

**45. Crassulaceae** J.St.-Hil. (6 genera and 17 taxa)

<i>Crassula aquatica</i> (L.) Schönland	[2, 4, 6, 10]
<i>Hylotelephium ewersii</i> (Ledeb.) H.Ohba [ $\equiv$ <i>Sedum ewersii</i> Ledeb.]	[6, 7, 14]
<i>Hylotelephium pallescens</i> (Freyn) H.Ohba	[2, 3, 4, 5]
<i>Hylotelephium telephium</i> (L.) H.Ohba [ $\equiv$ <i>Sedum telephium</i> L.]	[1–10]
<i>Orostachys fimbriata</i> (Turcz.) A.Berger [ $\equiv$ <i>Cotyledon fimbriata</i> Turcz.]	[2–6, 8–13]
<i>Orostachys malacophylla</i> (Pall.) Fisch. [ $\equiv$ <i>Cotyledon malacophylla</i> Pall.]	[1–5, 8, 9]
<i>Orostachys spinosa</i> (L.) Sweet [ $\equiv$ <i>Cotyledon spinosa</i> L.]	[1–4, 6–15]
<i>Orostachys thrysiflora</i> Fisch.	[1, 3–11, 13–15]
<i>Phedimus aizoon</i> (L.) 't Hart [ $\equiv$ <i>Sedum aizoon</i> L.]	[1–14]
<i>Phedimus hybridus</i> (L.) 't Hart [ $\equiv$ <i>Sedum hybridium</i> L.]	[1, 3–9, 12]
<i>Pseudosedum lievenii</i> A.Berger	[7, 14]
<i>Rhodiola algida</i> (Ledeb.) Fisch. & C.A.Mey.	[6, 7]
<i>Rhodiola coccinea</i> (Royle) Boriss.	[7]
<i>Rhodiola litwinowii</i> Boriss.	[3, 6, 7, 10]
<i>Rhodiola quadrifida</i> (Pall.) Fisch. & C.A.Mey.	[1–3, 6, 7, 13]
<i>Rhodiola rosea</i> L. [ $\equiv$ <i>Sedum roseum</i> (L.) Scop.]	[1–8, 13, 14]
<i>Rhodiola stephani</i> (Cham.) Trautv. & C.A.Mey. [= <i>Rhodiola krylovii</i> Polozhij & Revjakina = <i>Rhodiola pinnatifida</i> Boriss. = <i>Rhodiola subpinnata</i> (Krasnob.) Krasnob.]	[1, 2, 6, 7, 10]

**46. Cynomoriaceae** Endl. (1 genus and 1 species)

<i>Cynomorium songaricum</i> Rupr. [ $\equiv$ <i>Cynomorium coccineum</i> subsp. <i>songaricum</i> (Rupr.) J.Léonard]	[10–16]
--	---------

#### 47. Cyperaceae Juss. (10 genera and 131 taxa)

Note: To date, eight species of *Kobresia* Willd. have been recorded in Mongolia (Nyambayar 2009b; Urgamal et al. 2014), and they were recently transferred to the genus *Carex* L. (Global Carex Group 2015).

<i>Blysmus compressus</i> L. subsp. <i>brevifolius</i> (Decne.) Kukkonen	[2–5, 7–9, 11, 13, 14]
<i>Blysmus rufus</i> Link	[1, 3–15]
<i>Bolboschoenus maritimus</i> (L.) Palla [= <i>Scirpus maritimus</i> L.]	[14, 16]
<i>Bolboschoenus maritimus</i> subsp. <i>affinis</i> (Roth) T.Koyama [= <i>Bolboschoenus popovii</i> T.V.Egorova]	[10–16]
<i>Bolboschoenus planiculmis</i> (F.Schmidt) T.V.Egorova	[4, 5, 8–12, 15]
<i>Carex accrescens</i> Ohwi [= <i>Carex pallida</i> C.A.Mey.]	[2]
<i>Carex acuta</i> L.	[2, 3, 14]
<i>Carex alatauensis</i> S.R.Zhang [= <i>Kobresia humilis</i> (C.A.Mey.) Serg.]	[3, 6, 7, 11, 13]
<i>Carex alba</i> Scop.	[1, 2]
<i>Carex altaica</i> (Gorodkov) V.I.Krecz. [= <i>Carex orbicularis</i> subsp. <i>altaica</i> (Gorodkov) T.V.Egorova]	[3]
<i>Carex amgunensis</i> F.Schmidt [= <i>Carex chloroleuca</i> Meinsh.]	[1, 2, 3, 5, 7]
<i>Carex appendiculata</i> Kük.	[2, 4, 5, 8–10]
<i>Carex argunensis</i> Turcz. [= <i>Carex rupestris</i> subsp. <i>argunensis</i> (Turcz.) Vorosch.]	[2, 4, 5, 7, 9]
<i>Carex arnellii</i> Christ	[2, 3, 4, 5, 9]
<i>Carex aterrima</i> Hoppe	[1, 2, 3, 6, 7]
<i>Carex atherodes</i> Spreng.	[1–5, 7, 8, 9, 11]
<i>Carex atrofusca</i> Schkuhr	[1, 3, 7]
<i>Carex bigelowii</i> subsp. <i>ensifolia</i> (Gorodkov) Holub	[1, 2, 3, 6, 7]
<i>Carex bigelowii</i> subsp. <i>rigidiooides</i> (Gorodkov) T.V.Egorova	[2, 3]
<i>Carex bistaminata</i> (W.Z.Di & M.J.Zhong) S.R.Zhang [= <i>Kobresia bistaminata</i> W.Z.Di & M.J.Zhong = <i>Kobresia myosuroides</i> (Vill.) Fiori]	[1–3, 5–7, 10, 13, 14]
<i>Carex bohemica</i> Schreb.	[4]
<i>Carex borealis</i> (S.R.Zhang) Boeck. [= <i>Kobresia smirnovii</i> Ivanova]	[1, 2, 3, 6, 7, 14]
<i>Carex brunnescens</i> (Pers.) Poir. [= <i>Carex curta</i> var. <i>brunnescens</i> Pers.]	[2, 9]
<i>Carex canescens</i> L.	[2, 4, 7]
<i>Carex capillifolia</i> (Decne.) S.R.Zhang [= <i>Kobresia capilliformis</i> Ivanova]	[3, 7]
<i>Carex capitata</i> L.	[1, 2, 3, 7]
<i>Carex capricornis</i> Meinsh.	[10]
<i>Carex caryophyllea</i> Latourr.	[1–4, 6, 7, 8, 13]
<i>Carex cespitosa</i> L.	[1–10, 15]
<i>Carex chordorrhiza</i> L.f.	[2]
<i>Carex coriophora</i> Fisch. & C.A.Mey.	[1–5, 8, 9]

<i>Carex curaica</i> Kunth	[1–3, 6, 7, 10, 11, 14]
<i>Carex dahurica</i> Kük.	[1, 2]
<i>Carex delicata</i> C.B.Clarke	[1–10, 13]
<i>Carex diandra</i> Schrank	[2, 5, 9, 10]
<i>Carex diluta</i> M.Bieb.	[10]
<i>Carex distans</i> subsp. <i>aspratilis</i> (V.I.Krecz.) T.V.Egorova	[4, 10]
<i>Carex duriuscula</i> C.A.Mey.	[1–14, 16]
<i>Carex eleusinoides</i> Turcz.	[2, 3]
<i>Carex enervis</i> C.A.Mey.	[1–13, 15]
<i>Carex eremopyroides</i> V.I.Krecz.	[9, 11]
<i>Carex ericetorum</i> Pollich	[4]
<i>Carex globularis</i> L.	[1, 2, 7]
<i>Carex gotoi</i> Ohwi [= <i>Carex songorica</i> subsp. <i>gotoi</i> (Ohwi) Popov]	[4, 5, 9]
<i>Carex hancockiana</i> Maxim.	[2, 3, 5]
<i>Carex heterolepis</i> Bunge	[5]
SE <i>Carex iljinii</i> V.I.Krecz.	[1, 2, 3]
<i>Carex karoi</i> Freyn [= <i>Carex selengensis</i> N.A.Ivanova]	[2, 3, 4, 5, 9, 10]
<i>Carex korshinskii</i> Kom.	[1–6, 8–10, 12, 13]
<i>Carex lachenalii</i> Schkuhr	[1, 7, 14]
<i>Carex lanceolata</i> Boott	[2]
<i>Carex lasiocarpa</i> Ehrh.	[2]
<i>Carex laxa</i> Wahlenb.	[1, 2]
<i>Carex ledebouriana</i> C.A.Mey. [= <i>Carex capillaris</i> subsp. <i>ledebouriana</i> (C.A.Mey.) Vorosch.]	[1, 2, 3, 6, 7]
<i>Carex leporina</i> L.	[2, 3, 7]
<i>Carex limosa</i> L.	[2]
<i>Carex lithophila</i> Turcz. [= <i>Carex disticha</i> subsp. <i>lithophila</i> (Turcz.) D.Hämet-Ahti]	[2–5, 7, 9, 10]
<i>Carex loliacea</i> L.	[2, 5]
<i>Carex macrogyna</i> Turcz.	[1, 3, 6, 7, 13]
<i>Carex macrophylla</i> (Y.C.Yang) S.R.Zhang [= <i>Kobresia filifolia</i> (Turcz.) C.B.Clarke ≡ <i>Kobresia filifolia</i> (Turcz.) C.B.Clarke var. <i>macrophylla</i> Y.C.Yang]	[1–4, 6–9, 13]
<i>Carex magellanica</i> Lam. subsp. <i>irrigua</i> (Wahlenb.) Hiiitonen [= <i>Carex limosa</i> L. var. <i>irrigua</i> Wahlenb.]	[2, 3]
<i>Carex media</i> R.Br.	[1–4, 6, 7, 10]
<i>Carex melanantha</i> C.A.Mey.	[1–3, 6, 7, 9, 13]
<i>Carex melanocephala</i> Turcz.	[1, 3, 7]
<i>Carex meyeriana</i> Kunth	[1, 2, 3, 5]
<i>Carex microglochin</i> Wahlenb.	[1–4, 6–8, 11, 14]
<i>Carex nigra</i> subsp. <i>junccea</i> (Fr.) Soó [= <i>Carex juncella</i> T.M.Fries]	[1, 3]
<i>Carex norvegica</i> Retz.	[1, 2, 3, 4, 6, 7]
<i>Carex obtusata</i> Lilj.	[1–3, 6, 7, 9, 13]

	<i>Carex orbicularis</i> Boott	[1–8, 12–15]
	<i>Carex pamirica</i> (O.Fedtsch.) B.Fedtsch. subsp. <i>dichroa</i> (Freyn) T.V.Egorova [≡ <i>Carex pulla</i> Gooden. subsp. <i>dichroa</i> Freyn]	[1–4, 6, 7, 10]
	<i>Carex parallela</i> Laest. subsp. <i>redowskiana</i> (C.A.Mey.) T.V.Egorova [≡ <i>Carex redowskiana</i> C.A.Mey.]	[1, 7]
	<i>Carex parva</i> Nees	[3]
	<i>Carex pediformis</i> var. <i>macroura</i> (Meinsh.) Kük. [≡ <i>Carex macroura</i> Meinsh.]	[1, 2, 3, 5, 7]
	<i>Carex pediformis</i> subsp. <i>pediformis</i>	[1–9, 13, 14]
	<i>Carex praecox</i> Schreb.	[2, 4]
	<i>Carex pseudofoetida</i> Kük.	[7, 13]
	<i>Carex pycnostachya</i> Kar. & Kir. [≡ <i>Carex curaica</i> subsp. <i>pycnostachya</i> (Kar. & Kir.) T.V.Egorova ]	[3, 11, 14]
SE	<i>Carex raddei</i> Kük.	[2]
	<i>Carex relaxa</i> V.I.Krecz.	[3, 4, 8, 9]
	<i>Carex reptabunda</i> (Trautv.) V.I.Krecz.	[1, 3–5, 8, 9, 11, 12, 16]
	<i>Carex rhynchophysa</i> Fisch.	[2–5, 9, 10]
	<i>Carex rostrata</i> Stokes	[1–5, 8, 9, 10]
	<i>Carex rupestris</i> All.	[1, 2, 3, 6, 7, 13]
	<i>Carex sabulosa</i> Turcz.	[1–4, 7, 8, 10]
	<i>Carex sabyensis</i> Less.	[2, 3]
	<i>Carex sajanensis</i> V.I.Krecz.	[1–4, 6–9, 11, 12]
	<i>Carex sargentiana</i> (Hemsl.) S.R.Zhang [≡ <i>Kobresia sargentiana</i> Hemsl. = <i>Kobresia robusta</i> Maxim.]	[3]
	<i>Carex saxatilis</i> L. [= <i>Carex saxatilis</i> subsp. <i>laxa</i> (Trautv.) Kalela]	[1, 3]
	<i>Carex schmidtii</i> Meinsh.	[2, 3, 4, 5, 8, 9]
	<i>Carex sedakowii</i> C.A.Mey.	[1, 2, 4]
	<i>Carex simpliciuscula</i> Wahlenb. [= <i>Kobresia simpliciuscula</i> subsp. <i>subholarctica</i> T.V.Egorova]	[1, 2, 3, 7, 12, 14]
	<i>Carex songorica</i> Kar. & Kir.	[4, 5, 9, 10, 14]
	<i>Carex sordida</i> Van Heurck & Müll.Arg.	[1, 2, 3, 6, 7, 8, 13, 14]
	<i>Carex stenophylla</i> subsp. <i>stenophylloides</i> (V.I.Krecz.) T.V.Egorova	[3, 7–16]
	<i>Carex supermascula</i> V.I.Krecz.	[1, 2, 4, 5, 9, 13]
	<i>Carex tenuiflora</i> Wahlenb.	[1, 2, 3]
	<i>Carex tomentosa</i> L.	[2, 4, 8]
	<i>Carex tristis</i> subsp. <i>stenocarpa</i> (Turcz.) T.V.Egorova	[1–3, 6–8, 13, 14]
	<i>Carex vaginata</i> var. <i>petersii</i> (C.A.Mey.) Akiyama [= <i>Carex falcata</i> Turcz.]	[2, 3]
	<i>Carex vaginata</i> Tausch var. <i>vaginata</i>	[1, 2, 3]
	<i>Carex vesicata</i> Meinsh.	[1–6, 8, 9, 14]
	<i>Carex williamsii</i> Britton	[2]
	<i>Carex yamatsutana</i> Ohwi [= <i>Carex diplasiocarpa</i> V.I.Krecz.]	[2, 5, 9]
	<i>Cyperus fuscus</i> L.	[8, 9, 10]
	<i>Cyperus hamulosus</i> M.Bieb. [≡ <i>Mariscus hamulosus</i> (M.Bieb.) S.S.Hooper]	[10]

<i>Cyperus michelianus</i> (L.) Delile [= <i>Scirpus michelianus</i> L.]	[4]
<i>Cyperus pannonicus</i> Jacq. [= <i>Juncellus pannonicus</i> (Jacq.) C.B.Clarke]	[10, 11, 16]
<i>Eleocharis acicularis</i> (L.) Roem. & Schult.	[2–4, 6–9, 11, 14]
<i>Eleocharis mamillata</i> (H.Lindb.) H.Lindb.	[10]
<i>Eleocharis mitracarpa</i> Steud.	[7, 14, 15]
<i>Eleocharis palustris</i> (L.) Roem. & Schult. [= <i>Scirpus palustris</i> L.]	[1–16]
<i>Eleocharis quinqueflora</i> (Hartmann) O.Schwarz [= <i>Scirpus quinqueflorus</i> Hartmann]	[2–4, 7, 8, 10, 11, 15]
<i>Eleocharis uniglumis</i> Schult. [= <i>Eleocharis klingei</i> (Meinsh.) B.Fedtsch.]	[1, 3–5, 8–10, 12–15]
<i>Eleocharis yokoscensis</i> (Franch. & Sav.) Tang & F.T.Wang [= <i>Eleocharis acicularis</i> subsp. <i>yokoscensis</i> (Franch. & Sav.) T.V.Egorova]	[2–5, 8, 9, 14]
<i>Eriophorum altaicum</i> Meinsh.	[1, 2, 3, 6, 7]
<i>Eriophorum angustifolium</i> Honck.	[1–7, 9, 10, 11]
<i>Eriophorum angustifolium</i> subsp. <i>komarovii</i> (V.N.Vassil.) M.S.Novos.	[1–10]
<i>Eriophorum brachyantherum</i> Trautv. & C.A.Mey.	[1–4, 6, 7, 10]
<i>Eriophorum callitrix</i> C.A.Mey.	[1]
<i>Eriophorum chamissonis</i> C.A.Mey. [= <i>Eriophorum mandshuricum</i> Meinsh.]	[1, 3, 4, 5, 7]
<i>Eriophorum gracile</i> W.D.J.Koch [= <i>Eriophorum gracile</i> subsp. <i>asiaticum</i> (V.N.Vassil.) M.S.Novos.]	[3, 4]
<i>Eriophorum humile</i> Turcz.	[1, 2, 3, 6, 7]
<i>Eriophorum vaginatum</i> L.	[3]
<i>Schoenoplectiella supina</i> (L.) Lye [= <i>Schoenoplectus supinus</i> (L.) Pall.]	[10]
<i>Schoenoplectus lacustris</i> subsp. <i>hippolytii</i> (V.I.Krecz.) Kukkonen [= <i>Scirpus hippolyti</i> V.I.Krecz.]	[1–10, 12–15]
<i>Schoenoplectus tabernaemontani</i> (C.C.Gmel.) Pall.	[2, 3, 14]
<i>Schoenoplectus triquetus</i> (L.) Palla [= <i>Scirpus triquetus</i> L.]	[9]
<i>Scirpus orientalis</i> Ohwi [= <i>Scirpus sylvaticus</i> subsp. <i>orientalis</i> (Ohwi) Vorosch.]	[2, 3, 4, 5, 9]
<i>Scirpus radicans</i> Schkuhr	[2, 3, 4, 5, 9]
<i>Trichophorum pumilum</i> (Vahl) Schinz & Thell.	[1, 3, 4, 8, 9, 10, 14]

#### 48. **Droseraceae** Salisb. (2 genera and 3 species)

<i>Aldrovanda vesiculosa</i> L.	[10]
<i>Drosera anglica</i> Huds.	[2]
<i>Drosera rotundifolia</i> L.	[2]

#### 49. **Elaeagnaceae** Juss. (2 genera and 3 taxa)

<i>Elaeagnus angustifolia</i> L.	[15, 16]
----------------------------------	----------

<i>Hippophae rhamnoides</i> St.-Lag. subsp. <i>mongolica</i> Rousi [≡ <i>Hippophae mongolica</i> (Rousi) Tzvelev]	[3, 4, 6, 7, 10, 11, 13]
<i>Hippophae rhamnoides</i> St.-Lag. subsp. <i>turkestanica</i> Rousi [≡ <i>Hippophae turkestanica</i> (Rousi) Tzvelev]	[14]

## 50. Ericaceae Durande (12 genera and 27 taxa)

<i>Arctostaphylos uva-ursi</i> (L.) Spreng.	[2]
<i>Arctous alpina</i> (L.) Nied. [= <i>Arbutus alpina</i> L.]	[1, 3, 6, 10]
<i>Cassiope ericoides</i> D.Don	[4]
<i>Chamaedaphne calyculata</i> (L.) Moench	[1–3, 6, 7, 13]
<i>Empetrum nigrum</i> L. subsp. <i>nigrum</i>	[1, 2, 3, 4]
<i>Empetrum nigrum</i> subsp. <i>sibiricum</i> (V.N.Vassil.) Kuvaev [≡ <i>Empetrum sibiricum</i> V.N.Vassil.]	[1, 2, 3, 4, 6, 7]
<i>Moneses uniflora</i> A.Gray	[1, 2, 4, 7]
<i>Monotropa hypopitys</i> L.	[1, 3, 4]
<i>Orthilia obtusata</i> (Turcz.) H.Hara [≡ <i>Pyrola secunda</i> var. <i>obtusata</i> Turcz.]	[1, 2, 3, 4, 6, 7]
<i>Orthilia secunda</i> (L.) House [≡ <i>Pyrola secunda</i> L.]	[1, 2, 3, 4, 7]
<i>Phyllodoce caerulea</i> (L.) Bab. [= <i>Andromeda caerulea</i> L.]	[1]
<i>Pyrola asarifolia</i> subsp. <i>incarnata</i> (DC.) A.E.Murray [≡ <i>Pyrola rotundifolia</i> var. <i>incarnata</i> (DC.) A.P.Khokhr.]	[1–7, 9]
<i>Pyrola chlorantha</i> Sw.	[1, 4]
<i>Pyrola daurica</i> Kom.	[1, 4, 5, 7, 9]
<i>Pyrola media</i> Sw.	[3, 4]
<i>Pyrola minor</i> L.	[2]
<i>Pyrola rotundifolia</i> L.	[1, 2, 3, 4, 7]
<i>Rhododendron adamsii</i> Rehder	[1, 3]
<i>Rhododendron aureum</i> Georgi	[1, 2]
<i>Rhododendron dauricum</i> L.	[1, 2, 3, 4, 5]
<i>Rhododendron lapponicum</i> (L.) Wahlenb.	[1, 2, 3]
<i>Rhododendron ledebourii</i> Pojark. [≡ <i>Rhododendron dauricum</i> subsp. <i>ledebourii</i> (Pojark.) Alexandrova & P.A.Schmidt]	[1, 3]
<i>Rhododendron tomentosum</i> Harmaja [= <i>Ledum palustre</i> L., = <i>Ledum palustre</i> var. <i>decumbens</i> Aiton]	[1, 2, 3, 4]
<i>Vaccinium microcarpum</i> (Turcz.) Schmalh. [≡ <i>Oxycoccus microcarpus</i> Turcz.]	[1, 2]
<i>Vaccinium myrtillus</i> L. [≡ <i>Vitis-idaea myrtillus</i> (L.) Moench]	[1, 2]
<i>Vaccinium uliginosum</i> L.	[1, 2, 3, 4, 6]
<i>Vaccinium vitis-idaea</i> L. [≡ <i>Vitis-idaea vitis-idaea</i> (L.) Britton]	[1, 2, 3, 4, 5, 6]

## 51. Euphorbiaceae Juss. (1 genus and 15 species)

<i>Euphorbia alpina</i> C.A.Mey.	[7]
<i>Euphorbia caesia</i> Kar. & Kir.	[7]

<i>Euphorbia esula</i> L. [= <i>Euphorbia discolor</i> Ledeb.]	[1–5, 8, 9, 12]
<i>Euphorbia fischeriana</i> Steud.	[4, 5, 9]
<i>Euphorbia humifusa</i> Willd.	[3, 4, 7–16]
<i>Euphorbia kozlovi</i> Prokh.	[8, 10, 12, 13, 16]
<i>Euphorbia macrorhiza</i> Ledeb.	[6, 7]
<i>Euphorbia mongolica</i> Prokh.	[3, 6, 7, 10–13]
<i>Euphorbia pachyrhiza</i> Kar. & Kir.	[7]
<i>Euphorbia pilosa</i> L.	[7]
<i>Euphorbia potaninii</i> Prokh.	[3, 6, 7, 10, 13]
<i>Euphorbia soongarica</i> Boiss. [= <i>Galarhoeus soongaricus</i> (Boiss.) Prokh.]	[7]
<i>Euphorbia subcordata</i> C.A.Mey.	[3, 6, 7, 10, 14]
<i>Euphorbia tshuiensis</i> (Prokh.) Serg.	[6, 7, 10]
<i>Euphorbia virgata</i> Waldst. & Kit. [= <i>Galarhoeus virgatus</i> (Waldst. & Kit.) Prokh.]	[4, 8]

## 52. Fabaceae Lindl. (24 genera and 328 taxa)

	<i>Alhagi maurorum</i> Medik.	[14, 15, 16]
	<i>Alhagi pseudalhagi</i> subsp. <i>kirghisorum</i> (Schrenk) Yakovl. [= <i>Alhagi sparsifolia</i> Shap.]	[14, 15]
SE	<i>Ammopiptanthus mongolicus</i> (Maxim.) S.H.Cheng	[14, 15, 16]
SE	<i>Astragalus admirabilis</i> Pjak & E.Pjak	[7]
	<i>Astragalus adsurgens</i> Pall.	[1–13]
	<i>Astragalus agrestis</i> Douglas ex G.Don	[7, 10]
	<i>Astragalus aksaicus</i> Schischk.	[7]
SE	<i>Astragalus alaschanus</i> Bunge	[13]
	<i>Astragalus alberti</i> Bunge	[10, 11]
	<i>Astragalus albicans</i> Bong.	[14]
	<i>Astragalus alpinus</i> L.	[1, 2, 7, 9, 13]
	<i>Astragalus altaicola</i> Podlech	[6, 7, 14]
	<i>Astragalus ammodytes</i> Pall.	[10, 14]
	<i>Astragalus ankylostus</i> Fisch. & C.A.Mey.	[7, 14]
	<i>Astragalus arcuatus</i> Kar. & Kir.	[14]
	<i>Astragalus argutensis</i> Bunge	[4, 6, 7, 8]
	<i>Astragalus arkalyicensis</i> Bunge	[6, 7, 14]
	<i>Astragalus austro-sibiricus</i> Schischk.	[1, 3, 6, 7, 8, 13]
SE	<i>Astragalus baitagensis</i> Sanchir	[14]
	<i>Astragalus beketowii</i> (Krassn.) B.Fedtsch.	[6]
	<i>Astragalus borodinii</i> Krasnob.	[7, 14]
	<i>Astragalus brachybotrys</i> Bunge	[6, 7, 10–14]
	<i>Astragalus brevifolius</i> Ledeb.	[1–4, 6–8, 10–13]
SE	<i>Astragalus burtschumensis</i> Sumnev.	[6, 7, 10]
	<i>Astragalus candidissimus</i> Ledeb.	[7, 14]
E	<i>Astragalus chamonobrychis</i> Podlech	[7]

E	<i>Astragalus changaicus</i> Sanchir	[3, 7]
	<i>Astragalus chinensis</i> L.f.	[5, 9]
SE	<i>Astragalus chorinensis</i> Bunge [= <i>Astragalus pseudochorinensis</i> N.Ulziykh.]	[2, 3, 4]
E	<i>Astragalus chubsugulicus</i> Gontsch. ex N.Ulziykh.	[1]
	<i>Astragalus compressus</i> Ledeb.	[7]
	<i>Astragalus confertus</i> Benth.	[3, 7]
	<i>Astragalus consanguineus</i> Bong. & C.A.Mey.	[10]
	<i>Astragalus contortuplicatus</i> L.	[14]
	<i>Astragalus dahuricus</i> Patrin	[3–5, 8–10, 12]
	<i>Astragalus danicus</i> Retz.	[1, 4]
	<i>Astragalus depauperatus</i> Ledeb.	[3, 6, 7]
	<i>Astragalus dilutus</i> Bunge	[3, 6, 7, 10, 12–14]
	<i>Astragalus dschimensis</i> Gontsch.	[7, 14]
	<i>Astragalus ellipsoideus</i> Ledeb.	[3, 6, 7, 10, 12, 14–16]
	<i>Astragalus filiformis</i> (DC.) Poir. [= <i>Oxytropis filiformis</i> DC.]	[1–5, 7–11, 13]
	<i>Astragalus follicularis</i> Pall.	[3]
	<i>Astragalus frigidus</i> A.Gray	[1–3, 6, 7, 13]
	<i>Astragalus fruticosus</i> Pall.	[2, 3, 4, 9, 13]
	<i>Astragalus galactites</i> Pall.	[1–5, 8, 9]
	<i>Astragalus glomeratus</i> Ledeb.	[3, 7, 13]
E	<i>Astragalus gobicus</i> Hanelt & Davaz.	[14, 15]
E	<i>Astragalus granitovii</i> Sanchir	[7, 14]
SE	<i>Astragalus gregorii</i> B. Fedtsch. & Basil.	[7]
SE	<i>Astragalus grubovii</i> Sanchir [= <i>Astragalus alaschanensis</i> H.C.Fu]	[7, 10–16]
SE	<i>Astragalus grum-grshimailoi</i> Palib.	[7]
E	<i>Astragalus gubanovii</i> N.Ulziykh.	[7, 10]
SE	<i>Astragalus habaheensis</i> Y.X.Liou	[14]
SE	<i>Astragalus hamiensis</i> S.B.Ho [= <i>Astragalus banzragczii</i> N.Ulziykh.]	[14]
SE	<i>Astragalus hsinbaticus</i> P.Y.Fu & Y.A.Chen [= <i>Astragalus quasitesticulatus</i> Barratte & Z.Y.Chu]	[9]
	<i>Astragalus hypogaeus</i> Ledeb.	[3, 6, 7, 10]
	<i>Astragalus inopinatus</i> Boriss.	[1–5, 7–9, 13]
SE	<i>Astragalus junatovii</i> Sanchir	[12, 13, 15, 16]
	<i>Astragalus kasachstanicus</i> Golosk.	[7]
	<i>Astragalus kaufmannii</i> Krylov	[1, 3]
E	<i>Astragalus kenteicus</i> N.Ulziykh.	[2]
	<i>Astragalus klementzii</i> N.Ulziykh.	[3]
E	<i>Astragalus koslovi</i> B.Fedtsch. & N.Basil.	[13]
	<i>Astragalus kurtschumensis</i> Bunge	[7, 10]
	<i>Astragalus laguroides</i> Pall. [= <i>Astragalus gobi-altaicus</i> N.Ulziykh.]	[2–4, 6–8, 10–13]
	<i>Astragalus lasiopetalus</i> Bunge	[7, 14]

	<i>Astragalus laxmannii</i> Jacq.	[7, 10, 14]
	<i>Astragalus lepsensis</i> Bunge	[7]
	<i>Astragalus leptostachys</i> Pall. [= <i>Astragalus macropterus</i> DC. = <i>Astragalus multicaulis</i> Ledeb.]	[1, 3, 6, 7, 13]
SE	<i>Astragalus lupulinus</i> Pall.	[3, 4, 7, 11–14, 16]
SE	<i>Astragalus luxurians</i> Bunge	[7]
	<i>Astragalus macrolobus</i> M.Bieb. [= <i>Astragalus macrocerus</i> C.A.Mey.]	[7, 10, 11]
	<i>Astragalus macrotrichus</i> E.Peter	[7, 10, 12, 14–16]
	<i>Astragalus majevskianus</i> Krylov	[7]
	<i>Astragalus megalanthus</i> DC.	[8, 12]
	<i>Astragalus melilotoides</i> Pall.	[2, 3, 4, 8–13, 16]
	<i>Astragalus miniatus</i> Bunge	[3, 4, 8, 9, 11–13]
	<i>Astragalus mongholicus</i> Bunge [= <i>Astragalus membranaceus</i> Fisch. = <i>Astragalus propinquus</i> Schischk.]	[1–11, 13]
	<i>Astragalus monophyllus</i> Bunge	[6–16]
	<i>Astragalus norvegicus</i> Weber	[1, 2]
SE	<i>Astragalus ochrias</i> Bunge	[12, 14, 15, 16]
	<i>Astragalus onobrychis</i> L.	[10]
	<i>Astragalus ortholobus</i> Bunge	[7]
	<i>Astragalus oxyglottis</i> Steven	[7, 14]
	<i>Astragalus pallasi</i> Spreng. [= <i>Astragalus lasiophyllus</i> Ledeb.]	[14]
SE	<i>Astragalus pavlovii</i> B.Fedtsch. & Basil.	[13–16]
	<i>Astragalus peterae</i> Tsai & Yu	[6, 10]
	<i>Astragalus physocarpus</i> Ledeb.	[7]
SE	<i>Astragalus politovii</i> Krylov	[7]
SE	<i>Astragalus polozhiae</i> Timokhina	[6, 7]
SE	<i>Astragalus pseudoborodinii</i> S.B.Ho [= <i>Astragalus baischinticus</i> N.Ulziykh.]	[14]
	<i>Astragalus pseudobrachytropis</i> Gontsch.	[6]
E	<i>Astragalus pseudotesticulatus</i> Sanchir	[7]
E	<i>Astragalus pseudovulpinus</i> Sanchir	[14]
	<i>Astragalus puberulus</i> Ledeb. [= <i>Craccina puberula</i> (Ledeb.) Steven]	[7, 10, 11, 13, 14]
	<i>Astragalus roseus</i> Ledeb.	[7, 14]
	<i>Astragalus rudolfii</i> N.Ulziykh.	[7, 14]
	<i>Astragalus rytidocarpus</i> Ledeb.	[2, 3, 7]
	<i>Astragalus sabuletorum</i> Ledeb.	[7, 14, 15]
E	<i>Astragalus saichanensis</i> Sanchir	[7, 13]
E	<i>Astragalus sanczirii</i> N.Ulziykh.	[7, 14]
SE	<i>Astragalus saraleensis</i> Gontsch.	[1]
	<i>Astragalus scaberrimus</i> Bunge	[2–4, 8, 9, 12]
	<i>Astragalus scabrisetus</i> Bong.	[15]
	<i>Astragalus schanginianus</i> Pall.	[7]
	<i>Astragalus schrenkianus</i> Fisch. & C.A.Mey.	[7]

	<i>Astragalus scleropodius</i> Ledeb.	[7]
	<i>Astragalus secundus</i> DC. [≡ <i>Astragalus frigidus</i> subsp. <i>secundus</i> (DC.) Vorosch.]	[1, 3, 13]
	<i>Astragalus sphaerocystis</i> Bunge	[7]
	<i>Astragalus stenoceras</i> C.A.Mey.	[10]
	<i>Astragalus suffruticosus</i> DC.	[1–4, 7, 8, 13]
	<i>Astragalus sulcatus</i> L.	[7, 10, 11, 13, 14]
E	<i>Astragalus tamiricus</i> N.Ulziykh.	[3]
	<i>Astragalus tenuis</i> Turcz. [≡ <i>Astragalus melilotoides</i> Pall. var. <i>tenuis</i> (Turcz.) Ledeb.]	[1, 2, 3, 5, 8, 9]
	<i>Astragalus tephrolobus</i> Bunge	[7]
	<i>Astragalus tibetanus</i> Benth.	[7, 10, 11, 14, 15]
	<i>Astragalus tschujensis</i> Bunge	[7]
	<i>Astragalus tulinovii</i> B.Fedtsch.	[7]
SE	<i>Astragalus tuvinicus</i> Timokhina	[7, 14]
	<i>Astragalus uliginosus</i> L.	[1–5, 8]
E	<i>Astragalus ulziykhutagii</i> Sytin [= <i>Astragalus alexandrii</i> N.Ulziykh.]	[7]
	<i>Astragalus urunguensis</i> N.Ulziykh.	[14]
SE	<i>Astragalus vallestris</i> Kamelin	[3, 7, 10–14]
	<i>Astragalus variabilis</i> Bunge	[7, 11–16]
	<i>Astragalus versicolor</i> Pall. [= <i>Astragalus alexandrii</i> N.Ulziykh. nom. illegit.]	[1, 2, 3, 4, 6]
E	<i>Astragalus viridiflavus</i> N.Ulziykh.	[1, 2, 3, 4]
	<i>Astragalus xanthotrichos</i> Ledeb.	[7]
SE	<i>Astragalus yumenensis</i> S.B.Ho	[14, 15]
SE	<i>Astragalus zacharensis</i> Bunge	[9]
	<i>Astragalus zaissanensis</i> Sumnev.	[7]
	<i>Caragana arborescens</i> Lam.	[1, 3, 10]
	<i>Caragana brachypoda</i> Pojark.	[12, 13, 16]
	<i>Caragana bungei</i> Ledeb.	[3, 6, 7, 10, 11, 13–15]
SE	<i>Caragana davazamcii</i> Sanchir [≡ <i>Caragana korshinskii</i> var. <i>davazamcii</i> (Sanchir) Yakovlev]	[9, 11–13, 16]
E	<i>Caragana gobica</i> Sanczir	[7, 12, 13, 14]
	<i>Caragana halodendron</i> (Pall.) Dum.Cours. [≡ <i>Halimodendron halodendron</i> (Pall.) Voss.]	[7, 10, 14, 15]
	<i>Caragana jubata</i> Poir.	[1, 2, 3, 7, 13]
SE	<i>Caragana korshinskii</i> Kom.	[9, 11–13, 16]
	<i>Caragana leucophloea</i> Pojark.	[3, 4, 6–8, 10–16]
	<i>Caragana microphylla</i> Lam.	[2, 3, 4, 8, 9]
	<i>Caragana pygmaea</i> (L.) DC. [≡ <i>Robinia pygmaea</i> L.]	[1–14]
	<i>Caragana spinosa</i> (L.) Vahl	[4, 6–8, 10, 11, 14, 16]
	<i>Caragana stenophylla</i> Pojark.	[3–5, 8, 9, 12, 13]
SE	<i>Caragana tibetica</i> Kom.	[12, 13, 16]

	<i>Chesneya ferganensis</i> Korsh. [= <i>Chesniella ferganensis</i> (Korsh.) Boriss.]	[15]
SE	<i>Chesneya mongolica</i> Maxim.	[10–13, 15, 16]
SE	<i>Chesniella macrantha</i> (W.C.Cheng) L.Duan, J.Wen & Zhao Y.Chang [= <i>Spongiocarpella grubovii</i> (N.Ulziykh.) Yakovlev]	[15, 16]
	<i>Cicer songaricum</i> Steph.	[7]
	<i>Corethrodendron fruticosum</i> (Pall.) B.H.Choi & H.Ohashi [= <i>Hedysarum fruticosum</i> Pall.]	[3, 4, 5, 8–13, 16]
	<i>Corethrodendron scoparium</i> (Fisch. & C.A.Mey.) Fisch. & Basiner [= <i>Hedysarum scoparium</i> Fisch. & C.A.Mey. = <i>Hedysarum arbuscula</i> Maxim.]	[15, 16]
	<i>Glycyrrhiza aspera</i> Pall.	[10, 14]
	<i>Glycyrrhiza glabra</i> L. [= <i>Glycyrrhiza alaschanica</i> Grankina]	[10, 12, 14, 15, 16]
SE	<i>Glycyrrhiza inflata</i> Batalin	[7, 15, 16]
	<i>Glycyrrhiza pallidiflora</i> Maxim.	[9]
SE	<i>Glycyrrhiza squamulosa</i> Franch.	[12, 14]
	<i>Glycyrrhiza uralensis</i> Fisch. [= <i>Glycyrrhiza gobica</i> Grankina = <i>Glycyrrhiza soongorica</i> Grankina]	[2–5, 8–16]
	<i>Gueldenstaedtia monophylla</i> Fisch.	[6, 7, 10, 12, 13, 16]
	<i>Gueldenstaedtia verna</i> (Georgi) Boriss. [= <i>Gueldenstaedtia stenophylla</i> Bunge]	[1, 2, 4, 5, 9]
	<i>Hedysarum alpinum</i> L. [= <i>Echinolobium alpinum</i> (L.) Desv.]	[1–7, 9]
	<i>Hedysarum austrosibiricum</i> B.Fedtsch. [= <i>Hedysarum hedysaroides</i> subsp. <i>austrosibiricum</i> (B.Fedtsch.) Jurtzev]	[3, 6, 7]
SE	<i>Hedysarum brachypterum</i> Bunge	[2, 9]
	<i>Hedysarum chalchorum</i> N.Ulziykh.	[3, 4, 8]
	<i>Hedysarum consanguineum</i> DC.	[7]
	<i>Hedysarum dahuricum</i> Turcz. [= <i>Hedysarum gmelinii</i> var. <i>dahuricum</i> (Turcz.) R.Sha]	[1, 3–10, 13]
	<i>Hedysarum ferganense</i> Korsh.	[1–4, 6–11, 13, 14, 16]
	<i>Hedysarum gmelinii</i> Ledeb.	[1–4, 6, 7, 9, 10, 13]
	<i>Hedysarum hedysaroides</i> subsp. <i>arcticum</i> (B.Fedtsch.) P.W.Ball [= <i>Hedysarum arcticum</i> B.Fedtsch.]	[1, 3, 6, 7]
	<i>Hedysarum iliense</i> B.Fedtsch.	[7]
	<i>Hedysarum inundatum</i> Turcz.	[1–3, 6, 7, 10, 13]
	<i>Hedysarum kamelinii</i> N.Ulziykh.	[7]
	<i>Hedysarum krylovii</i> Sumn.	[7]
	<i>Hedysarum lintschevskyi</i> Bajtenov	[7, 13]
	<i>Hedysarum neglectum</i> Ledeb.	[1, 2, 3, 6, 7]
	<i>Hedysarum roseum</i> Sims	[2, 3, 4]
	<i>Hedysarum sajanicum</i> N.Ulziykh.	[1]
SE	<i>Hedysarum sangilense</i> Krasnob. & Timokhina	[1, 3]
	<i>Hedysarum setigerum</i> Turcz.	[2–7, 9]
	<i>Hedysarum theinum</i> Krasnob.	[7]

	<i>Lathyrus humilis</i> (Ser.) Fisch. [= <i>Orobus humilis</i> Ser.]	[1–5, 8, 9]
	<i>Lathyrus ledebourii</i> Trautv.	[7]
	<i>Lathyrus palustris</i> L. subsp. <i>pilosus</i> (Cham.) Hultén	[1–5, 8, 9, 10, 14]
	<i>Lathyrus pisiformis</i> L.	[5, 9]
	<i>Lathyrus pratensis</i> L.	[2, 3, 4, 10]
	<i>Lathyrus quinquenervius</i> (Miq.) Litv.	[3, 4, 5]
	<i>Lespedeza bicolor</i> Turcz.	[5]
	<i>Lespedeza daurica</i> (Laxm.) Schindl. [= <i>Trifolium dauricum</i> Laxm.]	[2–5, 8, 9, 11–13, 16]
	<i>Lespedeza juncea</i> (L.f.) Pers.	[2–5, 8, 9]
	<i>Lespedeza tomentosa</i> Siebold	[5, 9]
	<i>Lotus krylovii</i> Schischk. & Serg.	[7, 9, 10, 12, 14–16]
	<i>Medicago falcata</i> L.	[2–12, 14]
	<i>Medicago lupulina</i> L.	[2–5, 7–11, 13, 14]
	<i>Medicago platycarpa</i> (L.) Trautv.	[1, 2, 3, 4, 7, 9]
	<i>Medicago ruthenica</i> Trautv.	[1–5, 8–11, 13, 14]
	<i>Melilotus dentatus</i> (Waldst. & Kit.) Pers.	[2–5, 7–12, 14]
	<i>Melilotus officinalis</i> (L.) Lam.	[2, 3, 4]
	<i>Melilotus suaveolens</i> Ledeb. [= <i>Trigonella suaveolens</i> (Ledeb.) Coulot & Rabaute]	[1, 3–14]
	<i>Melilotus wolgicus</i> Poir.	[9, 12]
	<i>Onobrychis arenaria</i> (Kit.) DC. subsp. <i>arenaria</i>	[2, 3, 4, 8]
	<i>Onobrychis arenaria</i> subsp. <i>sibirica</i> (Turcz.) P.W.Ball [≡ <i>Onobrychis sibirica</i> (Sirj.) Turcz.]	[2, 3, 4, 13]
SE	<i>Oxytropis acanthacea</i> Jurtzev	[6, 7]
	<i>Oxytropis aciphylla</i> Ledeb.	[3, 6, 7, 10–16]
SE	<i>Oxytropis alpestris</i> Schischk.	[7]
	<i>Oxytropis alpicola</i> Turcz.	[2]
	<i>Oxytropis alpina</i> Bunge	[1, 2, 3, 6, 7, 13]
	<i>Oxytropis altaica</i> (Pall.) Pers.	[6, 7]
	<i>Oxytropis ambigua</i> (Pall.) DC.	[1, 2, 3, 4, 7, 13]
	<i>Oxytropis ampullata</i> (Pall.) Pers.	[2, 3, 7–9, 12, 13]
SE	<i>Oxytropis baicalia</i> (Pall.) Pers.	[1, 3, 4]
SE	<i>Oxytropis bicolor</i> Bunge	[9]
	<i>Oxytropis brachycarpa</i> Vassilcz.	[7]
E	<i>Oxytropis bungei</i> Kom.	[3, 6, 7, 8, 10–14]
	<i>Oxytropis caerulea</i> DC.	[1, 2, 4, 5, 9]
	<i>Oxytropis caespitosa</i> Pers.	[1, 2, 3, 4, 5, 8]
	<i>Oxytropis campanulata</i> Vassilcz.	[1, 3]
	<i>Oxytropis chionophylla</i> Schrenk	[3, 6, 7, 13]
	<i>Oxytropis deflexa</i> (Pall.) DC.	[1–4, 6, 7, 10, 13]
	<i>Oxytropis diantha</i> Bunge [= <i>Oxytropis changaica</i> B.Fedtsch. & Basil.]	[1, 3]
SE	<i>Oxytropis dubia</i> Turcz.	[2]

SE	<i>Oxytropis eriocarpa</i> Bunge	[6, 7]
	<i>Oxytropis falcata</i> Bunge	[6, 7]
E	<i>Oxytropis fragilifolia</i> N.Ulziykh.	[7, 13]
SE	<i>Oxytropis gebleri</i> Fisch.	[1, 3, 6, 7, 13]
	<i>Oxytropis glabra</i> DC.	[1–4, 6–16]
	<i>Oxytropis glandulosa</i> Turcz.	[1, 3]
	<i>Oxytropis glareosa</i> Vassilcz.	[3, 10]
	<i>Oxytropis gorbunovii</i> Boriss.	[3, 6, 7]
	<i>Oxytropis grandiflora</i> DC.	[2, 4, 5, 8, 9]
	<i>Oxytropis hailarensis</i> Kitag.	[5, 9]
SE	<i>Oxytropis heterophylla</i> Bunge	[6, 7, 10, 13, 14]
	<i>Oxytropis hirta</i> Bunge	[5]
SE	<i>Oxytropis intermedia</i> Bunge	[3, 6, 7, 10]
E	<i>Oxytropis junatovii</i> Sanchir	[13]
SE	<i>Oxytropis jurtzevii</i> Malyshev	[1]
E	<i>Oxytropis klementzii</i> N.Ulziykh.	[2, 3, 4, 8]
SE	<i>Oxytropis komarovii</i> Vassilcz.	[5, 9]
SE	<i>Oxytropis kossinskyyi</i> B.Fedtsch. & Basil.	[3, 4, 8, 11, 13]
	<i>Oxytropis krylovii</i> Schipcz.	[7]
SE	<i>Oxytropis kusnetzovii</i> Kryl. & Steinb.	[1, 7]
	<i>Oxytropis ladyginii</i> Krylov	[7]
	<i>Oxytropis lanata</i> DC.	[1, 3, 4, 8, 9]
SE	<i>Oxytropis lanuginosa</i> Kom.	[3, 10]
	<i>Oxytropis lapponica</i> Gaudin	[3, 6, 7]
	<i>Oxytropis lasiopoda</i> Bunge	[3, 4, 8, 9, 13]
SE	<i>Oxytropis latibracteata</i> Jurtzev	[3]
E	<i>Oxytropis lavrenkoi</i> N.Ulziykh.	[12]
	<i>Oxytropis leptophylla</i> DC.	[1, 3–5, 8, 9, 12]
SE	<i>Oxytropis leucotricha</i> Turcz.	[1, 2, 3, 8]
	<i>Oxytropis longirostra</i> DC.	[1, 2, 3]
	<i>Oxytropis macrosema</i> Bunge	[6, 7]
SE	<i>Oxytropis martjanovii</i> Krylov	[3, 6, 7, 10]
E	<i>Oxytropis micrantha</i> Bunge	[3, 6, 7, 10, 11]
	<i>Oxytropis microphylla</i> (Pall.) DC.	[3, 6, 7, 10, 12]
SE	<i>Oxytropis mixotricha</i> Bunge	[2, 3, 4, 8]
SE	<i>Oxytropis mongolica</i> Kom.	[6, 10]
SE	<i>Oxytropis monophylla</i> Grubov	[12, 13]
	<i>Oxytropis muricata</i> (Pall.) DC.	[1, 3, 4, 9, 13]
	<i>Oxytropis myriophylla</i> (Pall.) DC.	[1–5, 8, 9]
SE	<i>Oxytropis nitens</i> Turcz.	[1, 2, 3, 4, 8, 9]
SE	<i>Oxytropis ochrantha</i> Turcz.	[9]
	<i>Oxytropis oligantha</i> Bunge	[3, 6, 7, 10, 13]
	<i>Oxytropis oxyphylla</i> (Pall.) DC.	[1–5, 8, 9, 12]

	<i>Oxytropis pauciflora</i> Bunge	[1, 6, 7, 13]
E	<i>Oxytropis pavlovii</i> B.Fedtsch. & Basil.	[3, 8, 11, 12, 13]
SE	<i>Oxytropis physocarpa</i> Ledeb.	[7]
E	<i>Oxytropis potaninii</i> Bunge	[7, 10]
SE	<i>Oxytropis prostrata</i> (Pall.) DC.	[4, 8, 9]
SE	<i>Oxytropis pseudoglandulosa</i> Gontsch.	[1–4, 8, 9, 12, 13]
	<i>Oxytropis puberula</i> Boriss.	[7, 13, 14]
	<i>Oxytropis pumila</i> Fisch.	[3, 6, 7, 8, 10, 11, 13]
	<i>Oxytropis racemosa</i> Turcz. [= <i>Oxytropis gracillima</i> Bunge]	[3, 4, 6, 8–12]
	<i>Oxytropis recognita</i> Bunge	[6, 7]
SE	<i>Oxytropis reverdattoi</i> Jurtzev	[2, 3, 4]
SE	<i>Oxytropis rhizantha</i> Palib.	[6, 7, 10]
	<i>Oxytropis rhynchophysa</i> Schrenk	[6, 7]
SE	<i>Oxytropis sacciformis</i> H.C.Fu	[12]
SE	<i>Oxytropis sajanensis</i> Jurtzev	[1, 3]
	<i>Oxytropis saposhnikovii</i> Krylov	[7, 10]
SE	<i>Oxytropis selengensis</i> Bunge	[2, 3, 4, 8, 9]
SE	<i>Oxytropis setosa</i> (Pall.) DC.	[3, 6]
	<i>Oxytropis songorica</i> (Pall.) DC.	[7]
	<i>Oxytropis sordida</i> (Willd.) Pers.	[1]
	<i>Oxytropis squammulosa</i> DC.	[2–4, 6–10, 12, 13]
SE	<i>Oxytropis stenophylla</i> Bunge	[3, 13]
	<i>Oxytropis strobilacea</i> Bunge	[1–4, 6, 7, 13]
SE	<i>Oxytropis stukovii</i> Palib.	[3, 9]
	<i>Oxytropis sulphurea</i> Ledeb.	[7]
E	<i>Oxytropis sutaica</i> N.Ulziykh.	[3, 7]
E	<i>Oxytropis tenuis</i> Palib.	[6, 7]
	<i>Oxytropis teres</i> DC.	[7]
	<i>Oxytropis tragacanthoides</i> Fisch.	[1, 3, 6–8, 10, 11, 13–15]
	<i>Oxytropis trichophysa</i> Bunge	[3, 6, 7, 10, 11, 13, 14]
SE	<i>Oxytropis tschujae</i> Bunge	[1, 7]
SE	<i>Oxytropis turczaninovii</i> Jurtzev	[1, 3, 4]
E	<i>Oxytropis ulzijchutagii</i> Sanchir	[7]
SE	<i>Oxytropis varlakovi</i> Serg.	[4, 9]
E	<i>Oxytropis viridiflava</i> Kom.	[1–4, 7–9, 11, 13]
	<i>Sophora alopecuroides</i> L.	[12–16]
	<i>Sophora flavescens</i> Aiton	[5, 9]
	<i>Sphaerophysa salsula</i> (Pall.) DC.	[5, 7, 9–16]
	<i>Thermopsis alpina</i> Ledeb.	[1, 2]
SE	<i>Thermopsis dahurica</i> Czebr.	[2, 4, 5, 9, 12]
	<i>Thermopsis lanceolata</i> R.Br. [= <i>Thermopsis lanceolata</i> var. <i>glabra</i> (Czebr.) Yakovlev]	[1–5, 8, 9, 11, 13]
E	<i>Thermopsis longicarpa</i> N.Ulziykh.	[6, 10]

	<i>Thermopsis mongolica</i> Czebr. [= <i>Thermopsis lanceolata</i> var. <i>mongolica</i> (Czebr.) Q.R.Wang & X.Y.Zhu]	[6, 7, 10–14, 16]
SE	<i>Thermopsis przewalskii</i> Czebr.	[9, 13]
	<i>Trifolium eximium</i> Steph.	[1–4, 6–11, 13]
	<i>Trifolium lupinaster</i> L.	[1–9]
	<i>Trifolium pratense</i> L.	[1, 4]
	<i>Trifolium repens</i> L.	[2, 7]
	<i>Trigonella arcuata</i> C.A.Mey.	[7, 14]
	<i>Trigonella cancellata</i> Desf.	[7, 14]
	<i>Vicia amoena</i> Fisch. [= <i>Vicia amoena</i> subsp. <i>sericea</i> (Kitag.) Kamelin & Gubanov]	[1–5, 7, 8, 9]
	<i>Vicia amurensis</i> Oett.	[4, 5, 9]
	<i>Vicia costata</i> Ledeb.	[2–4, 6–14, 16]
	<i>Vicia cracca</i> L.	[1–10, 14]
	<i>Vicia geminiflora</i> Trautv.	[3, 4, 5]
	<i>Vicia japonica</i> A.Gray	[2, 5]
	<i>Vicia macrantha</i> Jurtzev [= <i>Vicia macrantha</i> subsp. <i>olchonensis</i> Peschkova]	[1, 2, 3]
	<i>Vicia megalotropis</i> Ledeb..	[1–5, 8, 9]
	<i>Vicia multicaulis</i> Ledeb. [= <i>Vicia nervata</i> Sipliv.]	[1–6, 8, 13]
SE	<i>Vicia olchonensis</i> (Peschkova) O.D.Nikif. [= <i>Vicia macrantha</i> subsp. <i>olchonensis</i> Peschkova]	[1, 4]
	<i>Vicia pseudorobus</i> Fisch. & C.A.Mey.	[5, 9]
	<i>Vicia ramuliflora</i> (Maxim.) Ohwi [= <i>Vicia baicalensis</i> (Turcz.) B.Fedtsch.]	[2, 3, 4, 5]
	<i>Vicia semenovii</i> B.Fedtsch.	[3, 13]
	<i>Vicia tenuifolia</i> Roth	[3, 6, 7]
SE	<i>Vicia tsydenii</i> Malyshев	[4]
	<i>Vicia unijuga</i> A.Braun	[1–8]
	<i>Vicia venosa</i> Maxim.	[1, 2, 3, 4, 5, 8]

**53. Frankeniaceae** Desv. (1 genus and 2 species)

	<i>Frankenia pulverulenta</i> L.	[10]
SE	<i>Frankenia tuvinica</i> Lomon.	[10]

**54. Gentianaceae** Juss. (8 genera and 32 taxa)

	<i>Centaurium pulchellum</i> subsp. <i>meyeri</i> (Bunge) Tzvelev	[7, 10, 13]
	<i>Centaurium pulchellum</i> (Sw.) Hayek subsp. <i>pulchellum</i>	[10, 11, 15]
	<i>Comastoma falcatum</i> (Turcz.) Toyokuni	[1, 6, 7, 13]
	<i>Comastoma malyschevii</i> (Zuev) Zuev [= <i>Gentianella malyschevii</i> Zuev]	[1, 3, 7]
	<i>Comastoma pulmonarium</i> (Turcz.) Toyokuni	[1, 2, 3, 6]
	<i>Comastoma tenellum</i> (Rottb.) Toyok. [= <i>Gentiana tenella</i> Rottb.]	[1–3, 6, 7, 13, 14]
	<i>Gentiana algida</i> Pall.	[1–3, 6, 7, 13]

<i>Gentiana aquatica</i> L. subsp. <i>aquatica</i>	[1, 2, 3, 6, 7, 8]
<i>Gentiana aquatica</i> var. <i>pseudoaquatica</i> (Kusn.) S.Agrawal [≡ <i>Gentiana pseudoaquatica</i> Kusnezow]	[1–4, 6–9, 13]
<i>Gentiana dahurica</i> Fisch. [≡ <i>Dasytaphana dahurica</i> (Fisch.) Zuev]	[4, 5, 8, 9]
<i>Gentiana decumbens</i> L.f. [≡ <i>Dasytaphana decumbens</i> (L.f.) Zuev]	[1–11, 13, 14]
<i>Gentiana grandiflora</i> Laxm.	[1, 2, 3, 7]
<i>Gentiana leucomelaena</i> Maxim. [≡ <i>Ciminalis leucomelaena</i> (Maxim.) Zuev]	[1–4, 7, 8, 10, 11, 13, 14]
<i>Gentiana macrophylla</i> Pall.	[1–7, 9, 13, 14]
<i>Gentiana prostrata</i> Haenke	[1–4, 6–8, 10, 11, 13, 14]
<i>Gentiana karelinii</i> Griseb. [≡ <i>Gentiana prostrata</i> var. <i>karelinii</i> (Griseb.) Kusn.]	[7]
<i>Gentiana riparia</i> Kar. & Kir.	[7, 14]
<i>Gentiana squarrosa</i> Ledeb. [≡ <i>Ciminalis squarrosa</i> (Ledeb.) Zuev]	[1–11]
<i>Gentiana triflora</i> Pall.	[2, 4, 5]
<i>Gentiana uniflora</i> Georgi	[1, 3, 6, 7]
<i>Gentianella amarella</i> L. subsp. <i>acuta</i> (Michx.) J.M.Gillett [≡ <i>Gentiana acuta</i> Michx.]	[1–4, 6–9, 13]
<i>Gentianella atrata</i> (Bunge) Holub	[5]
<i>Gentianella aurea</i> (L.) Harry Sm.	[2, 3, 6, 7, 13]
<i>Gentianella turkestanorum</i> (Gand.) Holub	[7, 14]
<i>Gentianopsis barbata</i> (Froel.) Ma	[1–11, 13, 14]
<i>Halenia corniculata</i> (L.) Cornaz [≡ <i>Swertia corniculata</i> L.]	[1–5, 8, 13]
<i>Lomatogonium carinthiacum</i> (Wulfen) Rchb. [≡ <i>Swertia carinthiaca</i> Wulfen]	[1–4, 6–8, 11, 13]
<i>Lomatogonium rotatum</i> Fr.	[1–8, 10, 13, 14]
<i>Swertia banzragczii</i> Sanchir	[6, 7]
<i>Swertia dichotoma</i> L. [≡ <i>Anagallidium dichotomum</i> (L.) Griseb.]	[1, 2, 3, 4, 8, 9]
<i>Swertia marginata</i> Schrenk [= <i>Swertia komarovii</i> Pissjauk.]	[1, 7]
<i>Swertia obtusa</i> Ledeb.	[2, 6, 7]

## 55. Geraniaceae Juss. (2 genera and 19 taxa)

<i>Erodium cicutarium</i> (L.) L'Hér.	[2, 4, 12]
<i>Erodium stephanianum</i> Willd.	[2–5, 7–16]
<i>Erodium tibetanum</i> Edgew. & Hook.f.	[4, 6, 7, 8, 10–16]
<i>Geranium affine</i> Ledeb.	[6, 7, 14]
<i>Geranium albiflorum</i> Ledeb.	[2, 3, 6, 7]
<i>Geranium amurense</i> Tsyren.	[3, 4, 9]
<i>Geranium collinum</i> Stephan	[7, 10, 14, 15, 16]
<i>Geranium dahuricum</i> DC.	[1, 5, 6, 9, 10, 12]
<i>Geranium krylovii</i> Tzvelev	[2, 3, 4]
<i>Geranium laetum</i> Ledeb.	[3, 7, 14]
<i>Geranium pamiricum</i> Ikonn.	[14]

<i>Geranium platyanthum</i> Duthie	[2, 3, 4, 5, 9, 12]
<i>Geranium pratense</i> L.	[1–4, 6–9, 12, 13]
<i>Geranium pseudosibiricum</i> J.Mayer	[1–8, 10, 14]
<i>Geranium saxatile</i> Kar. & Kir.	[7, 14]
<i>Geranium sibiricum</i> L.	[1–5, 7–10, 12–14, 16]
<i>Geranium transbaicalicum</i> Serg.	[1, 7, 9]
<i>Geranium transbaicalicum</i> subsp. <i>turczaninovii</i> (Serg.) Peschkova	[3, 4]
<i>Geranium wlassovianum</i> Fisch.	[1, 2, 3, 4, 5, 9]

**56. Grossulariaceae DC.** (1 genus and 12 taxa)

<i>Ribes aciculare</i> Sm. [= <i>Grossularia acicularis</i> (Sm.) Spach]	[2–4, 6–8, 10, 13, 14]
<i>Ribes diacanthum</i> Pall. [= <i>Ribes diacanthum</i> f. <i>weichangense</i> J.X.Huang & J.Z.Wang]	[1–5, 8, 9]
<i>Ribes fragrans</i> Pall.	[1, 2]
<i>Ribes graveolens</i> Bunge	[1, 7]
<i>Ribes heterotrichum</i> C.A.Mey.	[6, 7, 10, 14, 15]
<i>Ribes meyeri</i> Maxim.	[7, 14]
<i>Ribes nigrum</i> L.	[1–7, 10, 13]
<i>Ribes petraeum</i> Wulfen [= <i>Ribes altissimum</i> Turcz.]	[1–4, 6, 7, 14]
<i>Ribes procumbens</i> Pall.	[1, 3, 7]
<i>Ribes pulchellum</i> Turcz.	[1–5, 8, 9, 12, 14]
<i>Ribes rubrum</i> L.	[1–7, 9, 13]
<i>Ribes spicatum</i> E.Robson	[1–3, 6, 7, 9, 10]

**57. Haloragaceae R.Br.** (1 genus and 3 species)

<i>Myriophyllum sibiricum</i> Kom.	[5]
<i>Myriophyllum spicatum</i> L.	[1–11, 14]
<i>Myriophyllum verticillatum</i> L.	[1–10, 14]

**58. Hydrocharitaceae Juss.** (2 genera and 5 species)

<i>Hydrilla verticillata</i> (L.f.) Royle [= <i>Serpicula verticillata</i> L.f.]	[10]
<i>Najas flexilis</i> (Willd.) Rostk. & W.L.E.Schmidt [= <i>Caulinia flexilis</i> Willd.]	[10]
<i>Najas marina</i> L.	[10, 11]
<i>Najas minor</i> All.	[10]
<i>Najas tenuissima</i> (A.Braun) Magnus [= <i>Caulinia tenuissima</i> (A.Braun) Tzvelev]	[10]

**59. Hypericaceae Juss.** (1 genus and 4 taxa)

<i>Hypericum ascyron</i> L. subsp. <i>ascyon</i>	[2, 3, 4, 5]
<i>Hypericum ascyron</i> subsp. <i>gebleri</i> (Ledeb.) N.Robson [≡ <i>Hypericum gebleri</i> Ledeb.]	[2]
<i>Hypericum attenuatum</i> Choisy	[2, 3, 4, 5, 9]
<i>Hypericum perforatum</i> L.	[3, 6]

**60. Iridaceae** Juss. (1 genus and 21 taxa)

- SE *Iris bungei* Maxim. [= *Cryptobasis bungei* (Maxim.) M.B.Crespo] [3, 5, 8, 9, 11–13, 16]  
*Iris dichotoma* Pall. [2, 4, 5, 8, 9]  
*Iris glaucescens* Bunge [6]  
*Iris halophila* Pall. [= *Chamaeiris halophila* (Pall.) M.B.Crespo] [6, 14]  
*Iris humilis* Georgi [= *Iris flavissima* Pall.] [1–5, 8, 9, 12, 13]
- SE *Iris ivanovae* Doronkin [2, 3]  
SE *Iris kamelinii* Alexeeva [1, 7]  
*Iris lactea* Pall. [= *Eremiris lactea* (Pall.) Rodion.] [1–13, 15, 16]  
*Iris loczyi* Kanitz [7, 10]  
*Iris ludwigii* Maxim. [= *Xyridion ludwigii* (Maxim.) Rodion.] [7]  
*Iris potaninii* Maxim. [1–4, 6–13]  
*Iris psammocola* Y.T.Zhao [10]  
SE *Iris pseudothoroldii* Galanin [4]  
*Iris ruthenica* subsp. *brevituba* (Maxim.) Doronkin  
[= *Iris ruthenica* var. *brevituba* Maxim.] [1, 2]  
*Iris ruthenica* Ker Gawl. subsp. *ruthenica* [1, 2, 3, 4, 5]
- E *Iris schmakovii* Alexeeva [1]  
*Iris sibirica* L. [= *Iris sanguinea* Donn] [2, 4, 5, 9]  
*Iris tenuifolia* Pall. [7–15]  
*Iris tigridia* Bunge [1, 2, 3, 4, 8]  
*Iris uniflora* Pall. [= *Joniris uniflora* (Pall.) M.B.Crespo] [4, 5]  
*Iris ventricosa* Pall. [= *Cryptobasis ventricosa* (Pall.) M.B.Crespo] [5, 9]

**61. Juncaceae** Juss. (2 genera and 32 taxa)

- SE *Juncus alpinoarticulatus* subsp. *fischerianus* (V.I.Krecz.) Hämet-Ahti [1–10, 14]  
*Juncus arcticus* subsp. *grubovii* (Novikov) Novikov [= *Juncus grubovii* Novikov] [1, 2, 3]  
*Juncus articulatus* L. subsp. *articulatus* [4, 9, 14]  
*Juncus articulatus* subsp. *limosus* (Vorosch.) Vorosch. [= *Juncus limosus* Vorosch.] [3, 4, 5, 9]  
*Juncus biglumis* L. [1, 6, 7]  
*Juncus bufonius* L. [1–16]  
*Juncus castaneus* subsp. *leucochlamys* (V.I.Krecz.) Hultén  
[= *Juncus leucochlamys* V.I.Krecz.] [1–4, 6, 7, 9]  
*Juncus castaneus* subsp. *triceps* (Rostk.) Novikov [1–3, 6, 7, 13]  
*Juncus compressus* Jacq. [1, 3–5, 8–10, 13, 14]  
*Juncus filiformis* L. [7]  
*Juncus gerardi* Loisel [2–4, 6–10, 13–15]  
*Juncus gracillimus* (Buchenau) V.I.Krecz. & Gontsch. [2, 3, 5, 9]  
*Juncus hybridus* Brot. [= *Juncus bufonius* subsp. *ambiguus* (Guss.) Schinz & Thell.] [3, 4, 6, 7, 9–11, 13–15]

<i>Juncus orchonicus</i> Novikov	[2–5, 8, 9, 10]
<i>Juncus persicus</i> subsp. <i>libanoticus</i> (J.Thiébaut) Novikov & Snogerup [= <i>Juncus libanoticus</i> J.Thiébaut]	[3, 4, 6, 7]
<i>Juncus ranarius</i> Songeon & E.P.Perrier [= <i>Juncus bufonius</i> subsp. <i>nastanthus</i> (V.I.Krecz. & Gontsch.) Soó]	[10, 14]
<i>Juncus salsuginosus</i> Turcz.	[1–4, 6–8, 10–13, 15]
<i>Juncus soranthus</i> Schrenk	[3, 7]
<i>Juncus triglumis</i> L.	[1–4, 6, 7, 10, 13, 14]
<i>Juncus turkestanicus</i> V.I.Krecz. & Gontsch. [= <i>Juncus bufonius</i> subsp. <i>turkestanicus</i> (V.I.Krecz. & Gontsch.) Novikov]	[3–5, 7, 9, 10, 14]
<i>Juncus virens</i> Buchenau [= <i>Juncus papillosum</i> var. <i>virens</i> (Buchenau) Vorosch.]	[4]
<i>Luzula confusa</i> Lindeb.	[1, 2, 6, 7]
<i>Luzula multiflora</i> (Ehrh.) Lej.	[3]
<i>Luzula multiflora</i> subsp. <i>frigida</i> (Buchenau) V.I.Krecz.	[7]
<i>Luzula multiflora</i> subsp. <i>sibirica</i> V.I.Krecz. [= <i>Luzula sibirica</i> (V.I.Krecz.) V.I.Krecz.]	[1–4, 6, 7]
<i>Luzula nivalis</i> (Laest.) Spreng. [= <i>Luzula campestris</i> var. <i>nivalis</i> Laest.]	[1]
<i>Luzula pallescens</i> Sw.	[1–5, 7, 9]
<i>Luzula parviflora</i> Desv.	[1, 2, 3, 6, 7]
<i>Luzula pilosa</i> (L.) Willd. [= <i>Juncus pilosus</i> L.]	[2]
<i>Luzula rufescens</i> var. <i>macrocarpa</i> Buchenau [= <i>Luzula changaica</i> Novikov]	[3]
<i>Luzula spicata</i> subsp. <i>mongolica</i> Novikov	[1–3, 6, 7, 13]
<i>Luzula rufescens</i> Fisch. var. <i>rufescens</i>	[1, 2, 3, 4]

## 62. Juncaginaceae Juss. (1 genus and 2 species)

<i>Triglochin maritima</i> L.	[1–16]
<i>Triglochin palustris</i> L.	[1–9, 11–16]

## 63. Lamiaceae Martinov (22 genera and 103 taxa)

Note: The herbarium records of *Phlomis oreophila* in Mongolia was identified as *Phlmoides chinghoensis* by Lazkov (2011).

<i>Amethystea caerulea</i> L.	[2–11, 13]
<i>Caryopteris mongholica</i> Bunge	[2–4, 7–9, 11–13, 15, 16]
<i>Dracocephalum argunense</i> Fisch.	[5]
<i>Dracocephalum discolor</i> Bunge	[3, 7, 10]
<i>Dracocephalum foetidum</i> Bunge	[1–4, 6–13]
<i>Dracocephalum fragile</i> Turcz.	[1, 3, 6, 7]
<i>Dracocephalum fruticosum</i> Stephan	[3, 4, 6–8, 10–13, 16]
<i>Dracocephalum grandiflorum</i> L.	[1–3, 6, 7, 13]
<i>Dracocephalum heterophyllum</i> subsp. <i>heterophyllum</i> Benth.	[3]
<i>Dracocephalum heterophyllum</i> subsp. <i>ovalifolium</i> A.L.Budantzev [= <i>Dracocephalum ovalifolium</i> (A.L.Budantzev) Doronkin]	[3]

<i>Dracocephalum imberbe</i> Bunge	[1, 6, 7]
<i>Dracocephalum integrifolium</i> Bunge [= <i>Ruyschiana integrifolia</i> (Bunge) House]	[6, 7]
<i>Dracocephalum junatovii</i> A.L.Budantzev	[4, 9]
<i>Dracocephalum moldavicum</i> C.Morren	[12, 13, 15]
<i>Dracocephalum nodulosum</i> Rupr.	[14]
<i>Dracocephalum nutans</i> L.	[1, 2, 3, 4, 7]
<i>Dracocephalum olchonense</i> Peschkova	[4]
<i>Dracocephalum origanoides</i> Steph. subsp. <i>origanoides</i>	[1, 3, 4, 6–9, 13, 14]
<i>Dracocephalum origanoides</i> subsp. <i>bungeanum</i> (Schischk. & Serg. A.L.Budantzev [= <i>Dracocephalum bungeanum</i> Schischk. & Serg.])	[1, 6, 7, 13]
<i>Dracocephalum paulsenii</i> Briq.	[14]
<i>Dracocephalum peregrinum</i> L.	[6, 7]
<i>Dracocephalum pinnatum</i> L.	[6]
<i>Dracocephalum ruyschiana</i> L.	[2, 3, 4, 5, 6, 8]
<i>Elsholtzia ciliata</i> (Thunb.) Hyl.	[2, 4]
<i>Elsholtzia densa</i> Benth.	[4, 13]
<i>Galeopsis bifida</i> Boenn. [= <i>Galeopsis tetrahit</i> var. <i>bifida</i> (Boenn.) Lej. & Courtois]	[2, 3, 4, 9]
<i>Hyssopus ambiguus</i> (Trautv.) Iljin [= <i>Hyssopus officinalis</i> var. <i>ambiguus</i> Trautv.]	[7]
<i>Hyssopus cuspidatus</i> Boriss.	[7, 14]
<i>Lagochilus bungei</i> Benth.	[7, 14]
<i>Lagochilus diacanthophyllus</i> Benth.	[6, 7, 14]
<i>Lagochilus ilicifolius</i> Bunge	[3, 7, 8, 10–16]
<i>Lagopsis darwiniana</i> Pjak	[7]
<i>Lagopsis eriostachya</i> (Benth.) Ikonn.-Gal.	[1, 7, 10, 14]
<i>Lagopsis flava</i> Kar. & Kir.	[7]
<i>Lagopsis marrubiastrum</i> (Steph.) Ikonn.-Gal.	[3, 6, 7, 13, 14]
<i>Lagopsis supina</i> (Steph.) Ikonn.-Gal.	[2, 3, 4, 9]
<i>Lamium album</i> L.	[1, 2, 4, 5, 7, 9]
<i>Leonurus deminutus</i> V.I.Krecz. [= <i>Leonurus glaucescens</i> var. <i>deminutus</i> (V.I.Krecz.) Karav.]	[1–4, 7, 8, 9, 13]
<i>Leonurus glaucescens</i> Bunge	[6, 7, 8, 9]
<i>Leonurus mongolicus</i> V.I.Krecz. & Kuprian.	[2–4, 6–9]
<i>Leonurus pseudopanzerioides</i> Krestovsk. [= <i>Leonurus cardiaca</i> subsp. <i>turkestanicus</i> (V.I.Krecz. & Kuprian.) Rech.f.]	[7, 14]
<i>Leonurus sibiricus</i> L.	[1–5, 8, 9, 12]
<i>Leonurus turkestanicus</i> V.I.Krecz. & Kuprian.	[7]
<i>Lophanthus chinensis</i> Benth.	[1–4, 6–10, 12, 13]
<i>Lophanthus krylovii</i> Lipsky	[7]
<i>Lycopus lucidus</i> Turcz.	[9]

	<i>Mentha aquatica</i> L.	[4]
	<i>Mentha arvensis</i> L.	[2–10, 14]
	<i>Mentha canadensis</i> L.	[2]
	<i>Nepeta annua</i> Pall. [= <i>Schizonepeta annua</i> (Pall.) Schischk.]	[3, 6–16]
	<i>Nepeta densiflora</i> Kar. & Kir.	[7, 14]
	<i>Nepeta micrantha</i> Bunge	[7, 14]
	<i>Nepeta multifida</i> L.	[1–5, 7–9, 13]
	<i>Nepeta nuda</i> L.	[6]
	<i>Nepeta pungens</i> Benth.	[14]
	<i>Nepeta sibirica</i> L.	[2, 3, 6, 7, 10, 13, 14]
	<i>Origanum vulgare</i> L.	[1, 6, 9]
	<i>Panzerina canescens</i> (Bunge) Soják	[6, 7, 10, 13]
	<i>Panzerina lanata</i> (L.) Soják [= <i>Ballota lanata</i> Willd.]	[2, 3, 4, 6–14, 16]
	<i>Phlomoides agraria</i> (Bunge) Adylov [= <i>Phlomis agraria</i> Bunge]	[6, 7]
	<i>Phlomoides alpina</i> (Pall.) Adylov [= <i>Phlomis alpina</i> Pall.]	[7]
SE	<i>Phlomoides chinghoensis</i> (C.Y.Wu) Kamelin & Makhm. [= <i>Phlomis chinghoensis</i> C.Y.Wu]	[7, 10, 14]
	<i>Phlomoides molucelloides</i> (Bunge) Salmaki [= <i>Eremostachys molucelloides</i> Bunge]	[6, 14]
	<i>Phlomoides mongolica</i> (Turcz.) Kamelin & A.L.Budantzev [= <i>Phlomis mongolica</i> Turcz.]	[5, 9]
	<i>Phlomoides pratensis</i> (Kar. & Kir.) Adylov [= <i>Phlomis pratensis</i> Kar. & Kir.]	[6]
	<i>Phlomoides tuberosa</i> Moench [= <i>Phlomis tuberosa</i> L.]	[2–9]
	<i>Phlomoides tuvinica</i> (A.Schroet.) Kamelin [= <i>Phlomis tuvinica</i> A.Schroet.]	[6, 7, 8]
	<i>Salvia abrotanoides</i> (Kar.) Sytsma [= <i>Perovskia abrotanoides</i> Kar.]	[6]
	<i>Salvia deserta</i> Schangin	[6]
	<i>Scutellaria altaica</i> Ledeb.	[7]
	<i>Scutellaria baicalensis</i> Georgi	[1–5, 8, 9]
	<i>Scutellaria dependens</i> Maxim.	[2, 4]
	<i>Scutellaria galericulata</i> L.	[1–6, 9, 10, 14]
	<i>Scutellaria grandiflora</i> Sims subsp. <i>grandiflora</i>	[2–4, 6, 7, 10, 13, 14]
E	<i>Scutellaria grandiflora</i> subsp. <i>gymnosperma</i> Kamelin & Gubanov	[7, 13]
	<i>Scutellaria krasevii</i> Kom. & I.Schischk.	[3]
	<i>Scutellaria paulsenii</i> Briq.	[7]
	<i>Scutellaria regeliana</i> var. <i>ikonnikovii</i> (Juz.) C.Y.Wu & H.W.Li	[2, 4]
	<i>Scutellaria scordifolia</i> Fisch.	[1–9]
	<i>Scutellaria sieversii</i> Bunge	[6, 7]
	<i>Scutellaria supina</i> L. [= <i>Scutellaria alpina</i> subsp. <i>supina</i> (L.) I.Richardson]	[7, 14]
	<i>Scutellaria tuvensis</i> Juz. [= <i>Scutellaria grandiflora</i> subsp. <i>tuvensis</i> (Juz.) Kamelin & Gubanov]	[10]
	<i>Scutellaria viscidula</i> Bunge	[9]

	<i>Stachys aspera</i> subsp. <i>baicalensis</i> (Fisch.) Krestovsk. [≡ <i>Stachys baicalensis</i> Fisch.]	[2, 3, 4, 5]
	<i>Stachys palustris</i> L.	[2–6, 9, 10]
	<i>Thymus altaicus</i> Klokov & Des.-Shost.	[3, 6, 7, 10]
	<i>Thymus baicalensis</i> Serg.	[1, 2, 3, 4, 10]
	<i>Thymus bituminosus</i> Klokov	[1]
	<i>Thymus dahuricus</i> Serg.	[2, 4, 5, 8, 9]
E	<i>Thymus gobi-altaicus</i> (N.Ulziykh.) Kamelin & A.L.Budantzev	[13]
	<i>Thymus gobicus</i> Tscherneva	[2, 3, 4, 7–13]
	<i>Thymus komarovii</i> Serg.	[9]
	<i>Thymus michaelis</i> Kamelin & A.L.Budantzev	[2, 4, 8, 9]
	<i>Thymus minussinensis</i> Serg.	[10]
	<i>Thymus mongolicus</i> (Ronniger) Ronniger	[3, 7, 9, 13]
	<i>Thymus narymensis</i> Serg.	[7]
	<i>Thymus pavlovii</i> Serg.	[1, 3]
	<i>Thymus roseus</i> Schipcz.	[7]
	<i>Thymus sibiricus</i> Klokov & Des.-Shost.	[4]
	<i>Thymus turczaninovii</i> Serg.	[9]
	<i>Ziziphora clinopodioides</i> Lam. subsp. <i>clinopodioides</i>	[7, 14]
	<i>Ziziphora clinopodioides</i> subsp. <i>bungeana</i> (Juz.) Rech.f. [≡ <i>Ziziphora bungeana</i> Juz.]	[6, 7, 13]
	<i>Ziziphora pamiroalaica</i> Juz.	[7, 14]

#### 64. Lentibulariaceae Rich. (2 genera and 7 species)

	<i>Pinguicula alpina</i> L.	[1]
	<i>Pinguicula vulgaris</i> L.	[1]
	<i>Utricularia australis</i> R.Br.	[10, 14]
	<i>Utricularia intermedia</i> Hayne [≡ <i>Lentibularia intermedia</i> (Hayne) Nieuwl. & Lunell]	[1, 2, 3, 6]
	<i>Utricularia × japonica</i> Makino	[10]
	<i>Utricularia minor</i> L.	[3, 9, 10, 14]
	<i>Utricularia vulgaris</i> L. [≡ <i>Lentibularia vulgaris</i> (L.) Moench]	[1–11, 14, 15]

#### 65. Liliaceae Juss. (5 genera and 15 taxa)

SE	<i>Erythronium sibiricum</i> (Fisch. & C.A.Mey.) Krylov	[7]
	<i>Fritillaria dagana</i> Turcz.	[1, 2, 3]
	<i>Gagea brevistolonifera</i> Levichev	[7]
	<i>Gagea filiformis</i> Merckl.	[7]
	<i>Gagea granulosa</i> Turcz.	[7]
	<i>Gagea hiensis</i> Pasch. [= <i>Gagea terraccianoana</i> Pasch.]	[1, 2]
SE	<i>Gagea kuraiensis</i> Levichev	[7]
	<i>Gagea fragifera</i> (Vill.) Ehr.Bayer & G.López [= <i>Gagea liotardii</i> (Sternb.) Schult. & Schult. f.]	[7]

<i>Gagea pauciflora</i> Turcz.	[1–5, 7, 9, 14]
<i>Gagea serotina</i> (L.) Ker Gawl. [= <i>Lloydia serotina</i> (L.) Salisb.]	[1–3, 6, 7, 13]
<i>Lilium concolor</i> var. <i>partheneion</i> (Siebold & de Vriese) Baker [= <i>Lilium buschianum</i> G.Lodd.]	[5]
<i>Lilium martagon</i> L.	[1–7]
<i>Lilium pensylvanicum</i> Ker Gawl. [= <i>Lilium dauricum</i> Ker Gawl.]	[2, 4, 5]
<i>Lilium pumilum</i> Redouté [= <i>Lilium potaninii</i> Vrshcz]	[1–5, 8, 9, 12]
<i>Tulipa uniflora</i> (L.) Besser	[3, 5, 7–10, 14]

#### 66. Linaceae DC. (1 genus and 5 species)

<i>Linum altaicum</i> Ledeb.	[1, 6, 7, 8, 9]
<i>Linum baicalense</i> Juz.	[1–5, 7–9, 13]
<i>Linum pallescens</i> Bunge	[2–5, 7–10, 13, 14]
<i>Linum perenne</i> L.	[7, 14]
<i>Linum violascens</i> Bunge	[7]

#### 67. Lythraceae J.St.-Hil. (1 genus and 3 species)

<i>Lythrum salicaria</i> L.	[4]
<i>Lythrum virgatum</i> L.	[6, 14]
<i>Lythrum borysthenicum</i> (Schrank) Litv. [= <i>Peplis borysthenica</i> Schrank]	[10]

#### 68. Malvaceae Juss. (2 genera and 5 species)

<i>Abutilon theophrasti</i> Medik.	[15]
<i>Malva neglecta</i> Wallr..	[4, 7, 10, 11, 14]
<i>Malva pusilla</i> Sm.	[13, 14, 15]
<i>Malva sylvestris</i> L.	[13, 15]
<i>Malva verticillata</i> L.	[1–4, 7, 8, 10, 16]

#### 69. Mazaceae Reveal (3 genera and 3 species)

Note: Mazaceae was separated from Phrymaceae according to APG IV (2016).

<i>Dodartia orientalis</i> L.	[6, 7, 14]
<i>Lancea tibetica</i> Hook.f. & Thomson	[1, 3]
<i>Mazus stachydifolius</i> Maxim.	[5]

#### 70. Melanthiaceae Batsch (3 genera and 5 species)

<i>Anticlea sibirica</i> (L.) Kunth [= <i>Zigadenus sibiricus</i> (L.) A.Gray]	[1, 3]
<i>Paris quadrifolia</i> L.	[2, 3, 4, 5]
<i>Paris verticillata</i> M.Bieb.	[2, 4, 5]
<i>Veratrum lobelianum</i> Bernh.	[1, 3, 4, 5, 7, 9]
<i>Veratrum nigrum</i> L.	[3, 4, 5, 9]

- 71. Menispermaceae** Juss. (1 genus and 1 species)  
*Menispermum dauricum* DC. [2, 3, 4, 5]
- 72. Menyanthaceae** Dumort. (2 genera and 2 species)  
*Nymphoides peltata* (S.G.Gmel.) Kuntze [1, 3, 8–10, 14]  
*Menyanthes trifoliata* L. [1, 2, 3, 4]
- 73. Molluginaceae** Bartl. (1 genus and 1 species)  
*Hypertelis cerviana* (L.) Thulin [= *Mollugo cerviana* (L.) Ser.] [12, 14, 15]
- 74. Montiaceae** Raf. (1 genus and 1 species)  
*Claytonia joanneana* Roem. & Schult. [1, 2, 3, 4, 6, 7]
- 75. Nitrariaceae** Lindl. (2 genera and 5 species)  
*Nitraria roborowskii* Kom. [= *Nitraria schoberi* var. *roborowskii* (Kom.) Grubov] [7, 10, 13, 14, 15]  
*Nitraria sibirica* Poir. [4, 6–16]  
*Nitraria sphaerocarpa* Maxim. [13, 15, 16]  
*Peganum harmala* L. [= *Peganum multisectum* (Maxim.) Bobrov] [7, 10, 13, 14, 15]  
*Peganum nigellastrum* Bunge [3, 4, 8–13, 16]
- 76. Nymphaeaceae** Salisb. (2 genera and 2 species)  
Note: Recently, this family was revised based on field observations and extensive herbarium specimens in Mongolia (Baasanmunkh et al. 2022b). *Nymphaea tetragona* Georgi is not recorded in Mongolia, according to Baasanmunkh et al (2022b).  
*Nuphar pumila* (Timm) DC. [1, 3, 10, 11]  
*Nymphaea candida* J.Presl. [1, 3, 7, 10, 11]
- 77. Onagraceae** Juss. (2 genera and 12 taxa)  
*Circaeа alpina* L. subsp. *alpina* [2, 3, 4, 5]  
*Circaeа alpina* subsp. *caulescens* (Kom.) Tatew. [3]  
*Epilobium anagallidifolium* Lam. [= *Epilobium alpinum* L.] [7]  
*Epilobium angustifolium* L. [= *Chamaenerion angustifolium* (L.) Schur] [1–9, 14]  
*Epilobium ciliatum* Raf. [2]  
*Epilobium davuricum* Fisch. [2, 3, 4, 11]  
*Epilobium fastigiato-ramosum* Nakai [= *Epilobium baicalense* Popov] [3, 4, 9]  
*Epilobium hirsutum* L. [3, 6]  
*Epilobium latifolium* L. [= *Chamaenerion latifolium* (L.) Sweet] [1–3, 6, 7, 13, 14]  
*Epilobium minutiflorum* Hausskn. [7, 10, 15]  
*Epilobium nervosum* Boiss. & Buhse [7]  
*Epilobium palustre* L. [1–10, 13–15]

### 78. Orchidaceae Juss. (14 genera and 26 taxa)

Note: Orchids of Mongolia were recently revised by Baasanmunkh et al. (2021b).

<i>Calypso bulbosa</i> (L.) Oakes [= <i>Cypripedium bulbosum</i> L.]	[2, 4]
<i>Corallorrhiza trifida</i> Châtel.	[1, 2, 4]
<i>Cypripedium calceolus</i> L..	[1, 2, 4]
<i>Cypripedium guttatum</i> Sw.	[1, 2, 3, 4, 5]
<i>Cypripedium macranthos</i> Sw.	[1, 2, 4, 5]
<i>Cypripedium × ventricosum</i> Sw.	[2]
<i>Dactylorhiza fuchsii</i> (Druce) Soó	[2, 4]
<i>Dactylorhiza incarnata</i> (L.) Soó	[3, 5]
<i>Dactylorhiza incarnata</i> subsp. <i>cruenta</i> (O.F.Müll.) P.D.Sell	[3]
<i>Dactylorhiza salina</i> (Turcz.) Soó	[1–11, 14]
<i>Dactylorhiza umbrosa</i> (Kar. & Kir.) Nevski	[1, 3, 4, 7, 10, 14]
<i>Dactylorhiza viridis</i> (L.) R.M.Bateman [= <i>Coeloglossum viride</i> (L.) Hartm.]	[1–7]
<i>Epipogium aphyllum</i> Sw.	[1, 2, 3, 4]
<i>Goodyera repens</i> (L.) R.Br.	[1, 2, 3, 4, 6]
<i>Gymnadenia conopsea</i> (L.) R.Br. [= <i>Orchis conopsea</i> L.]	[1–5]
<i>Herminium alaschanicum</i> Maxim. [= <i>Peristylus alaschanicus</i> (Maxim.) N.Pearce & P.J.Cribb]	[16]
<i>Herminium monorchis</i> R.Br.	[1–5, 8, 9, 10]
<i>Malaxis monophyllos</i> (L.) Sw. [= <i>Ophrys monophyllos</i> L.]	[1, 2, 3, 4, 5]
<i>Neottia camtschatea</i> Sprengel	[1, 2, 3, 7]
<i>Neottia puberula</i> (Maxim.) Szlach. [= <i>Listera puberula</i> Maxim.]	[5]
<i>Orchis militaris</i> L.	[3, 4]
<i>Platanthera bifolia</i> (L.) Rich.	[1, 2, 3, 4]
<i>Platanthera fuscescens</i> Kraenzl.	[2, 3, 4, 5]
<i>Platanthera oligantha</i> Turcz.	[1, 3]
<i>Ponerorchis cucullata</i> (L.) X.H.Jin [= <i>Neottianthe cucullata</i> (L.) Schltr.]	[1, 2, 3, 4]
<i>Spiranthes australis</i> Lindl.	[2–5, 8, 9, 10]

### 79. Orobanchaceae Vent. (9 genera and 57 taxa)

<i>Boschniakia rossica</i> (Cham. & Schldl.) B.Fedtsch.	[2]
<i>Castilleja pallida</i> (L.) Spreng.	[1–9, 13]
<i>Cistanche deserticola</i> Ma	[7, 10–16]
<i>Cistanche feddeana</i> K.S.Hao	[9, 12, 13, 16]
<i>Cistanche lanzhouensis</i> Zhi Y.Zhang	[12]
<i>Cistanche salsa</i> (C.A.Mey.) Beck	[12–16]
<i>Cymbalaria daurica</i> L.	[2–5, 7–13]
<i>Euphrasia altaica</i> Serg.	[7]
<i>Euphrasia hirtella</i> Jord.	[2, 3, 4, 5]
<i>Euphrasia maximowiczii</i> Wettst.	[2, 4, 5, 9]
<i>Euphrasia pectinata</i> Ten.	[1–10, 13, 14]
<i>Euphrasia schischkinii</i> Serg.	[7]

	<i>Euphrasia syreitschikovii</i> Govor.	[1, 2, 3, 5–8]
	<i>Odontites vulgaris</i> Moench	[2–4, 7–11, 14]
	<i>Orobanche amoena</i> C.A.Mey.	[7, 14]
	<i>Orobanche caesia</i> Rchb. [= <i>Phelypaea lanuginosa</i> C.A.Mey. ≡ <i>Orobanche lanuginosa</i> (C.A.Mey.) Beck]	[1, 2, 3, 4]
	<i>Orobanche cernua</i> Loefl.	[7, 10, 13, 14]
	<i>Orobanche coerulescens</i> Steph. [= <i>Orobanchella coerulescens</i> (Steph.) Piwow.]	[5, 7–15]
	<i>Orobanche coerulescens</i> var. <i>albiflora</i> Kuntze [= <i>Orobanche korshinskyi</i> Novopokr.]	[1–15]
	<i>Orobanche pycnostachya</i> Hance	[5]
	<i>Pedicularis abrotanifolia</i> M.Bieb.	[1, 3, 6, 7, 13, 14]
	<i>Pedicularis achilleifolia</i> Steph.	[1, 3, 6–8, 10, 14]
	<i>Pedicularis altaica</i> Steph.	[6, 7, 10, 14]
	<i>Pedicularis amoena</i> Adams	[1–3, 6, 7, 13, 14]
	<i>Pedicularis anthemifolia</i> Fisch.	[1, 3, 6, 7, 13]
	<i>Pedicularis compacta</i> Steph.	[1, 2, 3, 6, 7]
	<i>Pedicularis dolichorrhiza</i> Schrenk	[7, 14]
	<i>Pedicularis elata</i> Willd.	[3, 6, 7]
SE	<i>Pedicularis fetisowii</i> Regel	[14]
	<i>Pedicularis fissa</i> Turcz.	[2, 7]
	<i>Pedicularis flava</i> Pall.	[2–4, 6–11, 13–15]
SE	<i>Pedicularis incarnata</i> L.	[1]
	<i>Pedicularis labradorica</i> Wirsing	[1, 2, 3, 4]
	<i>Pedicularis lapponica</i> L.	[1]
	<i>Pedicularis lasiostachys</i> Bunge	[3, 6, 7]
	<i>Pedicularis longiflora</i> Rudolph	[1, 2, 3, 7, 11]
	<i>Pedicularis moschata</i> Maxim.	[6, 7, 10]
	<i>Pedicularis myriophylla</i> Pall.	[1–4, 6–8, 13]
	<i>Pedicularis oederi</i> Vahl	[1, 2, 3, 6, 7]
	<i>Pedicularis palustris</i> L. subsp. <i>karoi</i> (Freyn) P.C.Tsoong [≡ <i>Pedicularis karoi</i> Freyn]	[1–6, 8, 9, 10, 14]
	<i>Pedicularis physocalyx</i> Bunge	[7]
	<i>Pedicularis proboscidea</i> Steven	[7]
	<i>Pedicularis resupinata</i> L.	[1–6, 8–10, 13]
	<i>Pedicularis rhinanthonoides</i> Schrenk	[7]
	<i>Pedicularis rubens</i> Steph.	[1, 2, 3, 4, 5]
	<i>Pedicularis sceptrum-carolinum</i> L.	[2, 3, 4, 5, 7]
	<i>Pedicularis sibirica</i> Vved.	[1, 3, 7]
	<i>Pedicularis spicata</i> Pall.	[3, 4, 5]
	<i>Pedicularis striata</i> Pall.	[1–5, 8, 9]
	<i>Pedicularis sudetica</i> Willd.	[1, 2]
	<i>Pedicularis tristis</i> L	[1, 2, 3, 6, 7, 8]

<i>Pedicularis uliginosa</i> Bunge	[1–4, 6, 7, 10, 13]
<i>Pedicularis venusta</i> Schangin	[1–4, 6–11]
<i>Pedicularis verticillata</i> L.	[1, 2, 3, 4, 5, 9]
<i>Pedicularis wlassoviana</i> Steven	[2]
<i>Rhinanthus serotinus</i> Oborny	[2]
<i>Rhinanthus songaricus</i> (Sterneck) B.Fedtsch. [= <i>Rhinanthus borbasii</i> (Dörf.) Soó subsp. <i>songaricus</i> (Sterneck) Soó]	[2, 3, 4]

**80. Oxalidaceae** R.Br. (1 genus and 1 species)

<i>Oxalis acetosella</i> L.	[1, 2]
-----------------------------	--------

**81. Paeoniaceae** Raf. (1 genus and 3 species)

<i>Paeonia anomala</i> L.	[1, 2, 3, 4, 6, 7]
<i>Paeonia intermedia</i> C.A.Mey.	[6, 7]
<i>Paeonia lactiflora</i> Pall.	[2, 4, 5, 9]

**82. Papaveraceae** Juss. (6 genera and 30 species)

<i>Chelidonium majus</i> L.	[1–5, 7, 9]
<i>Corydalis adunca</i> Maxim.	[6, 7, 13, 14, 15]
<i>Corydalis capnoides</i> Pers.	[2, 6, 7, 9, 14]
<i>Corydalis grubovii</i> Mikhailova	[6, 7]
<i>Corydalis impatiens</i> Fisch.	[1, 2, 3, 7]
<i>Corydalis inconspicua</i> Bunge	[1, 2, 6, 7]
<i>Corydalis pauciflora</i> Pers.	[1, 2, 6, 13]
<i>Corydalis sajanensis</i> Peschkova [= <i>Corydalis pauciflora</i> subsp. <i>sajanensis</i> (Peschkova) Mikhailova]	[1]
<i>Corydalis schanginii</i> (Pall.) B.Fedtsch.	[7, 14]
<i>Corydalis sibirica</i> Pers.	[1–4, 6–8, 10, 13]
<i>Corydalis stricta</i> Steph. [= <i>Corydalis grubovii</i> Mikhailova]	[6, 7, 13]
<i>Fumaria officinalis</i> L.	[7]
<i>Fumaria schleicheri</i> Soy.-Will.	[7, 14]
<i>Glaucium elegans</i> Fisch. & C.A.Mey.	[14]
<i>Glaucium squamigerum</i> Kar. & Kir.	[7, 14]
<i>Hypecoum erectum</i> L.	[2–5, 8, 9, 11, 12]
<i>Hypecoum lactiflorum</i> (Kar. & Kir.) Pazii	[3, 4, 6–16]
<i>Hypecoum leptocarpum</i> Hook.f. & Thomson	[3]
<i>Papaver baitagense</i> Kamelin & Gubanov	[6, 7, 14]
<i>Papaver canescens</i> Tolm.	[1–7, 13]
<i>Papaver chakassicum</i> Peschkova	[6, 7]
<i>Papaver lapponicum</i> (Tolm.) Nordh.	[7]
<i>Papaver nudicaule</i> L.	[1–7, 9, 13]
<i>Papaver pseudocanescens</i> Popov	[1, 3, 6, 7, 13]
<i>Papaver pseudotentellum</i> Grubov	[7, 10, 13, 14]

	<i>Papaver refractum</i> (DC.) K.-F.Günther [= <i>Roemeria refracta</i> DC.]	[3, 14]
	<i>Papaver rubroaurantiacum</i> (Fisch.) C.E.Lundstr.	[1–10, 13]
SE	<i>Papaver saichanense</i> Grubov [= <i>Papaver rubroaurantiacum</i> subsp. <i>saichanense</i> (Grubov) Kamelin & Cubanov]	[7, 13]
	<i>Papaver smirnovii</i> Peschkova [= <i>Papaver rubroaurantiacum</i> subsp. <i>smirnovii</i> (Peschkova) Kamelin & Cubanov]	[4, 9]
	<i>Papaver setosum</i> (Tolm.) Peschkova [= <i>Papaver rubroaurantiacum</i> subsp. <i>setosum</i> Tolm.]	[4]

**83. Phyllanthaceae** Martinov (1 genus and 1 species)

<i>Flueggea suffruticosa</i> Baill.	[5, 9]
-------------------------------------	--------

**84. Plantaginaceae** Juss. (7 genera and 47 species)

<i>Callitricha hermaphroditica</i> L.	[1, 3, 4, 5]
<i>Callitricha palustris</i> L	[1–5, 7, 9–11, 14]
<i>Hippuris vulgaris</i> L.	[1–11, 13, 14]
<i>Lagotis integrifolia</i> (Willd.) Schischk.	[1, 2, 3, 7, 13]
<i>Linaria acutiloba</i> Fisch. [= <i>Linaria vulgaris</i> Mill. subsp. <i>acutiloba</i> (Fisch. ) D.Y.Hong]	[1–4, 6–8, 13, 14]
<i>Linaria altaica</i> Fisch.	[3, 6, 7, 10, 14]
<i>Linaria buriatica</i> Turcz.	[1–6, 8, 9]
<i>Linaria debilis</i> Kuprian.	[7, 14]
<i>Linaria hepatica</i> Bunge	[6, 7, 11, 13, 14]
<i>Linaria incompleta</i> Kuprian.	[7]
<i>Linaria melampyroides</i> Kuprian.	[3, 4, 5, 7, 9]
<i>Linaria pedicellata</i> Kuprian.	[6, 7, 10, 13, 14]
<i>Plantago arachnoidea</i> Schrenk [= <i>Plantago lorata</i> (J.Z.Liu) Shipunov ≡ <i>Plantago arachnoidea</i> var. <i>lorata</i> J.Z.Liu]	[14]
<i>Plantago cornuti</i> Gouan	[2, 9, 10]
<i>Plantago depressa</i> Willd.	[1–10, 12, 13]
<i>Plantago komarovii</i> Pavlov	[1, 3, 6, 7, 13]
<i>Plantago major</i> L.	[2–14]
<i>Plantago maritima</i> subsp. <i>ciliata</i> Printz [= <i>Plantago salsa</i> Pall.]	[1–6, 8–11, 13, 14]
<i>Plantago minuta</i> Pall.	[3, 7, 8, 10–15]
<i>Plantago polysperma</i> Kar. & Kir.	[10, 13, 15]
<i>Plantago urvillei</i> Opiz	[2]
<i>Veronica anagallis-aquatica</i> L.	[1–4, 7–16]
<i>Veronica anagalloides</i> Guss.	[10]
<i>Veronica arenosa</i> (Serg.) Boriss. [= <i>Veronica laeta</i> auct. non Kar. & Kir.]	[2, 3, 6, 7, 8, 14]
<i>Veronica beccabunga</i> L.	[3, 7, 8]
<i>Veronica biloba</i> Schreb.	[3, 6, 7, 14, 15]
<i>Veronica ciliata</i> Fisch.	[1, 2, 3, 6, 7]

<i>Veronica daurica</i> Steven	[2, 4, 5, 8, 9]
<i>Veronica densiflora</i> Ledeb.	[2, 7]
<i>Veronica ferganica</i> Popov	[6, 7, 14]
<i>Veronica hispidula</i> Boiss. & Huet [= <i>Veronica pusilla</i> Hohen. & Boiss.]	[7]
<i>Veronica incana</i> L.	[1–11, 13]
<i>Veronica krylovii</i> Schischkin	[9]
<i>Veronica linariifolia</i> Link	[1–5, 8, 9]
<i>Veronica longifolia</i> L.	[1–4, 6, 7, 9, 10]
<i>Veronica macrostemon</i> Bunge	[1, 7]
<i>Veronica oxycarpa</i> Boiss. [= <i>Veronica anagallis-aquatica</i> subsp. <i>oxycarpa</i> (Boiss.) A. Jelen]	[7]
<i>Veronica pinnata</i> subsp. <i>nana</i> Polozhij	[7]
<i>Veronica pinnata</i> L. subsp. <i>pinnata</i>	[2–4, 6–8, 10, 14]
<i>Veronica porphyriana</i> Pavlov	[7, 14]
<i>Veronica sajanensis</i> Printz	[7]
E <i>Veronica sapozhnikovii</i> Kosachev	[7, 14]
<i>Veronica scutellata</i> L.	[7]
<i>Veronica × schmakovii</i> Kosachev	[7]
<i>Veronica × smirnovii</i> Kosachev & D.A.German	[7]
<i>Veronicastrum sibiricum</i> (L.) Pennell	[2, 4, 5, 6, 7, 9]
<i>Veronicastrum tubiflorum</i> (Fisch. & C.A.Mey.) Soják [≡ <i>Veronica tubiflora</i> Fisch. & C.A.Mey.]	[4]

### 85. Plumbaginaceae Juss. (4 genera and 20 taxa)

<i>Armeria maritima</i> subsp. <i>sibirica</i> (Turcz.) Nyman [= <i>Armeria sibirica</i> Turcz.]	
	[3, 7]
<i>Goniolimon callicomum</i> Boiss.	[7, 14]
<i>Goniolimon eximium</i> Boiss.	[7]
<i>Goniolimon krylovii</i> A.V.Grebenjuk	[7, 14]
<i>Goniolimon speciosum</i> Boiss.	[1–4, 6–15]
<i>Limonium aureum</i> (L.) Hill	[3–13, 15, 16]
<i>Limonium bicolor</i> Kuntze	[4, 6, 8, 9, 11–14]
<i>Limonium chrysocomum</i> (Kar. & Kir.) Kuntze	[3, 7, 10, 11, 13–15]
<i>Limonium chrysocomum</i> subsp. <i>semenovii</i> (Herder) Kamelin	[7, 11, 13]
<i>Limonium congestum</i> Kuntze	[6, 7, 10]
<i>Limonium coralloides</i> (Tausch) Lincz.	[6, 10, 14]
<i>Limonium flexuosum</i> Kuntze	[1–4, 6–9, 12, 13]
<i>Limonium gmelini</i> Kuntze	[6, 10, 14]
E <i>Limonium gobicum</i> Ikonn.-Gal.	[12]
E <i>Limonium grubovii</i> Lincz.	[9]
E <i>Limonium klementzii</i> Ikonn.-Gal.	[7, 10, 15]
<i>Limonium myrianthum</i> Kuntze	[14]
<i>Limonium suffruticosum</i> Kuntze	[14]

<i>Limonium tenellum</i> Kuntze	[4, 8, 9, 11–13, 15, 16]
<i>Plumbagella micrantha</i> (Ledeb.) Spach	[3, 4]

## 86. Poaceae Barnhart (58 genera and 229 taxa)

Note: The genus *Stipa* L. was recently revised which included a taxonomic key and species synopsis by Zhao et al (2019).

<i>Achnatherum caragana</i> (Trin.) Nevski [= <i>Stipa conferta</i> Poir.]	[7]
<i>Achnatherum confusum</i> (Litv.) Tzvelev [= <i>Stipa confusa</i> Litv.]	[2, 3, 4]
SE <i>Achnatherum inebrians</i> (Hance) Keng [= <i>Stipa inebrians</i> Hance]	[7, 12, 13, 16]
SE <i>Achnatherum pelliotii</i> (Danguy) Röser & Hamasha [= <i>Stipa pelliotii</i> Danguy]	[7, 11–16]
<i>Achnatherum sibiricum</i> (L.) Keng [= <i>Stipa sibirica</i> (L.) Lam.]	[1–13]
<i>Aeluropus littoralis</i> (Gouan) Parl.	[10, 14, 15]
<i>Agropyron cristatum</i> (L.) Gaertn. [= <i>Bromus cristatus</i> L.]	[1–7, 9–15]
<i>Agropyron desertorum</i> Schult.	[3, 6–9, 11–13, 15]
<i>Agropyron fragile</i> (Roth) P.Candargy	[6, 8, 9, 11, 12, 14]
<i>Agropyron krylovianum</i> Schischk. [= <i>Kengyilia kryloviana</i> (Schischk.) C.Yen, J.L.Yang & B.R.Baum]	[4, 6, 7]
<i>Agropyron michnoi</i> Roshev. [= <i>Agropyron cristatum</i> (L.) Gaertn. subsp. <i>michnoi</i> (Roshev.) Å.Löve]	[1, 3–5, 8–10, 13, 14]
<i>Agropyron pumilum</i> (Steud.) P.Candargy	[6, 10, 14]
SE <i>Agrostis clavata</i> Trin. [= <i>Agrostis exarata</i> subsp. <i>clavata</i> (Trin.) T.Koyama]	[1, 2, 3, 4, 5, 9]
<i>Agrostis divaricatissima</i> Mez [= <i>Agrostis mongolica</i> Roshev.]	[2–13, 15]
<i>Agrostis gigantea</i> Roth	[1–7, 9, 10, 13, 14]
<i>Agrostis stolonifera</i> L.	[1, 3, 4, 7–10, 14]
SE <i>Agrostis tuvinica</i> Peschkova	[1, 2]
<i>Agrostis vinealis</i> Schreb. [= <i>Agrostis trinii</i> Turcz.]	[1–10, 13]
<i>Alopecurus aequalis</i> Sobol.	[1–10, 12]
<i>Alopecurus arundinaceus</i> Poir.	[1–10, 12–14]
<i>Alopecurus brachystachys</i> M.Bieb.	[1–10, 13]
SE <i>Alopecurus pratensis</i> L.	[3, 4, 6, 8–10, 13, 14]
<i>Alopecurus turczaninovii</i> O.D.Nikif.	[1–4, 6–8, 13]
<i>Anthoxanthum glabrum</i> (Trin.) Veldkamp [= <i>Hierochloe glabra</i> Trin.]	[1–10]
<i>Anthoxanthum monticola</i> (Bigelow) Veldkamp [= <i>Holcus monticola</i> Bigelow]	[1–3, 6, 7, 10]
<i>Anthoxanthum nitens</i> (Weber) Y.Schouten & Veldkamp [= <i>Poa nitens</i> Weber = <i>Hierochloe odorata</i> (L.) P.Beauv.]	[1–4, 6–10, 14]
<i>Anthoxanthum odoratum</i> L.	[2, 3, 4, 5, 7]
<i>Arctagrostis latifolia</i> Griseb.	[1, 2, 3]
<i>Arctopoa schischkinii</i> (Tzvelev) Prob. [= <i>Poa schischkinii</i> Tzvelev]	[1, 7, 8]
<i>Arctopoa tibetica</i> (Munro) Prob. [= <i>Poa tibetica</i> Munro]	[1, 3, 6–8, 10–15]

	<i>Aristida adscensionis</i> L.	[3, 8–13, 15, 16]
	<i>Arundinella hirta</i> (Thunb.) Tanaka [= <i>Poa hirta</i> Thunb.]	[4, 5, 9]
	<i>Beckmannia syzigachne</i> Fernald	[1–6, 8–16]
	<i>Brachypodium pinnatum</i> (L.) P.Beauv. [= <i>Bromus pinnatus</i> L.]	[3, 4]
	<i>Bromus inermis</i> Leyss.	[1–10, 12–15]
	<i>Bromus japonicus</i> Thunb.	[6, 7, 10, 12, 14]
	<i>Bromus oxyodon</i> Schrenk	[7, 11, 14]
	<i>Bromus pumpellianus</i> Scribn. [= <i>Bromus korotkiji</i> Drobow]	[3, 4, 7–11, 13]
	<i>Bromus scoparius</i> L.	[14]
	<i>Bromus squarrosum</i> L.	[10]
	<i>Bromus tectorum</i> L.	[7, 14]
	<i>Calamagrostis angustifolia</i> Kom. [= <i>Calamagrostis angustifolia</i> subsp. <i>tenuis</i> (V.N.Vassil.) Tzvelev]	[1–11, 13, 14]
	<i>Calamagrostis epigejos</i> (L.) Roth [= <i>Calamagrostis epigejos</i> subsp. <i>glomerata</i> (Boiss. & Buhse) Tzvelev]	[2, 4, 5, 9]
	<i>Calamagrostis inexpansa</i> A.Gray [= <i>Calamagrostis inexpansa</i> subsp. <i>micrantha</i> (Kearney) Stebbins]	[2, 3, 4, 5]
	<i>Calamagrostis korotkyi</i> Litv. [= <i>Deyeuxia korotkyi</i> (Litv.) S.M.Phillips & W.L.Chen]	[1, 2, 3, 4, 6]
SE	<i>Calamagrostis × kuznetzovii</i> Tzvelev	[4]
	<i>Calamagrostis lapponica</i> (Wahlenb.) Hartm. [= <i>Arundo lapponica</i> Wahlenb.]	[1–4, 6–8, 10, 13–15]
	<i>Calamagrostis macilenta</i> Litv.	[1–4, 7–13, 15]
	<i>Calamagrostis macrolepis</i> Litv. [= <i>Calamagrostis epigejos</i> subsp. <i>macrolepis</i> (Litv.) Tzvelev]	[2, 3, 4]
	<i>Calamagrostis obtusata</i> Trin.	[3, 4]
	<i>Calamagrostis pavlovii</i> (Roshev.) Roshev.	[1–4, 7, 8, 10, 11]
	<i>Calamagrostis pseudophragmites</i> (Haller f.) Koeler [= <i>Arundo pseudophragmites</i> Haller f.]	[1–7, 9]
	<i>Calamagrostis purpurea</i> (Trin.) Trin. [= <i>Arundo purpurea</i> Trin.]	[2, 3, 4, 10]
SE	<i>Calamagrostis sajanensis</i> Malyschev	[8, 9, 10, 13]
	<i>Calamagrostis salina</i> Tzvelev	[1–5, 8, 9, 13]
	<i>Calamagrostis stricta</i> (Timm) Koeler [= <i>Arundo stricta</i> Timm]	[2]
	<i>Catabrosa aquatica</i> (L.) P.Beauv.	[2–4, 8–10, 13]
	<i>Cenchrus flaccidus</i> (Griseb.) Morrone [= <i>Pennisetum flaccidum</i> Griseb.]	[10, 12]
SE	<i>Cinna latifolia</i> (Trevir.) Griseb. [= <i>Agrostis latifolia</i> Trevir.]	[2, 3, 4]
	<i>Cleistogenes caespitosa</i> Keng	[12]
SE	<i>Cleistogenes festucacea</i> Honda [= <i>Cleistogenes foliosa</i> Keng]	[4, 12, 13]
	<i>Cleistogenes kitagawae</i> Honda [= <i>Kengia kitagawae</i> (Honda) Packer]	[2–5, 8, 9]
	<i>Cleistogenes songorica</i> (Roshev.) Ohwi	[4, 7–16]
	<i>Cleistogenes squarrosa</i> (Trin.) Keng	[2–13]

	<i>Colpodium altaicum</i> Trin.	[1, 7]
	<i>Deschampsia caespitosa</i> P.Beauv. subsp. <i>caespitosa</i>	[1–4, 6, 7, 8]
	<i>Deschampsia caespitosa</i> subsp. <i>orientalis</i> Hultén	[1, 2, 3, 4, 5]
	<i>Deschampsia caespitosa</i> subsp. <i>pamirica</i> (Roshev.) Tzvelev [≡ <i>Deschampsia pamirica</i> Roshev.]	[1, 3]
	<i>Deschampsia koelerioides</i> Regel	[1, 2, 3, 6, 7]
	<i>Echinochloa crus-galli</i> (L.) P.Beauv.	[4, 7, 9, 12, 13]
	<i>Elymus bungeanus</i> (Trin.) Melderis	[2, 4, 6, 7, 12, 14, 15]
	<i>Elymus confusus</i> (Roshev.) Tzvelev [≡ <i>Roegneria confusa</i> (Roshev.) Nevski]	[1–4, 7, 8, 13]
	<i>Elymus dahuricus</i> Turcz.	[1–10, 13]
	<i>Elymus fedtschenkoi</i> Tzvelev [≡ <i>Roegneria fedtschenkoi</i> (Tzvelev) J.L.Yang & C.Yen]	[7]
SE	<i>Elymus gmelinii</i> (Ledeb.) Tzvelev	[1–5, 7, 8, 9, 13]
	<i>Elymus karakabinicus</i> Kotukhov	[7]
	<i>Elymus macrorurus</i> (Turcz.) Tzvelev [= <i>Elymus kronokensis</i> (Kom.) Tzvelev, <i>Elymus kronokensis</i> subsp. <i>subalpinus</i> (Neuman) Tzvelev]	[1, 4, 7]
	<i>Elymus mutabilis</i> (Drobow) Tzvelev [= <i>Elymus transbaicalensis</i> (Nevski) Tzvelev = <i>Elymus praecaespitosus</i> (Nevski) Tzvelev]	[1, 2, 3, 4, 6, 7]
	<i>Elymus nutans</i> Griseb.	[3, 4, 9, 13]
	<i>Elymus pendulinus</i> (Nevski) Tzvelev [= <i>Agropyron vernicosum</i> Nevski = <i>Elymus brachypodioides</i> (Nevski) Peschkova]	[4, 5, 7, 9, 12, 13, 16]
	<i>Elymus reflexiaristatus</i> (Nevski) Melderis [= <i>Elymus aegilopoides</i> (Drobow) Vorosch.]	[1–7, 12, 13, 14]
	<i>Elymus repens</i> (L.) Gould [= <i>Elytrigia repens</i> (L.) Nevski]	[2–11, 13–15]
	<i>Elymus schrenkianus</i> (Fisch. & C.A.Mey.) Tzvelev [= <i>Elymus pamiricus</i> Tzvelev]	[3, 4, 7, 13]
	<i>Elymus sibiricus</i> L.	[1–10, 12–16]
	<i>Elymus uralensis</i> (Nevski) Tzvelev [= <i>Elymus uralensis</i> subsp. <i>komarovii</i> (Nevski) Tzvelev]	[2, 3, 4, 6, 13]
	<i>Elymus varius</i> (Keng) Tzvelev	[4]
	<i>Enneapogon desvauxii</i> P.Beauv.	[3, 4, 6–15]
	<i>Eragrostis ciliaris</i> (All.) Vignolo	[10]
	<i>Eragrostis minor</i> Host	[2, 3, 4, 6–16]
	<i>Eragrostis pilosa</i> (L.) P.Beauv. [= <i>Eragrostis pilosa</i> subsp. <i>imberbis</i> (Franch.) Tzvelev]	[2, 4, 7–10, 12]
	<i>Eremopyrum distans</i> (K.Koch) Nevski	[14]
	<i>Festuca altaica</i> Trin.	[1, 3, 4, 6, 7]
	<i>Festuca brachyphylla</i> Schult. & Schult.f.	[1, 3, 6, 7, 13]
	<i>Festuca dahurica</i> V.I.Krecz. & Bobr.	[2, 4, 5, 9]
	<i>Festuca extremiorientalis</i> Ohwi	[2]
	<i>Festuca hubsugulica</i> Krivot. [= <i>Festuca sumneviczii</i> Serg.]	[1]
SE	<i>Festuca jacutica</i> Drobow	[4, 9]

	<i>Festuca komarovii</i> Krivot.	[1, 2]
	<i>Festuca kryloviana</i> Reverd.	[1–4, 6, 7, 9, 13]
	<i>Festuca kurtschumica</i> E.B.Alexeev	[7]
	<i>Festuca lenensis</i> Drobow	[1–9, 13, 15]
	<i>Festuca litwinovii</i> (Tzvelev) E.B.Alexeev	[9]
	<i>Festuca oreophila</i> Markgr.-Dann. [= <i>Festuca valesiaca</i> subsp. <i>hypspila</i> (St.-Yves) Tzvelev]	[1–4, 6, 7, 13, 14]
	<i>Festuca ovina</i> L. [= <i>Festuca ovina</i> subsp. <i>sphagnicola</i> (B.Keller) Tzvelev]	[1–7, 9, 13]
SE	<i>Festuca pseudosulcata</i> Drobow	[4]
	<i>Festuca rubra</i> L.	[1–9, 13]
	<i>Festuca sibirica</i> Hack.	[1–5, 8–10, 13]
SE	<i>Festuca tristis</i> Krylov & Ivanitzk.	[3, 6, 7]
	<i>Festuca tschujensis</i> Reverd.	[3, 6, 7, 10]
	<i>Festuca valesiaca</i> Schleich.	[1–10, 13]
	<i>Festuca venusta</i> St.-Yves	[1, 2, 3, 4]
	<i>Glyceria arundinacea</i> Kunth	[1–10, 14]
	<i>Glyceria lithuanica</i> (Gorski) Gorski	[3, 5]
	<i>Glyceria spiculosa</i> Roshev. [= <i>Glyceria longiglumis</i> Hand.-Mazz.]	[4, 5, 9]
	<i>Helictochloa dahurica</i> (Kom.) Romero Zarco [≡ <i>Helictotrichon dahuricum</i> (Kom.) Kitag.]	[1, 4, 5]
	<i>Helictochloa hookeri</i> (Scribn.) Romero Zarco [≡ <i>Helictotrichon schellianum</i> (Hack.) Kitag.]	[1–9, 13]
	<i>Helictotrichon desertorum</i> (Less.) Pilg. [= <i>Helictotrichon altaicum</i> Tzvelev]	[3, 6, 7]
	<i>Helictotrichon mongolicum</i> (Roshev.) Henrard [≡ <i>Avenastrum mongolicum</i> (Roshev.) Roshev.]	[1–3, 6, 7, 13]
	<i>Helictotrichon pubescens</i> (Huds.) Pilg. [= <i>Hordeum brevisubulatum</i> subsp. <i>turkestanicum</i> (Nevski) Tzvelev]	[3, 4, 7, 9]
	<i>Hordeum bogdanii</i> Wilensky	[3, 7, 9, 10, 14–16]
	<i>Hordeum brevisubulatum</i> Link [= <i>Hordeum brevisubulatum</i> subsp. <i>turkestanicum</i> (Nevski) Tzvelev]	[1–16]
	<i>Hordeum roshevitzii</i> Bowden [≡ <i>Critesion roshevitzii</i> (Bowden) Tzvelev]	[1, 2, 4, 8–11, 13]
	<i>Koeleria altaica</i> (Domin) Krylov	[1–10, 13]
	<i>Koeleria asiatica</i> Domin	[2, 8]
	<i>Koeleria glauca</i> DC.	[4]
	<i>Koeleria litvinowii</i> Domin	[1, 3, 7, 13]
	<i>Koeleria macrantha</i> (Ledeb.) Schult.	[1–5, 8–13]
	<i>Koeleria spicata</i> subsp. <i>mongolica</i> (Hultén) Barberá, Quintanar, Soreng, & P.M.Peterson [≡ <i>Trisetum spicatum</i> subsp. <i>mongolicum</i> Hultén]	[1–4, 6, 7, 13]

SE	<i>Koeleria thonii</i> Domin	[5]
	<i>Leymus angustus</i> (Trin.) Pilg. [= <i>Elymus angustus</i> Trin.]	[3, 6, 7, 10–14, 16]
	<i>Leymus chinensis</i> (Trin.) Tzvelev	[1–6, 8–14]
SE	<i>Leymus ordensis</i> Peschkova	[15]
	<i>Leymus paboanus</i> (Claus) Pilg.	[3, 6–8, 10–14]
	<i>Leymus racemosus</i> (Lam.) Tzvelev	[3, 5, 6, 8–13]
	<i>Leymus ramosus</i> (K.Richt.) Tzvelev [= <i>Agropyron ramosum</i> K.Richt.]	[5, 9]
	<i>Leymus secalinus</i> (Georgi) Tzvelev [= <i>Leymus secalinus</i> var. <i>mongolicus</i> (Meld.) Tzvelev = <i>Leymus ovatus</i> (Trin.) Tzvelev]	[1–4, 6–16]
	<i>Melica nutans</i> L.	[1, 2, 4]
	<i>Melica transsilvanica</i> Schur	[7]
	<i>Melica turczaninowiana</i> Ohwi	[2, 3, 4, 5, 9]
	<i>Melica virgata</i> Turcz.	[1–5, 8–10, 12, 13]
	<i>Milium effusum</i> L.	[2]
	<i>Nardus stricta</i> L.	[3]
	<i>Neotrinia splendens</i> (Trin.) M.Nobis [≡ <i>Achnatherum splendens</i> (Trin.) Nevski ≡ <i>Stipa splendens</i> Trin.]	[2–5, 7–16]
	<i>Phalaris arundinacea</i> L.	[1–5, 7, 9, 10, 14]
	<i>Phleum alpinum</i> L.	[7]
	<i>Phleum phleoides</i> (L.) H.Karst.	[2–4, 6–8, 10]
	<i>Phragmites australis</i> (Cav.) Steud.	[1–16]
	<i>Piptatherum songaricum</i> (Trin. & Rupr.) Roshev.	[7, 14]
	<i>Poa alpina</i> L.	[1, 3, 6, 7, 15]
	<i>Poa alta</i> Hitchc. [= <i>Poa mongolica</i> (Rendle) Keng]	[2, 5]
	<i>Poa altaica</i> Trin. [= <i>Poa glauca</i> subsp. <i>altaica</i> (Trin.) Olonova & G.H.Zhu]	[1–3, 6, 7, 10, 13]
	<i>Poa angustifolia</i> L. [= <i>Poa pratensis</i> subsp. <i>angustifolia</i> (L.) Dumort.]	[2, 3, 4, 5, 9]
	<i>Poa annua</i> L. [= <i>Ochlopoa annua</i> (L.) H.Scholz]	[2, 7, 9]
	<i>Poa argunensis</i> Roshev.	[1–11, 13, 15]
	<i>Poa attenuata</i> Trin. subsp. <i>attenuata</i> .	[1–4, 6–10, 12–14]
	<i>Poa attenuata</i> subsp. <i>botryoides</i> (Trin.) Tzvelev	[1, 3–9, 13]
	<i>Poa attenuata</i> subsp. <i>dahurica</i> (Trin.) Cubanov [= <i>Poa dahurica</i> Trin.]	[1–4, 7–13]
	<i>Poa attenuata</i> subsp. <i>tshuensis</i> (Serg.) Olonova [≡ <i>Poa argunensis</i> f. <i>tshuensis</i> Serg.]	[1–4, 6–10, 12–15]
	<i>Poa glauca</i> Vahl	[1, 3, 6, 7, 13]
	<i>Poa ircutica</i> Roshev.	[4]
SE	<i>Poa kenteica</i> Ivanova	[2, 3]
	<i>Poa krylovii</i> Reverd. [= <i>Poa urssulensis</i> subsp. <i>krylovii</i> (Reverd.) Olonova]	[2, 3, 4, 7, 8]
	<i>Poa nemoralis</i> L.	[1–5, 8, 9]

	<i>Poa palustris</i> L.	[1–5, 7, 8, 9, 13]
	<i>Poa pratensis</i> L. [= <i>Poa pratensis</i> subsp. <i>sabulosa</i> (Turcz.) Tzvelev]	[1–10, 13, 14]
	<i>Poa raduliformis</i> Prob.	[2, 7]
	<i>Poa sibirica</i> Roshevitz	[1–7, 10, 13]
	<i>Poa smirnowii</i> Roshev.	[1, 3, 4, 6, 7]
	<i>Poa subfastigiata</i> Trin.	[1–6, 8–10, 13]
	<i>Poa supina</i> Schrad.	[3, 4, 7]
	<i>Poa tianschanica</i> Hack.	[1–10, 13]
	<i>Poa trivialis</i> L.	[1]
	<i>Poa urssulensis</i> Trin.	[2, 3, 5]
	<i>Poa veresczaginii</i> Tzvelev	[7]
	<i>Poa versicolor</i> subsp. <i>reverdattoi</i> (Roshev.) Olonova & G.H.Zhu [≡ <i>Poa reverdattoi</i> Roshev.]	[1, 3, 5–10, 13, 14]
	<i>Poa versicolor</i> Besser subsp. <i>versicolor</i> [= <i>Poa versicolor</i> subsp. <i>stepposa</i> (Krylov) Tzvelev]	[1–9, 11, 13]
SE	<i>Polypogon maritimus</i> Willd.	[11, 15]
	<i>Polypogon monspeliensis</i> (L.) Desf.	[7, 10, 11, 13, 15, 16]
	<i>Psammochloa villosa</i> (Trin.) Bor	[3, 9–13, 16]
	<i>Psathyrostachys juncea</i> (Fisch.) Nevski	[3, 6, 7, 10, 12–15]
	<i>Psathyrostachys lanuginosus</i> (Trin.) Nevski	[7, 14]
	<i>Ptilagrostis mongholica</i> (Turcz.) Griseb. [≡ <i>Stipa mongholica</i> Turcz.]	[1, 2, 3, 4, 7]
	<i>Puccinellia altaica</i> Tzvelev	[14]
SE	<i>Puccinellia distans</i> (Jacq.) Parl.	[11, 13, 14]
	<i>Puccinellia filifolia</i> (Trin.) Tzvelev	[8, 12]
	<i>Puccinellia hackeliana</i> (V.I.Krecz.) V.I.Krecz.	[7]
	<i>Puccinellia hauptiana</i> (V.I.Krecz.) Kitag.	[3–11, 13, 14]
	<i>Puccinellia macranthera</i> V.I.Krecz.	[2–5, 7–10, 12, 14]
	<i>Puccinellia manchuriensis</i> Ohwi	[3]
	<i>Puccinellia nudiflora</i> (Hack.) Tzvelev	[7, 13]
SE	<i>Puccinellia przewalskii</i> Tzvelev	[10]
	<i>Puccinellia schischkinii</i> Tzvelev	[5, 8, 10, 12, 13, 15]
	<i>Puccinellia tenuiflora</i> Scribn. & Merr. [= <i>Puccinellia kreczetoviczii</i> Bubnova]	[1–16]
	<i>Schismus arabicus</i> Nees	[4, 7, 11, 14]
	<i>Schizachne purpurascens</i> subsp. <i>callosa</i> (Turcz.) T.Koyama & Kawano	[2, 3, 4, 5]
	<i>Scolochloa festucacea</i> Link	[4]
	<i>Sibirotrisetum sibiricum</i> (Rupr.) Barberá [≡ <i>Trisetum sibiricum</i> Rupr.]	[1–5, 7–10, 13]
	<i>Spodiopogon sibiricus</i> Trin.	[2, 3, 4, 5, 8, 9]
	<i>Sporobolus aculeatus</i> (L.) P.M.Peterson [≡ <i>Crypsis aculeata</i> (L.) Aiton]	[10–13, 15]
	<i>Sporobolus schoenoides</i> (L.) P.M.Peterson [≡ <i>Crypsis schoenoides</i> Lam.]	[3, 10, 14]

E	<i>Stipa austromongolica</i> M.Nobis	[10]
	<i>Stipa baicalensis</i> Roshev.	[1–5, 7, 8, 9, 12]
	<i>Stipa breviflora</i> Griseb.	[7, 8, 9, 12, 13]
	<i>Stipa capillata</i> L.	[3, 4, 7, 10, 12]
	<i>Stipa caucasica</i> Schmalh. subsp. <i>caucasica</i>	[7, 10–14, 16]
	<i>Stipa caucasica</i> subsp. <i>desertorum</i> (Roshev.) Tzvelev	[10]
	<i>Stipa consanguinea</i> Trin. & Rupr.	[2, 7, 10]
	<i>Stipa glareosa</i> f. <i>pubescens</i> P.A.Smirn.	[4–12]
	<i>Stipa glareosa</i> P.A.Smirn. [= <i>Stipa caucasica</i> subsp. <i>glareosa</i> (P.A.Smirn.) Tzvelev]	[3, 6–16]
	<i>Stipa gobica</i> Roshev.	[3, 4, 6–16]
	<i>Stipa grandis</i> P.A.Smirn.	[3, 4, 5, 8, 9]
E	<i>Stipa khovdensis</i> L.Q.Zhao	[3, 6]
	<i>Stipa kirghisorum</i> P.A.Smirn.	[3, 7, 10, 14]
	<i>Stipa klemenzii</i> Roshev.	[3, 4, 6, 8, 9–13]
	<i>Stipa krylovii</i> Roshev.	[1–14]
	<i>Stipa mongolorum</i> Tzvelev	[7, 8, 10, 11, 12]
	<i>Stipa orientalis</i> Trin.	[1, 3, 6, 7, 10, 11, 14]
	<i>Stipa pennata</i> L. subsp. <i>pennata</i>	[3, 4]
	<i>Stipa pennata</i> subsp. <i>sabulosa</i> (Pacz.) Tzvelev	[10]
	<i>Stipa sareptana</i> A. Beck.	[3, 7, 10, 11, 13]
	<i>Stipa sczerbakovii</i> Kotuch.	[7]
	<i>Stipa tianschanica</i> Roshev. [= <i>Stipa tianschanica</i> subsp. <i>gobica</i> (Roshev.) D.F. Cui = <i>Stipa tianschanica</i> var. <i>klemenzii</i> (Roshev.) Norl.]	[3, 4, 6–16]
	<i>Stipa zalesskii</i> Wilensky	[3, 7, 10]
	<i>Timouria saposhnikowii</i> Roshev. [= <i>Stipa saposhnikowii</i> (Roshev.) Kitag.]	[13, 15]
	<i>Tragus mongolorum</i> Ohwi	[12, 13]
	<i>Tripogon chinensis</i> Hack.	[2, 4, 5, 8, 9, 12]
	<i>Tripogon purpurascens</i> Duthie	[12, 13, 16]
	<i>Trisetum altaicum</i> Roshev.	[1, 2, 3, 6, 7]
	<i>Zizania latifolia</i> Turcz.	[9]

#### 87. Polemoniaceae Juss. (2 genera and 4 species)

<i>Phlox sibirica</i> L.	[1, 4]
<i>Polemonium boreale</i> Adams	[1, 3, 6]
<i>Polemonium chinense</i> Brand	[1–7, 9]
<i>Polemonium pulchellum</i> Bunge	[1, 3, 6]

#### 88. Polygalaceae Hoffmanns. & Link (1 genus and 3 species)

<i>Polygala comosa</i> Schkuhr [= <i>Polygala hybrida</i> DC.]	[1–4, 6, 7, 14]
<i>Polygala sibirica</i> L.	[1, 2, 3, 4, 5, 9]
<i>Polygala tenuifolia</i> Willd.	[1–5, 8, 9, 12, 13]

89. **Polygonaceae** Juss. (11 genera and 63 taxa)

	<i>Atraphaxis bracteata</i> Losinsk.	[3, 6, 7, 9, 10–16]
	<i>Atraphaxis compacta</i> Ledeb.	[13, 14, 15]
E	<i>Atraphaxis frutescens</i> (L.) K.Koch	[3, 5, 6, 7, 9–16]
	<i>Atraphaxis kamelinii</i> Yurtseva	[14]
	<i>Atraphaxis pungens</i> Jaub. & Spach	[2–16]
	<i>Atraphaxis spinosa</i> L.	[14, 15]
	<i>Atraphaxis virgata</i> (Regel) Krassn.	[7, 10, 12, 14–16]
SE	<i>Bistorta elliptica</i> (Willd.) V.V.Petrovsky	[1, 2, 3, 6, 7]
	<i>Calligonum ebinuricum</i> Ivanova	[14, 15]
	<i>Calligonum junceum</i> (Fisch. & C.A.Mey.) Litv.	[14, 15]
	<i>Calligonum litwinowi</i> Drobow [= <i>Calligonum gobicum</i> Losinsk.]	[14, 15]
	<i>Calligonum mongolicum</i> Turcz.	[7, 10–16]
	<i>Fallopia convolvulus</i> (L.) Á.Löve [= <i>Polygonum convolvulus</i> L.]	[2 – 5 , 8–10, 12, 15]
	<i>Fallopia dumetorum</i> (L.) Holub	[5]
	<i>Knorringia sibirica</i> (Laxm.) Tzvelev subsp. <i>sibirica</i>	[1–16]
	<i>Knorringia sibirica</i> subsp. <i>ubsunurica</i> Tzvelev	[10]
	<i>Koenigia islandica</i> L.	[1–3, 6, 7, 10, 13]
	<i>Oxyria digyna</i> Hill	[1, 2, 3, 6, 7, 13]
	<i>Persicaria alpina</i> Gross.	[1–4, 6, 7, 8, 14]
	<i>Persicaria amphibia</i> (L.) Delarbre	[1, 3–12, 14]
	<i>Persicaria bistorta</i> Samp.	[2]
	<i>Persicaria bungeana</i> Nakai	[1–3, 5, 7–10, 13]
	<i>Persicaria hydropiper</i> (L.) Delarbre [= <i>Polygonum hydropiper</i> L.]	[2 – 4 , 6–8, 10, 13, 14]
	<i>Persicaria lapathifolia</i> (L.) Delarbre [= <i>Polygonum ochreatum</i> L.]	[1–16]
	<i>Persicaria longiseta</i> var. <i>rotundata</i> (A.J.Li) B.Li [	
	≡ <i>Polygonum longisetum</i> var. <i>rotundatum</i> A.J.Li ]	[2, 4, 8]
	<i>Persicaria minor</i> (Huds.) Opiz	[10, 14]
	<i>Persicaria sagittata</i> (L.) H.Gross	[2, 3, 4, 5, 8, 9]
	<i>Persicaria vivipara</i> (L.) Ronse Decr.	[1–4, 6–8, 10, 13, 14]
	<i>Polygonum abbreviatum</i> Kom.	[1, 2, 7]
	<i>Polygonum alopecuroides</i> Turcz.	[1–6, 8]
	<i>Polygonum angustifolium</i> Pall.	[1–5, 7–9, 11, 13]
	<i>Polygonum arenastrum</i> Boreau	[1, 3, 7, 10, 14]
	<i>Polygonum argyrocoleon</i> Steud.	[7, 10, 11, 13–15]
	<i>Polygonum aviculare</i> L.	[1–5, 7–14, 16]
	<i>Polygonum cognatum</i> Meisn.	[3, 4, 6, 7, 8, 10]
	<i>Polygonum divaricatum</i> L.	[1–5, 8, 9]
	<i>Polygonum ellipticum</i> Willd.	[1, 2, 3, 6, 7]
	<i>Polygonum humifusum</i> C.Merck	[3]
	<i>Polygonum intramongolicum</i> Borodina	[12, 13]

<i>Polygonum novoascanicum</i> Klokov	[14]
<i>Polygonum patulum</i> M.Bieb.	[3, 7, 9, 10, 14, 15]
<i>Polygonum polycnemoides</i> Jaub. & Spach	[7, 14]
<i>Polygonum sericeum</i> Pall.	[2, 3, 4, 8, 9]
<i>Polygonum tenuissimum</i> A.I.Baranov & Skvortsov	[9]
<i>Polygonum valerii</i> A.K.Skvortsov	[2, 4, 5, 8]
<i>Polygonum volchovense</i> Tzvelev	[7]
<i>Rheum compactum</i> L.	[1–4, 6, 7, 12–14]
<i>Rheum nanum</i> Siev.	[7, 8, 10–16]
<i>Rheum rhabarbarum</i> L. [= <i>Rheum undulatum</i> L.]	[1–5, 7–9, 12–14]
SE <i>Rheum uninerve</i> Maxim.	[13]
<i>Rumex acetosa</i> L.	[1, 2, 3, 6, 7]
<i>Rumex acetosella</i> L.	[1–5, 8, 9]
<i>Rumex aquaticus</i> L.	[1–10, 14]
<i>Rumex crispus</i> L.	[1, 7, 9, 10, 14]
<i>Rumex gmelinii</i> Turcz.	[2, 3, 4, 5, 8, 9]
<i>Rumex maritimus</i> L.	[2–5, 8–11, 14]
<i>Rumex marschallianus</i> Rchb.	[6, 8, 9, 10, 11]
<i>Rumex patientia</i> L.	[7, 9, 13, 14]
<i>Rumex popovii</i> Pachom.	[10, 13]
<i>Rumex pseudonatronatus</i> (Borbás) Murb.	[11, 13]
<i>Rumex similans</i> Rech.f.	[2–4, 6, 7, 10–12, 14, 15]
<i>Rumex stenophyllus</i> Ledeb.	[2–4, 6, 7, 9, 10, 14, 15]
<i>Rumex thysiflorus</i> Fingerh.	[1–14]

## 90. Potamogetonaceae Bercht. & J.Presl (3 genera and 18 taxa)

<i>Potamogeton angustifolius</i> Bercht. & J.Presl	[10]
<i>Potamogeton alpinus</i> subsp. <i>tenuifolius</i> (Raf.) Hulten	[3, 4, 9, 10, 12]
<i>Potamogeton berchtoldii</i> Fieber	[3, 6, 7, 9]
<i>Potamogeton compressus</i> L.	[1, 8, 9, 10, 14]
<i>Potamogeton crispus</i> L.	[1, 8, 10]
<i>Potamogeton friesii</i> Rupr.	[5, 8, 9]
<i>Potamogeton gramineus</i> L.	[1–10, 14]
<i>Potamogeton lucens</i> L.	[1, 8, 10]
<i>Potamogeton mandschuriensis</i> A.Benn.	[5]
<i>Potamogeton natans</i> L.	[1, 5–7, 10, 11]
<i>Potamogeton obtusifolius</i> Mert. & W.D.J.Koch	[1, 3, 5]
<i>Potamogeton perfoliatus</i> L.	[1–12, 14, 16]
<i>Potamogeton praelongus</i> F.Muell.	[1–5, 8, 9]
<i>Potamogeton pusillus</i> L.	[1–6, 8–11, 13, 14]
<i>Stuckenia filiformis</i> (Pers.) Börner [= <i>Potamogeton filiformis</i> Pers.]	[1, 3, 6–11, 13, 15]

<i>Stuckenia pectinata</i> (L.) Börner [= <i>Potamogeton pectinatus</i> L.]	[1–10, 11]
<i>Stuckenia vaginata</i> Holub	[1, 3–8, 10, 11, 13]
<i>Zannichellia palustris</i> L. [= <i>Zannichellia palustris</i> subsp. <i>pedicellata</i> (Rosén & Wahlenb.) Hook.f.]	[3, 4, 7–11, 15]

## 91. Primulaceae Batsch (3 genera and 28 taxa)

Note: The genus *Primula* L. was recently revised by Baasanmunkh et al. (2020a).

<i>Androsace fedtschenkoi</i> Ovcz.	[1, 6, 7, 13]
<i>Androsace filiformis</i> Retz.	[1, 2, 3, 4, 5, 9]
<i>Androsace gmelinii</i> Gaertn.	[2, 3, 9, 11]
<i>Androsace incana</i> Lam.	[1–9, 13]
<i>Androsace lactiflora</i> Fisch. [= <i>Androsace amurensis</i> Prob.]	[1–4, 6, 7, 9, 14]
<i>Androsace lemanniana</i> Spreng. [= <i>Androsace bungeana</i> Schischk. & Bobrov]	[1, 2, 3, 6, 7, 9]
<i>Androsace longifolia</i> Turcz.	[5, 9]
<i>Androsace maxima</i> L.	[2–4, 6–10, 13–15]
<i>Androsace ovczinnikovii</i> Schischk. & Bobrov	[3, 6, 7]
<i>Androsace septentrionalis</i> L.	[1–9, 12–14]
<i>Androsace villosa</i> L. var. <i>dasyphylla</i> (Bunge) Kar. & Kir. [= <i>Androsace dasyphylla</i> Bunge]	[1, 2, 3, 6, 7, 13]
<i>Lysimachia davurica</i> Ledeb.	[2, 3, 4, 5, 9, 14]
<i>Lysimachia europaea</i> (L.) U.Manns & Anderb. [= <i>Trientalis europaea</i> L.]	[1, 2, 3, 4, 5]
<i>Lysimachia maritima</i> (L.) Galasso [= <i>Glaux maritima</i> L.]	[1–16]
<i>Lysimachia thyrsiflora</i> L. [= <i>Naumburgia thyrsiflora</i> (L.) Rchb.]	[2–5, 9, 10, 14]
<i>Primula algida</i> Adams	[1–3, 6, 7, 11, 13]
<i>Primula bukukunica</i> Kovt.	[7, 11, 13]
<i>Primula cortusoides</i> L.	[3]
<i>Primula farinosa</i> L.	[1–4, 6, 7, 10, 13, 15]
<i>Primula longiscapa</i> Ledeb.	[3, 6, 7, 10, 13, 14]
<i>Primula matthioli</i> subsp. <i>altaica</i> (Losinsk.) Kovt. [= <i>Cortusa altaica</i> Losinsk.]	[1, 2, 3, 7, 13, 14]
<i>Primula matthioli</i> subsp. <i>brotheri</i> (Pax) Kovt. [= <i>Cortusa matthioli</i> f. <i>brotheri</i> Pax]	[7]
<i>Primula maximowiczii</i> Regel	[5]
<i>Primula nivalis</i> subsp. <i>nivalis</i> Pall.	[1, 2, 3, 6, 7, 10]
<i>Primula nivalis</i> subsp. <i>turkestanica</i> (J.N.Haage & E.Schmidt) Kovt.	[1, 3]
<i>Primula nivalis</i> subsp. <i>xanthobasis</i> (Fed.) Halda	[1, 2, 3]
<i>Primula nutans</i> Georgi	[1–7, 9, 10, 11]
<i>Primula serrata</i> Georgi	[1–5, 7–10, 13]

## 92. Ranunculaceae Juss. (20 genera and 156 taxa)

Note: In regards to the phylogeny and position of *Actaea* and *Cimicifuga*, we follow Compton et al. (1998). *Anemone* is considered here according to Hoot (2012) with some changes leaving *Pulsatilla* at generic level (Sramkó et al. 2019). The taxonomy of the tribe Ranunculae is given according to Emadzade (2010) and Wang et al. (2014). The taxonomical positions of some taxa have been changed according to taxonomic works (Solovjev 1998; Erst 2007). Previously, four species of *Batrachium* were recorded in Mongolia which are treated as a synonym of *Ranunculus* by Wieglob et al. (2017). Taxonomic revision of the genus *Aquilegia* L. was carried out by Erst et al. (2016).

	<i>Aconitum ambiguum</i> Rchb.	[1, 2, 3, 4, 10]
	<i>Aconitum anthoroideum</i> DC.	[3, 7]
	<i>Aconitum baicalense</i> Turcz. [= <i>Aconitum ambiguum</i> Rchb. subsp. <i>baicalense</i> (Turcz.) Vorosch.]	[2, 3, 4, 5, 9]
	<i>Aconitum barbatum</i> Patr.	[1–4, 6–8, 10, 11, 13]
	<i>Aconitum biflorum</i> Fisch.	[3, 7]
	<i>Aconitum coreanum</i> (H.Lév.) Rapaics	[5]
	<i>Aconitum decipiens</i> Vorosch. & Anfalov	[3, 7]
	<i>Aconitum glandulosum</i> Rapaics [= <i>Aconitum altaicum</i> Steinb. = <i>Aconitum smirnovii</i> Steinb.]	[1–4, 6, 7, 13, 14]
E	<i>Aconitum gubanovii</i> Luferov & Vorosch.	[7, 14]
E	<i>Aconitum kamelinii</i> A.A.Solovjev [= <i>Aconitum chasmanthum</i> Stapf]	[3, 13]
SE	<i>Aconitum khanminthunii</i> A.A.Solovjev & Shmakov	[3, 6, 7, 11, 13]
	<i>Aconitum kusnezoffii</i> Rchb. [= <i>Aconitum birobidshanicum</i> Vorosch.]	[5, 9]
	<i>Aconitum leucostomum</i> Vorosch.	[6, 7, 8, 10]
	<i>Aconitum macrorhynchum</i> Turcz.	[4]
SE	<i>Aconitum paskoi</i> Vorosch.	[2, 3]
	<i>Aconitum ranunculoides</i> Turcz.	[4]
SE	<i>Aconitum rubicundum</i> (Ser.) Fisch. [= <i>Aconitum septentrionale</i> subsp. <i>rubicundum</i> (Ser.) Vorosch.]	[1]
	<i>Aconitum septentrionale</i> Koelle	[1–4, 6, 7, 8]
SE	<i>Aconitum turczaninowii</i> Vorosch.	[2, 3, 4, 5, 9]
	<i>Aconitum volubile</i> Koelle	[2, 3, 7]
	<i>Actaea cimicifuga</i> L. [= <i>Cimicifuga foetida</i> L.]	[1, 2, 3, 4]
	<i>Actaea dahurica</i> (Turcz.) Franch. [= <i>Cimicifuga dahurica</i> (Turcz.) Maxim.]	[2, 5, 9]
	<i>Actaea erythrocarpa</i> (Fisch.) Kom.	[1, 2, 3, 4]
	<i>Actaea simplex</i> Prantl	[5]
	<i>Adonis apennina</i> L. [= <i>Adonis sibirica</i> Patrin]	[1, 2, 3, 4, 6, 8]
E	<i>Adonis mongolica</i> Simonovich	[1, 2, 3, 4, 8]
	<i>Anemonastrum crinitum</i> (Juz.) Holub [= <i>Anemone narcissiflora</i> subsp. <i>crinita</i> (Juz.) Kitag.]	[1, 2, 3, 4, 6, 7]
	<i>Anemonastrum dichotomum</i> (L.) Mosyakin [= <i>Anemone dichotoma</i> L.]	[2, 3, 4, 5, 9]

	<i>Anemonastrum obtusilobum</i> (D.Don) Mosyakin [= <i>Anemone obtusiloba</i> Lindl.]	[3]
	<i>Anemonastrum sibiricum</i> (L.) Holub [= <i>Anemone sibirica</i> L.]	[1, 2, 4]
	<i>Anemone reflexa</i> Steph.	[2, 3, 4]
	<i>Anemone sylvestris</i> L. [= <i>Anemonoides sylvestris</i> (L.) Galasso]	[1–7, 9]
E	<i>Aquilegia amurensis</i> Kom.	[2]
	<i>Aquilegia aradanica</i> Shaulo & Erst	[4]
	<i>Aquilegia daingolica</i> Erst & Shaulo	[7]
	<i>Aquilegia ganboldii</i> Kamelin & Gubanov	[5]
	<i>Aquilegia glandulosa</i> Fisch.	[1, 4, 6, 7]
E	<i>Aquilegia grubovii</i> Erst	[1, 2, 3, 4]
	<i>Aquilegia jucunda</i> Fisch. & Lallem.	[1]
	<i>Aquilegia sibirica</i> Lam.	[1, 2, 3, 4, 6, 7]
	<i>Aquilegia viridiflora</i> Pall.	[2–5, 7–10, 12, 13]
	<i>Aquilegia xinjiangensis</i> Erst	[7]
	<i>Callianthemum angustifolium</i> Witasek	[7]
	<i>Callianthemum isopyroides</i> Witasek	[1, 2, 3]
	<i>Callianthemum sajanense</i> Witasek	[1, 3, 7]
	<i>Caltha membranacea</i> (Turcz.) Schipcz.	[5]
	<i>Caltha natans</i> (Pall.) Deyl & Sojak	[1–5, 8, 10, 11]
	<i>Caltha palustris</i> L.	[1–5, 9–11]
	<i>Ceratocephala testiculata</i> (Crantz) Besser	[7]
	<i>Clematis aethusifolia</i> Turcz.	[9]
	<i>Clematis brevicaudata</i> DC.	[5, 9]
SE	<i>Clematis fruticosa</i> Turcz.	[11–13, 15, 16]
	<i>Clematis glauca</i> Willd.	[3, 7, 10, 14, 15]
	<i>Clematis hexapetala</i> Pall.	[2, 4, 5, 8, 9]
	<i>Clematis intricata</i> Bunge	[7–14, 16]
	<i>Clematis macropetala</i> Ledeb.	[9]
	<i>Clematis ochotensis</i> (Pall.) Poir. & Lam.	[4]
	<i>Clematis orientalis</i> L.	[15]
	<i>Clematis sibirica</i> (L.) Mill.	[1–4, 6–8, 10, 13]
	<i>Clematis songarica</i> Siev.	[7, 11–16]
SE	<i>Clematis tangutica</i> subsp. <i>mongolica</i> Grey-Wilson	[2]
	<i>Clematis tangutica</i> Korsh. subsp. <i>tangutica</i>	[2–4, 7, 8, 10, 13–15]
	<i>Delphinium altaicum</i> Nevski	[6, 7, 13, 14]
	<i>Delphinium barlykense</i> Lomon. & Khanm.	[1, 6, 7]
E	<i>Delphinium changaicum</i> N.Friesen	[3, 13]
	<i>Delphinium cheilanthum</i> Fisch.	[1–4, 6, 7, 8, 13]
	<i>Delphinium crassifolium</i> Schrad.	[1, 2, 3, 5, 6, 7]
	<i>Delphinium dictyocarpum</i> DC.	[7]
SE	<i>Delphinium dissectum</i> Huth	[1, 3, 4, 8]
	<i>Delphinium elatum</i> L.	[1, 3, 6, 7]

	<i>Delphinium grandiflorum</i> L.	[1–5, 9, 13]
E	<i>Delphinium gubanovii</i> N.Friesen	[7]
	<i>Delphinium iliense</i> Huth	[14]
	<i>Delphinium inconspicuum</i> Serg. subsp. <i>inconspicuum</i>	[3, 6, 7, 14]
E	<i>Delphinium inconspicuum</i> subsp. <i>mongolicum</i> A.L.Ebel	[7]
SE	<i>Delphinium malyschevii</i> N.Friesen	[1]
	<i>Delphinium mirabile</i> Serg.	[6, 7]
SE	<i>Delphinium sajanense</i> Jurtzev	[1]
	<i>Delphinium triste</i> Fisch.	[1, 2, 3, 4, 8, 13]
	<i>Delphinium ukokense</i> Serg.	[6, 7]
	<i>Halerpestes salsuginosa</i> Greene	[1–4, 6–15]
	<i>Halerpestes sarmentosa</i> (Adams) Kom. & Klob.-Alis	[3, 4, 6–16]
	<i>Isopyrum anemonoides</i> Kar. & Kir.	[7]
	<i>Leptopyrum fumariooides</i> Rchb.	[1–4, 6–9, 13]
	<i>Oxygraphis glacialis</i> (Fisch.) Bunge	[1, 2, 3, 6, 7, 13]
	<i>Paraquilegia anemonoides</i> Ulbr.	[1, 6, 7]
	<i>Pulsatilla ambigua</i> Turcz.	[1–4, 6, 7, 13]
SE	<i>Pulsatilla bungeana</i> C.A.Mey. [= <i>Pulsatilla bungeana</i> var. <i>astragalifolia</i> (Pobed.) Grubov]	[1–11, 13]
	<i>Pulsatilla campanella</i> Fisch.	[1, 3, 6, 7, 14]
	<i>Pulsatilla dahurica</i> (Fisch.) Spreng.	[3, 9]
	<i>Pulsatilla multifida</i> (G.Pritz.) Juz. [= <i>Pulsatilla patens</i> subsp. <i>multifida</i> (G.Pritz.) Zämelis]	[3, 4, 6, 7]
	<i>Pulsatilla patens</i> (L.) Mill. subsp. <i>flavescens</i> (Zucc.) Zämelis [= <i>Pulsatilla flavescens</i> (Zucc.) Juz.]	[1, 2, 3, 4, 5, 7]
	<i>Pulsatilla tenuiloba</i> (Hayek) Juz.	[2, 3, 4, 9]
	<i>Pulsatilla turczaninovii</i> Krylov & Serg.	[1–6, 8, 9]
	<i>Ranunculus acris</i> L.	[1–5, 7, 8, 9, 10]
	<i>Ranunculus altaicus</i> Laxm.	[1, 2, 3, 6, 7]
	<i>Ranunculus aquatilis</i> L.	[3, 5, 9]
E	<i>Ranunculus archantynicus</i> Kamelin, Shmakov & S.V.Smirn.	[7, 14]
	<i>Ranunculus chinensis</i> Bunge	[2, 3, 4, 6, 10]
	<i>Ranunculus circinatus</i> Sibth.	[3, 4, 5, 9, 10]
	<i>Ranunculus confervoides</i> (Fr.) Fr. [= <i>Ranunculus trichophyllus</i> subsp. <i>eradicatus</i> (Laest.) C.D.K.Cook]	[2, 5–10]
	<i>Ranunculus gmelinii</i> DC.	[1, 2, 4, 5, 9]
	<i>Ranunculus gobicus</i> Maxim.	[13]
	<i>Ranunculus grandifolius</i> C.A.Mey.	[3]
	<i>Ranunculus kauffmannii</i> Clerc [= <i>Batrachium kauffmannii</i> (Clerc) Krecz.]	[2, 7]
SE	<i>Ranunculus lapponicus</i> L.	[1, 2, 3, 6, 7, 13]
	<i>Ranunculus lasiocarpus</i> C.A.Mey.	[1, 3, 6, 7]
	<i>Ranunculus lingua</i> L.	[14]

	<i>Ranunculus longicaulis</i> C.A.Mey.	[1–3, 6, 7, 11, 14]
	<i>Ranunculus mongolicus</i> (Krylov) Serg. [= <i>Batrachium mongolicum</i> Serg.]	[3, 6, 7, 10, 11]
	<i>Ranunculus monophyllus</i> Ovcz.	[1–7]
	<i>Ranunculus natans</i> C.A.Mey.	[1–4, 6–10, 13, 15]
	<i>Ranunculus pedatifidus</i> Sm. [= <i>Ranunculus rigescens</i> Turcz.]	[1–7, 9, 13, 14]
	<i>Ranunculus polyanthemos</i> L.	[6]
	<i>Ranunculus propinquus</i> C.A.Mey. subsp. <i>propinquus</i>	[1–4, 6–8, 10]
	<i>Ranunculus propinquus</i> var. <i>subborealis</i> (Tzvel.) Luferov	[1, 2, 3, 4]
SE	<i>Ranunculus pseudohirculus</i> Schrenk	[1–3, 6, 7, 13, 14]
	<i>Ranunculus pseudomonophyllus</i> Timokhina	[1, 2]
	<i>Ranunculus pulchellus</i> C.A.Mey.	[1–4, 6–10, 13]
E	<i>Ranunculus radicans</i> C.A.Mey.	[1–8, 10, 13]
	<i>Ranunculus repens</i> L.	[2–9]
	<i>Ranunculus reptans</i> L.	[1, 2, 4, 7, 10]
	<i>Ranunculus sapozhnikovii</i> Schegol.	[7]
	<i>Ranunculus sceleratus</i> L.	[1–9, 11, 13, 14]
SE	<i>Ranunculus schmakovii</i> Erst	[7]
	<i>Ranunculus smirnovii</i> Ovcz.	[2]
	<i>Ranunculus sulphureus</i> subsp. <i>exaltatus</i> Erst	[7]
	<i>Ranunculus tanguticus</i> (Maxim) Ovcz.	[3, 6]
	<i>Ranunculus trautvetterianus</i> Regel	[7]
	<i>Ranunculus trichophyllum</i> Chaix [= <i>Batrachium trichophyllum</i> (Chaix) Bosch = <i>Batrachium divaricatum</i> (Schrank) Schur]	[1–11, 14]
SE	<i>Ranunculus turczaninovii</i> (Luferov) Vorosch.	[2]
SE	<i>Ranunculus tuvinicus</i> Erst	[7]
	<i>Thalictrum alpinum</i> L.	[1–4, 6, 7, 13]
	<i>Thalictrum baicalense</i> Turcz.	[2, 5]
	<i>Thalictrum contortum</i> L. [= <i>Thalictrum aquilegiifolium</i> var. <i>sibiricum</i> Regel & Tiling]	[5, 9]
	<i>Thalictrum foetidum</i> L.	[1–4, 6–10, 13, 14]
	<i>Thalictrum isopyroides</i> C.A.Mey.	[7, 14]
SE	<i>Thalictrum minus</i> subsp. <i>appendiculatum</i> (C.A.Mey.) Gubanov [= <i>Thalictrum appendiculatum</i> C.A.Mey.]	[3]
	<i>Thalictrum minus</i> subsp. <i>elatum</i> (Jacq.) Stoj. & Stef. [= <i>Thalictrum minus</i> subsp. <i>kemense</i> (Fries) Cajander]	[3]
	<i>Thalictrum minus</i> L. subsp. <i>minus</i>	[1–10, 13, 14]
	<i>Thalictrum petaloideum</i> L.	[1, 2, 3, 4, 5, 9]
SE	<i>Thalictrum schischkinii</i> N.Friesen [= <i>Thalictrum altaicum</i> (Schischk.) Serg.]	[7]
	<i>Thalictrum simplex</i> L.	[1–10, 13, 14]
	<i>Thalictrum squarrosum</i> Steph.	[2, 3, 4, 5, 8, 9]
	<i>Trollius altaicus</i> C.A.Mey.	[6, 7, 14]
	<i>Trollius asiaticus</i> L.	[1–4, 6, 7, 9, 13]

<i>Trollius austrosibiricus</i> Erst & Luferov	[7]
<i>Trollius chinensis</i> Bunge	[7]
<i>Trollius dschungaricus</i> Regel	[14]
<i>Trollius ledebourii</i> Rchb.	[2, 3, 4, 5, 9]
<i>Trollius lilacinus</i> Bunge	[1, 2, 6, 7]
<i>Trollius sajanensis</i> (Malyschev) Sipliv.	[1]
<i>Trollius sibiricus</i> Schipcz.	[5]
<i>Trollius vicarius</i> Sipliv.	[5]

**93. Rhamnaceae** Juss. (1 genus and 5 species)

<i>Rhamnus davurica</i> Pall.	[2, 4]
<i>Rhamnus erythroxylon</i> Pall.	[2–4, 8, 9, 12, 13]
<i>Rhamnus maximovicziana</i> J.J.Vassil.	[13, 16]
<i>Rhamnus parvifolia</i> Bunge	[2, 4, 5, 9]
<i>Rhamnus utilis</i> Decne.	[4, 5, 9]

**94. Rosaceae** Juss. (28 genera and 168 taxa)

<i>Agrimonia pilosa</i> Ledeb.	[1–6, 9]
<i>Alchemilla argutiserrata</i> H.Lindb.	[7]
E <i>Alchemilla changaica</i> V.N.Tikhom.	[1, 3]
<i>Alchemilla circularis</i> Juz.	[7]
<i>Alchemilla cyrtopleura</i> Juz.	[3, 7]
<i>Alchemilla flavescentia</i> Buser	[3]
<i>Alchemilla gracilis</i> Pax	[3]
<i>Alchemilla gubanovii</i> V.N.Tikhom.	[2, 3]
<i>Alchemilla hebescens</i> Juz.	[2, 3, 7]
<i>Alchemilla krylovii</i> Juz.	[7]
<i>Alchemilla murbeckiana</i> Buser	[7]
SE <i>Alchemilla pavlovii</i> Juz.	[2, 3, 6]
<i>Argentina anserina</i> (L.) Rydb. [= <i>Potentilla anserina</i> L.]	[1–11, 13–15]
<i>Aruncus sylvester</i> Kostel.	[5]
SE <i>Chamaerhodos altaica</i> Bunge	[1–4, 6–8, 10, 11, 13]
<i>Chamaerhodos corymbosa</i> Murav.	[5, 9]
<i>Chamaerhodos erecta</i> (L.) Bunge	[1–13]
<i>Chamaerhodos grandiflora</i> Ledeb.	[5]
<i>Chamaerhodos sabulosa</i> Bunge	[3, 6–16]
<i>Chamaerhodos trifida</i> Ledeb.	[4, 5, 8, 9, 12, 13]
<i>Coluria geoides</i> (Pall.) Ledeb. [= <i>Dryas geoides</i> Pall.]	[3, 6]
<i>Comarum palustre</i> L.	[1, 2, 3, 4, 6]
<i>Cotoneaster megalocarpus</i> Popov	[7]
<i>Cotoneaster melanocarpus</i> Lodd.	[1–10, 13, 14]
<i>Cotoneaster mongolicus</i> Pojark.	[2–5, 7–9, 12, 13]
<i>Cotoneaster neopopovii</i> Czerep.	[4]

	<i>Cotoneaster uniflorus</i> Bunge	[1, 2, 3, 7, 8, 13]
	<i>Crataegus dahurica</i> Koehne	[2, 4, 5, 9]
	<i>Crataegus maximowiczii</i> C.K.Schneid.	[5]
	<i>Crataegus sanguinea</i> Pall.	[2, 3, 4, 5, 9]
	<i>Dasiphora fruticosa</i> (L.) Rydb. [= <i>Potentilla fruticosa</i> L.]	[1–9, 11, 13]
	<i>Dasiphora parvifolia</i> (Fisch.) Juz. [= <i>Potentilla parvifolia</i> Fisch.]	[2, 3, 4, 6, 8]
	<i>Dryas grandis</i> Juz.	[1, 7]
	<i>Dryas incisa</i> Juz.	[1]
	<i>Dryas oxyodonta</i> Juz.	[1, 2, 3, 4, 6, 7]
	<i>Dryas punctata</i> Juz.	[1, 3]
SE	<i>Dryas sumneviczii</i> Serg.	[1]
	<i>Farinopsis salesoviana</i> (Steph.) Chrtek & Soják [≡ <i>Comarum salesovianum</i> (Steph.) Ledeb]	[6, 7, 10, 11, 13, 14]
	<i>Filipendula angustiloba</i> Maxim.	[5, 9]
	<i>Filipendula palmata</i> Maxim.	[2, 3, 4, 5, 9]
	<i>Filipendula ulmaria</i> (L.) Maxim.	[2, 3, 4]
	<i>Fragaria orientalis</i> Losinsk.	[2, 3, 4, 5]
	<i>Fragaria viridis</i> Weston	[2]
	<i>Geum aleppicum</i> Jacq.	[2, 3, 4, 5, 9]
	<i>Geum rivale</i> L.	[7]
	<i>Malus baccata</i> (L.) Borkh.	[2, 3, 4, 5, 8, 9]
SE	<i>Potaninia mongolica</i> Maxim.	[11, 12, 13, 16]
	<i>Potentilla acaulis</i> L.	[1–11, 13]
	<i>Potentilla acervata</i> Soják [= <i>Potentilla chenteica</i> Soják]	[2–5, 8, 9, 13]
	<i>Potentilla agrimonoides</i> M.Bieb. [= <i>Potentilla lydiae</i> Kurbatski]	[7, 14]
	<i>Potentilla altaica</i> Bunge [= <i>Potentilla nivea</i> L. var. <i>pinnatifida</i> Lehm.]	[7]
	<i>Potentilla angustiloba</i> T.T.Yu & C.L.Li	[7, 14]
	<i>Potentilla aphanes</i> Soják	[3, 6, 7, 10, 13, 14]
	<i>Potentilla arenosa</i> (Turcz.) Juz. [= <i>Potentilla nivea</i> var. <i>arenosa</i> Turcz.]	[2, 3, 4, 6]
	<i>Potentilla asiatica</i> (Th.Wolf) Juz.	[7]
	<i>Potentilla astragalifolia</i> Bunge	[3, 6, 7, 10, 11]
SE	<i>Potentilla × burjatica</i> Soják	[2]
	<i>Potentilla chalchorum</i> Soják	[2, 3, 7, 9]
SE	<i>Potentilla × chamaeleo</i> Soják	[6, 7, 14]
	<i>Potentilla chinensis</i> Ser.	[4, 5, 9]
	<i>Potentilla chionea</i> Soják	[1–4, 10, 13, 14]
	<i>Potentilla chrysantha</i> Trevir.	[7, 14]
	<i>Potentilla conferta</i> Bunge	[1–9, 12, 13, 14]
E	<i>Potentilla coriacea</i> Soják	[3]
	<i>Potentilla crantzii</i> (Crantz) Fritsch	[7]
	<i>Potentilla crebridens</i> Juz. [= <i>Potentilla nivea</i> var. <i>elongata</i> Th.Wolf]	[1, 2, 3, 10]
	<i>Potentilla desertorum</i> Bunge	[1, 6, 7, 9, 12–14]

SE	<i>Potentilla × drymeja</i> Soják	[2, 3, 13]
E	<i>Potentilla ekaterinae</i> Kamelin ex Kechaykin	[13]
	<i>Potentilla elegans</i> Cham. & Schltdl.	[1]
	<i>Potentilla elegantissima</i> Polozhij	[3]
	<i>Potentilla evestita</i> Th.Wolf	[1–4, 6, 7, 13, 14]
	<i>Potentilla exuta</i> Soják	[3, 7, 13, 14]
	<i>Potentilla flagellaris</i> D.F.K.Schltdl.	[2, 3, 4, 5, 9]
	<i>Potentilla fragarioides</i> L.	[2, 3, 4, 5, 7]
	<i>Potentilla gelida</i> C.A.Mey.	[1–3, 6, 7, 9, 13, 14]
E	<i>Potentilla gobica</i> Soják	[14]
SE	<i>Potentilla gracillima</i> Kamelin	[3, 7, 10]
E	<i>Potentilla hilbigii</i> Soják	[3]
E	<i>Potentilla hubsugulica</i> Soják	[1]
E	<i>Potentilla ikonnikovii</i> Juz.	[7, 13]
E	<i>Potentilla inopinata</i> Soják	[6, 7]
	<i>Potentilla jenissejensis</i> Polozhij & W.Smirnova [= <i>Potentilla agrimonoides</i> M.Bieb. var. <i>kobdoensis</i> Soják]	[6, 7, 10]
	<i>Potentilla kryloviana</i> Th.Wolf	[3, 7, 14]
E	<i>Potentilla laevipes</i> Soják	[7]
E	<i>Potentilla laevissima</i> Kamelin	[7]
	<i>Potentilla leucophylla</i> Pall. [= <i>Potentilla betonicifolia</i> Poir.]	[2, 3, 4, 5, 8, 9]
	<i>Potentilla longifolia</i> D.F.K.Schltdl.	[1–13]
E	<i>Potentilla mongolica</i> Krasch.	[3, 8]
	<i>Potentilla multicaulis</i> Bunge	[1, 3]
	<i>Potentilla multifida</i> L. [= <i>Potentilla tenella</i> Turcz.]	[1–14]
	<i>Potentilla nivea</i> L.	[1–4, 6, 7, 13, 14]
	<i>Potentilla norvegica</i> L. [= <i>Potentilla monspeliensis</i> L.]	[1, 2, 3, 4]
	<i>Potentilla nudicaulis</i> D.F.K.Schltdl. [= <i>Potentilla strigosa</i> Pall.]	[12]
SE	<i>Potentilla × olchonensis</i> Peschkova	[6]
	<i>Potentilla ornithopoda</i> Tausch	[1–4, 6, 7, 10, 14]
SE	<i>Potentilla ozjorensis</i> Peschkova	[1, 3, 4, 7]
	<i>Potentilla pamirica</i> Th.Wolf	[6, 7, 10, 14]
	<i>Potentilla pamiroalaica</i> Juz.	[14]
	<i>Potentilla pensylvanica</i> L. [= <i>Pentaphyllum pennsylvanicum</i> (L.) Lunell]	[1–11, 13, 14]
	<i>Potentilla regelianae</i> Th.Wolf	[6]
SE	<i>Potentilla × rhipidophylla</i> Soják	[3]
SE	<i>Potentilla rigidula</i> Th.Wolf	[6, 10]
	<i>Potentilla sanguisorba</i> D.F.K.Schltdl.	[1, 2, 3, 4, 12, 13]
E	<i>Potentilla schmakovii</i> Kechaykin	[7, 14]
SE	<i>Potentilla sergievskajae</i> Peschkova	[5, 8]
	<i>Potentilla sericea</i> L.	[1–4, 6–13, 15]
SE	<i>Potentilla serrata</i> Soják	[3]

SE	<i>Potentilla sischanensis</i> Bunge	[4, 9]
	<i>Potentilla songorica</i> Bunge	[10, 14]
SE	<i>Potentilla stepposa</i> Soják	[7, 10]
	<i>Potentilla subdigitata</i> T.T.Yu & C.L.Li [= <i>Potentilla junatovii</i> Rudaya & A.L.Ebel]	[7]
	<i>Potentilla supina</i> L.	[1–12, 14–16]
	<i>Potentilla tanacetifolia</i> D.F.K.Schltdl.	[2–9, 12, 13]
	<i>Potentilla tergemina</i> Soják	[2, 3, 4, 5, 9]
SE	<i>Potentilla tericholica</i> Sobolevsk.	[6, 7]
	<i>Potentilla tetrandra</i> (Bunge) Hook.f. [= <i>Sibbaldia tetrandra</i> Bunge]	[1, 3, 6, 7]
	<i>Potentilla turczaninowiana</i> Stschegl.	[6, 7, 14]
	<i>Potentilla turkestanica</i> Soják	[7, 14]
E	<i>Potentilla tytthantha</i> (Soják) Kechaykin	[6, 7]
E	<i>Potentilla × vanzhilii</i> Gundegmaa & Kechaykin	[3]
	<i>Potentilla verticillaris</i> Stephan	[2, 3, 4, 5, 8, 9]
	<i>Potentilla virgata</i> Lehm.	[1, 3, 4, 6–15]
	<i>Prunus mongolica</i> Maxim. [= <i>Amygdalus mongolica</i> (Maxim.) Ricker]	[12, 13, 16]
	<i>Prunus padus</i> L.	[1, 2, 3, 4, 5, 9]
	<i>Prunus pedunculata</i> (Pall.) Maxim. [= <i>Amygdalus pedunculata</i> Pall.]	[2, 3, 4, 6–13, 16]
	<i>Prunus sibirica</i> L. [= <i>Armeniaca sibirica</i> (L.) Lam.]	[2, 3, 4, 5, 9]
	<i>Rosa acicularis</i> Lindl.	[1–9, 13]
	<i>Rosa albertii</i> Regel	[7]
E	<i>Rosa baitagensis</i> Kamelin & Gubanov	[14]
	<i>Rosa beggeriana</i> Schrenk	[14]
	<i>Rosa davurica</i> Pall.	[2, 4, 5, 9]
	<i>Rosa kokanica</i> (Regel) Regel	[7]
	<i>Rosa laxa</i> var. <i>kaschgarica</i> (Rupr.) Y.L.Han [= <i>Rosa kaschgarica</i> Rupr.]	[14, 15]
	<i>Rosa laxa</i> Lindl. var. <i>laxa</i>	[6, 7, 13, 14, 15]
	<i>Rosa oxyacantha</i> M.Bieb.	[1, 2, 3, 7]
	<i>Rosa platyacantha</i> Schrenk	[14]
	<i>Rosa spinosissima</i> L.	[3, 7, 14]
	<i>Rosa xanthina</i> Lindl.	[9]
	<i>Rubus arcticus</i> L.	[1, 2, 3, 4]
	<i>Rubus chamaemorus</i> L.	[2]
	<i>Rubus humilifolius</i> C.A.Mey.	[1, 2]
	<i>Rubus sachalinensis</i> H.Lév.	[1, 2, 3, 4, 7, 8]
	<i>Rubus saxatilis</i> L.	[1, 2, 3, 4, 5, 9]
	<i>Sanguisorba alpina</i> Bunge	[3, 6, 7]
	<i>Sanguisorba officinalis</i> L.	[1–11]
	<i>Sanguisorba parviflora</i> (Maxim.) Takeda	[9]
	<i>Sanguisorba tenuifolia</i> Fisch.	[4, 9]
	<i>Sibbaldia procumbens</i> L.	[2, 7]

	<i>Sibbaldianthe adpressa</i> (Bunge) Juz. [= <i>Sibbaldia adpressa</i> Bunge]	[1–13, 15, 16]
	<i>Sibbaldianthe bifurca</i> (L.) Kurtto & T.Erikss. [= <i>Potentilla bifurca</i> L.]	[1–14]
	<i>Sibbaldianthe imbricata</i> (Kar. & Kir.) Mosyakin & Shiyan [≡ <i>Potentilla imbricata</i> Kar. & Kir.]	[6, 7, 10, 14]
	<i>Sibbaldianthe orientalis</i> (Soják) Mosyakin & Shiyan [= <i>Potentilla bifurca</i> var. <i>major</i> Ledeb.]	[7, 8, 9, 14]
	<i>Sibbaldianthe semiglabra</i> (Soják) Mosyakin & Shiyan [≡ <i>Potentilla semiglabra</i> Juz.]	[5, 9]
SE	<i>Sibbaldianthe sericea</i> Grubov	[7, 8, 12, 13]
	<i>Sibiraea laevigata</i> (L.) Maxim.	[7]
	<i>Sorbaria sorbifolia</i> (L.) A.Braun	[4, 5]
	<i>Sorbus aucuparia</i> L. subsp. <i>glabrata</i> (Wimm. & Grab.) Hedl. [= <i>Sorbus sibirica</i> Hedl.]	[1, 2, 3, 4, 5]
	<i>Spiraea alpina</i> Pall.	[1, 2, 3, 6, 7]
	<i>Spiraea aquilegiifolia</i> Pall.	[1–5, 8, 9, 12, 13]
	<i>Spiraea chamaedryfolia</i> L.	[5]
	<i>Spiraea dahurica</i> (Rupr.) Maxim.	[2, 4]
	<i>Spiraea elegans</i> Pojark.	[4]
	<i>Spiraea flexuosa</i> Fisch.	[1–6, 8, 9, 13]
	<i>Spiraea hypericifolia</i> L.	[2–4, 6, 7, 9, 10, 12, 14]
	<i>Spiraea media</i> F.Schmidt subsp. <i>media</i> [= <i>Spiraea sericea</i> Turcz.]	[1 – 8 , 13]
	<i>Spiraea pubescens</i> Turcz.	[4, 5, 9]
	<i>Spiraea salicifolia</i> L.	[2, 3, 4, 5, 9]

### 95. Rubiaceae Juss. (3 genera and 13 taxa)

E	<i>Asperula gobicola</i> Grubov [= <i>Asperula saxicola</i> Grubov]	[13, 16]
	<i>Galium amblyophyllum</i> Schrenk	[1, 2, 14]
	<i>Galium boreale</i> L.	[1–10, 13, 14]
	<i>Galium dahuricum</i> Turcz.	[2]
	<i>Galium densiflorum</i> Ledeb.	[3, 6, 7, 14]
	<i>Galium humifusum</i> M.Bieb.	[3, 7, 13]
	<i>Galium songaricum</i> Schrenk	[1, 2, 3]
	<i>Galium spurium</i> L.	[1–8, 10, 13, 14]
	<i>Galium trifidum</i> L.	[2–5, 7, 10, 14]
	<i>Galium uliginosum</i> L.	[1, 2, 3, 4, 10]
	<i>Galium verum</i> L. subsp. <i>verum</i> [= <i>Galium densiflorum</i> Ledeb.]	[1–10, 13, 14]
	<i>Galium verum</i> subsp. <i>wirtgenii</i> (F.W.Schultz) Oborny	[7]
	<i>Rubia cordifolia</i> L. [≡ <i>Galium cordifolium</i> (L.) Kuntze]	[2–5, 8, 9, 12, 13]

### 96. Ruppiaceae Horan. (1 genus and 1 species)

	<i>Ruppia maritima</i> L.	[10]
--	---------------------------	------

97. **Rutaceae** Juss. (2 genera and 2 species)

- Haplophyllum dauricum* (L.) G.Don [2–6, 8, 9, 11–14, 16]  
*Dictamnus albus* L. [5, 9]

98. **Salicaceae** Mirb. (2 genera and 47 species)

- |  |                         |
|--|-------------------------|
| <i>Populus euphratica</i> Olivier  | [12–16]                 |
| <i>Populus laurifolia</i> Ledeb. [= <i>Populus pilosa</i> Rehder]                                  | [2–4, 6, 7, 10, 13, 14] |
| <i>Populus simonii</i> Carrière  | [9]                     |
| <i>Populus suaveolens</i> Fisch.   | [1, 2, 3, 4]            |
| <i>Populus tremula</i> L.  | [1–5, 7–9, 11]          |
| <i>Salix abscondita</i> Laksch.  | [1, 2, 4, 5, 13]        |
| <i>Salix alatavica</i> Kar.  | [6, 7]                  |
| <i>Salix arctica</i> Pall.   | [1, 3, 6, 7]            |
| <i>Salix bebbiana</i> Sarg.  | [1–10, 12, 13]          |
| <i>Salix berberifolia</i> Pall.  | [1, 2, 3, 6, 7, 13]     |
| <i>Salix brachypoda</i> (Trautv. & C.A.Mey.) Kom.  | [2, 4, 5, 9]            |
| <i>Salix caesia</i> Vill.  | [1–4, 6, 7, 10, 14]     |
| <i>Salix divaricata</i> Pall.  | [1–4, 6, 7, 13]         |
| <i>Salix glauca</i> L.   | [1, 2, 3, 6, 7, 13]     |
| <i>Salix gmelinii</i> Pall. [= <i>Salix dasyclados</i> Wimmer]                                     | [2, 4, 6, 7, 8, 10]     |
| <i>Salix gorderevii</i> Y.L.Chang & Skvortsov  | [5, 8, 9]               |
| <i>Salix hastata</i> L.  | [1, 6, 7, 10, 13]       |
| <i>Salix jenisseensis</i> (F.Schmidt) Flod.  | [1, 6, 7]               |
| <i>Salix kochiana</i> Trautv.  | [1–5, 7, 10]            |
| <i>Salix ledebouriana</i> Trautv.  | [1–4, 6, 7, 9–15]       |
| <i>Salix microstachya</i> Turcz.   | [2–6, 8, 9, 10]         |
| <i>Salix miyabeana</i> Seemen  | [1–5, 8, 9]             |
| <i>Salix myrtilloides</i> L.   | [1, 2, 3]               |
| <i>Salix nasarovii</i> A.K.Skvortsov   | [1]                     |
| <i>Salix nipponica</i> Franch. & Sav.  | [9]                     |
| <i>Salix nummularia</i> Andersson  | [2, 3, 6, 7]            |
| <i>Salix polaris</i> Wahlenb.  | [1]                     |
| <i>Salix pseudopentandra</i> (Flod.) Flod. [= <i>Salix pentandra</i> var. <i>intermedia</i> Nakai] | [1–10, 12, 13]          |
| <i>Salix pyrolifolia</i> Ledeb.  | [1–4, 6, 7, 10]         |
| <i>Salix rectijulis</i> Ledeb.   | [1, 2, 3, 6, 7]         |
| <i>Salix recurvигemmata</i> A.K.Skvortsov [= <i>Salix recurvигemmis</i> A.K.Skvortsov]             | [1, 3, 6]               |
| <i>Salix reticulata</i> L.   | [1, 3, 6, 7]            |
| <i>Salix rhamnifolia</i> Pall.   | [1–6, 9]                |
| <i>Salix rorida</i> Laksch.  | [2, 3, 4, 5, 9]         |
| <i>Salix rosmarinifolia</i> L.   | [1–4, 6, 10, 14]        |
| <i>Salix sajanensis</i> Nasarow  | [1, 6, 7]               |

SE

<i>Salix saposhnikovii</i> A.K.Skvortsov	[1, 3, 6, 7]
<i>Salix saxatilis</i> Turcz.	[1, 2, 3]
<i>Salix schwerinii</i> E.L.Wolf	[2, 3, 4, 5, 9]
<i>Salix taraikensis</i> Kimura	[1–5, 7, 13]
<i>Salix tenuijulis</i> Ledeb.	[3, 7, 14, 15]
<i>Salix triandra</i> L.	[1, 3, 4, 14]
<i>Salix turanica</i> Nasarow	[6, 7, 10, 14]
<i>Salix turczaninowii</i> Laksch.	[1, 2, 6, 7]
<i>Salix udensis</i> Trautv. & C.A.Mey.	[9]
<i>Salix vestita</i> Pursh	[1, 3, 6, 7]
<i>Salix viminalis</i> L.	[3, 6, 7, 10, 11, 14]

### 99. Santalaceae R.Br. (1 genus and 6 species)

<i>Thesium chinense</i> Turcz.	[9]
<i>Thesium longifolium</i> Turcz.	[2, 3, 4]
<i>Thesium refractum</i> C.A.Mey.	[1–10, 13]
<i>Thesium repens</i> Ledeb.	[1, 2, 3, 4]
SE <i>Thesium saxatile</i> Turcz.	[1, 3–6, 8–10]
SE <i>Thesium tuvense</i> Krasnob.	[5, 10]

### 100. Saxifragaceae Juss. (5 genera and 21 taxa)

<i>Bergenia crassifolia</i> (L.) Fritsch	[1, 2, 3, 4, 7]
<i>Chrysosplenium nudicaule</i> Bunge	[6]
SE <i>Chrysosplenium peltatum</i> Turcz.	[1, 3]
SE <i>Chrysosplenium sedakowii</i> Turcz.	[2, 3, 8]
<i>Chrysosplenium serreanum</i> Hand.-Mazz. [= <i>Chrysosplenium alternifolium</i> subsp. <i>sibiricum</i> (Ser.) Hultén]	[1, 2, 4]
<i>Micranthes davurica</i> (Willd.) Small [= <i>Saxifraga davurica</i> Willd.]	[2]
<i>Micranthes foliolosa</i> (R.Br.) Gornall [= <i>Saxifraga foliolosa</i> R.Br.]	[1, 2]
<i>Micranthes hieraciifolia</i> (Waldst. & Kit.) Haw. [= <i>Saxifraga hieraciifolia</i> Waldst. & Kit.]	[1, 2, 3, 6, 7]
<i>Micranthes melaleuca</i> (Fisch.) Losinsk. [= <i>Saxifraga melaleuca</i> Fisch.]	[1, 2, 6, 7]
<i>Micranthes nelsoniana</i> subsp. <i>aestivalis</i> (Fisch. & C.A.Mey.) Elven & D.F.Murray [= <i>Saxifraga aestivalis</i> Fisch. & C.A.Mey.]	[1, 2, 3, 6]
<i>Micranthes nivalis</i> (L.) Small [= <i>Saxifraga nivalis</i> L.]	[1]
<i>Mitella nuda</i> L.	[1, 2, 4]
<i>Saxifraga bronchialis</i> L. [= <i>Saxifraga caulescens</i> Sipliv., = <i>Saxifraga spinulosa</i> Adams]	[2, 3, 4, 8]
<i>Saxifraga cernua</i> L.	[1–3, 6–9, 13]
<i>Saxifraga hirculus</i> L.	[1–7, 13, 14]
<i>Saxifraga macrocalyx</i> Tolm. [= <i>Saxifraga flagellaris</i> Willd.]	[1, 3, 6, 7, 13, 14]

<i>Saxifraga oppositifolia</i> L. subsp. <i>oppositifolia</i> [= <i>Saxifraga asiatica</i> Hayek]	[1, 3, 6, 7]
<i>Saxifraga setigera</i> Pursh	[1, 2, 3, 6, 7, 13]
<i>Saxifraga sibirica</i> L.	[1–3, 6–8, 10, 13, 14]
<i>Saxifraga terekensis</i> Bunge	[1, 3, 6, 7]

**101. Scheuchzeriaceae F.Rudolphi (1 genus and 1 species)**

<i>Scheuchzeria palustris</i> L.	[2]
----------------------------------	-----

**102. Scrophulariaceae Juss. (3 genera and 6 species)**

<i>Limosella aquatica</i> L.	[1–4, 6, 7, 9–11, 13, 14]
<i>Scrophularia altaica</i> Murray	[1, 3, 6, 7]
<i>Scrophularia canescens</i> Bong. [= <i>Scrophularia hilbigii</i> Jäger]	[13, 14]
<i>Scrophularia incisa</i> Weinm.	[2–4, 6–15]
<i>Scrophularia umbrosa</i> Dumort.	[10]
<i>Verbascum thapsus</i> L.	[4]

**103. Solanaceae Juss. (4 genera and 9 taxa)**

<i>Hyoscyamus niger</i> L.	[2–5, 7–10, 12, 13]
<i>Hyoscyamus pusillus</i> L.	[6, 7, 10, 14, 15]
<i>Lycium chinense</i> var. <i>potaninii</i> (Pojark.) A.M.Lu [= <i>Lycium potaninii</i> Pojark.]	[16]
<i>Lycium ruthenicum</i> Murray	[10, 11, 13–16]
<i>Lycium truncatum</i> Y.C.Wang	[10, 12, 15, 16]
<i>Physoclaina albiflora</i> Grubov	[3, 4]
<i>Physoclaina physalooides</i> (L.) G.Don	[1, 3–9, 11–13]
<i>Solanum kitagawae</i> Schönb.-Tem.	[3, 6, 9, 14]
<i>Solanum septemlobum</i> Bunge	[4, 8, 9, 12]

E

**104. Tamaricaceae Link (3 genera and 13 taxa)**

<i>Myricaria bracteata</i> Royle	[1, 7, 11–14, 16]
<i>Myricaria longifolia</i> Ehrenb.	[2–4, 6, 7, 11]
<i>Reaumuria soongarica</i> Maxim.	[3, 6–16]
<i>Tamarix arceuthoides</i> Bunge	[10, 14, 15]
<i>Tamarix elongata</i> Ledeb.	[11, 14]
<i>Tamarix gracilis</i> Willd.	[13, 15, 16]
<i>Tamarix hispida</i> Willd.	[13]
<i>Tamarix × karelinii</i> Bunge	[10, 14–16]
<i>Tamarix kasahorum</i> Gorschk.	[15, 16]
<i>Tamarix laxa</i> Willd.	[11, 12, 15]
<i>Tamarix leptostachya</i> Bunge	[10, 13–16]
<i>Tamarix ramosissima</i> Ledeb.	[10, 12–16]
<i>Tamarix smyrnensis</i> Bunge	[12, 14, 16]

**105. Thymelaeaceae** Juss. (2 genera and 3 species)

<i>Diarthron altaicum</i> (Thiéb.-Bern.) Kit Tan [= <i>Stellera altaica</i> Thiéb.-Bern.]	[7]
<i>Diarthron linifolium</i> Turcz.	[2, 3, 4, 5]
<i>Stellera chamaejasme</i> L.	[2, 3, 4, 5, 9]

**106. Tofieldiaceae** Takht. (1 genus and 1 species)

<i>Tofieldia coccinea</i> Richardson	[1]
--------------------------------------	-----

**107. Typhaceae** Juss. (2 genera and 12 species)

<i>Sparganium emersum</i> Rehm	[3, 5, 6, 7, 9, 10]
<i>Sparganium glomeratum</i> (Laest.) Beurl.	[2, 4]
<i>Sparganium natans</i> L.	[7, 10]
<i>Sparganium stoloniferum</i> (Graebn.) Buch.-Ham.	[1, 4, 8, 9, 10]
<i>Typha angustifolia</i> L.	[1, 10]
<i>Typha domingensis</i> Pers.	[4, 10]
<i>Typha joannis</i> Mavrodiev	[9]
<i>Typha latifolia</i> L.	[2, 5]
<i>Typha laxmannii</i> Lepech.	[3–6, 8, 9, 11]
<i>Typha minima</i> Funck	[9, 10]
<i>Typha orientalis</i> C.Presl	[5]
<i>Typha tzvelevii</i> Mavrodiev	[4]

**108. Ulmaceae** Mirb. (1 genus and 3 taxa)

<i>Ulmus davidiana</i> var. <i>japonica</i> (Rehder) Nakai	[4, 5, 8]
<i>Ulmus macrocarpa</i> Hance	[2, 4, 5, 8, 9, 12]
<i>Ulmus pumila</i> L.	[2–5, 7–9, 11, 13, 16]

**109. Urticaceae** Juss. (2 genera and 4 taxa)

<i>Parietaria debilis</i> G.Forst.	[2–4, 6, 9, 10]
<i>Urtica angustifolia</i> Fisch.	[1–5, 7, 9, 10]
<i>Urtica cannabina</i> L.	[2–10, 12–14]
<i>Urtica dioica</i> L. subsp. <i>sondonii</i> (Simm.) Hyl. [= <i>Urtica sondenii</i> (Simm.) Avror]	[7, 14]

**110. Violaceae** Batsch. (1 genus and 27 taxa)

Note: Recently, Esput (2020) critically revised the genus *Viola* L in the Russian Far East and adjacent territories. In this study, accepted species and nomenclature mostly follow Esput (2020).

SE	<i>Viola acuminata</i> Ledeb.	[4, 5, 9]
	<i>Viola alexandrowiana</i> (W.Becker) Juz.	[4]
	<i>Viola altaica</i> Ker Gawl.	[3, 7]

	<i>Viola arvensis</i> Murray	[4]
	<i>Viola biflora</i> L.	[1, 2, 3, 7]
	<i>Viola brachyceras</i> Turcz.	[2]
	<i>Viola collina</i> Besser	[5, 7]
	<i>Viola dactyloides</i> Schult.	[2, 3, 4, 5, 6]
	<i>Viola disiuncta</i> W.Becker	[7]
	<i>Viola dissecta</i> Ledeb.	[1–5, 7, 9, 13]
	<i>Viola epipsilooides</i> Å.Löve & D.Löve	[1, 2]
	<i>Viola gmeliniana</i> Schult.	[1, 2, 3, 4, 5]
	<i>Viola incisa</i> Turcz.	[2, 4]
SE	<i>Viola ircutiana</i> Turcz.	[2]
	<i>Viola macroceras</i> Bunge	[7]
	<i>Viola mauritii</i> Teplouchow	[1, 4, 5, 7, 9, 13]
	<i>Viola mirabilis</i> L.	[2, 5]
	<i>Viola nemoralis</i> Kuetz.	[2]
	<i>Viola patrinii</i> Ging.	[2, 5, 7, 9, 10, 14]
	<i>Viola rudolfii</i> Vl.V.Nikitin	[4, 5]
	<i>Viola rupestris</i> F.W.Schmidt	[2, 3, 4, 6, 7]
	<i>Viola sachalinensis</i> H.Boissieu	[2, 4, 5]
	<i>Viola × schauloi</i> Vl.V.Nikitin	[2, 4]
	<i>Viola selkirkii</i> Pursh	[2]
	<i>Viola tenuicornis</i> subsp. <i>trichosepala</i> W.Becker	[4]
	<i>Viola uniflora</i> L.	[1, 2, 3, 4, 7]
	<i>Viola variegata</i> Fisch.	[1, 4, 5]

### 111. Zygophyllaceae R.Br. (2 genera and 13 taxa)

	<i>Tribulus terrestris</i> L.	[3, 5–16]
	<i>Zygophyllum brachypterum</i> Kar. & Kir.	[6, 10]
	<i>Zygophyllum gobicum</i> Maxim.	[14, 15]
	<i>Zygophyllum kaschgaricum</i> Boriss. [≡ <i>Sarcozygium kaschgaricum</i> (Boriss.) Y.X.Liou]	[12–16]
	<i>Zygophyllum macropterum</i> C.A.Mey. [= <i>Zygophyllum pinnatum</i> Cham. & Schltdl.]	[7, 14]
SE	<i>Zygophyllum melongena</i> Bunge	[3, 6, 7, 10, 11, 13, 14]
SE	<i>Zygophyllum mucronatum</i> Maxim.	[15, 16]
E	<i>Zygophyllum neglectum</i> Grubov	[10, 13, 14, 16]
	<i>Zygophyllum potaninii</i> Maxim.	[6, 7, 12–16]
	<i>Zygophyllum pterocarpum</i> Bunge	[6, 7, 10–16]
	<i>Zygophyllum rosowii</i> var. <i>latifolium</i> (Schrenk) Popov	[13–16]
	<i>Zygophyllum rosowii</i> Bunge var. <i>rosowii</i>	[3, 7, 8, 10–16]
	<i>Zygophyllum xanthoxylon</i> (Bunge) Maxim.	[7, 8, 10–16]

## Acknowledgements

Our study was supported by a research project (A study on the distribution of vascular plants in Mongolia; Grant KNA1-2-38, 20-5) of the Korea National Arboretum, Republic of Korea and partially supported by the National University of Mongolia, Mongolia (Grant No. P2021-4186). The research of Alexander Sukhorukov, which included the taxonomy of Chenopodiaceae and revisions to specimens of Chenopodiaceae in Mongolia in the herbaria MW and MHA, is in accordance with the scientific programme 12-2-21 of the Department of Higher Plants, Moscow State University. The work on the revision of the Ranunculaceae was carried out with financial support from the Russian Science Foundation (Grant 19-74-10082). The work on revising Rosaceae was carried out with the financial support of the Russian Foundation for Basic Research (Grant 20-04-00183A). We thank our colleagues for cross-checking the following family and genera: D.A. German (Altai State University, Russia) for the Brassicaceae family, N. Friesen (Osnabrueck University, Germany) for the genus *Allium*, E.V. Boltenkov (Botanical Garden-Institute of the Far Eastern Branch of the Russian Academy of Sciences, Russia) for the genus *Iris* as well as Lorenzo Peruzzi and other anonymous reviewers for valuable comments. The first author greatly thanks Jun-Ho Song (Korea Institute of Oriental Medicine, Korea) for his valuable comments and suggestions on an earlier draft of the paper.

## References

- Akan H, Ekici M, Aytac Z (2020) The synopsis of the genus *Trigonella* L. (Fabaceae) in Turkey. Turkish Journal of Botany 44(6): 670–693. <https://doi.org/10.3906/bot-2004-63>
- Alexeeva NB (2018) New species of *Iris* L. (Iridaceae) from Mongolia. Turczaninowia 21: 145–149. [in Russian] <https://doi.org/10.14258/turczaninowia.21.4.14>
- Al-Shehbaz IA (2021) Nomenclatural Adjustments in *Eutrema*, *Ceratocnemum*, *Rhamphospermum*, and *Sinapsis* (Brassicaceae, Cruciferae). Harvard Papers in Botany 26(1): 1–4. <https://doi.org/10.3100/hpib.v26iss1.2021.n1>
- Al-Shehbaz IA, Ozudogru B, German DA (2021) *Iljinckaea* (Brassicaceae), a new genus based on *Conringia planisiliqua*. Phytotaxa 500(2): 142–146. <https://doi.org/10.11646/phytotaxa.500.2.7>
- APG IV (2016) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. Botanical Journal of the Linnean Society 181(1): 1–20. <https://doi.org/10.1111/boj.12385>
- Baasanmunkh S, Oyunsetseg B, Oyundelger K, Khaliunaa K, Urgamal M, Batkhuu N, Shiga T, Chung GY, Choi HJ (2019a) Contribution to the knowledge on the flora of northern Mongolia. Journal of Asia-Pacific Biodiversity 12(4): 643–660. <https://doi.org/10.1016/j.japb.2019.08.009>
- Baasanmunkh S, Takashi S, Oyunsetseg B, Wesche K, Ritz CM, Khaliunaa K, Kim JY, Jo HJ, Batkhuu N, Chung GY, Choi HJ (2019b) Contribution to the knowledge on the flora of Numrug Strictly Protected Area and some parts of East Mongolia. Journal of Asia-Pacific Biodiversity 12: 284–301. <https://doi.org/10.1016/j.japb.2019.08.009>

- Baasanmunkh S, Oyunsetseg B, Lazkov G, Chung GY, Choi HJ (2019c) A new record and new distribution points of vascular plants from Dzungarian Gobi, Mongolia. *Turczaninowia* 22(1): 132–136. <https://doi.org/10.14258/turczaninowia.22.1.12>
- Baasanmunkh S, Kovtonyuk NK, Oyunsetseg B, Tsegmed Z, Han IV, Choi HJ (2020a) Diversity and distribution of the genus *Primula* L. (Primulaceae) in Mongolia. *Journal of Asia-Pacific Biodiversity* 13(4): 687–700. <https://doi.org/10.1016/j.japb.2020.09.002>
- Baasanmunkh S, Nyamgerel N, Bayarmaa G, Oyunsetseg B, Oyundelger K, Choi HJ (2020b) A new record of critically endangered *Saussurea bogedaensis* (Asteraceae) from Dzungarian Gobi, Mongolia. *PhytoKeys* 160: 109–121. <https://doi.org/10.3897/phytokeys.160.55603>
- Baasanmunkh S, Oyunsetseg B, Oyundari C, Oyundelger K, Urgamal M, Darikhand D, Soninkhishig N, Nyambayar D, Khaliunaa K, Tsegmed Z, Kechaykin AA, Shmakov AI, Erst AS, Friesen N, Ritz CM, Wesche K, Choi HJ (2021a) The vascular plant diversity of Dzungarian Gobi in western Mongolia, with an annotated checklist. *Phytotaxa* 501(1): 1–55. <https://doi.org/10.11646/phytotaxa.501.1.1>
- Baasanmunkh S, Oyunsetseg B, Efimov P, Tsegmed Z, Vandendorj S, Oyundelger K, Urgamal M, Undruul A, Khaliunaa K, Namuulin T, Choi HJ (2021b) Orchids of Mongolia: Taxonomy, species richness, and conservation status. *Diversity (Basel)* 13(7): e302. <https://doi.org/10.3390/d13070302>
- Baasanmunkh S, Urgamal M, Oyunsetseg B, Grabovskaya-Borodina A, Oyundelger K, Tsegmed Z, Gundemgaa V, Kechaykin AA, Pyak AI, Zhao LQ, Choi HJ (2021c) Updated checklist of vascular plants endemic to Mongolia. *Diversity (Basel)* 301(12): e619. <https://doi.org/10.3390/d13120619>
- Baasanmunkh S, Oyunsetseg B, Khaliunaa K, Kosachev P, Choi HJ (2021d) *Pedicularis incarnata* L. (Orobanchaceae) – New species to the flora of Mongolia. *Mongolian Journal of Biological Sciences* 19(1): 51–54. <https://doi.org/10.22353/mjbs.2021.19.14>
- Baasanmunkh S, Oyunsetseg B, Urgamal M, Norris J, Shiga T, Choi HJ (2022a) Notes on the taxonomy of Nymphaeaceae and Menyanthaceae in Mongolia. *Journal of Asia-Pacific Biodiversity* 15(1): 129–137. <https://doi.org/10.1016/j.japb.2021.09.011>
- Baasanmunkh S, Oyunsetseg B, Tsegmed Z, Oyundelger K, Urgamal M, Gantuya B, Javzandolgor C, Nyambayar N, Kosachev P, Choi HJ (2022b) Distribution of vascular plants in Mongolia –I part. *Mongolian Journal of Biological Sciences* 20(1): 3–28. <https://doi.org/10.22353/MJBS.2022.20.01>
- Barberá P, Quintanar A, Peterson PM, Soreng RJ, Romaschenko K, Aedo C (2019) New combinations, new names, typifications, and a new section, sect. *Hispanica*, in *Koeleria* (Poeae, Poaceae). *Phytoneuron* 46: 1–13.
- Barberá P, Soreng RJ, Peterson PM, Romaschenko K, Quintanar A, Aedo C (2020) Molecular phylogenetic analysis resolves *Trisetum* (Poaceae: Pooideae: Koeleriinae) polyphyletic: Evidence for a new genus, *Sibirotrisetum* and resurrection of *Acrospelion*. *Journal of Systematics and Evolution* 58(4): 517–526. <https://doi.org/10.1111/jse.12523>
- Batsaikhan N, Buuveibaatar B, Chimed B, Enkhtuya O, Galbrakh D, Ganbaatar O, Lkhagvasuren B, Nandintsetseg D, Berger J, Calabrese JM, Edwards A, Fagan WF, Fuller TK, Heiner M, Ito T, Kaczensky P, Leimgruber P, Lushchekina A, Milner-Gulland EJ, Mueller T, Murray MG, Olson KA, Reading R, Schaller GB, Stubbe A, Stubbe M, Walzer

- C, Von Wehrden H, Whitten T (2014) Conserving the world's finest grassland amidst ambitious national development. *Conservation Biology* 28(6): 1736–1739. <https://doi.org/10.1111/cobi.12297>
- Bazarragchaa B, Batdelger G, Shagdar D, Paek WK, Lee J (2019) *Scutellaria krasevii* Kom. & I. Schischk. ex Juz. (Lamiaceae): A new record species from Mongolia. *Korean Journal of Plant Taxonomy* 49(2): 198–201. <https://doi.org/10.11110/kjpt.2019.49.2.198>
- Biazrov LG, Ganbold E, Gubanov IA, Ulziikhutag N (1989) Flora of Khangay. Nauka, Leningrad, 191 pp. [in Russian]
- Boltenkov EV (2018) Taxonomic notes on *Iris* ser. *Lacteae* (Iridaceae) with typifications of fifteen names and one new combination. *Phytotaxa* 383(3): 283–292. <https://doi.org/10.11646/phytotaxa.383.3.5>
- Christenhusz MJM, Reveal JL, Farjon A, Gardner MF, Mill RR, Chase MW (2011) A new classification and linear sequence of extant gymnosperms. *Phytotaxa* 19(1): 55–70. <https://doi.org/10.11646/phytotaxa.19.1.3>
- Compton JA, Culham A, Jury SL (1998) Reclassification of *Actaea* to include *Cimicifuga* and *Souliea* (Ranunculaceae): Phylogeny inferred from morphology, nrDNA ITS, and cpDNA trnLF sequence variation. *Taxon* 47(3): 593–634. <https://doi.org/10.2307/1223580>
- Dariimaa S (2014) Flora of Mongolia. (Asteraceae). Bembi san, Ulaanbaatar, 275 pp. [in Mongolian]
- Dariimaa S (2021) Flora of Mongolia. (Asteraceae: Artemisiinae). Udam Soyol, Ulaanbaatar, 199 pp. [in Mongolian]
- Dariimaa S, Saruul N (2017) Flora of Mongolia. (Asteraceae: Asteroideae). Udam Soyol, Ulaanbaatar, 220 pp. [in Mongolian]
- Dorofeyev VI (2019) *Lepidium gobicum* (Cruciferae, Brassicaceae), a new species form Gobi Desert of Mongolia and China. *Turczaninowia* 22(4): 37–41. <https://doi.org/10.14258/turczaninowia.22.4.5>
- Dorofeyev VI, Ekhmaa U (2020) The genus *Camelina* (Cruciferae) in Mongolia and China reviewed on the basis of herbarium materials from the Institute of General and Experimental Biology of the ASM (UBA) and the Komarov Botanical Institute (LE). Proceeding on Applied Botany. Genetics and Breeding 181(3): 163–165. <https://doi.org/10.30901/2227-8834-2020-3-163-165>
- Doronkin V, Shaulo D, Han I, Vlasova N, Ivleva V, Enkhtuya L, Munkh-Erdene T, Ochgerel N, Munkhjargal B (2015) New records for the flora of Selenge Aimag (Mongolia). *Skvortsovia* 2(1): 8–27.
- Drew BT, González-Gallegos JG, Xiang CL, Kriebel R, Drummond CP, Walked JB, Sytsma KJ (2017) *Salvia* united: The greatest good for the greatest number. *Taxon* 66(1): 133–145. <https://doi.org/10.12705/661.7>
- Du C, Liao S, Boufford DE, Ma JS (2020) Twenty years of Chinese vascular plant novelties, 2000 through 2019. *Plant Diversity* 42(5): 393–398. <https://doi.org/10.1016/j.pld.2020.08.004>
- Duan L, Yang X, Liu P, Johnson G, Wen J, Chang ZY (2016) A molecular phylogeny of Caraganeae (Leguminosae, Papilionoideae) reveals insights into new generic and infrageneric delimitations. *PhytoKeys* 70: 111–137. <https://doi.org/10.3897/phytokeys.70.9641>

- Ebel AL, Strelnikova TO, Kupriyanov AN, Anenkhonov OA, Ankipovich ES, Antipova EM, Verkhozina AV, Efremov AN, Zykova EY, Mikhailova SI, Plikina NV, Ryabovol SV, Silanteva MM, Stepanov NV, Terekhina TA, Chernova OD, Shaulo DN (2014) Invasive and potentially invasive species in Siberia. *Bulletin Botanical Garden* 200: 52–61. [in Russian]
- Emadzade K, Lehnebach C, Lockhart P, Hörndl EA (2010) molecular phylogeny, morphology and classification of genera of Ranunculeae (Ranunculaceae). *Taxon* 59(3): 809–828. <https://doi.org/10.1002/tax.593011>
- Erst A (2007) New tax of buttercup (*Ranunculus*) from Altai mountains. *Turczaninowia* 10: 5–11. [in Russian]
- Erst A, Sukhorukov AP, Shaulo D, Kuznetsov A (2015) Chorological and taxonomic notes on *Aquilegia ganboldii* Kamelin & Gubanov (Ranunculaceae) previously considered to be a Mongolian endemic. *Acta Botanica Gallica* 162(3): 168–175. <https://doi.org/10.1080/12538078.2015.1040998>
- Erst A, Luferov AN, Xiang K, Wang W (2016) A review of the genus *Aquilegia* L. (Ranunculaceae) in Mongolia. Systematic notes on the materials of P.N. Krylov Herbarium of Tomsk State University 114: 37–48. [in Russian] <https://doi.org/10.17223/20764103.114.5>
- Erst A, Luferov A, Troshkina V, Shaulo D, Kuznetsov A, Xiang K, Wang W (2019) *Trollius austrosibiricus* (Ranunculaceae), a new species from South Siberia. *PhytoKeys* 115: 83–92. <https://doi.org/10.3897/phytokeys.115.30863>
- Esput M (2020) Revision of the genus *Viola* L. (Violaceae) in the Russian Far East with notes on adjacent territories. *Botanica Pacifica* 9(1): 3–52. <https://doi.org/10.17581/bp.2020.09113>
- Friesen N, Smirnov S, Shmakov AI, Herden T, Oyuntsetseg B, Hurka H (2020) *Allium* species of section *Rhizomatosa*, early members of the Central Asian steppe vegetation. *Flora* 263: e151536. <https://doi.org/10.1016/j.flora.2019.151536>
- Galasso G, Conti F, Luisa Peruzzi NMG (2018) An updated checklist of the vascular flora alien to Italy. *Plant Biosystems* 152(3): 556–592. <https://doi.org/10.1080/11263504.2018.1441197>
- GBIF (2021) Global Biodiversity Information Facility. <https://www.gbif.org> [accessed: 10 October 2021]
- German DA (2015) Cruciferae (Brassicaceae): Alternative treatment for the “Conspectus of the vascular plants of Mongolia”. *Turczaninowia* 18(2): 39–67. <https://doi.org/10.14258/turczaninowia.18.2.4>
- German DA (2020) Taxonomic remarks on selected Asian *Lepidium* (Brassicaceae). *Phytotaxa* 452(1): 33–45. <https://doi.org/10.11646/phytotaxa.452.1.3>
- Gillespie LJ, Soreng RJ, Cabi E, Amiri N (2018) Phylogeny and taxonomic synopsis of *Poa* subgenus *Pseudopoaa* (including *Eremopoa* and *Lindbergella*) (Poaceae, Poeae, Poinae). *PhytoKeys* 111: 69–101. <https://doi.org/10.3897/phytokeys.111.28081>
- Global Carex Group (2015) Making *Carex* monophyletic (Cyperaceae, tribe Cariceae): A new broader circumscription. *Botanical Journal of the Linnean Society* 179(1): 1–42. <https://doi.org/10.1111/boj.12298>
- Govaerts R, Nic Lughadha E, Black N, Turner R, Paton A (2021) The World Checklist of Vascular Plants, a continuously updated resource for exploring global plant diversity. *Scientific Data* 8(1): e215. <https://doi.org/10.1038/s41597-021-00997-6>

- Grubov VI (1955) Conspectus of Flora of People's Republic of Mongolia. Series of Mongolian Commission of Academy of Sciences of USSR. vol. 67. Moscow, Leningrad. [in Russian]
- Grubov VI (1982) Key to the vascular plants of Mongolia. Nauka, Leningrad, 443 pp. [in Russian]
- Grubov VI, Yunatov AA (1952) Main peculiarities of the flora of Mongolian People's Republic in relation to its division. Botanicheskii Zhurnal 37(1): 45–64. [in Russian]
- Gubanov IA (1996) Conspectus of flora in Outer Mongolia. Valang, Moscow, 136 pp. [in Russian]
- Gundegmaa V, Kechaykin AA (2018) A new intersectional hybrid in the genus *Potentilla* (Rosaceae) from Northern Mongolia. Turczaninowia 21(1): 174–179. <https://doi.org/10.14258/turczaninowia.21.1.17>
- Gunin PD, Vostokova EA, Dorofeyuk NI, Tarasov PE, Black CC (1999) Vegetation dynamics of Mongolia. Kluwer Academic, Dordrecht-Boston-London, 238 pp. <https://doi.org/10.1007/978-94-015-9143-0>
- He Y, Xu XM, Zhou Y, Liu QR (2020) *Onosma fuyunensis* (Boraginaceae), a new species from Xinjiang, China. PhytoKeys 144: 11–22. <https://doi.org/10.3897/phytokeys.144.33287>
- Hilbig W (1995) The vegetation of Mongolia. SPB Academic, Amsterdam, 258 pp.
- Hoot SB, Meyer KM, Manning JC (2012) Phylogeny and reclassification of *Anemone* (Ranunculaceae), with an emphasis on austral species. Systematic Botany 37(1): 139–152. <https://doi.org/10.1600/036364412X616729>
- Hurka H, Friesen N, Bernhardt KG, Neuffer B, Smirnov SV, Shmakov AI, Blattner FR (2019) The Eurasian steppe belt. Status quo, origin and evolutionary history. Turczaninowia 22(3): 5–71. <https://doi.org/10.14258/turczaninowia.22.3.1>
- IPNI (2021) International Plant Names Index. The Royal Botanic Gardens, Kew, Harvard University Herbaria & Libraries and Australian National Botanic Gardens. <http://www.ipni.org> [accessed: 24 August 2021]
- Javzandolgor C, Baasanmunkh S, Tsegmed Z, Oyuntsetseg B, Gundegmaa V, Choi HJ (2021) *Arctium tomentosum* (Asteraceae): A new report of a native genus in the flora of Mongolia. Korean Journal of Plant Taxonomy 51(4): 391–394. <https://doi.org/10.11110/kjpt.2021.51.4.391>
- Kechaykin AA, Kutsev M (2015) Notes on *Potentilla* L. (Rosaceae) from the Altai. 2. New species from South Siberia and West Mongolia. Feddes Repertorium 126(3–4): 73–76. <https://doi.org/10.1002/fedr.201500017>
- Knyazev MS (2020) *Astragalus habaheensis* (sect. *Erioceras*, Fabaceae) in Mongolia and China. Novitates Systematicae Plantarum Vascularium 51: 39–42. <https://doi.org/10.31111/novitates/2020.51.39>
- Korea National Arboretum (2021) Checklist of vascular plants in Korea (Alien plants). Korea National Arboretum, Pocheon, 305 pp.
- Kosachev PA (2017) Check-list of Scrophulariaceae Juss. s. l. of North Asia. Acta Biologica Sibirica 3(4): 31–76. [in Russian] <https://doi.org/10.14258/abs.v3i4.3631>
- Lazkov GA (2011) The genus *Phlomoides* (Lamiaceae) in Kirghizia. Komarovia 7: 1–64.
- Li WJ, Tojibaev KS, Hisoriev H, Shomurodov KF, Luo MF, Feng Y, Ma KP (2020) Mapping Asia Plants: Current status of floristic information for Central Asian flora. Global Ecology and Conservation 24: e01220. <https://doi.org/10.1016/j.gecco.2020.e01220>
- Liu DH, Xu XM, He Y, Liu QR (2021) A new combination in *Pseudolappula* (Boraginaceae, Rochelieae) based on morphological, molecular and palynological evidence. PhytoKeys 187: 77–92. <https://doi.org/10.3897/phytokeys.187.75346>

- Madhani H, Rabeler R, Pirani A, Oxelman B, Heubl G, Zarre S (2018) Untangling phylogenetic patterns and taxonomic confusion in tribe Caryophylleae (Caryophyllaceae) with special focus on generic boundaries. *Taxon* 67(1): 83–112. <https://doi.org/10.12705/671.6>
- Moore AJ, Dillenberger MS (2017) A conspectus of the *Cherleria* (*Minuartia* s.l., Caryophyllaceae). *Willdenowia* 47(1): 5–14. <https://doi.org/10.3372/wi.47.47101>
- Munkhnast D, Chuluunjav C, Urgamal M, Wong L J, Pagad S (2020) GRIIS Checklist of Introduced and Invasive Species - Mongolia. Version 2.7. Invasive Species Specialist Group ISSG.
- Murakami S, Takayama K, Fuse S, Hirota SK, Koi S, Ideno T, Yamamoto T, Tamura MN (2020) Recircumscription of sections of *Hemerocallis* (Asphodelaceae) from Japan and adjacent regions based on MIG-seq data. *Acta Phytotaxonomica et Geobotanica* 71: 1–11. <https://doi.org/10.18942/apg.201913>
- Nesom GL (2020) *Tibetiodes* Nesom, gen. nov. (Asteraceae: Astereae), including 27 species of Himalayan Aster. *Phytoneuron* 63: 1–30.
- Nobis M (2014) Taxonomic revision of the Central Asian *Stipa tianschanica* complex (Poaceae) with particular reference to the epidermal micromorphology of the lemma. *Folia Geobotanica* 49(2): 283–308. <https://doi.org/10.1007/s12224-013-9164-2>
- Nobis M, Ebel AL, Nowak A, Turginov OT, Kupriyanov AN, Nobis A, Olonova MV, Paszko B, Piwowarczyk R, Chen WL, Gudkova PD, Klichowska E, Nowak S, Pujadas-Salvà AJ (2014) Contribution to the flora of Asian and European countries: New national and regional vascular plant records, 2. *Acta Botanica Gallica* 161(2): 209–221. <https://doi.org/10.1080/12538078.2014.921643>
- Nobis M, Klichowska E, Terlević A, Wróbel A, Erst A, Hrvnák R, Ebel AL, Tikhomirov VN, Byalt VV, Gudkova PD, Király G, Kipriyanova LM, Olonova M, Piwowarczyk R, Pliszko A, Rosadziński S, Seregin AP, Honcharenko V, Marciniuk J, Marciniuk P, Oklejewicz K, Wolanin M, Batlai O, Bubíková K, Choi HJ, Dzhus MA, Kochjarová J, Molnár AV, Nobis A, Nowak A, Otáheľová H, Óvári M, Shimko II, Shukherdorj B, Sramkó G, Troshkina VI, Verkhozina AV, Wang W, Xiang K, Zykova EY (2019a) Contribution to the flora of Asian and European countries: New national and regional vascular plant records, 8. *Botany Letters* 166(2): 163–188. <https://doi.org/10.1080/23818107.2019.1600165>
- Nobis M, Gudkova PD, Nowak A (2019b) *Neotrinia* gen. nov. and *Pennatherum* sect. nov. in *Achnatherum* (Poaceae: Stipeae). *Turczaninowia* 22(1): 37–41. <https://doi.org/10.14258/turczaninowia.22.1.5>
- Nosov NN, Punina EO, Rodionov AV (2017) Synopsis of the grasses (Poaceae) of Altai Krai and Altai Republic. I. Genera *Arctopoa*, *Eremopoa* and *Poa*. *Turczaninowia* 20(2): 31–55. [in Russian] <https://doi.org/10.14258/turczaninowia.20.2.4>
- Nyambayar D (2009a) Flora of Mongolia. (Cyperaceae). Udam Soyol Publishing, Ulaanbaatar, 277 pp. [in Mongolian]
- Nyambayar D (2009b) The genus *Kobresia* Willd. (Cyperaceae Juss.) in the flora of Mongolia. *Mongolian Journal of Biological Sciences* 7: 7–13. <https://doi.org/10.22353/mjbs.2009.07.02>
- Ovczinnikova V, Korolyuk (2016). New species of the genus *Craniospermum* (Boraginaceae) from Mongolia. *Plant Life of Asian Russia* 3: 33–40. [in Russian] [https://doi.org/10.21782/RMAR1995-2449-2016-3\(33-40\)](https://doi.org/10.21782/RMAR1995-2449-2016-3(33-40)).
- Ovczinnikova V (2019a) Addition to the species composition of Boraginaceae of Outer Mongolia. *Turczaninowia* 22(3): 97–110. [in Russian] <https://doi.org/10.14258/turczaninowia.22.3.5>

- Ovczinnikova V (2019b) *Craniospermum volkovae* (Boraginaceae), a new species from Mongolia. Novitates Systematicae Plantarum Vascularium 50: 148–153. [in Russian] <https://doi.org/10.31111/novitates/2019.50.148>
- Ovczinnikova V (2020) Three new species of the genus *Craniospermum* (Boraginaceae) from Mongolia. Turczaninowia 23(2): 108–119. [in Russian] <https://doi.org/10.14258/turczaninowia.23.2.15>
- Oyunmaa J, de Priest P (2011) New records of the plant distribution of the phytogeographical region in Khuvsgul. Proceedings of Institute of Botany. Mongolian Academy of Sciences 23: 40–44. [in Mongolian]
- Pagad S, Genovesi P, Carnevali L, Schigel D, McGeoch MA (2018) Introducing the global register of introduced and invasive species. Scientific Data 5(1): e170202. <https://doi.org/10.1038/sdata.2017.202>
- Park I, Yang S, Song JH, Moon BC (2020) Dissection for floral micromorphology and plastid genome of valuable medicinal Borages *Arnebia* and *Lithospermum* (Boraginaceae). Frontiers in Plant Science 11: e606463. <https://doi.org/10.3389/fpls.2020.606463>
- Pfeiffer M, Dulamsuren C, Wesche K (2020) Grassland and shrubland of Mongolia. Reference Module in Earth Systems and Environmental Sciences, 759–772. <https://doi.org/10.1016/B978-0-12-409548-9.12057-3>
- Pimenov MG (2017) Updated checklist of Chinese Umbelliferae: Nomenclature, synonymy, typification, distribution. Turczaninowia 20(2): 106–239. <https://doi.org/10.14258/turczaninowia.20.2.9>
- Podlech D, Zarre SH [with collaboration of Ekici M, Maassoumi AAR, Sytin A] (2013) A taxonomic revision of the genus *Astragalus* L. (Leguminosae) in the Old World. Naturhistorisches Museum Wien, Vienna, 1039–1150.
- POWO (2021) Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. <http://www.plantsoftheworldonline.org/>
- PPG I (2016) A community-derived classification for extant lycophytes and ferns. Journal of Systematics and Evolution 54(6): 563–603. <https://doi.org/10.1111/jse.12229>
- Pyak AI, Pyak EA (2019) A new species of *Astragalus* section *Laguropsis* (Fabaceae: Galegeae) from a cross border highland region of Altai Mountains in Russia and Mongolia. Phytotaxa 414(4): 194–198. <https://doi.org/10.11646/phytotaxa.414.4.6>
- Pyak EA, Pyak AI, Madyka VV (2020) *Saussurea odorata* (Asteraceae), a new species from western Mongolia. Phytotaxa 470(3): 235–242. <https://doi.org/10.11646/phytotaxa.470.3.3>
- Pyšek P, Hulme PE, Simberloff D, Bacher S, Blackburn TM, Carlton JT, Dawson W, Essl F, Foxcroft LC, Genovesi P, Jeschke JM, Kühn I, Liebhold AM, Mandrak NE, Meyerson LA, Pauchard A, Pergl J, Roy HE, Seebens H, van Kleunen M, Vilà M, Wingfield MJ, Richardson DM (2020) Scientist' warning on invasive alien species. Biological Reviews of the Cambridge Philosophical Society 95(6): 1511–1534. <https://doi.org/10.1111;brv.12627>
- Ren C, Wang L, Illarionava ID, Yang QE (2020) Circumscription and phylogenetic position of *Ligularia* sect. *Stenostegia* (Asteraceae: Senecioneae) based on morphological, cytological, and molecular phylogenetic evidence. Taxon 69(4): 1–17. <https://doi.org/10.1002/tax.12280>
- Rilke S, Najmi U, Schnittler M (2012) Contributions to 'E-Taxonomy'—A virtual approach to the flora of Mongolia (Flora GREIF). Feddes Repertorium 123(3): 219–232. <https://doi.org/10.1002/fedr.201200013>

- Sennikov AN, Lazkov GA (2021) The first checklist of alien vascular plants of Kyrgyzstan, with new records and critical evaluation of earlier data. Contribution 1. Biodiversity Data Journal 9: e75590. <https://doi.org/10.3897/BDJ.9.e75590>
- Sennikov AN, Lazkov GA, Wong LJ, Pagad S (2021) Global register of introduced and invasive species – Kyrgyzstan. Invasive Species Specialist Group ISSG. <https://doi.org/10.15468/b8ro4l>
- Shiga T, Khaliunaa K, Baasanmunkh S, Oyuntsetseg B, Midorkawa S, Choi HJ (2020) New Mongolian records of two genera, seven species, and two hybrid nothospecies from Khar-US Lake and its associated wetlands. Journal of Asia-Pacific Biodiversity 13(3): 443–453. <https://doi.org/10.1016/j.japb.2020.06.008>
- Sinitsyna TA, Herden T, Friesen N (2018) Dated phylogeny and biogeography of the Eurasian *Allium* section *Rhizirideum* (Amaryllidaceae). Plant Systematics and Evolution 302(9): 1311–1328. <https://doi.org/10.1007/s00606-016-1333-3>
- Solovjev A (1998) A new species of the genus *Aconitum* L. (Ranunculaceae) from Mongolia. Turczaninowia 1: 5–8. [in Russian]
- Sramkó G, Laczkó L, Volkova PA, Bateman RM, Mlinarec J (2019) Evolutionary history of the Pasque-flowers (*Pulsatilla*, Ranunculaceae): Molecular phylogenetics, systematics and rDNA evolution. Molecular Phylogenetics and Evolution 135: 45–61. <https://doi.org/10.1016/j.ympev.2019.02.015>
- Sukhorukov AP, Uotila P, Zhang M, Zhang HX, Speranskaya AS, Krinitsyna AA (2013) New combinations in Asiatic *Oxybasis* (Amaranthaceae s.l.): Evidence from morphological, carpological and molecular data. Phytotaxa 144(1): 1–12. <https://doi.org/10.11646/phytotaxa.144.1.1>
- Sukhorukov AP, Liu PL, Kushunina M (2019) Taxonomic revision of Chenopodiaceae in Himalaya and Tibet. PhytoKeys 116: 1–146. <https://doi.org/10.3897/phytokeys.116.27301>
- Thiers B (2021) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih/> [accessed 15 Sep 2021]
- Troshkina VI (2021) Taxonomic revision of the family Geraniaceae in the flora of Mongolia. Turczaninowia 24(1): 9–20. [in Russian] <https://doi.org/10.14258/turczaninowia.24.1.2>
- Tungalag R (2020) Flora of Mongolia. (Amaranthaceae). Bembi San, Ulaanbaatar, 1–133. [in Mongolian]
- Ulziikhutag N, Dariimaa S, Ganbold E, Nyambayar D, Urgamal M, Zumberelmaa D, Enkhmaa U (2015) Flora of Mongolia. (Huperziaceae-Ephedraceae). Munkhiin Useg, Ulaanbaatar, 219 pp. [in Mongolian]
- Urgamal M (2009) Flora of Mongolia. (Apiaceae) Bembi San, Ulaanbaatar, 130 pp. [in Mongolian]
- Urgamal M, Oyuntsetseg B, Nyambayar D, Dulamsuren C (2014) Conspectus of the Vascular Plants of Mongolia. In: Sanchir C, Jamsran T (Eds) Ulaanbaatar, Mongolia. Admon, 1–334.
- Urgamal M, Oyuntsetseg B, Gundegmaa V, Munkh-Erdene T, Solongo K (2016) Additions to the vascular flora of Mongolia-IV. Proceeding of the Mongolian Academy of Sciences 56(2): 32–38. <https://doi.org/10.5564/pmas.v56i4.840>
- Urgamal M, Gundegmaa V, Baasanmunkh S, Oyuntsetseg B, Darikhant D, Munkh-Erdene T (2019) Additions to the vascular flora of Mongolia-IV. Proceeding of the Mongolian Academy of Sciences 59(2): 14–53. <https://doi.org/10.5564/pmas.v59i2.1218>

- Urgamal M, Munkh-Erdene T, Solongo K, Gundegmaa V, Amartuvshin N, Altantsetseg G (2020) Flora of Mongolia. (Ceratophyllaceae-Zygophyllaceae). Bembi San, Ulaanbaatar, 181 pp. [in Mongolian]
- Urgamal M, Munkh-Erdene T, Enkhmaa U, Zumberelmaa D, Ganbold E, Saruul N, Gantuya B (2021) Flora of Mongolia. (Nymphaeaceae-Asphodelaceae). Bembi San, Ulaanbaatar, 190 pp. [in Mongolian]
- van Kleunen M, Pyšek P, Dawson W, Essl F, Kreft H, Pergl J, Weigelt P, Stein A, Dullinger S, König C, Lenzner B, Maurel N, Moser D, Seebens H, Kartesz J, Nishino M, Aleksanyan A, Ansong M, Antonova LA, Barcelona JF, Breckle SW, Brundu G, Cabezas FJ, Cádenas D, Cádenas-Toro J, Castaño N, Chacón E, Chatelain C, Conn B, de Sá Dechoum M, Dufour-Dror JM, Ebel AL, Figueiredo E, Fragman-Sapir O, Fuentes N, Groom QJ, Henderson L, Inderjit, Jogan N, Krestov P, Kupriyanov A, Masciadri S, Meerman J, Morozova O, Nickrent D, Nowak A, Patzelt A, Pelser PB, Shu WS, Thomas J, Uludag A, Velayos M, Verkhosina A, Villaseñor JL, Weber E, Wieringa JJ, Yazlık A, Zeddam A, Zykova E, Winter M (2019) The Global Naturalized Alien Flora (GloNAF) database. *Ecology* 100: 1–2. <https://doi.org/10.1002/ecy.2542>
- Vinogradova YK, Aistova EV, Antonova LA, Chernyagina OA, Chubar EA, Darman GF, Devyatova EA, Khoreva MG, Kotenko OV, Marchuk EA, Nikolin EG, Prokopenko SV, Rubtsova TA, Shieko VV, Kudryavtseva EP, Krestov PV (2020) Invasive plants in flora of the Russian Far East: The checklist and comments. *Botanica Pacifica* 9(1): 103–129. <https://doi.org/10.17581/bp.2020.09107>
- Wang W, Li HL, Xiang XG, Chen ZD (2014) Revisiting the phylogeny of Ranunculeae: Implications for divergence time estimation and historical biogeography. *Journal of Systematics and Evolution* 52(5): 551–565. <https://doi.org/10.1111/jse.12101>
- Wang HF, Soejima A, Chang KS, Ma KP (2020) Mapping Asia Plants: Current status of floristic information for Northeast Asia. *Global Ecology and Conservation* 24: e01321. <https://doi.org/10.1016/j.gecco.2020.e01321>
- Wesche K, Ambarlı D, Kamp J, Török P, Treiber J, Dengler J (2016) The Palaearctic steppe biome: A new synthesis. *Biodiversity and Conservation* 25(12): 2197–2231. <https://doi.org/10.1007/s10531-016-1214-7>
- Wiegleb G, Bobrov AA, Zalewska-Gałoz J (2017) A taxonomic account of *Ranunculus* section *Batrachium* (Ranunculaceae). *Phytotaxa* 319(1): 1–55. <https://doi.org/10.11646/phytotaxa.319.1.1>
- Xu H, Qiang S, Genovesi P, Ding H, Wu J, Meng L, Han Z, Miao J, Hu B, Guo J, Sun H, Huang C, Lei J, Le Z, Zhang X, He S, Wu Y, Zheng Z, Chen L, Jarošík V, Pyšek P, Pysek P (2012) An inventory of invasive alien species in China. *NeoBiota* 15: 1–26. <https://doi.org/10.3897/neobiota.15.3575>
- Yano O, Shiga T, Khaliunaa K, Baasanmunkh S, Oyuntsetseg B, Choi HJ (2021) A new record of *Carex capricornis* (Cyperaceae) from Mongolia. *Japanese Journal of Botany* 96(4): 238–241.
- Yurtseva OV, Kuznetsova OI, Mavrodiev EV (2016) A broadly sampled 3-loci plastid phylogeny of *Atraphaxis* (Polygoneae, Polygonoideae, Polygonaceae) reveals new taxa: I. *Atraphaxis kamelinii* spec. nov. from Mongolia. *Phytotaxa* 268(1): 1–24. <https://doi.org/10.11646/phytotaxa.268.1.1>

- Zaika MA, Kilian N, Jones K, Krinitina AA, Nilova MV, Speranskaya AS, Sukhorukov AP (2020) *Scorzonera* sensu lato (Asteraceae, Cichorieae) – taxonomic reassessment in the light of new molecular phylogenetic and carpological analyses. *PhytoKeys* 137: 1–85. <https://doi.org/10.3897/phytokeys.137.46544>
- Zemmrich A, Schnittler M, Hartleib J, Busch M, Bobertz B, Zolitz R (2013) FloraGREIF – An Internet-based data repository for biogeographical research in Mongolia. *Folia Geobotanica* 48(4): 523–536. <https://doi.org/10.1007/s12224-012-9149-6>
- Zhang GJ, Nie B, Gao TG (2018) *Aster lingii*, a new name for *Rhinactinidia limoniifolia* (Astereae, Asteraceae). *Phytotaxa* 344(3): 297–299. <https://doi.org/10.11646/phytotaxa.344.3.12>
- Zhao LQ, Ri X, Alatanzhula, Gundegmaa V, Qiao X-G, Mungunchimeg C (2019) *Stipa khovdensis* (Poaceae), a new species, and a checklist of *Stipa* s. stricto from Mongolia. *Annales Botanici Fennici* 56(1–3): 95–99. <https://doi.org/10.5735/085.056.0114>

## Supplementary material 1

### Appendix 1

Authors: Baasanmunkh et al.

Data type: Checklist

Explanation note: List of new species (marked by red color) and new records to the flora of Mongolia, since Urgamal et al (2014).

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/phytokeys.192.79702.suppl1>

## Supplementary material 2

### Appendix 2

Authors: Baasanmunkh et al.

Data type: Checklist

Explanation note: Changes of accepted names compared with Urgamal et al (2014).

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/phytokeys.192.79702.suppl2>