

Spiradiclis pengshuiensis (Ophiorrhizeae, Rubioideae), a new species from Chongqing, China

Bo Pan¹, Hu-Sheng Ma¹, Rui-Jiang Wang²

1 Guangxi Institute of Botany, Guangxi Zhuangzu Autonomous Region and Chinese Academy of Sciences, Guilin 541006, China **2** Key Laboratory of Plant Resources Conservation and Sustainable Utilization, Guangdong Provincial Key Laboratory of Applied Botany, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China

Corresponding author: Rui-Jiang Wang (wangrj@scbg.ac.cn)

Academic editor: Pavel Stoev | Received 3 February 2016 | Accepted 14 May 2016 | Published 18 May 2016

Citation: Pan B, Ma H-S, Wang R-J (2016) *Spiradiclis pengshuiensis* (Ophiorrhizeae, Rubioideae), a new species from Chongqing, China. *PhytoKeys* 63: 41–45. doi: 10.3897/phytokeys.63.8016

Abstract

Spiradiclis pengshuiensis Bo Pan & R. J. Wang (Rubiaceae) is described as a new species from Chongqing in SW China. It is morphologically compared with *S. pauciflora* L. Wu & Q. R. Liu because of their similarities in habit, pubescent surface, small leaf laminas and subglobose capsules. Its conservation status is evaluated as “VU” according to the IUCN categories and criteria.

Keywords

China, New taxon, Rubiaceae, *Spiradiclis*

Introduction

The genus *Spiradiclis* Blume (Rubiaceae) mainly distributed in the tropical and sub-tropical limestone areas of Southern China and Northern Vietnam. Geographically, the localities of *Spiradiclis* species can extend northward to Mt. Emei (29°33'N) in Sichuan province of China and southward to Central Java (ca. 7°10'S) of Indonesia. The genus comprise about 50 species and ca. 91% species can be found in China, the center of species diversity (Chen and Taylor 2011, Deng et al. 2014, Wang et al. 2015, Wen et al. 2015, Wu et al. 2015, Wang 2016). *Spiradiclis* species are usually similar to *Ophiorrhiza* L. because of their similar inflorescence and flower characters,

but the former is characterized by globose or ovoid (vs. strongly laterally compressed in *Ophiorrhiza*) capsules.

During a botanical inventory in Pengshui County, the east of Chongqing Municipality, in 2013, the senior author found a striking *Spiradiclis* species growing on the dry cliffs. In order to get enough materials for morphological observation and comparison, we successively collected the vouchers during the flowering and fruiting seasons in recent years. This new species can be readily distinguished from other *Spiradiclis* species in habit and flowers and herein described and illustrated.

Material and methods

All materials were collected by ourselves and deposited at the herbarium of South China Botanical Garden, Chinese Academy of Sciences (IBSC). The morphological data were collected by a Digimatic Caliper (Mitutoyo, Japan).

Taxonomy

Spiradiclis pengshuiensis B. Pan & R. J. Wang, sp. nov.

urn:lsid:ipni.org:names:77154905-1

Figure 1

Diagnosis. *Spiradiclis pengshuiensis* is similar to *S. pauciflora* L. Wu & Q. R. Liu, from which it differs in having linear (vs. triangular) stipules, 4–6 (vs. 3–4) secondary veins each side of the leaf laminae, salverform (vs. funnellform) corolla with 9–15 mm (vs. 7–9 mm) long tubes (Table 1).

Type. CHINA. Chongqing Municipality: Pengshui County, Hanxia Town, Baixiang Village, 29°8'N, 108°6'E, 360 m alt., 29 Nov 2014, Rui-Jiang Wang 2931, long-styled flower (holotype IBSC; isotypes IBSC).

Description. Perennial herbs, 4–9 cm tall, densely pubescent on plant surfaces; stems erect or prostrate at base and then ascending, terete, rooting at nodes; internodes 2–15 mm long. Stipules linear, 0.9–1.2 mm long. Petiole (3.5–)7–12(–23.5) mm long. Leaf blades opposite, ovate, (4–)7–15(–25) × (2.5–)5–10(–15) mm; base cuneate or late cuneate, asymmetry, decurrent; apex obtuse to rounded, papery, green adaxially and pale green abaxially; secondary veins 4–6 on each side; margin revolute slightly. Inflorescence terminal, cymose, 1–7(–12)-flowered but only 1–3 blooming simultaneously; peduncles 1–2.1 cm long; bracts and bracteoles linear, 1–2 mm long. Flowers distylous; pedicels 0–2 mm long; hypanthium obconical, 1–2 mm long; calyx lobes 5, lanceolate, 0.8–1.5 × ca. 0.4 mm. Corolla salverform, white adaxially, pinkish to white abaxially, pubescent both sides; tube 9–15 × 1.5–2 mm; lobes (4–)5, 5–7.5 × 2.5–3.5 mm. Stamens 5; anthers linear-oblong, 1.5–2 mm long, dorsifixed. Stigma bilobed, lobes ovoid; ovary 2-celled, ovules many, axile. Long-styled flowers:

Table 1. Morphological comparison of *Spiradiclis pengshuiensis* and *S. pauciflora*.

Characters	<i>S. pengshuiensis</i>	<i>S. pauciflora</i>
Habit	Erect or prostrate at base	Creeping or with upper parts ascending
Indument	Pubescent whole plant surfaces	Pubescent whole plant surfaces
Stem	Terete	Terete
Stipules	Linear, 0.9–1.2 mm long	Triangular, less than 1 mm
Petioles length (mm)	3.5–23.5	3–10
Leaf blades	Ovate; base cuneate or late cuneate, asymmetry, decurrent; apex obtuse to rounded	Ovate to elliptic-ovate; base obtuse to broadly cuneate; apex obtuse to acute
Leaf texture	Papery	Papery
Leaf size (mm)	(4–) 7–15(–25) × (2.5–) 5–10 (–15)	5–20 × 5–15
Leaf color	Green adaxially and pale green abaxially	Olive-green adaxially, pale or sometime purplish red abaxially
Secondary veins on each side	4–6	3–4
Inflorescence	Terminal, 1–7(–12)-flowered	Terminal, 3–7(–9)-flowered
Peduncle length (cm)	1–2.1	1–3
Pedicel length (mm)	0–2	0.3–3
Calyx lobes	Lanceolate, 0.8–1.5 mm long	Ovate-triangular, 1.2–1.6 mm long
Corolla	Salverform; corolla tube 9–15 cm long; lobes oval, 5–7.5 mm long	Funnelform; corolla tube 7–9 mm long; lobes ovate-triangular, 2.5–3 mm long
Pin flowers	Corolla pubescent adaxially, without ring hairs; styles 9–12 mm long	Ring hairs present in adaxial side of corolla; styles 7.5–8.5 mm long
Style length in thrum flowers (mm)	ca. 5	2.7–3
Capsules	Subglobose, 3–5 mm in diam.	Subglobose, ca. 2 mm in diam.
Phenology	Flowering in November to next January, fruiting in December to February	Flowering in March to June, fruiting in May to August

stamens included; filaments adnate to the lower corolla tube, ca. 1.5 mm long; styles 9–12 mm long; stigma ca. 0.5 mm long, red to brown, extended to corolla throat, not exerted. Short-styled flowers: stamens extended to corolla throat, not exerted; filaments adnate to upper corolla tube, ca. 0.5 mm long; styles ca. 5 mm long; stigmas ca. 1.5 mm long, included. Capsules subglobose, 3–5 mm in diam., with persistent calyx lobes, dehiscent septically and loculicidally; valves 4, ovate, 3–5 mm long. Seeds 15–20 per capsule, ca. 0.5 mm long, brown, irregular pyramid to cuboid, foveolate on surface.

Phenology. Flowers from November to next January; fruits from December to next February.

Distribution and habitat. *Spiradiclis pengshuiensis* is known only from the type locality. Its habitat is on the cliffs nearby the A-Yi-He river but with very poor conditions.

Conservation status. Only three small populations and less than 500 individuals of *Spiradiclis pengshuiensis* within 5 km² were found in the A-Yi-He Scenic Area. The tourist activity is the primary factor in the decline of populations. In addition, the




Figure 1. *Spiradiclis pengshuiensis* sp. nov. **A** habitat **B** habit **C** linear stipules **D** visiting insects **E** bracts and hypanthium **F–I** longitudinal section of long- (**F, H**) and short-styled flowers (**G, I**), respectively, showing the induments, relative positions and morphology of the stigmas and anthers and the developing capsules in distylous flowers **J** infructescence **K–L** transverse and longitudinal section of young capsules, respectively **M** mature capsule **N** dehiscent capsules **O** seeds. Photos by Ruijiang Wang.

plants always grow on the cliffs with little soil and insufficient water, which limited the development and dispersal of the species. Therefore we assign a preliminary IUCN threat status of Vulnerable [VU, B2ab(ii, iii, iv)] to *S. pengshuiensis* (IUCN 2001).

Specimens examined. CHINA. **Chongqing Municipality:** Pengshui County, Hanxia Town, Baixiang Village, 29°8'N, 108°6'E, 350 m alt., 29 Nov 2014, short-styled flowers, Rui-Jiang Wang 2932, 2937 (IBSC); same locality, 12 Jan 2016, fruiting, Rui-Jiang Wang 3095, 3099 (IBSC).

Acknowledgements

This work was supported by National Natural Science Foundation of China (31070177) and Natural Science Foundation of Guangxi (2013GXNSFBA019078).

References

- Chen T, Taylor CM (2011) *Spiradiclis* Blume. In: Wu ZY, Raven PH (Eds) Flora of China, vol. 19. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 330–339.
- Deng SJ, Wen HZ, Huang XX, Wang RJ (2014) *Spiradiclis coriaceifolia* and *S. tonglingensis* spp. nov. (Rubiaceae, Ophiorrhizeae) from Guangxi, China. Nordic Journal of Botany 32: 594–601. doi: 10.1111/njb.00461
- IUCN (2001) IUCN Red List Categories and Criteria (Version 3.1). IUCN Species Survival Commission, Gland, Switzerland and Cambridge. http://www.iucnredlist.org/info/categories_criteria2001
- Wang RJ, Wen HZ, Deng SJ, Zhou LX (2015) *Spiradiclis danxiashanensis* (Rubiaceae), a new species from South China. Phytotaxa 206(1): 30–36. doi: 10.11646/phytotaxa.206.1.5
- Wang RJ (2016) *Spiradiclis yangchunensis* (Rubiaceae), a new species from Guangdong, China. Plant Science Journal 34(1): 13–17. doi: 10.11913/PSJ.2095–0837.2016.10013
- Wen HZ, Wang RJ, Deng SJ (2015) *Spiradiclis longanensis*, a new species of Rubiaceae from China. PhytoKeys 55: 113–117. doi: 10.3897/phytokeys.55.4975
- Wu L, Wang JL, Liu QR (2015) *Spiradiclis pauciflora* (Rubiaceae), a new species from limestone areas in Guangxi, China. Annales Botanici Fennici 52(3–4): 257–261. doi: 10.5735/085.052.0318