

***Thylacopteris minuta* (Polypodiaceae), a new fern species from Myanmar**

Kiyotaka Hori¹, Phyto Kay Khine², Tao Fujiwara^{2,3}, Thant Shin⁴, Harald Schneider²

1 The Kochi Prefectural Makino Botanical Garden 4200-6 Godaisan, Kochi 781-8125, Japan **2** Center for Integrative Conservation, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Menglin, Mengla, Yunnan 666303, China **3** Makino Herbarium, Tokyo Metropolitan University, 1-1 Minami-osawa, Hachioji, Tokyo 192-0397, Japan **4** Forest Research Institute, Yezin, Nay Pyi Taw, Myanmar

Corresponding author: Kiyotaka Hori (katori@makino.or.jp)

Academic editor: Joel Nitta | Received 4 March 2022 | Accepted 26 May 2022 | Published 9 June 2022

Citation: Hori K, Khine PK, Fujiwara T, Shin T, Schneider H (2022) *Thylacopteris minuta* (Polypodiaceae), a new fern species from Myanmar. *PhytoKeys* 199: 141–153. <https://doi.org/10.3897/phytokeys.199.83107>

Abstract

The genus *Thylacopteris* is a small, phylogenetically isolated genus belonging to the fern family Polypodiaceae. This study describes a new species, *Thylacopteris minuta*, based on collections obtained during field surveys of Shan State, Myanmar. This new species is distinct from other species of *Thylacopteris* in its small size and presence of sclerenchyma strands in the rhizome. This species is also distinct from the only other species of *Thylacopteris* with molecular data available, *T. papillosa*, in a plastid *rbcL* phylogeny of Polypodiaceae. This new discovery of *Thylacopteris* from Myanmar suggests that this genus is still overlooked in Southeast Asia.

Keywords

Myanmar, new species, Polypodiaceae, *Thylacopteris*

Introduction

With about 1,652 species, Polypodiaceae is the second largest family of pteridophytes (ferns and lycophytes) (PPGI 2016). Species of this family share the presence of some characteristics such as yellowish or greenish spores, round or flat, yellowish-brown, ex-indusiate sori, and creeping rhizomes usually covered by scales (Zhang et al. 2013). Despite the progress in our understanding of the generic classification of these ferns as summarized in PPGI (2016), some issues are still unresolved. Based on recent molecular studies, Testo et al. (2019) segregated the genus *Bosmania* Testo, *Dendroconche* Copel.

and *Zealandia* Testo from the paraphyletic genus *Microsorum*. Link, and Zhao et al. (2020) expanded the definition of the genus *Lepisorus* (J.Sm.) Ching. In addition, new species are still being discovered in Polypodiaceae in recent studies (Khine et al. 2016; Zhao et al. 2017) around Southeast Asia. To improve the pteridophyte flora of Southeast Asia, collections and molecular studies of Polypodiaceae are still needed in this area. In this study, we address the relationships of *Thylacopteris* specimens obtained in Shan State, Myanmar. The genus *Thylacopteris* includes currently only two species (Rödl-Linder 1994) of which one, *Thylacopteris diaphana* (Brause) Copel., is endemic to New Guinea. The other species, *T. papillosa* (Blume) Kunze ex J.Sm, is known to occur throughout Malesia but has not been found in the north of the Isthmus of Kra (Rödl-Linder 1994).

The genus *Thylacopteris* Kunze ex J.Sm. was established by Smith (1875) with *T. papillosa* as the type species. This species was originally described as *Polypodium pilosum* Blume based on an accession collected in Java (Blume 1828). As distinct characteristics of *Thylacopteris* from *Polypodium* sensu stricto, Blume (1828) mentioned the deeply sunken sori and the articulation of the lateral segments to the rachis. Subsequently, Copeland (1947) added a second species by introducing the combination *T. diaphana* (Brause) Copel. based on *P. diaphanum* Brause, which was based on an accession collected in New Guinea (Brause 1912). Some studies pointed out the close relationships of the genus *Thylacopteris* with either *Goniophlebium* or *Polypodium* (Christensen and Holttum 1934; Ching 1978; Tryon and Tryon 1982). Subsequently, *Thylacopteris* was treated as a group of *Polypodium* (Holttum 1966; Tryon and Lugard 1991), or of uncertain systematic position in Polypodiaceae (Hennipman et al. 1990). Finally, utilizing DNA-based phylogenetics, *Thylacopteris* was found to be sister to a clade including *Goniophlebium*, *Lepisorus*, and *Microsorum* (Schneider et al. 2004). Considering the reported results of phylogenetic studies focusing on Polypodiaceae, *Thylacopteris* was included in a broadly defined Microsoroideae in PPG I (2016). Reflecting its isolated phylogenetic position, Chen et al. (2019, 2021) introduced a tribe Thylacopteraceae only including *Thylacopteris*.

In the recent years, some studies have reported new species (Khine et al. 2016) or new records of ferns (Nwe et al. 2016; Khine et al. 2017; Hori et al. 2019; Khine and Schneider 2020) from Myanmar. The discovery of new species and new records reflect the lack of comprehensive fern floristic studies of Myanmar (Khine et al. 2017; Khine and Schneider 2020), which limit the ability to report and manage the conservation of Myanmar's unique biodiversity (Khine and Schneider 2020). Here, we describe a new species of the genus *Thylacopteris* from Myanmar and the first record of this genus in the country based on morphological characteristics and molecular phylogenetic analysis of Polypodiaceae.

Materials and methods

We collected three specimens in Shan State, Myanmar during the inventories conducted under the leadership of the Makino Botanical Garden team on 13th September 2015, 16th September 2015, and 30th September 2019 together with the team from the

Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences. To identify the new species *Thylacopteris minuta* sp. nov., the following characteristics were studied carefully and compared to the description and specimens of previously described species: the size and shape of plants, morphology of leaves including the shape of segments, stipe, anatomy of rhizome, scales, and the morphology of reproductive organs including sori, sporangia, and spores. Voucher specimens were deposited at the herbarium of the Kochi Prefectural Makino Botanical Garden (MBK), Xishuangbanna Tropical Botanical Garden (HITBC), Queen Sirikit Botanic Garden (QBG) and the Forest Research Institute, Myanmar (RAF). Herbarium codes follow Thiers (2021).

The plastid *rbcL* gene was employed for phylogenetic analysis. Total DNA was extracted from silica-dried leaves using cetyltrimethylammonium bromide (CTAB) solution according to the method of Doyle and Doyle (1990). PCR amplification was performed using the primers af3 and cr3 (Hori et al. 2018) and PrimeSTAR Max DNA Polymerase (Takara, Kyoto, Japan). PCR involved an initial denaturation step at 95 °C for 10 min, followed by 35 cycles of denaturation, annealing, and elongation steps at 98 °C for 10 s, 55 °C for 5 s, and 72 °C for 8 s, respectively (Model 9700 Thermal Cycler, Applied Biosystems, Foster City, CA, USA). The PCR products were purified using Illustra ExoStar 1-Step (GE Healthcare, Wisconsin, USA) and used as templates for direct sequencing. Reaction mixtures for sequencing were prepared using the BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Foster City, CA, USA). The reaction mixtures were analyzed using an ABI 3130 Genetic Analyzer (Applied Biosystems, Foster City, CA, USA).

To estimate the phylogenetic position of the accession of interest, plastid *rbcL* sequences of Polypodiaceae were obtained from Genbank (<https://www.ncbi.nlm.nih.gov/genbank/>), covering all genera accepted in PPG I (2016) as far as data availability enabled. To reflect recent progress in our understanding of the natural classification of Polypodiaceae, the treatment of some taxa deviated from PPGI by adapting new concepts (Testo et al. 2019; Zhao et al. 2020; Chen et al. 2021). In the Genbank database, some accessions contained indels, which should not be present in *rbcL* since it is a protein-coding gene; we removed such low-quality accessions. The final data set included 94 accessions of Polypodiaceae, three samples of *Thylacopteris minuta* from Myanmar, and a set of outgroup taxa including *Davallia*, *Oleandra*, *Nephrolepis* and *Tectaria* (Table 1). The *rbcL* sequences were aligned using MUSCLE (Edgar 2004) and analyzed with Bayesian inference (BI) using MrBayes 3.2.6 (Ronquist et al. 2012) and maximum likelihood (ML) using MEGA X software (Kumar et al. 2018). Based on BIC values, GTR + G + I model was selected as the best-fit model of sequence evolution for BI analysis by jModelTest 2.1.10 (Darriba et al. 2012), and Tamura 3-parameter + G + I model was selected for the ML analysis by MEGA X software. Four chains of Markov chain Monte Carlo were run simultaneously and sampled every 100 generations for 1 million generations in total. Tracer 1.7.1 (Rambaut et al. 2018) was used to examine the posterior distribution of all parameters and their associated statistics, including estimated sample sizes. The first 2,500 sample trees from each run were discarded as burn-in. In ML analysis, initial trees for the heuristic search were obtained automatically by applying the Neighbor-Join and BioNJ algorithms to a ma-

Table 1. Accessions of rbcL sequences in this study.

Accessions of rbcL sequences	Species
AF468205	<i>Adenophorus montanus</i>
AY529147	<i>Aglaomorpha acuminata</i>
AF470349	<i>Aglaomorpha coronans</i>
AY529150	<i>Aglaomorpha heraclea</i>
MW138159	<i>Alansmia smithii</i>
MT215977	<i>Archigrammitis marquesensis</i>
JQ685380	<i>Arthromeris lehmannii</i>
MG948938	<i>Ascogrammitis anfractuosa</i>
EU482962	<i>Bosmania membranacea</i>
MT215995	<i>Calymmodon cucullatus</i>
MF318061	<i>Campyloneurum brevifolium</i>
MF317971	<i>Campyloneurum lorentzii</i>
MF318013	<i>Campyloneurum rigidum</i>
MW138183	<i>Ceradenia kalbreyeri</i>
KM218797	<i>Chrysogrammitis musgraveana</i>
EF178615	<i>Cochlidium serrulatum</i>
MT657584	<i>Ctenopterella blechnoides</i>
KM218775	<i>Dasygrammitis brevivenosa</i>
MZ957125	<i>Davallia pulchra</i>
KM114198	<i>Davallia solida var. fejeensis</i>
MN018180	<i>Dendroconche annabellae</i>
MN018176	<i>Dendroconche sayeri</i>
DQ227292	<i>Dictymia brownii</i>
DQ164441	<i>Dictymia mckeei</i>
MW138254	<i>Enterosora trifurcata</i>
KM218794	<i>Galactodenia parrisiae</i>
MN017598	<i>Goniophlebium amoenum</i>
AB043100	<i>Goniophlebium formosanum</i>
DQ078627	<i>Goniophlebium microrhizoma</i>
AB043098	<i>Goniophlebium niponicum</i>
MT657640	<i>Goniophlebium percussum</i>
AB043099	<i>Goniophlebium persicifolium</i>
MT657642	<i>Goniophlebium subauriculatum</i>
MT216033	<i>Grammitis cincta</i>
AB232409	<i>Gymnogrammitis dareiformis</i>
AF470322	<i>Lecanopteris carnosia</i>
AF470329	<i>Lecanopteris crustacea</i>
AF470325	<i>Lecanopteris luzonensis</i>
GU387043	<i>Lellingeria dissimulans</i>
MT169815	<i>Lepisorus carnosus</i>
MN623364	<i>Lepisorus hederaceus</i>
MT169813	<i>Lepisorus longifolius</i>
MT169824	<i>Lepisorus normalis</i>
AY362564	<i>Lepisorus nudus</i>
EU482971	<i>Lepisorus superficialis</i>
GQ256304	<i>Lepisorus thunbergianus</i>
GQ256310	<i>Lepisorus uchiyamae</i>
MH768462	<i>Leptochilus decurrens</i>
MH768470	<i>Leptochilus digitatus</i>
MH768471	<i>Leptochilus saxicola</i>
GU376488	<i>Leucotrichum mitchellae</i>
KF992501	<i>Loxogramme lanceolata</i>

Accessions of rbcL sequences	Species
DQ227294	<i>Loxogramme salicifolia</i>
GU476898	<i>Melpomene anazalea</i>
MW138194	<i>Microgramma lycopodioides</i>
AY362579	<i>Microgramma squamulosa</i>
MF317960	<i>Microgramma vacciniifolia</i>
AY362344	<i>Micropolypodium hyalinum</i>
LC496693	<i>Microsorum cuspidatum</i>
KY099830	<i>Microsorum membranifolium</i>
DQ179633	<i>Microsorum scolopendria</i>
MW620392	<i>Moranopteris taenifolia</i>
MT216066	<i>Nephrolepis cordifolia</i>
MT216068	<i>Nephrolepis hirsutula</i>
EF463254	<i>Niphidium crassifolium</i>
MF317999	<i>Niphidium longifolium</i>
JQ904094	<i>Notogrammitis billardierei</i>
EF463242	<i>Oleandra articulata</i>
AB232405	<i>Oleandra pistillaris</i>
MT657589	<i>Oreogrammitis forbesiana</i>
EF463255	<i>Pecluma eurybasis</i>
AY362588	<i>Pecluma ptilodon</i>
KT780748	<i>Pecluma sicca</i>
MW138202	<i>Phlebodium pseudoaureum</i>
MN623367	<i>Platycerium bifurcatum</i>
AY362591	<i>Pleopeltis fructuosa</i>
EF463258	<i>Pleopeltis sanctae-rosae</i>
KF909057	<i>Pleurosoriopsis makinoi</i>
KF909059	<i>Polypodium scouleri</i>
KF186527	<i>Polypodium virginianum</i>
AB044899	<i>Polypodium vulgare</i>
MT657600	<i>Prosaptia alata</i>
EF463259	<i>Pyrrosia polydactyla</i>
AY362558	<i>Pyrrosia rupestris</i>
EF463260	<i>Pyrrosia serpens</i>
KM218771	<i>Radiogrammitis holttumii</i>
AY096199	<i>Selliguea feei</i>
AF470347	<i>Selliguea hastata</i>
AY529171	<i>Selliguea laciniata</i>
MW138195	<i>Stenogrammitis limula</i>
DQ168808	<i>Synammia intermedia</i>
KF667652	<i>Tectaria griffithii</i>
EF463274	<i>Tectaria trifoliata</i>
KM218802	<i>Terpsichore asplenifolia</i>
KM218758	<i>Themelium decrescens</i>
LC685475	<i>Thylacopteris minuta</i> sp.nov., Baba et al. 103191
LC685476	<i>Thylacopteris minuta</i> sp.nov., Baba et al. 103361
LC685054	<i>Thylacopteris minuta</i> sp.nov., Hori et al. 108601
AY459175	<i>Thylacopteris papillosa</i>
MH665089	<i>Thylacopteris papillosa</i>
KM218780	<i>Tomophyllum macrum</i>
MG452028	<i>Zealandia powellii</i>
DQ401117	<i>Zealandia pustulatum</i>
DQ179635	<i>Zealandia vieillardii</i>
KM218793	<i>Zygophlebia devoluta</i>

trix of pairwise distances estimated using the Tamura 3-parameter model, and then selecting the topology with superior log likelihood value. A discrete Gamma distribution was used to model evolutionary rate differences among sites (5 categories (+G, parameter = 0.9922)). The rate variation model allowed for some sites to be evolutionarily invariable ([+I], 57.20% sites). The tree was drawn to scale, with branch lengths measured in the number of substitutions per site. The bootstrap method with 1,000 replications was employed in ML analysis.

Results and discussions

The aligned matrix included 1209 bp of *rbcL*, of which 329 bp (27%) were parsimony-informative. The ML (the highest log likelihood = -11294.22) tree showed that the three accessions of *Thylacopteris minuta* sp.nov (Baba et al. 103191, 103361, Hori et al. 108601) comprised a clade with two accessions of *T. papillosa* (Fig. 1). The *rbcL* sequence of Myanmar accessions of *Thylacopteris* had 25 substitutions relative to *T. papillosa*. The phylogenetic placement of *T. diaphana* is unresolved because DNA sequences were not available in this study. However, at least, morphological characteristics of the new species *T. minuta* can be differentiated from those of *T. diaphana* and *T. papillosa* as described below.

Taxonomic treatment

Thylacopteris minuta K.Hori & Khine, sp. nov.

urn:lsid:ipni.org:names:77299336-1

Diagnosis. *Thylacopteris minuta* is similar to *T. papillosa* with 20–40 sclerenchyma strands per rhizome in cross-section. However, *T. minuta* is distinct from *T. papillosa* with sori shallowly sunken vs. *T. papillosa* sori deeply sunken. In addition, the lamina of *T. minuta* has a maximal length of 15 cm vs. a maximal length of 59 cm in *T. papillosa*. *Thylacopteris minuta* is distinct from the New Guinea endemic *T. diaphana*, which lacks sclerenchyma strands in the rhizome, has superficial sori, and lamina with a maximal length of 45 cm.

Type. MYANMAR: Shan State; Ah Lel Chaung reserve forest, Ywangan Township. 20°59'44.8"N, 96°34'26.81"E, ca.1325 m, 30 Sep. 2019, K. Hori, P.K. Khine ["Kine"], T. Fujiwara, M. Nagashima, P.P. Shwe & A.K. Moe 108601 (holotype: MBK 0328421 (herbarium barcode), Figs 2–5 isotype: HITBC, RAF).

Epilithic. **Rhizome** long-creeping, weakly branched, 1.0–2.0 mm in diam. (without scales), light brown, densely clothed with scales, phylloodia sometimes prominent, these 1.0–2.0 mm high, 5.0–10.0 mm apart; 20–40 black sclerenchyma strands in rhizome, longitudinal, scattered in the ground tissue. **Rhizome scales** evenly inserted, dull brown, fragile, adpressed or apically spreading, quite densely set, deciduous, deltoid or ovate, 1.0–1.5 mm long, 0.5–1.0 mm wide, gradually narrowed from base to apex, sometimes with wavy margins, apex acute or rounded. **Cell walls of**




Figure 1. Phylogenetic hypothesis selected by applying maximum likelihood (the highest log-likelihood = -11294.22) to *rbcL* sequences. Posterior probabilities (> 0.90) and bootstrap percentages (> 70%) of Bayesian inference/maximum likelihood analyses are depicted at each node.




Figure 2. *Thylacopteris minuta* K. Hori & P.K.Khine (holotype, Hori et al. 108601 = MBK0328421, illustration by K.Hori) **A** habit **B** abaxial view of a middle segment **C** adaxial view of a middle segment **D** cross-section of rhizome **E** spore **F** rhizome scales **G** sporangium. Scale bars: 5 cm (**A1–A3**); 1 cm (**B–C**); 0.5 mm (**D**); 10 µm (**E**); 20 µm (**F**); 200 µm (**G**).




Figure 3. Warty, thickened cell walls of rhizome scales (MBK 0328421; photo taken using DP20 microscope camera, OLYMPUS, Japan). Scale bar: 2.5 μm .




Figure 4. The distribution of *Thylacopteris minuta* sp.nov.



Figure 5. *Thylacopteris minuta* in its natural habitat (MBK 0328421). Scale bar: 5 cm. (Photo by P.K. Khine).

rhizome scales dark brown, jigsaw-puzzle-shaped and wavy at basal and central part of scales, thickened, densely warty, in a single layer or double layers in basal scales. **Fronds** monomorphic, articulate to rhizome, petiolate. **Stipes** glabrous, 3.0–5.0 cm long, 0.7–1.0 mm diam, yellowish green. **Blades** membranous, lanceolate, 7.0–15.0 cm long, 2.0–4.0 cm wide, equally wide all along or rather wider above the basal part, pinnatisect, yellowish green. **Segments** glabrous, 20–30 pairs, lanceolate, ascending at an angle of 90°, 0.8–2.3 cm long, 0.3–0.5 cm wide, apically obtuse, entire at basal margin, crenate at apical margin, lower segments sometimes reduced, apical segments continuously reduced, terminal segments adnate or caudate. **Veins** free, once-forked, excurrent with terminal hydathodes. **Sori** exindusiate, uniserial on each side of costa, placed medially between costa and margin, shallowly sunken, 0.5–1.0 mm in diam., depth of papillae 0.2–0.5 mm, paraphyses absent. **Sporangium** globe-shaped, long stalked, 200–250 µm in diam., annulus vertical, indurated cells 10–13. **Spores** bilateral, oblong, light yellow, 40–60 µm long, 25–35 µm wide in lateral view, laesura 20–25 µm long, exospore smooth, perispore thin, surface shallowly wrinkled, globules absent.

Distribution. Myanmar.

Habitat. Epilithic, growing on shady surfaces of limestones (1–3 m high) in semi-evergreen or evergreen forest; altitude 940–1450 m.

Etymology. The name refers to the relatively small size of this species compared to other species of *Thylacopteris*.

Additional specimens examined. Myanmar: Shan State; Phaya Taung, Lein Le village, Paunglang Reserve Forest, Pinlaung Township; 19°59'41.0"N, 96°39'3.0"E, ca.947 m alt., 13 Sep. 2015, Y. Baba, K. Kertsawang, C. Kilgour, C. Puglisi, M. Rodda, P. Srisanga, T. Shin & P.P. Hnin 103191 (MBK0306471, duplicates on RAF, QBG). ibid., road between Nyaung Phyue village and Pinglaung village, Paunglang Reserve Forest, Pinlaung Township; 20°02'56.1"N, 96°46'00.1"E, ca.1448 m alt., 16 Sep. 2015, ibid., 103361 (MBK0313746, duplicates in RAF, QBG).

Notes. The genus *Thylacopteris* is sometimes confused with *Goniophlebium* and *Polypodium* (Rödl-Linder 1994; Fraser-Jenkins 2020). Warty cell walls of rhizome scales (Fig. 3) can be used to conclusively identify the genus *Thylacopteris* (Rödl-Linder 1994).

Key to species of the genus *Thylacopteris*

1	Blades 7–15 cm long.....	<i>T. minuta</i>
–	Blades 30–60 cm long.....	2
2	Sori deeply sunken; sclerenchyma present in cross-section of rhizome.....	
	<i>T. papillosa</i>
–	Sori not sunken sclerenchyma; absent in cross-section of rhizome.....	
	<i>T. diaphana</i>

Acknowledgements

This study was conducted under a Memorandum of Understanding between the Forest Department, Ministry of Natural Resources and Environmental Conservation, Myanmar and the Kochi Prefectural Makino Botanical Garden, Japan. We express our cordial thanks to Dr. Nyi Nyi Kyaw (Director General (retired) of the Forest Department), Dr. Thaung Naing Oo (Deputy Director General of the Forest Department), Dr. Naing Zaw Htun (Director (retired) of the Nature and Wildlife Conservation Division), Myanmar Forest Department and Ministry of Natural Resources and Environmental Conservation, for their help in coordinating the expeditions. We are grateful to the Myanmar-Japanese Cooperative Program for allowing us to study the specimens collected in Shan State. Field work was financially supported through the JICA grassroots program, the Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, and the Post-doctoral orientation training in Yunnan Province in 2018 (Y7YN021B14). This research received partial financial support through the Kochi Prefectural Government. We are grateful to Dr. Thais Almeida, Dr. Weston Testo for reviewing and Dr. Joel Nitta for management of this article.

References

- Blume CL (1828) *Enumeratio Plantarum Javae et Insularum adjacentium*, fasc. II. Filices. Leiden, 131 pp.

- Brause G (1912) Neue Flora Papuasiens, nebst allgemeinen Bemerkungen fiber das Vorkommen der Pteridophyten in Neu-Guinea von R. Schlechter. Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie 49: 1–42.
- Chen CC, Hyvönen J, Schneider H (2019) Exploring phylogeny of the microsoroid ferns (Polypodiaceae) based on six plastid DNA markers. Molecular Phylogenetics and Evolution 143: e106665. <https://doi.org/10.1016/j.ympev.2019.106665>
- Chen CC, Liu HY, Chen CW, Schneider H, Hyvönen J (2021) On the spore ornamentation of the microsoroid ferns (microsoroideae, Polypodiaceae). Journal of Plant Research 134(1): 55–76. <https://doi.org/10.1007/s10265-020-01238-4>
- Ching RC (1978) The Chinese fern families and genera: Systematic arrangement and historical origin. Acta Phytotaxonomica Sinica 16(3): 1–19.
- Christensen C, Holtum RE (1934) The ferns of Mount Kinabalu. Gardens' bulletin. Straits Settlements 7(3): 191–324.
- Copeland EB (1947) Genera Filicum: The Genera of Ferns. Waltham, Mass, 181–182.
- Darriba D, Taboada GL, Doallo R, Posada D (2012) jModelTest 2: More models, new heuristics and parallel computing. Nature Methods 9(8): e772. <https://doi.org/10.1038/nmeth.2109>
- Doyle JA, Doyle JL (1990) Isolation of plant DNA from fresh tissue. Focus [San Francisco, Calif.] 12(1): 13–15.
- Edgar RC (2004) MUSCLE: Multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Research 32(5): 1792–1797. <https://doi.org/10.1093/nar/gkh340>
- Fraser-Jenkins CR (2020) An Annotated Checklist of Indian Pteridophytes, Part 3: Lomariopsidaceae to Salviniaceae. Bishen Singh Mahendra Pal Singh, Uttarakhand, 450 pp.
- Hennipman E, Veldhoen P, Kramer KU, Price MG (1990) Polypodiaceae. In: Kubitzki K (Ed.) Families and genera of vascular plants. Vol 1. Pteridophytes and gymnosperms. Springer, Berlin, 203–230. https://doi.org/10.1007/978-3-662-02604-5_41
- Holtum RE (1966) Flora of Malaya 2, Ferns of Malaya, Vol. 2, Government Printing Office, Singapore, 203 pp.
- Hori K, Zhou XL, Shao W, Yan YH, Ebihara A, Wang RX, Ishikawa H, Watano Y, Murakami N (2018) Hybridization of the Dryopteris erythrosora complex (Drypteridaceae, Polypodiidae) in Japan. Hikobia 17(4): 299–313.
- Hori K, Ebihara A, Aung ZW (2019) New Records of Pteridophytes from Myanmar (1). Shokubutsu Kenkyu Zasshi 94(1): 35–38.
- Khine PK, Schneider H (2020) Data on pteridophyte species diversity and status of the International Union for Conservation of Nature in each political unit of Myanmar. Data in Brief 30: e105503. <https://doi.org/10.1016/j.dib.2020.105503>
- Khine PK, Lindsay S, Fraser-Jenkins CR, Kluge J, Kyaw M, Hovenkamp P (2016) Selliguea kachinensis (Polypodiaceae), a new fern species of uncertain affinity from northern Myanmar. PhytoKeys 62: 73–81. <https://doi.org/10.3897/phytokeys.62.8101>
- Khine PK, Fraser-Jenkins C, Lindsay S, Middleton D, Miehe G, Thomas P, Kluge J (2017) A Contribution Toward the Knowledge of Ferns and Lycophytes from Northern and Northwestern Myanmar. American Fern Journal 107(4): 219–256. <https://doi.org/10.1640/0002-8444-107.4.219>

- Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGAX: Molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution* 35(6): 1547–1549. <https://doi.org/10.1093/molbev/msy096>
- Nwe TY, Moon MO, Hwang IC, Lee SH, Sun B-Y (2016) New records of ferns in the flora of Myanmar found in Natma Taung National Park in the Chin State. *Korean Journal of Plant Taxonomy* 46(3): 283–287. <https://doi.org/10.11110/kjpt.2016.46.3.283>
- PPG I (2016) A community-derived classification for extant lycophytes and ferns. *Journal of Systematics and Evolution* 54(6): 563–603. <https://doi.org/10.1111/jse.12229>
- Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA (2018) Posterior summarisation in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* 67(5): 901–904. <https://doi.org/10.1093/sysbio/syy032>
- Rödl-Linder G (1994) A monograph of the fern genus *Thylacopteris* (Polypodiaceae). *Blumea* 39: 351–364.
- Ronquist F, Teslenko M, Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Schneider H, Janssen T, Hovenkamp P, Smith AR, Cranfill R, Haufler CH, Ranker TA (2004) Phylogenetic relationships of the enigmatic Malesian fern *Thylacopteris* (Polypodiaceae, Polypodiidae). *International Journal of Plant Sciences* 165(6): 1077–1087. <https://doi.org/10.1086/423882>
- Smith J (1875) *Historia Filicum*, Macmillan, London, 87 pp.
- Testo WL, Field AR, Sessa EB, Sundue M (2019) Phylogenetic and morphological analyses support the resurrection of *Dendroconche* and the recognition of two new genera in Polypodiaceae subfamily Microsoroideae. *Systematic Botany* 44(4): 1–16. <https://doi.org/10.1600/036364419X15650157948607>
- Thiers B (2021) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih/> [accessed 6 May 2022]
- Tryon AF, Lugardon B (1991) Spores of the Pteridophyta. Springer, New York, 302–361. https://doi.org/10.1007/978-1-4613-8991-0_21
- Tryon RM, Tryon AF (1982) Ferns and fern allies, with special reference to tropical America. Springer, New York, 684–758. <https://doi.org/10.1007/978-1-4613-8162-4>
- Zhang XC, Lu SG, Lin YX, Qi XP, Moore S, Xing FW, Wang FG, Hovenkamp PH, Gilbert MG, Nooteboom HP, Parris BS, Haufler C, Kato M, Smith AR (2013) Polypodiaceae. In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China*, vol. 2–3: (Pteridophytes). Science Press, Beijing & Missouri Botanical Garden, St. Louis, 758–850.
- Zhao XF, Xiang JY, Geekiyanage N, Sun HY, Myo MK, Dian EN, Tang FL (2017) *Leptochilus mengsongensis* (Polypodiaceae), a new species of fern from SW China. *Phytotaxa* 317(2): e144. <https://doi.org/10.11646/phytotaxa.317.2.7>
- Zhao CF, Wei R, Zhang XC, Xiang QP (2020) Backbone phylogeny of *Lepisorus* (Polypodiaceae) and a novel infrageneric classification based on the total evidence from plastid and morphological data. *Cladistics* 36(3): 235–258. <https://doi.org/10.1111/cla.12403>