

***Gastrochilus heminii* (Orchidaceae, Epidendroideae), a new species from Sichuan, China, based on molecular and morphological data**

Min Liao^{1,2*}, Yue-Hong Cheng^{3*}, Jun-Yi Zhang^{1,4}, Yu Feng¹,
Gui-Ying Liu³, Ping Ye³, Sen-Long Jin³, Hong-Qiang Lin³, Bo Xu^{1,2}

1 China-Croatia “Belt and Road” Joint Laboratory on Biodiversity and Ecosystem Services, CAS Key Laboratory of Mountain Ecological Restoration and Bioresource Utilization & Ecological Restoration and Biodiversity Conservation Key Laboratory of Sichuan Province, Chengdu Institute of Biology, Chinese Academy of Sciences, Chengdu 610041, China **2** University of Chinese Academy of Sciences, Beijing 10049, China **3** Wolong National Natural Reserve Administration Bureau, Wenchuan 623006, Sichuan, China **4** College of Life Sciences, Chongqing Normal University, Chongqing 401331, China

Corresponding author: Bo Xu (xubo@cib.ac.cn)

Academic editor: Timothée LePéchon | Received 30 July 2022 | Accepted 1 December 2022 | Published 16 December 2022

Citation: Liao M, Cheng Y-H, Zhang J-Y, Feng Y, Liu G-Y, Ye P, Jin S-L, Lin H-Q, Xu B (2022) *Gastrochilus heminii* (Orchidaceae, Epidendroideae), a new species from Sichuan, China, based on molecular and morphological data. *PhytoKeys* 215: 95–106. <https://doi.org/10.3897/phytokeys.215.91061>

Abstract

Gastrochilus heminii (Orchidaceae), a new orchid species from Sichuan Province, Southwest China, is described and illustrated. It morphologically resembles *G. affinis* and *G. yei*, but differs markedly from the former in having a thinner and slightly rolled downwards reniform epichile and the central thickened purple-red mat with irregular folds (vs. subtriangular epichile curves upwards, with 2 thick, brown to purplish-brown median ridges from base to apex), and can be clearly distinguished from the latter by having reniform epichile with lobed apex and subconical hypochile with bilobed apex that splits into two conical protrusions (vs. semi-rounded epichile not lobed and subconical hypochile not bilobed). The results of molecular phylogenetic analysis based on nuclear ribosome internal transcribed spacer (nrITS) and four chloroplast DNA fragments (*matK*, *psbA-trnH*, *psbM-trnD*, and *trnL-F*) of 36 *Gastrochilus* species showed that *G. heminii* was closely related to *G. affinis* and *G. distichus*.

Keywords

Gastrochilus, morphological, Orchidaceae, phylogeny, taxonomy

* These authors contributed equally to this work as co-first authors.

Introduction

Gastrochilus D.Don (Orchidaceae, Epidendroideae; Don 1825) is diagnosed by a subdivided labellum with a recurved epichile and a saccate hypochile attached to the column at the base and two subglobose pollinia attached to a slender and filiform stipe (Tsi 1999; Pridgeon et al. 2014). The genus comprises more than 70 accepted species, most of which are distributed in Southeast Asia (Kumar et al. 2014; Liu et al. 2016, 2019; Liu and Gao 2018; Rao et al. 2019; Govaerts et al. 2021; Nguyen et al. 2021; Zhou et al. 2021a, b; Zhang et al. 2022). Liu et al. (2019) first constructed the phylogenetic relationships of *Gastrochilus* and divided them into five clades. Recently, the phylogenetic studies by Liu et al. (2020), Li et al. (2022), and Zhang et al. (2022) also supported the monophyly of *Gastrochilus*.

An unusual arboreal miniature orchid was recently discovered for the first time in Wolong Nature Reserve (Wenchuan County, Sichuan Province, China) during a routine survey. It was tentatively identified as a species of *Gastrochilus* sect. *Microphyllae* Bentham & Hooker (Bentham and Hooker 1883), characterized by plants with extremely small flowers, distichous and alternate leaves compared to plants in the other sections (larger flowers or clustered leaves). However, a morphological examination revealed that this species shows an unusual combination of characters that does not match any known species of *G.* sect. *Microphyllae*. The phylogenetic analysis using five DNA markers (nrITS, *matK*, *psbA-trnH*, *psbM-trnD*, and *trnL-F*) confirmed the monophyly of this taxon. The objectives of this study are (1) to describe, (2) to examine both molecular and morphological affinities of this new *Gastrochilus* species, *Gastrochilus heminii* M.Liao, B.Xu & Yue.H.Cheng, sp. nov.

Materials and methods

Morphological analyses

The measurements and description of *Gastrochilus heminii* were based on two living plant individuals and two herbarium specimens (voucher information: *Min Liao & Yue-Hong Cheng ZJY143*; *Min Liao & Yue-Hong Cheng ZJY167*), respectively. The taxonomic description follows the terminology used by Beentje (2012). Voucher specimens and additional silica-gel dried leaves are deposited in CDBI Herbarium (herbarium follows Thiers 2021).

DNA extraction, amplification and sequencing

The sequences of the two individuals of the new species newly generated in this study, and the sequences of the remaining 42 species used in the molecular phylogenetic analysis, were retrieved from GenBank. The information on the DNA fragments and four complete plastid genomes were listed in Appendix 1. Total DNA was

extracted exclusively from silica-gel dried leaves via a Plant DNA Isolation Kit (Cat. No.DE-06111). We used the same primers as Liu et al. (2019) to amplify the nuclear ribosome internal transcribed spacer (nrITS) and the four chloroplast DNA fragments (i.e., *matK*, *psbA-trnH*, *psbM-trnD*, and *trnL-F*) through polymerase chain reaction (PCR). All DNA samples were sent to TSINGKE Biotech Co. Ltd (Chengdu, China) for sequencing. The sequences were then deposited with GenBank, with the following accession numbers: *G. heminii*, nrITS (ON286752, ON286753), *matK* (ON331126, ON331127), *psbA-trnH* (ON331128, ON331129), *psbM-trnD* (ON331130, ON331131), and *trnL-F* (ON331132, ON331133).

Phylogenetic analyses

All sequences were edited via Sequencher v4.1.4 (Gene Codes, Ann Arbor, Michigan, USA) and aligned via MAFFT v7.475 (Katoh and Standley 2013) with default parameters. We performed phylogenetic analyses based on combined nuclear ribosome internal transcribed spacer (nrITS) and the four chloroplast DNA fragments. The nucleotide substitution model for the data matrix was estimated using jModeltest v2.1.6 (Posada 2008) software and the best-fit model (GTR+I+G) was selected using the corrected Akaike Information Criterion (AICc). Two different methods, Maximum likelihood (ML) and Bayesian inference (BI), were employed. The ML analysis was performed using IQ-TREE v1.4.2 (Nguyen et al. 2014) with branch support estimated by 2,000 replicates of ultrafast bootstrapping algorithm (UFboot) (Minh et al. 2013). The BI analysis was conducted using MrBayes v3.2.7a (Ronquist and Huelsenbeck 2003), with two separate Markov-chain Monte Carlo (MCMC) chains (1,000,000 generations and sampled every 1,000 generations). The first 25% of the trees were discarded as burn-in, and the remaining trees were used to generate a majority-rule consensus tree.

Results

The molecular phylogenetic tree showed that the 36 taxa of *Gastrochilus* formed a well-supported monophyletic group (BI/ML = 1/97, Fig. 1). The two individuals of *G. heminii* were resolved as sisters to each other (BI/ML = 1/99, Fig. 1). Our data recovered a sister relationship between *G. affinis* (King & Pantl.) Schltr. (King and Pantling 1898; Schlechter 1913) and *G. distichus* (Lindl.) Kuntze (Lindley 1859; Kuntze 1891); these two species formed a monophyletic group with *G. heminii* (BI/ML = 0.90/89, Fig. 1), which formed a subclade of section *Microphyllae* together with *G. fargesii* (Kraenzl.) Schltr. (Kraenzlin 1903; Schlechter 1919), *G. alatus* X.H.Jin & S.C.Chen (Jin et al. 2007), *G. fuscopunctatus* (Hayata) Hayata (Hayata 1912, 1917), and *G. pseudodistichus* (King & Pantl.) Schltr. (King and Pantling 1895; Schlechter 1913).

Morphologically, *G. heminii* is closest to *G. affinis* and *G. yei* JianW.Li & X.H.Jin (Li et al. 2021). A thorough morphological comparison among *G. heminii*, *G. affinis*, *G. yei* and *G. distichus* is summarized in Table 1 and Fig. 2.




Figure 1. Maximum likelihood tree of *Gastrochilus* from phylogenetic analysis of combined nrITS and plastid DNA markers. Numbers before and after the slash indicate Bayesian posterior probabilities and ML bootstrap supports, respectively. The position of *Gastrochilus heminii* is indicated in red.

Taxonomic treatment

***Gastrochilus heminii* M.Liao, B.Xu & Yue H.Cheng, sp. nov.**

[urn:lsid:ipni.org:names:77310051-1](https://nri.org/names/77310051-1)

Figs 2A, B, 3

Type. CHINA. Sichuan: Wenchuan, coniferous and broadleaf mixed forest, on tree trunk, elev. ca. 2640 m, in flowering and fruiting, 15 March 2022, *Min Liao & Yue-Hong Cheng ZJY143* (holotype CDBI!).

Diagnosis. *Gastrochilus heminii* is morphologically related to *G. affinis* and *G. yei* based on vegetative and floral characteristics such as similar habit, distichous and alternate leaves, epichile surface smooth and glabrous, sepals and petals with purplish-red




Figure 2. Comparison of three taxa of *Gastrochilus* **A, B** *Gastrochilus heminii* **C, D** *G. affinis* **E, F** *G. distichus* **G, H** *G. yei*. [Images **C, D** cited from Jalal et al. 2020; image **E** reproduced from website (<http://www.orchidspecies.com/gastdistichus.htm>); image **F** cited from Kumar et al. 2014, which was photographed by Xiao-Hua Jin; images **G** and **H** cited from Li et al. 2021].




Figure 3. *Gastrochilus heminii* **A** flowering plant **B** young leaves (B1: front view; B2: reverse view) **C** old leaves (C1: front view; C2: reverse view) **D** raceme and capsule (side view) **E–I** flowers (different views) **J** anatomy of flower (J1: dorsal sepals; J2 and J3: lateral sepals; J4 and J5: petals; J6: labellum) **K** hypochile (dorsal view) **L** stamens.

patches. However, it can be differentiated from *G. affinis* on the basis of flower numbers (1–2 in the former vs. 1–4 in the latter), peduncle length (0.4–0.7 cm in the former vs. 1.5–2.0 cm in the latter) and an additional morphological characteristic: young leaves are densely covered with purple-red spots and old leaves have hardly any purple-red spots in the former (both have purple-red spots in the latter); the reniform epichile is rolled downwards, smooth and glabrous above, and central thickened purple-red mat with irregular folds in the former (subtriangular epichile curves upwards, with 2 thick, brown to purplish-brown median ridges from base to apex in the latter). It differs from *G. yei* by having reniform and lobed epichile (not lobed in the latter), apex of hypochile bilobed and splits into two conical protrusions (not bilobed in the latter), apex of the leaf with 1–2 lobules, lobes setaceous (apex of leaf with a tine in the latter).

Description. Miniature trunk epiphyte. Roots vermiform, 2.0–7.0 cm long and ca. 1.5 mm thick. Stem pendulous, slender, purplish-red spots, 3.0–6.5 cm long and ca. 1.8 mm thick. Leaves alternate, distichous, narrowly oblong or falcate oblong, 0.9–2.3 × 0.3–0.5 cm, apex acute and with 1–2 lobules, lobes setaceous, young leaves with conspicuous purplish-red spots, old leaves with hardly any purplish-red spots. Raceme with 1 or 2 flowers; inflorescence stalk curved upward and thickened, 4.0–9.0 mm long, proximally covered with two sheaths; floral bracts ovate-lanceolate, 0.7–1.0 mm long, apex acute; pedicel and ovary connate, 4.0–5.5 mm long; flowers spreading, ca. 6.0 × 5.0 mm; sepals and petals heterochromatic on both surfaces, outside yellow-green with purplish-red spots, inside purplish-red with yellow-green margin; dorsal sepals and lateral sepals similar and equal in size, elliptic-oblong, ca. 2.4 × 1.5 mm, apex obtuse; petals narrowly oblong, ca. 2.6 × 1.3 mm, apex acute, base narrowed; epichile reniform, yellow-green with purplish-red spots, 4.2–6.5 × 2.0–3.0 mm, margin erose, smooth and glabrous above, central thickened purple-red mat with irregular folds; hypochile subconical, yellow-green with purple-red spots, 2–2.4 × 1.6–2 mm, dorsally compressed, slightly bent outward, the apex splits into two conical protrusions; column cylindrical, ca. 1.0 mm; anther cap subhemispheric, with two chambers, 0.7 × 0.4 mm, hanging from both ends of the stipe; pollinia 2, 0.4 × 0.3 mm, yellow, full and nearly spherical, with a depression in the center; stigma deeply sunken, inverted V-shaped, ca. 0.6 mm long, yellow, apically forked, forked in a subtriangular outline. Capsule shuttle-shaped with six ribs, green with sparse purplish-red spots, ca. 1.1 cm long, inflated to ca. 0.6 cm in the middle, persistent and growing for one year until maturity.

Additional specimens examined. —CHINA. Sichuan: Wenchuan, coniferous and broadleaf mixed forest, on tree trunk, elev. ca. 2640 m, 18 April 2022, *Min Liao & Yue-Hong Cheng* ZJY167 (CDBI).

Distribution, habitat and phenology. The new species is currently known only from Wenchuan County, Sichuan Province, Southwest China (Fig. 4). It is found epiphytic on the trunk of *Tsuga chinensis* (Franch.) Pritz. in a subalpine mixed coniferous forest at elevation ca. 2640 m. *Gastrochilus heminii* flowers from March to April.




Figure 4. Distribution map of *Gastrochilus heminii*.

Table 1. Morphological comparison of *Gastrochilus heminii* with three related species, *G. affinis*, *G. distichus* and *G. yei*.

Character	<i>G. heminii</i>	<i>G. affinis</i>	<i>G. distichus</i>	<i>G. yei</i>
Plant length	3.0–6.5 cm	4.0–15.0 cm	1.5–20.0 cm	3.0–8.0 cm
Leaf shape	narrowly oblong or falcate oblong	oblong-lanceolate to subspathulate	lanceolate or falcate lanceolate	blade lanceolate
No. of flowers per inflorescence	1 or 2 flowers	1–4 flowers	2–4 flowers	2–6 flowers
Peduncle length	0.4–0.7 cm	1.5–2.0 cm	1.0–1.5 cm	0.7–1.0 cm
Dorsal sepal	elliptic-oblong, ca. 2.4 × 1.5 mm, concave, apex obtuse	elliptic-oblong, 3–5 × 1–1.3 mm, concave, apex obtuse	oblong-elliptic, 4.5–0.5 × 2.5–3 mm, concave, apex obtuse	oblong, 3.3 × 1.9–2.0 mm, apex rounded
Lateral sepals	similar to dorsal sepal, equal in size	elliptic-ovate, 3.5–4 × 0.7–1.3 mm, slightly oblique and incurved, apex obtuse	similar to dorsal sepal, equal in size	oblong, 3.9–4.0 × 1.8–1.9 mm, apex obtuse
Petals	narrowly oblong, ca. 2.6 × 1.3 mm, apex acute, base narrowed	ovate-elliptic to elliptic, 3–4 × 1–1.3 mm, apex obtuse	subobovate, slightly smaller than sepals, apex obtuse	oblong, 3.5 × 1.8 mm, apex rounded
Epichile	reniform, 4.2–6.5 × 2.0–3.0 mm, margin erose, smooth and glabrous above, central thickened purple-red mat with irregular folds	subtriangular, decurved, subacute at apex, margin finely erose at base, disk with 2 thick, brown to purplish-brown median ridges from base to apex.	subcircular, 5 × 3 mm, apex obtuse, margin entire, smooth and glabrous above and thickened cushion-like centrally, with 2 conical callosities near base	semi-rounded, 2.0–2.2 × 4.0–4.2 mm, glabrous, with a thickened central, rugose cushion, tint with purple, margin irregularly denticulate
Hypochile	subconical or helmet-shaped, ca. 2–2.4 × 1.6–2 mm, dorsally compressed, slightly bent outward, the end splits into two conical protrusions	obconical, 3–4 × 2–3 mm, dorsally compressed, slightly bent outward, subacute to obtuse and shortly bifid at apex	subcupular, 4 × 2–3 mm, rounded at end, dorsally compressed, slightly bent outward	subconical, 3 mm tall, 3 mm in diameter, apex rounded

Etymology. Named in honor of Mr. He-Min Zhang, the advocate of the panda “Release to the Forest”, one of the pioneers of panda research in China, renowned as the “father” of pandas, in recognition of his contribution to the conservation of flora and fauna in the region which was one of the earliest reserves dedicated to preserving the habitat of wild pandas. A Chinese name, *he min peng ju lan* (和民盆距兰), is suggested here.

Conservation status. Three populations with a total of ca. 200 individuals of *Gastrochilus heminii* have been documented during our investigation. However, similar habitat of this new species is widespread in the Wolong Natural Reserve. Therefore, we assessed the conservation status of *Gastrochilus heminii* as DD (Data Deficient) according to the IUCN (2022).

Acknowledgements

Our work was funded by the National Key Research and Development Program of China (Grant No. 2020YFE0203200), the Second Tibetan Plateau Scientific Expedition and Research (STEP) program (Grant No. 2019QZKK0502), 2022 Central Finance Forestry Grassland Ecological Protection and Restoration National Park Subsidy Project, 2022 Subsidy Projects of Prohibited Developmental Areas from the Transfer Payment of the National Key Ecological Functional Areas, and Wild Plants Sharing and Service Platform of Sichuan Province.

References

- Beentje H (2012) The Kew Plant Glossary, an illustrated dictionary of plant terms. Kew Publishing, Richmond, 164 pp.
- Bentham G, Hooker JD (1883) *Gastrochilus* section *Microphyllae*. In: Genera plantarum 3. Reeve, London, 579 pp.
- Don D (1825) Prodrromus Florae Nepalensis. J. Gale, London, 256 pp.
- Govaerts R, Campacci MA, Baptista DH, Baptista PJ, George A, Kreutz K, Wood JJ (2021) World Checklist of Orchidaceae. The Board of Trustees of the Royal Botanic Gardens, Kew. <http://www.kew.org/wcps/monocots/>
- Hayata B (1912) Icones Plantarum Formosanarum, nec non et contributions ad floram Formosanam, vol. 2 (suppl.). Government of Formosa, Taihoku [Taipei], 143 pp.
- Hayata B (1917) Icones Plantarum Formosanarum, nec non et contributions ad floram Formosanam, vol. 6 (suppl.). Government of Formosa, Taihoku [Taipei], 78 pp.
- IUCN (2022) Guidelines for Using the IUCN Red List Categories and Criteria, prepared by the Standards and Petitions Committee, version 15. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Jalal JS, Rai ID, Bhattacharjee A, Singh G (2020) New distributional record of *Gastrochilus affinis* (Orchidaceae) from Western Himalaya with notes on a new synonym and typification. Nelumbo 62(1): 12–15. <https://doi.org/10.20324/nelumbo/v62/2020/153741>

- Jin XH, Li H, Li DZ (2007) Additional notes on Orchidaceae from Yunnan, China. *Acta Phytotaxonomica Sinica* 45(6): 796–807. <https://doi.org/10.1360/aps06216>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- King G, Pantling R (1895) New Orchids from Sikkim. *Journal of the Asiatic Society of Bengal* 64(2): 341.
- King G, Pantling R (1898) The Orchid of the Sikkim-Himalaya. *Annals of the Royal Botanic Garden, Calcutta*, vol. VIII. Bengal Secretariat Press, London, 1–133.
- Kraenzlin FWL (1903) Deux Orchidees Nouvelles. *Le Journal de Botanique* 17(12): 423.
- Kumar P, Gale SW, Kocyan A, Fischer GA, Averyanov LV, Borosova R, Bhattacharjee A, Li JH, Pang KS (2014) *Gastrochilus kadooriei* (Orchidaceae), a new species from Hong Kong, with notes on allied taxa in section Microphyllae found in the region. *Phytotaxa* 164(2): 91–103. <https://doi.org/10.11646/phytotaxa.164.2.3>
- Kuntze O (1891) Dicotyledones II. *Revisio generum plantarum: vascularium omnium atque cellularium multarum secundum leges nomenclaturae internationales cum enumeratione plantarum exoticarum in itinere mundi collectarum, Pars II*. Arthur Felix, Leipzig, 661 pp. <https://doi.org/10.5962/bhl.title.124064>
- Li JW, Ya JD, Ye DP, Liu C, Liu Q, Pan R, He ZX, Pan B, Cai J, Lin DL, Jin XH (2021) Taxonomy notes on Vandaeae (Orchidaceae) from China: Five new species and two new records. *Plant Diversity* 43(05): 379–389. <https://doi.org/10.1016/j.pld.2021.01.009>
- Li Y, Jin WT, Zhang LG, Zhou P, Luo Y, Zhu ZW, Xiang XG (2022) Biogeography and diversification of the Tropical and Subtropical Asian genus *Gastrochilus* (Orchidaceae, Aeridinae). *Diversity* 14(5): 396. <https://doi.org/10.3390/d14050396>
- Lindley J (1859) The Orchidology of India. *Journal of the Proceedings of the Linnean Society, Botany*, vol. III. Williams and Norgate, London, 36 pp.
- Liu Q, Gao JY (2018) *Gastrochilus dulongjiangensis* (Aeridinae, Vandaeae, Epidendroideae, Orchidaceae), a new species from Yunnan Province, China. *Phytotaxa* 340(3): 293–296. <https://doi.org/10.11646/phytotaxa.340.3.11>
- Liu Q, Tan YH, Gao JY (2016) A new species of *Gastrochilus* (Aeridinae, Vandaeae, Orchidaceae) and a new record species from Yunnan, China. *Phytotaxa* 282: 066–070. <https://doi.org/10.11646/phytotaxa.282.1.8>
- Liu Q, Song Y, Jin XH, Gao JY (2019) Phylogenetic relationships of *Gastrochilus* (Orchidaceae) based on nuclear and plastid DNA data. *Botanical Journal of the Linnean Society* 189(3): 228–243. <https://doi.org/10.1093/botlinnean/boy084>
- Liu D, Tu XD, Zhao Z, Zeng MY, Zhang S, Ma L, Zhang GQ, Wang MM, Liu ZJ, Lan SR, Chen SP, Li MH (2020) Plastid phylogenomic data yield new and robust insights into the phylogeny of *Cleisostoma* – *Gastrochilus* clades (Orchidaceae, Aeridinae). *Molecular Phylogenetics and Evolution* 145: 106729. <https://doi.org/10.1016/j.ympev.2019.106729>
- Minh BQ, Nguyen MAT, von-Haeseler A (2013) Ultrafast approximation for phylogenetic bootstrap. *Molecular Biology and Evolution* 30(5): 1188–1195. <https://doi.org/10.1093/molbev/mst024>

- Nguyen LT, Schmidt HA, Von-Haeseler A, Minh BQ (2014) IQ-TREE: A fast and effective stochastic algorithm for estimating Maximum-Likelihood phylogenies. *Molecular Biology and Evolution* 32(1): 268–274. <https://doi.org/10.1093/molbev/msu300>
- Nguyen DH, Averyanov LV, Dang VS, Bui VH, Maisak TV, Truong BV (2021) *Gastrochilus dresslerii* (Orchidaceae), a new species from northern Vietnam. *Phytotaxa* 498(2): 145–149. <https://doi.org/10.11646/phytotaxa.498.2.8>
- Posada D (2008) jModelTest: Phylogenetic model averaging. *Molecular Biology and Evolution* 25(7): 1253–1256. <https://doi.org/10.1093/molbev/msn083>
- Pridgeon AM, Cribb PJ, Chase MW, Rasmussen FN (2014) *Genera orchidacearum: Epidendroideae*, volume 6, Part 3. Oxford University Press, Oxford, 544 pp.
- Pritzel GA (1901) Pinaceae. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 29(2): 217.
- Rao WH, Liu ZJ, Zhang GQ, Chen XH, Huang J, Chen GZ, Chen LJ (2019) A new epiphytic species of *Gastrochilus* (Orchidaceae: Epidendroideae) from Yunnan, China. *Phytotaxa* 340: 293–296. <https://doi.org/10.11646/phytotaxa.413.4.5>
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19(12): 1572–1574. <https://doi.org/10.1093/bioinformatics/btg180>
- Schlechter FRR (1913) Die Gattungen *Gastrochilus* Don. Und *Gastrochilus* Wall. *Repertorium specierum novarum regni vegetabilis*. Selbstverlag des Herausgebers, Berlin, 313–315. <https://doi.org/10.1002/fedr.19130121713>
- Schlechter FRR (1919) *Orchideologiae Sino-Japonicae prodromus*. Eine kritische Besprechung der Orchideen Ost-Asien. *Repertorium Specierum Novarum Regni Vegetabilis*, Beihefte 4: 1–319.
- Thiers B (2021) Index Herbariorum: a global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih>
- Tsi ZH (1999) *Gastrochilus* D. Don. In: Tsi ZH (Ed.) *Flora Reipublicae Popularis Sinicae*, vol. 19. Science Press, Beijing, 399–420.
- Zhang JY, Cheng YH, Liao M, Jin SL, Qu CM, Tan YC, Plenković-Moraj A, Xu B (2022) *Gastrochilus wolongensis* (Orchidaceae): A new species from Sichuan, China, based on molecular and morphological data. *Ecosystem Health and Sustainability* 8(1): 2101546. <https://doi.org/10.1080/20964129.2022.2101546>
- Zhou ZH, Shi RH, Zhang Y, Jin XH (2021a) Orchid diversity in China: Recent discoveries. *Plant Diversity* 43(5): 341–342. <https://doi.org/10.1016/j.pld.2021.07.004>
- Zhou ZH, Shi RH, Zhang Y, Xing XK, Jin XH (2021b) Orchid conservation in China from 2000 to 2020: Achievements and perspectives. *Plant Diversity* 43(5): 343–349. <https://doi.org/10.1016/j.pld.2021.06.003>

Appendix I

Table A1. The GenBank accession numbers for DNA sequences used in this study.

Taxa	nrITS	matK	psbA-trnH	trnL-F	psbM-trnD
<i>G. acaulis</i> (Lindl.) Kuntze	KM583455	KM583465	—	—	—
<i>G. acinacifolius</i> Z.H.Tsi	KJ733412	KJ733569	KJ733492	KJ733649	MK357216
<i>G. acutifolius</i> (Lindl.) Kuntze	MT225573	MW433889	—	—	MK357230
<i>G. affinis</i> (King & Pantl.) Schltr.	—	MK357141	—	—	MK357141
<i>G. alatus</i> X.H.Jin & S.C.Chen	—	—	—	—	MK357228
<i>G. bellinus</i> (Rchb.f.) Kuntze	KY966597	KY966884	MK357164	MK357202	MK357240
<i>G. bigibbus</i> (Rchb.f. ex Hook.f.) Kuntze	—	MN124439	MN124439	MN124439	MN124439
<i>G. calceolaris</i> (Buch.-Ham. ex Smith) D.Don	MN517123	MK357144	MK357169	MK357204	MK357232
<i>G. changjiangensis</i> Q.Liu & M.Z.Huang	MK357124	—	MK357166	—	MK357236
<i>G. ciliaris</i> F.Mackawa	—	MK357148	MK357173	—	MK357225
<i>G. dasypogon</i> (Lindl.) Kuntze	DQ091685	MK357149	MK357181	MK357197	MK357219
<i>G. deminutus</i> J.M.H.Shaw	KY966600	KY966887	—	—	—
<i>G. distichus</i> (Lindl.) Kuntze	KJ733414	KJ733571	KJ733494	KJ733651	—
<i>G. fargesii</i> (Kraenzl.) Schltr.	KJ733415	KJ733572	—	KJ733652	—
<i>G. formosanus</i> (Hayata) Hayata	KJ733416	KJ733573	KJ733495	KJ733653	MK357226
<i>G. fuscopunctatus</i> (Hayata) Hayata	MK317970	MK357150	MK357171	MK357192	MK357231
<i>G. gongshanensis</i> Z.H.Tsi	—	MN124438	MN124438	MN124438	MN124438
<i>G. guangtungensis</i> Z.H.Tsi	KJ733417	KJ733574	KJ733496	KJ733654	KJ733654
<i>G. heminii</i> M.Liao, B.Xu & Yue.H.Cheng	ON286752	ON331126	ON331128	ON331130	ON331132
<i>G. heminii</i> M.Liao, B.Xu & Yue.H.Cheng	ON286753	ON331127	ON331129	ON331131	ON331133
<i>G. intermedius</i> (Griff. ex Lindl.) Kuntze	MK357121	MK357151	MK357172	MK357190	MK357213
<i>G. japonicus</i> (Makino) Schltr.	KJ733418	KF545886	KF545866	KF545897	KX871236
<i>G. linearifolius</i> Z.H.Tsi & Garay	MK357133	MK357136	MK357187	MK357194	MK357229
<i>G. linii</i> Ormerod	—	MK357152	MK357176	MK357198	MK357224
<i>G. malipoensis</i> X.H.Jin & S.C.Chen	—	MK357147	MK357177	MK357200	MK357235
<i>G. minutiflorus</i> Aver.	—	MK357153	MK357179	—	MK357215
<i>G. obliquus</i> (Lindl.) Kuntze	MK357131	MK357137	KJ733498	KJ733656	MK357211
<i>G. platycalcaratus</i> (Rolf.) Schltr.	MK357122	—	MK357175	—	MK357222
<i>G. pseudodistichus</i> (King & Pantl.) Schltr.	MK357132	—	MK357170	—	MK357221
<i>G. rantabunensis</i> C.Chow ex T.P.Lin	—	MK357155	MK357184	MK357193	MK357223
<i>G. nanaensis</i> Fukuyama	KJ733420	KJ733577	KJ733499	KJ733657	MK357239
<i>G. sinensis</i> Z.H.Tsi	OM985813	OK042953	OK172399	OK172401	—
<i>G. somai</i> (Hayata) Hayata	MK357128	MN124436	MK357180	MN124436	MK357220
<i>G. tianbaoensis</i> Q.Liu & Y.H.Tan	MK357120	MK357157	MK357186	MK357207	MK357214
<i>G. wolongensis</i> Jun.Y.Zhang, B.Xu & Yue.H.Cheng	OM985810	OK172400	OK172402	OK172404	OK172403
<i>G. wolongensis</i> Jun.Y.Zhang, B.Xu & Yue.H.Cheng	OM985811	OM974209	OM974211	OM974210	—
<i>G. yunnanensis</i> Schltr.	MK165469	MK357158	MK357185	—	MK357212
<i>G. zhenyuanensis</i> Q.Liu & D.P.Ye	MK357127	MK357146	MK357168	MK357199	MK357237
<i>Holcoglossum amesianum</i> (Rchb.f.) Christenson	HQ404389	JF763779	HQ404439	—	—
<i>Holcoglossum kimballianum</i> (Rchb.f.) Garay	HQ452901	JF763787	HQ404452	—	—
<i>Luisia magniflora</i> Z.H.Tsi & S.C.Chen	KJ733426	KJ733583	KJ733505	—	KJ733663
<i>Pomatocalpa diffusum</i> Breda	AB217576	AB217752	—	—	EF670432
<i>Pomatocalpa spicatum</i> Breda	DQ091706	KJ733595	KJ733518	—	KJ733675
<i>Saccolabium pusillum</i> Blume	AB217580	AB217756	—	—	—