

Primulina jiulianshanensis, a new species of Gesneriaceae from Jiangxi Province, China

Guo-Liang Xu^{1*}, Li-Fen Liang^{2*}, Di-Ya Chen^{3,4}, Zhi-Fang Jing¹,
Xiao-Hai Zuo¹, Zheng-Yu Zuo⁵, Fang Wen^{3,6}

1 Jiulianshan National Nature Reserve Administrative Bureau, Longnan, CN-341700, China **2** Jiangxi Environmental Engineering Vocational College, Ganzhou, CN-341000, China **3** Guangxi Key Laboratory of Plant Conservation and Restoration Ecology in Karst Terrain, Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences, CN-541006, Guilin, China **4** College of Tourism and Landscape Architecture, Guilin University of Technology, Guilin, CN-541006, China **5** Germplasm Bank of Wild Species, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, CN-650201, China **6** National Gesneriaceae Germplasm Resources Bank of GXIB, Gesneriad Committee of China Wild Plant Conservation Association, Gesneriad Conservation Center of China (GCCC), Guangxi Institute of Botany, Guilin Botanical Garden, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences, CN-541006, Guilin, China

Corresponding authors: Fang Wen (wenfang760608@139.com); Zheng-Yu Zuo (zuozhengyu@mail.kib.ac.cn)

Academic editor: Alan Paton | Received 15 October 2022 | Accepted 2 April 2023 | Published 9 May 2023

Citation: Xu G-L, Liang L-F, Chen D-Y, Jing Z-F, Zuo X-H, Zuo Z-Y, Wen F (2023) *Primulina jiulianshanensis*, a new species of Gesneriaceae from Jiangxi Province, China. *PhytoKeys* 226: 1–16. <https://doi.org/10.3897/phytokeys.226.96351>

Abstract

Primulina jiulianshanensis F.Wen & G.L.Xu, a new species of Gesneriaceae from Jiulianshan National Nature Reserve of Jiangxi Province, China, is described and illustrated here. Molecular evidence showed it was sister to *P. wenii* Jian Li & L.J.Yan, while the morphological observation found clear differences between them, petiole, both sides of leaf blades, adaxial surface of the calyx lobes, corolla inside toward the bottom, bract margins covered glandular-pubescent hairs in *P. jiulianshanensis* (*vs.* no glandular-pubescent hairs in *P. wenii*); lateral bracts 4–9 × ca. 2 mm, the central one 2–5 × 1–1.5 mm, adaxially glabrous but sparsely pubescent at apex (*vs.* lateral bracts 14–16 × 2.5–3.0 mm, the central one 10–12 × 1.3–1.6 mm, all adaxially pubescent); calyx lobes 8–11 × ca. 2 mm, each side with several brown serrate teeth at apex (*vs.* 14–15 × ca. 2.5 mm, margin entire); filaments and staminodes sparsely yellow glandular-puberulent (*vs.* white, glabrous).

Keywords

Flora of Jiangxi, Jiulianshan National Nature Reserve, new taxon, *Primulina wenii*, taxonomy

* The authors contributed equally to this work.

Introduction

The genus *Primulina* Hance (Hance 1883) in Gesneriaceae is mainly distributed in the mountainous areas of southern and southwestern China to northern Vietnam, especially in Karst landforms (Wei 2018; Xu et al. 2020b). Since it was redefined (Wang et al. 2011; Weber et al. 2011), many new species have been discovered and published. For example, ten new taxa of *Primulina* from China were reported in 2020 (Du et al. 2021). As of December 2022, about 240 species (including infraspecies, the same below) have been confirmed throughout the world, of which 224 species are distributed in China. So far, this genus is the largest genus of Gesneriaceae in China (POWO 2022).

Jiangxi Province is located in the mid-subtropical region of East China; the species number of *Primulina* is not very rich. After consulting references, checking herbarium specimens, and excluding the species that have been mistakenly identified, only 13 species of *Primulina* have been confirmed in Jiangxi Province (Peng et al. 2021). Three of them were discovered and published after 2011, which are *P. lepingensis* Z.L.Ning & M.Kang (Ning et al. 2014), *P. suichuanensis* X.L.Yu & J.J.Zhou (Zhou et al. 2016) and *P. inflata* Li H.Yang & M.Z.Xu (Xu et al. 2020a).

In April 2021, an interesting population of *Primulina* was found on a cliff of Danxia landform under the evergreen broad-leaved forest in Jiulianshan National Nature Reserve, Longnan City, Jiangxi Province. Morphologically, this species is similar to *P. wenii* Jian Li & L.J.Yan (Li et al. 2017) in some characteristics. For example, leaf blades are oblong or broadly rounded, corolla purple, and so on.

We collected the plants at the flowering and fruiting stage in the type locality to make specimens, and at the same time carried out botanical fine anatomical photography to observe carefully. We saw that the indumentum characters of petiole, leaf, bract, calyx, corolla tube, filaments, staminodes of this unknown species were obviously different from *P. wenii*, and the differences of two species' calyx lobes and bracts characters can help us easily distinguish them. In order to understand the phylogenetic placements of this unknown species in *Primulina*, ITS and *trnL-F* sequences of this species were amplified and included for phylogenetic analysis to examine the relationships of the putative new species.

Materials and methods

Morphological observation

All available specimens of *Primulina* were used and compared (i. e. those stored in the following herbaria ANU, HITBC, IBK, IBSC, KUN, PE), as was the material of *Primulina* from recent fieldwork by the authors' team in South and Southwest China. All the morphological characters, such as leaves, inflorescences, flowers and capsules, were observed and measured in the field. The description, measurements, shape, color

and other details given in this description are based on living plants and specimens. We examined distinguished morphological characters of the presumed new species and the compared one, *P. wenii*, under a dissecting microscope. We described this presumed new species using the terminology of Wang et al. (1990, 1998).

Sampling and DNA sequencing

We randomly selected one plant from the population to collect its leaves for a DNA experiment. Fresh leaf materials were preserved in silica gel for quick drying. Total genomic DNA was extracted from dried leaves using modified cetyltrimethylammonium bromide (CTAB) protocol (Doyle and Doyle 1987). ITS and *trnL-F* were amplified and sequenced following the methods of Möller et al. (2009) and Smitsen et al. (2004), respectively. Besides, we downloaded the ITS and *trnL-F* sequences from GenBank for 188 *Primulina* species and two *Petrocodon* species. Species and GenBank accession numbers employed in this study are listed in Table 1.

Phylogenetic analysis

We assembled and aligned the newly obtained sequences and those from GenBank using MAFFT v.7.017 (Kato et al. 2002) and subsequently corrected and combined the ITS and *trnL-F* sequences in Geneious 9.1.4 (Kearse et al. 2012). We used the Maximum likelihood (ML) and Bayesian inference (BI) analyses to do the phylogenetic analysis of the ITS and *trnL-F* matrixes, and the combined ITS + *trnL-F* sequences data-set. The two best supported tree topologies from maximum likelihood (ML) analyses of ITS and *trnL-F* were visually compared for topological incongruence. A conflict in tree topologies of each tree was considered significant when incongruent topologies both received bootstrap values $\geq 80\%$ (Fu et al. 2022). The ML analyses were conducted using IQ-TREE 1.6.12 (Nguyen et al. 2015) with the GTR+R6 model and 1000 ultrafast bootstrap replicates. For BI analysis, we employed MrBayes v.3.2.6 (Ronquist et al. 2012) to obtain a maximum clade credibility (MCC) tree. Bayesian inference was performed using one million generations, four runs, four chains, a temperature of 0.001, 25% trees discarded as burn-in, and trees sampled every 1,000 generations (1,000 trees sampled in total) with GTR+I+G model.

Results

Phylogenetic analysis

The ITS matrix had a length of 782 characters, with 449 (57.4%) variable characters and 355 (45.4%) parsimony-informative. In comparison, the *trnL-F* matrix had a length of 836 characters, with 198 (23.6%) variable characters and 93 (11.1%) were parsimony-informative. The comparison of trees for ITS and *trnL-F* revealed no

Table 1. Species names and GenBank accession numbers of ITS and *trnL-F* DNA sequences used in this study.

Species name	Voucher number	ITS	<i>trnL-F</i>
<i>Petrocodon ainsliifolius</i>	CWH88	KF202291	KF202298
<i>Petrocodon hancei</i>	CIPeng22903	KY796057	KY796059
<i>Primulina alutacea</i>	YD07	KY394847	KY393441
<i>Primulina argentea</i>	YMBC	KY394848	KY393442
<i>Primulina baishouensis</i>	GXLG05	KY394849	KY393443
<i>Primulina balansae</i>	BALAN	MK747141	MK746274
<i>Primulina beiliuensis</i>	GXBLBC	KY394850	KY393444
<i>Primulina beiliuensis</i> var. <i>fimbribracteata</i>	SGQJ04	KY394851	KY393445
<i>Primulina bicolor</i>	SLHLCB	KY394852	KY393446
<i>Primulina bipinnatifida</i>	GXLG04	KY394853	KY393447
<i>Primulina bobaiensis</i>	BBGL01	KY394854	KY393448
<i>Primulina bogneriana</i>	WF7	MK747166	MK746225
<i>Primulina brachytricha</i>	DWDMCZ	KF498048	KY393450
<i>Primulina brachytricha</i> var. <i>magnibracteata</i>	KFC4193	MK369979	MK369994
<i>Primulina brunnea</i>	BRUN	MK747142	MK746275
<i>Primulina bullata</i>	GXJX06	KF498071	KY393451
<i>Primulina cangwuensis</i>	GXLG04	KY394853	KY393447
<i>Primulina cardaminifolia</i>	GXLB	MK747131	MK746255
<i>Primulina carinata</i>	NTBC	KY394858	KY393452
<i>Primulina cataractarum</i>	N1	MW900263	MW960358
<i>Primulina chizhouensis</i>	JXFY01	KY394860	KY393454
<i>Primulina colaniae</i>	WF8	MK747167	MK746224
<i>Primulina confertiflora</i>	GDYS05	MK747101	MK746253
<i>Primulina cordata</i>	HYH010	KC190200	KC190207
<i>Primulina cordifolia</i>	GXRA02	KY394863	KY393457
<i>Primulina cordistigna</i>	GDYCXZ	MK747118	MK746251
<i>Primulina crassirhizoma</i>	CJGL01	KY394864	KY393458
<i>Primulina crassituba</i>	HNSP	MK747147	MK746230
<i>Primulina curvituba</i>	GXHJ01	MK747137	MK746242
<i>Primulina danxiaensis</i>	P22865	JX506886	JX506778
<i>Primulina debaoensis</i>	DBGL01	KY394868	KY393462
<i>Primulina depressa</i>	DXS02	KY394869	KY393463
<i>Primulina diffusa</i>	PJGL01	KY394871	KY393465
<i>Primulina dongguanica</i>	DGBC	KY394872	KY393466
<i>Primulina drakei</i>	YNCP01	KY394873	KY393467
<i>Primulina dryas</i>	HKDMS	KY394875	KY393469
<i>Primulina duanensis</i>	DABC	KY394877	KY393471
<i>Primulina eburnea</i>	P22908	JX506891	JX506783
<i>Primulina effusa</i>	KFC4167	MK369976	MK369991
<i>Primulina fengkaiensis</i>	KFC4130	MK369975	MK369990
<i>Primulina fengshanensis</i>	KFC4195	MK369970	MK369985
<i>Primulina fimbrisepala</i>	P22863	JX506894	JX506786
<i>Primulina fimbrisepala</i> var. <i>mollis</i>	GXIB	JX506895	JX506787
<i>Primulina flavimaculata</i>	KFC3988	MK369974	MK369989
<i>Primulina floribunda</i>	DHGL01	KY394886	KY393480
<i>Primulina fordii</i>	LJM1207202	MG727881	MG727878
<i>Primulina fordii</i> var. <i>dolichotricha</i>	DHS01	MK747125	MK746247
<i>Primulina gemella</i>	GEME	MK747146	MK746254
<i>Primulina glabrescens</i>	GZLBSM	MK747132	MK746278

Species name	Voucher number	ITS	<i>trnL-F</i>
<i>Primulina glandaceistriata</i>	GXLCHW	MK747114	MK746256
<i>Primulina glandulosa</i>	GXPCLG	KY394887	KY393481
<i>Primulina gongchengensis</i>	GCGLO1	KY394889	KY393483
<i>Primulina grandibracteata</i>	YNHK	MK747121	MK746266
<i>Primulina guigangensis</i>	GXGGBBC	KY394892	KY393486
<i>Primulina guihaiensis</i>	GXLG036	KY394893	KY393487
<i>Primulina guizhongensis</i>	GXGZBC	KY394894	KY393488
<i>Primulina halongensis</i>	HLW01	KY394895	KY393489
<i>Primulina hedyotidea</i>	XWB	JX506905	JX506797
<i>Primulina heterochroa</i>	GXMES01	KY394898	KY393492
<i>Primulina heterotricha</i>	HNBT01	KY394899	KY393493
<i>Primulina hezhouensis</i>	HZXH	MK747143	MK746258
<i>Primulina hiepii</i>	WF2	MK747144	MK746223
<i>Primulina hochiensis</i>	GXIB	JX506903	JX506795
<i>Primulina huaijiensis</i>	GDHJ02	KF498127	KY393495
<i>Primulina huangii</i>	WF12	MK747138	MK746231
<i>Primulina hunanensis</i>	Xu11697	KU220602	KU220608
<i>Primulina jiangyongensis</i>	HNJY01	KY394902	KY393496
<i>Primulina jingxiensis</i>	LZXHGL01	KY394903	KY393497
<i>Primulina jiuwanshanica</i>	JWS	MK747116	MK746260
<i>Primulina juliae</i>	LJM1210011	MG727889	MG727873
<i>Primulina langshanica</i>	LSCZ	KY394907	KY393501
<i>Primulina latinervis</i>	XIN1	KY394908	KY393502
<i>Primulina laxiflora</i>	P22927	JX506910	JX506802
<i>Primulina lechangensis</i>	GDLC12	KY394910	KY393504
<i>Primulina leei</i>	LSGL01	KY394911	KY393505
<i>Primulina leiophylla</i>	GXJX07	KY394912	KY393506
<i>Primulina lepingensis</i>	JXLP01	KY394913	KY394913
<i>Primulina leprosa</i>	GXMS055	KY394914	KY393508
<i>Primulina lianpingensis</i>	CHLT016	MH3443910	MH344542
<i>Primulina liboensis</i>	GXJX08	KY394917	KY393511
<i>Primulina liguliformis</i>	GXIB	JX506912	JX506804
<i>Primulina lijiangensis</i>	GLS01	KY394919	KY393513
<i>Primulina linearicalyx</i>	KFC4141	MH032854	MH032841
<i>Primulina linearifolia</i>	GXNN01	KY394921	KY393515
<i>Primulina lingchuanensis</i>	LCXHGL01	KY394922	KY393516
<i>Primulina linglingensis</i>	LLBC	KY394923	KY393517
<i>Primulina linglingensis</i> var. <i>fragrans</i>	XHLLBC2	MK746285	MK746285
<i>Primulina liujiangensis</i>	LJGL01	KY394924	KY393518
<i>Primulina lobulata</i>	GDQX04	KF498054	KY393519
<i>Primulina longgangensis</i>	P22948	JX506916	JX506808
<i>Primulina longicalyx</i>	GXGL01	KY394927	KY393521
<i>Primulina longii</i>	XWB	JX506917	JX506809
<i>Primulina longzhouensis</i>	P22963	JX506918	JX506810
<i>Primulina lunglinensis</i>	GZXY04	KY394930	KY393524
<i>Primulina lunglinensis</i> var. <i>amblyosepala</i>	LCDE	MK747105	MK746281
<i>Primulina lungzhouensis</i>	GXJX10	KY394931	KY393525
<i>Primulina luochengensis</i>	LCWCGL01	KY394932	KY393526
<i>Primulina lutea</i>	1844	JX506921	JX506813
<i>Primulina lutescens</i>	PBLS01	MK747135	MK746263
<i>Primulina lutvittata</i>	KFC4149	MK369978	MK369993

Species name	Voucher number	ITS	<i>trnL-F</i>
<i>Primulina luzhaiensis</i>	HYH019	KC190197	KC190204
<i>Primulina mabaensis</i>	SZY02	KY394937	KY393531
<i>Primulina macrodonta</i>	GXIB	JX506923	JX506815
<i>Primulina maculata</i>	Xu11916	KU220604	KU220609
<i>Primulina maguanensis</i>	YNMG	MK747127	MK746267
<i>Primulina malipoensis</i>	YNMLP01	MK747123	MK746240
<i>Primulina medica</i>	GXPLCM	KY394940	KY393534
<i>Primulina melanofilamenta</i>	GXXA	MK747158	MK746277
<i>Primulina minor</i>	WXXH1	MK747160	MK746290
<i>Primulina minutimaculata</i>	GXLZ10	KY394941	KY393535
<i>Primulina moi</i>	SGWY03	KF498115	KY393536
<i>Primulina mollifolia</i>	GXESWC	KY394943	KY393537
<i>Primulina multifida</i>	DLXHGL01	KY394946	KY393540
<i>Primulina nandanensis</i>	GXJX02	KY393541	KY393541
<i>Primulina napoensis</i>	GXIB	JX506930	JX506821
<i>Primulina ningmingensis</i>	NMGL01	KY394949	KY393543
<i>Primulina obtusidentata</i>	GZJK01	KF498096	KY393544
<i>Primulina ophiopogoides</i>	GXFS01	KF498062	KY393545
<i>Primulina orthandra</i>	ZRBC2	MK747128	MK746286
<i>Primulina parvifolia</i>	GGSL01	KY394952	KY393546
<i>Primulina pengii</i>	W0397	KU220603	KU220610
<i>Primulina petrocosomeoides</i>	SHDBC	KY394953	KY393547
<i>Primulina pinnatifida</i>	MS02	KY394954	KY393548
<i>Primulina polycephala</i>	GDLZ06	KY394955	KY393549
<i>Primulina porphyrea</i>	DNGL01	KU173793	KU173799
<i>Primulina pseudoeburnea</i>	KY394958	KY394958	KY393552
<i>Primulina pseudoglandulosa</i>	GXYS06	KF498138	KY393482
<i>Primulina pseudobeterotricha</i>	XWB	JX506933	JX506824
<i>Primulina pseudolinearifolia</i>	JXY	MK747140	MK746280
<i>Primulina pseudomollifolia</i>	JMMXH1	MK747134	MK746244
<i>Primulina pseudoroseoalba</i>	JFHGL01	KY394959	KY393553
<i>Primulina pteropoda</i>	HNCJ01	KY394960	KY393554
<i>Primulina pungentisepala</i>	JEGL01	KY394962	KY393556
<i>Primulina purpurea</i>	ZHGL01	KY394964	KY393558
<i>Primulina qingyuanensis</i>	GDQX01	KY394965	KY394965
<i>Primulina renifolia</i>	GXDA02	KY394966	KY393560
<i>Primulina repanda</i>	GXBM03	KY394968	KY393562
<i>Primulina ronganensis</i>	GXRA01	KF498135	KY393564
<i>Primulina rongshuiensis</i>	GXRS01	KF498088	KY393565
<i>Primulina roseoalba</i>	LDGL01	KY394972	KY393566
<i>Primulina rosulata</i>	GXPL05	KU528874	KU528884
<i>Primulina rotundifolia</i>	OO3	KY394975	KY393569
<i>Primulina rubibracteata</i>	JH01R	KU173791	KU173797
<i>Primulina sclerophylla</i>	GXDA01	KY394979	KY393573
<i>Primulina secundiflora</i>	GZQZ	MK747119	MK746279
<i>Primulina shouchengensis</i>	GXYF02	KY394980	KY393574
<i>Primulina sichuanensis</i>	SCBC	MK747162	MK746264
<i>Primulina dryas</i>	HKDMS	KY394875	KY393469
<i>Primulina sinovietnamica</i>	Peng21956	MK369973	MK369988

Species name	Voucher number	ITS	<i>trnL-F</i>
<i>Primulina spinulosa</i>	GXFS02	KF498063	KY393576
<i>Primulina subrhomboidea</i>	GXYS02	KY395018	KY393577
<i>Primulina subulata</i>	GDYA01	KY395020	KY393579
<i>Primulina subulata</i> var. <i>guilinensis</i>	GXHYXH	KY394967	KY393561
<i>Primulina subulatisepala</i>	CQAYH01	MK747122	MK746246
<i>Primulina suichuanensis</i>	GDLC07	KY395021	KY393580
<i>Primulina swinglei</i>	GXXR01	KY395022	KY393581
<i>Primulina tabacum</i>	LZ01	KY395023	KY393582
<i>Primulina tenuifolia</i>	GXBM01	KY395024	KY393583
<i>Primulina tenuituba</i>	GZGY01	KY395025	KY393584
<i>Primulina tiandengensis</i>	GXTD03	KY395027	KY393586
<i>Primulina tribracteata</i>	GXFS04	KY395028	KY393587
<i>Primulina tribracteata</i> var. <i>zhuana</i>	1877	JX506952	JX506843
<i>Primulina tsoongii</i>	ZSGL01	KY395029	KY393588
<i>Primulina varicolor</i>	GXNP01	KF498086	KY393589
<i>Primulina verecunda</i>	LBJX01	KY395031	KY393590
<i>Primulina versicolor</i>	GDYD01	MK747155	MK746252
<i>Primulina vestita</i>	QZXT	MK747156	MK746282
<i>Primulina villosissima</i>	QXY01	KY395032	KY393591
<i>Primulina wenii</i>	WENI	MK747148	MK746284
<i>Primulina wentsaii</i>	GXLZ047	KY395033	KY393592
<i>Primulina wuae</i>	WSBC	MK747159	MK746265
<i>Primulina xinningensis</i>	GGL01	KY394891	KY393485
<i>Primulina xiziae</i>	ZJHZ01	KY395038	KY393597
<i>Primulina yangchunensis</i>	GDYC01	KY395039	KY393598
<i>Primulina yangshanensis</i>	GDNX01	KY395040	KY393599
<i>Primulina yangshuoensis</i>	GXYS07	KY395042	KY393601
<i>Primulina yingdeensis</i>	YD03	KU528876	KU528886
<i>Primulina yungfuensis</i>	GXIB	JX506957	JX506848
<i>Primulina zhoui</i>	WF18	MK747104	MK746222
<i>Primulina jiulianshanensis</i>	WF217	OP243287	OP243283

significant incongruence topology and both indicated that *Primulina jiuyishanensis* is closely related to *P. wenii* (Suppl. materials 1, 2). Because the combined dataset resulted in a better-resolved tree with higher support values, we use the combined dataset to do the further molecular studies. The combined data-set was 1628 characters, with 653 (40.1%) variable characters and 452 (27.8%) parsimony-informative, including the indels in all matrixes. The undescribed species and *P. wenii* were sister groups [Bayesian posterior probabilities (BIPP) = 1.00, ML ultrafast bootstrap support values (UFBoot) = 100%], and we found 10 and 1 different sites in the ITS (totally 615 bp) and the *trnL-F* (totally 750 bp) sequences between them, respectively (Table 2). belonging to a strongly supported clade (BIPP = 0.99, UFBoot = 98%) included *Primulina crassituba* (W.T.Wang) Mich.Möller & A.Weber, *P. effusa* F.Wen & B.Pan and *P. lobulata* (W.T.Wang) Mich.Möller & A.Weber (Wang 1982, 1989; Weber et al. 2011; Pan et al. 2017) (Fig. 1).




Figure 1. Phylogenetic tree of *Primulina* generated from maximum likelihood (ML) of combined *trnL-F* and ITS data-set. Numbers of each branch are support values in the order of UFBoot/BIPP. Stars indicate UFBoot = 100% or BIPP = 1.0. The dash (–) indicates a node with UFBoot or < 50% or BIPP < 0.5.

Table 2. Sequence differences of ITS and *trnL-F* regions between *Primulina jiulianshanensis* and *P. wenii*.

Species & marker	Sequences
<i>P. jiulianshanensis</i> ITS	1-CCCGAGAACATGTTTAAACACGCTTGCGT-30
<i>P. wenii</i> ITS	1-CCCGAGAACATGTTTAAAGACACGCTTGCGT-30
<i>P. jiulianshanensis</i> ITS	141-CGAGCGCCTCTCCGTCCTGGCTAAGTTCGC-170
<i>P. wenii</i> ITS	141-CGAGCGCCTCTCCGTACCGGTGAAGTTCGC-170
<i>P. jiulianshanensis</i> ITS	396-CGTTTTTTCCACGCTCAAAAGGTGTC-GGGGACGA-430
<i>P. wenii</i> ITS	396-CGTCTTTTCCACGCTCCAAAGGTGTCGGGGGAAGA-430
<i>P. jiulianshanensis trnL-F</i>	501-GTTCAAAAGTCCTTTATCTT-520
<i>P. wenii trnL-F</i>	501-GTTCAAAATTCCTTTATCTT-520

Taxonomic treatment

Primulina jiulianshanensis F.Wen & G.L.Xu, sp.nov.

urn:lsid:ipni.org:names:77318694-1

Figs 2, 3

Diagnosis. This new species differs from *P. wenii* by the combination of the following characteristics: petiole, both sides of leaf blades, adaxial surface of the calyx lobes, corolla inside toward the bottom, bract margins glandular-pubescent (*vs.* what above-mentioned eglandular-pubescent in *P. wenii*); lateral bracts 4–9 × ca. 2 mm, the central one 2–5 × 1–1.5 mm, adaxially glabrous but sparsely pubescent at apex (*vs.* lateral bracts 14–16 × 2.5–3.0 mm, the central one 10–12 × 1.3–1.6 mm, all adaxially pubescent); calyx lobes 8–11 × ca. 2 mm, each side with several brown serrate teeth at apex (*vs.* 14–15 × ca. 2.5 mm and entire); filaments and staminodes sparsely glandular-puberulent (*vs.* glabrous). Detailed morphological comparisons with *P. wenii* are provided in Table 3.

Holotype. CHINA, Jiangxi Province: Ganzhou City, Longnan County, Jiulianshan National Nature Reserve, growing in shady and moist cliffs in the forest, 24°34'8.9"N, 114°25'49.7"E, altitude ca. 440 m, April 20, 2021, *Guo-Liang Xu, JLSXGL-20210420* (Holotype IBK!; Isotype: KUN!)

Description. Herbs perennial, acaulescent, rhizome terete, 1–4 cm long, 0.8–1.5 cm in diam., leaves all basal, 4–8, petiole 10–40 × 4–13 mm, densely villous and very sparsely glandular-pubescent. Leaf blade oblong-elliptic or broadly elliptic, 4–13 × 4–8 cm, thickly chartaceous, more and less fleshly, adaxially pale green to dark green, densely white to light red villous and very sparsely glandular-pubescent, abaxially pale green, densely white villous and pubescent, and very sparsely glandular-pubescent, base apparently asymmetric but cuneate, margin irregularly obtuse-serrate, apex slightly obtuse, lateral veins 4–6 on each side, adaxially inconspicuously sunken, abaxially prominently raised. Inflorescence dichotomous cymes 2–4, axillary, 1–3-branched, 3–9-flowered or more; peduncle and pedicle green, erectly white or light red pubescent; peduncle 5–10 cm long, 2–3 mm in diam.; pedicle 5–15 mm long, 1–2 mm in diam. Bracts 3, lanceolate or spatulate, a pair on either side in same size, opposite, 4–9 mm long, ca. 2 mm wide, the middle one smaller, 2–5 mm long, 1–1.5 mm




Figure 2. *Primulina jiulianshanensis* sp. nov. **A** habitat **B** population **C** plant in blooming **D** habit **E** adaxial surface of mature leaf blades and petiole **F** abaxial surface of mature leaf blade and petiole.

wide, all three abaxially white or reddish pubescent, adaxially glabrous but sparsely pubescent at apex, margin entire, ciliate and very sparsely glandular-puberulent, apex acute; bracteoles 3, shape and indumentum same as bracts, 2–5 mm long, 1–1.5 mm wide. Calyx 5-parted, lobes lanceolate, 8–11 mm long, ca. 2 mm wide, nearly equal, abaxially densely white or light red villous, adaxially sparsely white puberulent and glandular-puberulent, margin entire but each side of calyx lobes with 1–3 purplish brown crenate at the apex. Corolla pinkish purple to bluish purple, 3.6–4.0 cm long; corolla tube funnellform, 2.6–3 cm long, mouth 1.3–1.6 cm in diam., base ca. 5 mm in diam., outside densely glandular-pubescent, inside from the middle to the base sparsely glandular-puberulent, and the upper part of the corolla tube glabrous; corolla tube abdomen with two obviously longitudinal ridges, the upper part (close to the mouth) of the longitudinal ridge dark bluish purple, and the lower part (close to the bottom)




Figure 3. *Primulina jiulianshanensis* sp. nov. **A** cyme **B** adaxial surfaces of bracts **C** abaxial surfaces of bracts **D** adaxial surfaces of calyx lobes **E** abaxial surfaces of calyx lobes **F** opened corolla **G** stamens and anthers **H** one of lateral staminodes **I** pistil **J** immature capsules **K** transverse section of capsule.

changing into yellowish brown; a dark reddish-brown lump on the upper throat of the corolla tube inside and between upper lip lobes, ovate to spatulate, extending to the middle of the corolla tube, the lump densely glandular-puberulent; a narrow triangular thickened dark reddish-brown stripe extending to the middle of the corolla tube inside at each side of corolla tube and at the junction of the abaxial and adaxial lip;

limb distinctly 2-lipped, adaxial lip 2-parted to the middle, lobes broadly ovate to semicircular, apex round, 6–8 mm long, 6–9 mm wide at the bottom; abaxial lip 3-parted to near the base, lobes elliptical to oblong, 8–12 mm long, 7–9 mm wide at the bottom. Stamens 2, adnate to 1.8 cm above the base of corolla tube; filaments linear, yellow from middle to base but white upper half, 8–11 mm long, geniculate near the base, glandular-puberulent; anthers reniform, slightly constricted at the middle, densely villous and fewer glandular-puberulent; staminodes 3, yellowish, lateral ones ca. 6 mm long, adnate to 14 mm above the base of corolla tube, straight, linear, very sparsely glandular-puberulent, apex capitate, the central one ca. 1 mm long, adnate to 5–6 mm above the base of corolla tube, glabrous. Disc annular, ca. 1 mm high, margin undulate, glabrous, white. Pistil pale green, 2.8–3.2 cm long; style linear, 1.9–2.3 cm long, ca. 1 mm in diam., upper part densely glandular-puberulent, lower part densely glandular-puberulent and eglandular-puberulent, ovary oblong, ca. 10 mm long, ca. 2 mm in diam., densely villous and glandular-pubescent, parietal placenta. Stigma acute triangle to narrowly obtrapeziform, 2-lobed, ca. 3 mm long. Capsule linear, 2–4 cm long, parietal placenta, densely villous and glandular-pubescent.

Phenology. Flowering from April to May, fruiting from June to September.

Etymology. The specific epithet '*jiulianshanensis*' is derived from the type locality, Jiulianshan National Natural Reserve, Jiangxi Province, China.

Vernacular name. 九连山报春苣苔 (Chinese name); Jiǔ Lián Shān Bào Chūn Jù Tái (Chinese pronunciation).

Distribution and habitat. We found three small subpopulations in Jiulianshan National Nature Reserve in Jiangxi Province, which are distributed in the shady and wet place on the cliffs under the evergreen broad-leaved forest in the reserve. And the new species is mainly accompanied by *Begonia palmata* D. Don, *Utricularia striatula* J. Smith, *Selaginella moellendorffii* Hieron., *S. involvens* (Sw.) Spring, etc.

Conservation status. At present, only three small subpopulations with total ca. 300 mature individuals of the new species are known in the type locality, Jiulianshan National Natural Reserve, Jiangxi Province, China. The three subpopulations are stable because they are in the reserve. The known AOO and EOO of the new species are about 0.2 km² and 25 m², respectively. Thus, if considering its fewer individuals of three subpopulations, it should be temporarily assessed as Near Threatened (NT), following the IUCN Red List Categories and Criteria (IUCN Standards and petitions committee 2022).

Notes. The mainly morphological differences are showed in diagnosis and Table 3. In addition, the insides of the corolla tube are also somewhat different. For example, at the junction of the abaxial and adaxial lip, there are two narrow triangular thickened dark reddish-brown stripes inside the corolla tube in *Primulina jiulianshanensis*, but there are only two bluish purple spots at the same places in *P. wenii*; at corolla tube abdomen of *P. jiulianshanensis*, the upper parts of the longitudinal ridges are dark bluish purple, and the lower parts are yellowish brown, but the longitudinal ridges inside corolla tube of *P. wenii* are all dark bluish purple; the lump on the upper throat inside corolla tube of *P. jiulianshanensis* is dark reddish-brown, but *P. wenii* is dark bluish brown (Li et al. 2017).

Table 3. Comparisons between the characters of *Primulina jiulianshanensis* and *P. wenii*.

Characters	<i>Primulina jiulianshanensis</i>	<i>P. wenii</i>
Petiole indumentum	densely villous and very sparsely glandular pubescent	densely villous
Leaf blades indumentum	adaxially densely white to light red villous and very sparsely glandular pubescent, abaxially densely white villous and pubescent, and very sparsely glandular-pubescent	adaxially densely pubescent and villous, abaxially densely appressed pubescent
Bract	lateral ones 4–9 × ca. 2 mm, the middle one 2–5 × 1–1.5 mm, all abaxially white or reddish pubescent, adaxially glabrous but sparsely pubescent at apex, margin ciliate and very sparsely glandular-puberulent	lateral ones 14–16 × 2.5–3.0 mm, the central one 10–12 × 1.3–1.6 mm; outside white pubescent and villous, adaxially white pubescent, margin entire and ciliate
Calyx lobes	adaxially sparsely white puberulent and glandular puberulent, margin entire but each side of calyx lobes with 1–3 brown crenate teeth at apex; lobes 8–11 × ca. 2 mm	adaxially sparsely shortly pubescent to nearly glabrous, margin entire, lobes 14–15 × ca. 2.5 mm
Filament	yellow from middle to base but white upper half, glandular-puberulent	white, glabrous
Staminodes	yellowish, lateral ones very sparsely glandular-puberulent	white, glabrous

The type locality of *Primulina wenii* is Rixi Township, Fuzhou, Fujian Province, while the type locality of *P. jiulianshanensis* is Jiulian Mountain, Jiangxi Province. Their type localities are separated by the Wuyi Mountains, and the two places are more than 500 kilometers apart. *P. wenii* has a narrow distribution range and is only recorded in Rixi Township, Fuzhou, Fujian Province (Li et al. 2017). This area is coastal and the climate zone of this region belongs to the typical subtropical marine monsoon climate. *P. wenii* grows in the limestone evergreen broad-leaved forest area with stable morphology. *P. jiulianshanensis*, grows on the rocks under the evergreen broad-leaved forest in Jiulianshan Nature Reserve, which is a typical Danxia landform. The soil forming rock are mainly sandstone and conglomerate in Jiulianshan Nature Reserve, which has a subtropical monsoon humid climate. The morphology of the *P. jiulianshanensis* in three subpopulations is also relatively stable. Therefore, considering the differences in molecular, morphological, and habitats between *P. jiulianshanensis* and *P. wenii*, they should be classified as two different species.

Except for a few species, such as *Primulina fimbrisepala* (Hand.-Mazz.) Yin Z. Wang, *P. eburnea* (Hance) Yin Z. Wang, *P. tenuituba* (W.T. Wang) Yin Z. Wang, *P. juliae* (Hance) Mich. Möller & A. Weber, most of the species of *Primulina* are narrowly distributed and endemic. Among the known species worldwide, more than 170 species are endemic to Karst areas in southern to southwestern China and to northern Vietnam (Wei 2018; Xu et al. 2020b). However, the diversity of *Primulina* in the Danxia landform has not been well understood so far (Yu et al. 2019). For example, the new species, *P. suichuanensis*, was found in the Danxia landform in Jiangxi Province (Zhou et al. 2016). It should be noted that a new provincial record for *Primulina wenii* from Jiangxi Province was discovered in Jiulianshan Nature Reserve (Liao et al. 2020). However, we carefully examined the voucher specimen (No. PVHJX-05557, stored in GNNU), and we noted that the voucher was collected from the same subpopulation of *P. jiulianshanensis* from the same site in Jiulianshan Nature Reserve. Further, we found there are many incon-

sistencies between the morphological description from the article (Liao et al. 2020) and the corresponding voucher specimens. Thus, this species' new provincial record of *P. wenii* in Jiangxi province is a mistaken identification. Lastly, none of the close relatives of *Primulina* are morphologically similar to this new species found in Jiangxi (Fig. 1). Thus, the new taxon is not easily confused with the others in this province.

Acknowledgements

We want to thank Stephen Maciejewski, the Gesneriad Society, and Michael LoFurno, Associate Professor, Temple University, Philadelphia, USA, for their editorial assistance. This study was financially supported by the Basic Research Fund of Guangxi Academy of Sciences (CQZ-C-1901), the Key Sci. & Tech. Research and Development Project of Guangxi (Guike AD20159091 & ZY21195050), the capacity-building project of SBR of CAS (KFJ-BRP-017-68), Basal Research Fund of GXIB (Guizhifa010) and the Fund of Technology Innovation Alliance of Flower Industry (2020hhlm005).

References

- Doyle JJ, Doyle JL (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* 19: 11–15.
- Du C, Liu J, Ye W, Liao S, Ge BJ, Liu B, Ma JS (2021) Annual report of new taxa and new names for Chinese plants in 2020. *Shengwu Duoyangxing* 29(8): 1011–1020. <https://doi.org/10.17520/biods.2021122>
- Fu LF, Wen F, Maurin O, Rodda M, Gardner EM, Xin ZB, Wei YG, Monro AK (2022) A revised delimitation of the species-rich genus *Pilea* (Urticaceae) supports the resurrection of *Achudemia* and a new infrageneric classification. *Taxon* 71(4): 796–813. <https://doi.org/10.1002/tax.12711>
- Hance HF (1883) New Chinese Cyrtandreae. *Journal of Botany. British and Foreign* 21: 165–170.
- IUCN Standards and petitions committee (2022) Guidelines for Using the IUCN Red List Categories and Criteria. Version 15. Prepared by the Standards and petitions committee. <https://www.iucnredlist.org/resources/redlistguidelines> [Accessed on 4 Mar. 2022]
- Katoh K, Misawa K, Kuma K, Miyata T (2002) MAFFT: A novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research* 30(14): 3059–3066. <https://doi.org/10.1093/nar/gkf436>
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A (2012) Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28(12): 1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>
- Li J, Chen X, Li S, Yan LJ (2017) *Primulina wenii* (Gesneriaceae), a new species from China. *Bangladesh Journal of Plant Taxonomy* 24(2): 173–182. <https://doi.org/10.3329/bjpt.v24i2.35113>

- Liao HH, Li L, Wei Y, Liang YL (2020) *Primulina wenii* (Gesneriaceae), a new record species from Jiangxi Province. South China Forestry Science 48(3): 45–46, 71. <https://doi.org/10.16259/j.cnki.36-1342/s.2020.03.009>
- Möller M, Pfosser M, Jang CG, Mayer V, Clark A, Hollingsworth ML, Barfuss MHJ, Wang YZ, Kiehn M, Weber A (2009) A preliminary phylogeny of the ‘Didymocarpoideae’ based on three molecular data sets: Incongruence with available tribal classifications. American Journal of Botany 96(5): 989–1010. <https://doi.org/10.3732/ajb.0800291>
- Nguyen LT, Schmidt HA, Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular Biology and Evolution 32(1): 268–274. <https://doi.org/10.1093/molbev/msu300>
- Ning ZL, Wang J, Tao JJ, Kang M (2014) *Primulina lepingensis* (Gesneriaceae), a new species from Jiangxi, China. Annales Botanici Fennici 51(5): 322–325. <https://doi.org/10.5735/085.051.0508>
- Pan B, Zou LL, Zhang RL, Kang M, Wen F (2017) *Primulina effusa* F.Wen & B.Pan, a new species of Gesneriaceae from limestone areas of South China. Guangxi Zhi Wu 37(10): 1250–1256. <https://doi.org/10.11931/guihaia.gxzw201706006>
- Peng YS, Tang ZB, Xie YF (2021) Inventory of species diversity of Jiangxi vascular plants. China Forestry Publishing House, Beijing, 434 pp.
- POWO (2022) Plants of the World Online. Royal Botanic Gardens, Kew. [Accessed 31 October 2022] <http://www.plantsoftheworldonline.org/>
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Smissen RD, Breitwieser I, Ward JM (2004) Phylogenetic implications of trans-specific chloroplast DNA sequence polymorphism in New Zealand Gnaphalieae (Asteraceae). Plant Systematics and Evolution 249(1–2): 37–53. <https://doi.org/10.1007/s00606-004-0209-0>
- Wang WT (1982) Notulae de Gesneriaceis Sinensibus (IV). Bulletin of Botanical Research 2(4): 37–64.
- Wang WT (1989) Tres Species Novae *Chirita* (Gesneriaceae) e Hunan. Bulletin of Botanical Research 9(4): 21–28.
- Wang WT, Pan KY, Li ZY (1990) Gesneriaceae. In: Wang WT (Ed.) Flora Republicae Popularis Sinicae, Vol. 69. Science Press, Beijing, 331–418.
- Wang WT, Pan KY, Li ZY, Weitzman AL, Skog LE (1998) Gesneriaceae. In: Wu ZY, Raven PH (Eds) Flora of China Vol. 18. Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis.
- Wang YZ, Mao RB, Liu Y, Li JM, Dong Y, Li ZY, Smith JF (2011) Phylogenetic reconstruction of *Chirita* and allies (Gesneriaceae) with taxonomic treatments. Journal of Systematics and Evolution 49(1): 50–64. <https://doi.org/10.1111/j.1759-6831.2010.00113.x>
- Weber A, Middleton DJ, Forrest A, Kiew R, Lim CL, Rafidah AR, Sontag S, Triboun P, Wei YG, Yao TL, Möller M (2011) Molecular systematics and remodeling of *Chirita* and associated genera (Gesneriaceae). Taxon 60(3): 767–790. <https://doi.org/10.1002/tax.603012>
- Wei YG (2018) The Distribution and Conservation Status of Native Plants in Guangxi, China. China Forestry Publishing House, Beijing, 876 pp.

- Xu MZ, Kong HH, Kang M, Yang LH (2020a) A new species of *Primulina* (Gesneriaceae) from Danxia landform in Jiangxi, China. *Taiwania* 65(2): 163–166. <https://doi.org/10.6165/tai.2020.65.163>
- Xu MZ, Yang LH, Kong HH, Wen F, Kang M (2020b) Congruent spatial patterns of species richness and phylogenetic diversity in karst flora: Case study of *Primulina* (Gesneriaceae). *Journal of Systematics and Evolution* 59(2): 251–261. <https://doi.org/10.1111/jse.12558>
- Yu XY, Li JM, Ren MX (2019) Adaptive differentiation of Gesneriaceae on karst and Danxia landforms in Southern China. *Guangxi Sciences* 26(1): 132–140. <https://doi.org/10.13656/j.cnki.gxkx.20190321.003>
- Zhou DS, Zhou JJ, Li M, Yu XL (2016) *Primulina suichuanensis* sp. nov. (Gesneriaceae) from Danxia landform in Jiangxi, China. *Nordic Journal of Botany* 34(2): 148–151. <https://doi.org/10.1111/njb.00956>

Supplementary material 1

The tree of the *trnL-F* plastid marker for *Primulina* species

Author: Zheng-Yu Zuo

Data type: (phylogenetic, genomic, tree)

Explanation note: The *trnL-F* plastid marker and the ITS nuclear marker separately and discuss any incongruences between the two markers, particularly with respect to the position of the new species.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/phytokeys.226.96351.suppl1>

Supplementary material 2

The tree of the ITS nuclear marker for *Primulina* species

Author: Zheng-Yu Zuo

Data type: (phylogenetic, genomic, tree)

Explanation note: The *trnL-F* plastid marker and the ITS nuclear marker separately and discuss any incongruences between the two markers, particularly with respect to the position of the new species.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/phytokeys.226.96351.suppl2>