

***Lithocarpus orbicarpus* (Fagaceae), a new species of Stone Oak from Phang Nga province, Thailand**

Joeri S. Strijk¹, Sukontip Sirimongkol², Sukid Rueangruea²,
Nikom Ritphet², Voradol Chamchumroon²

1 Ecological Evolution Group, Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Menglun, Mengla, Yunnan, 666303, PR China **2** Forest Herbarium (BKF), Department of National Parks, Wildlife and Plant Conservation, 61 Phahonyothin Rd., Chatuchak, Bangkok 10900, Thailand

Corresponding author: Joeri S. Strijk (joeristrijk@xtbg.org.cn)

Academic editor: H. Schaefer | Received 14 October 2013 | Accepted 2 February 2014 | Published 11 February 2014

Citation: Strijk JS, Sirimongkol S, Rueangruea S, Ritphet N, Chamchumroon V (2013) *Lithocarpus orbicarpus* (Fagaceae), a new species of Stone Oak from Phang Nga province, Thailand. PhytoKeys 34: 33–45. doi: 10.3897/phytokeys.34.6429

Abstract

A new species is described, *Lithocarpus orbicarpus* Strijk, collected from Ton Pariwat Wildlife Sanctuary, Mueang district, Phang Nga Province in the Peninsular Floristic Region of Thailand. We provide the first technical illustrations and colour photographs of the new species, as well as a description of its conservation status and the collecting locality. The species can be easily distinguished by its unique orbicular acorns, each covered with a dense pattern of irregularly placed scales, which completely conceal the nut, except for a tiny apical pore, and which are arranged in a dense cluster on an erect woody spike. We also provide an amendment to the existing diagnostic key to *Lithocarpus*, and discuss important differences with morphologically similar species found in Thailand and the surrounding region.

Keywords

Lithocarpus orbicarpus, new species, Stone Oaks, Fagaceae, Thailand, Ton Pariwat Wildlife Sanctuary

Introduction

Lithocarpus Blume is the second largest genus within Fagaceae, with over 300 species described (Camus 1948, 1954; Phengklai 2008; Soepadmo 1972; Wu et al. 1999). Members of the genus, commonly called Stone Oaks, can be found in (sub-)

tropical broad-leaved evergreen forests near sea level to mixed forests at altitudes of over 3200 m. The geographic distribution of *Lithocarpus* roughly covers southern and south-eastern Asia from eastern India to southern Japan, to the Philippines and southward to New Guinea. *Lithocarpus densiflorus* (Hook. & Arn.) Rehder, the only North American member of the genus was recently moved to a new monotypic genus (Manos et al. 2008). All species within *Lithocarpus* are trees, ranging from small understory elements, to very large emergent trees. Many species have a geographically wide distribution and in some locations may constitute the main canopy forming elements together with other Fagaceae (*Lithocarpus*, *Quercus*, *Castanopsis*), Lauraceae and Theaceae. Leaves are simple, entire, rarely serrate, generally glabrous and mostly spirally arranged. Male and female flowers are white to pale whitish-yellow. Genders can be either on separate inflorescences, on the same inflorescence mixed throughout, or with female flowers below and male flowers terminal, arranged with dense indumentum on erect spikes. Male flowers are solitary or in clusters of three or more, with the perianth campanulate or cup-shaped, usually 6-lobed, partially united; stamens generally 12. Female flowers solitary or in clusters of three, perianth like male flowers but smaller; 12 staminodes; 3(-4) styles (Camus 1948, 1954; Phengklai 2008).

Previous studies and the most recent treatment for Thailand have recovered 57 species of *Lithocarpus* (Barnett 1940; Phengklai 2004, 2008). During fieldwork in remote Ton Pariwat Wildlife Sanctuary, carried out as part of ongoing research on the genomics, systematics, biogeography and evolution of Asian Fagaceae, we made collections of an individual tree with unique features that could not be matched with any previously described taxa in Fagaceae. After careful examination of herbaria and literature, comparison of other specimens collected during the fieldwork and consultation of specialists on the regional flora, we report this collection here as a new species, placed within the genus *Lithocarpus*.

Taxonomy

Lithocarpus orbicarpus Strijk, sp. nov.

urn:lsid:ipni.org:names:77135982-1

http://species-id.net/wiki/Lithocarpus_orbicarpus

Fig. 1, 2

Type. THAILAND, Ton Pariwat Wildlife Sanctuary, Mueang district, Phang Nga Province, 8°37'25"N; 98°33'14"E; alt. 455 m, 16 July 2013, Chamchumroon et al. 5823 (Holotype: BKF; Isotypes: E, K, L, SING).

Diagnosis. *Lithocarpus orbicarpus* is a small-medium sized tree. It differs from similar species by its unique orbicular acorns, each covered with a dense pattern of irregularly placed scales, which completely conceal the nut, except for a tiny apical pore, and which are arranged in a dense cluster on an erect woody spike. Unique for Thai species of *Lithocarpus*, almost the entire surface of the round nut is covered with scar area (re-

ceptacle tissue), leaving only the topmost part of the nut covered with a thin vestigial exocarp layer. Pending discovery of additional individuals, the species appears to be locally restricted to low-mid-elevation forests in the peninsular region of Thailand.

Description. Small-medium sized tree, up to 15 m tall. *Bark* smooth to slightly rough grey-green, with superficial horizontal lines. Sapwood white to yellow, with inner bark ridges forming light brown longitudinal slits in sapwood surface. *Branches* dark brown to grey brown, mostly glabrous, densely lenticellate; young twigs, leaf buds and old fruits with short, soft (occasionally long) gray indumentum. Leaf buds tiny and terminal buds solitary. *Leaves* simple; lamina elliptic to oblanceolate with (strongly) acuminate tip, 11.0–22.3 × 4.4–7.1 cm. Margin entire. Leaves often with slightly asymmetric lamina. Leaf apex acuminate to strongly acuminate, leaf base cuneate to slightly attenuate. Both surfaces generally glabrous except emerging leaf buds, terminal shoots and young leaves, which have soft grey indumentum. Young leaves light green, but turning dark green above and glaucous below when older. *Venation*. Pinnately veined; secondary venation discretely anastomosing near the leaf margin. Pairs of secondary veins 9–13, slightly raised and clearly visible on underside of leaf. *Peduncles* carrying fruits 5–11 cm long, up to 1 cm thick at the base, glabrescent, grey-brown and densely lenticellate. *Male and female inflorescences* not seen. *Infructescence* a woody spike, terminal, up to 15–21 cm long. Fruits sessile on thick woody peduncle, closely pressed against each other, but walls of individual units not fused. Number of fruits per infructescence very variable, ranging from 9–20 units. *Acorn*. Orbicular, globose, 2.7–3.4 by 2.9–3.5 cm (including cupule) and covered with glabrous, semi-concentric interlocking ridges when young, which transform over time into ridges with irregularly placed scales. Cupule nearly completely enclosing the nut, indehiscent, but showing small cracks when mature; fruit wall up to 4–6 mm thick, apical pore very small, 1–4 mm wide, exposing the persistent punctiform styles (3) and a tiny fraction of vestigial exocarp. Young cupule walls light green, ridges light to dark brown. Old cupule walls turning light brown to yellow-brown and pubescent with short (occasionally long), greyish-yellow indumentum. *Nut* 1 in each cupule, ball shaped, globose, 2.4–2.9 by 2.6–3.0 cm. Up to 95% of surface area of the nut made up of scar area (receptacle tissue), upper 5% of surface area of the nut slightly raised and made up of vestigial exocarp layer. Nut scar pale yellow-whitish, tiny exocarp layer light brown. Scar area covered with deep grooves and red-brown to purplish vein-like lines, stretching down to the base of the nut. Up to 5/6 of the scar area of the young nut (from the base upward) covered with dotted pattern of small depressions. Cotyledons black when dried.

Flowering and fruiting. Flowering unknown, but thought to be around November-January. Infructescences on the ground in July, fruiting commencing possibly as early as April-May.

Distribution. This species is only known from Thailand, and has not been recorded outside Ton Pariwat Wildlife Sanctuary, Mueang district, Phang Nga Province. During our field survey, we found only one individual tree, located on a gentle sloping section of closed dense forest. Additional survey work will have to be undertaken to determine the actual population size of this species within the wildlife sanctuary.




Figure 1. *Lithocarpus orbicarpus* Strijk, sp. nov. Chamchumroon et al. 5823 (BKF). Technical illustration. **A** Habit **B** Detail of glabrous young fruit with ridges and apical pore **C** Detail of interior of young fruit, showing nearly complete fruit scar, covered umbo and ‘pitted pattern’ on the nut surface **D** Infructescence with ripe fruits showing highly irregular scaly patterns on the fruit exterior **E** Details of seed, from left to right: bottom view, side view, top view. Note venation and crevice pattern on surface of fruit, and cover of the umbo section. All drawings by J.S. Strijk.

Ecology. This species grows in dense hillside forest at an elevation of around 450 m.

Vernacular name. Unknown.

Etymology. *Lithocarpus orbicarpus* is named after its unique orbicular acorns, arranged densely clustered on an erect spike, of which the cup almost completely encloses the nut (except for a tiny apical pore). The exterior of the cup is covered with a dense pattern of ridges, transforming with age into horizontal and vertical lines with irregularly placed scales. Apart for a tiny portion of the umbo which is covered with vestigial exocarp, the exterior surface of the nut is completely covered by scar. Although the cupule nearly completely covers the nut, the latter is free and not fused to the wall of the cupule – a condition which occurs throughout the genus (Cannon 2001; Cannon and Manos 2001). Upon drying, the shrinking of the cotyledons inside the nut causes the whole infructescence to make a rattling sound when shaken.

Discussion. Thailand has a total of 121 species, 2 subspecies and 2 varieties of Fagaceae in four genera indigenous to the country. Of these, nine species (*Castanopsis*: 4; *Quercus*: 1; *Lithocarpus*: 4) and 1 subspecies (*Quercus*: 1) are national endemics (Bunpha et al. 2011; Phengklai 2008). Three of the endemic *Lithocarpus* species (including *L. orbicarpus*) are restricted to small ranges in the peninsular region. Of the 58 *Lithocarpus* known to occur in Thailand (including this newly described species), 43 species (74%) can be found in the Peninsular floristic province making it the most diverse floristic region (in terms of *Lithocarpus* species) followed by the North (25 species – 43%). While the Northern (and North-eastern) floristic province is characterized by species that reach high elevation habitat (>75% can be found between 1200–2500 m), the Peninsular complement includes species that are restricted to lower elevations (21 species restricted <500 m asl). With two exceptions (*L. falconeri* (Kurz) Rehder and *L. erythrocarpus* (Ridl.) A.Camus), none of these species are shared with the North (or North-east). In contrast, four of the Peninsular lowland species (*L. lucidus* (Roxb.) Rehder; *L. maingayi* (Benth.) Rehder; *L. reinwardtii* (Korth.) A.Camus; *L. tubulosus* (Hickel & A.Camus) A.Camus) are shared with the South-eastern floristic province, in addition to *L. elephantum* (Hance) A.Camus and *L. pierrei* (Hickel & A.Camus) A.Camus, which can only be found in the South-eastern region.

Lithocarpus in Thailand can be further characterized based on their acorn (cupule+nut) properties. The open- or semi-open type, in which the cupule shape ranges from saucer shaped-flat to covering up to 4/5th of the nut is present in 37 species (64%), while the remaining 18 species have cupules that entirely cover the nut, in most cases leaving a tiny portion of the umbo uncovered. With one exception (*L. truncates* (King ex Hook.f.) Rehder), all closed-type species occur in the Peninsula, and eight are restricted to it (two shared with the Southeast). *L. orbicarpus* resembles species in this group, with its indehiscent and near-closed cupules, restricted geographical distribution and presence in lower elevation habitat, but is clearly distinct from them based on the structure and shape of the fruit, the infructescence and the properties of the nut.

During our field survey, we encountered and collected additional Fagaceae species, e.g. *Lithocarpus reinwardtii* (Korth.) A.Camus (Burma, Cambodia, Malaysia, Indonesia), *L. sundaicus* (Blume) Rehder (Malaysia, Indonesia, Brunei), *L. cantleyanus* (King ex Hook.f.)




Figure 2. *Lithocarpus orbicarpus* Strijk, sp. nov. Chamchumroon et al. 5823 (BKF). Pictures from field collection. **A** Branch with leaves **B** Young infructescences **C** Bark and sapwood **D** Fresh fruit – top view **E** Fresh fruit – side view **F** Fresh fruit – bottom view **G** Dried fruit – top view **H** Dried fruit – side view **I** Dried fruit – bottom view (**G–J**: blue lines demarcate vestigial exocarp; red lines demarcate scar area (receptacle tissue)) **K** Dried infructescence **L** Young acorn, opened up to show dotted pattern of small depressions and surface structure of the umbo **M** Cross-section of dried nut, showing seed coat and black cotyledons **N** Terminal leaf, twig and very small terminal bud, showing grey indumentum **O** Young emerging leaf with soft grey indumentum. All pictures by S. Sirimongkol and J. S. Strijk.

Rehder (Burma, Malaysia, Singapore) and several, as of yet, unidentified collections. Additional species encountered in previous surveys in the wildlife sanctuary are: *Castanopsis javanica* (Blume) A.DC. (Vietnam, Malaysia, Singapore, Indonesia); *Castanopsis nephelioides* King ex Hook. f. (Malaysia, Singapore); *Castanopsis purpurea* Barnett (Thailand (endemic)); *Castanopsis wallichii* King ex Hook. f. (Malaysia, Singapore, Indonesia); *Castanopsis inermis* (Lindl.) Benth. & Hook. f. (Burma, Malaysia, Singapore, Indonesia, Philippines); *Lithocarpus bennettii* (Miq.) Rehder (Malaysia, Singapore, Indonesia); *Lithocarpus clementianus* (King) A.Camus (Malaysia, Indonesia); *Lithocarpus eucalyptifolia* (Hickel et A.Camus) A.Camus (Burma, Vietnam, Cambodia); *Lithocarpus falconeri* (Kurz) Rehder (Burma, Malaysia); *Lithocarpus garrettianus* (Craib) A.Camus (China, Burma, Laos, Vietnam); *Lithocarpus lucidus* (Roxb.) Rehder (India, Malaysia, Singapore, Indonesia, Brunei); *Lithocarpus macphailii* (Henders.) Barnett (Malaysia, Indonesia); *Lithocarpus wrayi* (King) A.Camus (Vietnam, Malaysia, Indonesia) and *Quercus oidocarpa* Korth. (Burma, Vietnam, Malaysia, Indonesia). Out of these 17 species, at least 14 have a geographic distribution that is primarily or completely located in the Sundaland biogeographic region. Conversely, only three species have a distribution that is mostly or entirely contained within the Indochinese biogeographic region. Despite the paucity of detailed biological records on this remote area, the distributional data shows us that the Fagaceae flora in Ton Pariwat Wildlife Sanctuary is distinctly Sundaic in composition, and differs substantially from forests in the more northern Indochinese region. This is in fact not surprising, as the Ton Pariwat Wildlife Sanctuary is located near the southern edge of one of the world's major phyto-geographic and zoogeographic transition zones: the ~500km stretch between the biogeographically distinct and well-known Isthmus of Kra – the narrowest part of the connection between mainland Southeast Asia and the Malay Peninsula (10°30'N) – and the line between Kangar (Malaysia) - 6°51'N, and Pattani (Thailand) 6°87'N. Within this relatively short distance, an abrupt and major shift takes place from northern Indochinese flora and fauna, to those distinct for the southern Sundaland region (Hughes et al. 2003; Meijaard 2009; Parnell 2013; Van Steenis 1950; Woodruff 2003; Woodruff and Turner 2009).

When looking at the Fagaceae flora in the wider region, some properties of *L. orbicarpus* resemble species in Peninsular Malaysia, such as *Castanopsis nephelioides* King ex Hook.f., but can easily be distinguished from species within *Castanopsis* (D.Don) Spach, based on the presence of bark ridges that penetrate the sapwood, the nut which is not fused to the cup and the orbicular shape of the fruit. Congeneric species from the Indo-Chinese zone that share some characters with *L. orbicarpus* are *L. rouletii* (Hickel & A.Camus) A.Camus (but fruit with basal scar, flattened, dehiscent; South Vietnam); *L. pachycarpus* (Hickel & A.Camus) A.Camus (but leaves with yellow indumentum, cupules pear-shaped, acorns not orbicular; Vietnam-Laos); *L. kontumensis* A.Camus (but cupules truncate, higher than acorn, cupules sometimes fused; Vietnam-Laos); *L. lepidocarpus* (Hayata) Hayata (but cupules truncate, sometimes fused, fruit not orbicular; central and south Taiwan); and *L. laoticus* (Hickel & A.Camus) A.Camus (but cupule ovoid, high elevation habitat (Tibet, south and central China, Vietnam) (Wu et al. 1999). Within Thailand, *L. orbicarpus* is unique in its combination of properties, and we outline some of the defining differences with Thai species in Table 1 below.

Table 1. Morphological differences between *L. orbicarpus* and other Thai species of Fagaceae.

Characters	<i>L. orbicarpus</i> Strijk	<i>L. encleisocarpus</i> A.Camus	<i>L. wrayi</i> (King) A.Camus	<i>C. nephelioides</i> King ex Hook.f.
1. Nut wall	Free from the cup	Free from the cup	Free from the cup	Fused to the cup
2. Cupule enclosure	Almost complete, but small apical pore showing flat umbo remains ($\leq 5\%$). Indehiscent.	Almost complete, but raised umbo free ($\pm 5\text{--}10\%$). Easily dehiscent in irregular parts.	Almost complete, but raised umbo free ($\pm 5\text{--}15\%$). Indehiscent.	Enclosure complete. Indehiscent.
3. Nut shape	Orbicular.	Ovoid to globose.	Broadly conical.	Ovoid, usually depressed to one longitudinal side.
4. Cup surface	Spines absent. Small, flattened scales present. Irregularly intersecting lines present. Old acorns pubescent with short (occasionally long), greyish-yellow indumentum.	Spines and scales absent. Wall smooth, densely greenish-brown hairy.	Alternate pseudo-spines and free scales present; pseudo-spines incurved or erect.	Sparsely covered with short, woody spines, 2-3 branched reclining and decurved.
5. Acorn shape	Orbicular, symmetric; young fruits occasionally slightly skewed in young and dense infructescences	Ovoid or turbinate.	Broadly ovoid.	Obovoid, always asymmetric, usually flattened adaxially.
6. Leaf margin	Entire throughout.	Entire throughout.	Entire throughout.	Entire or serrate in the upper half.
7. Scar position, shape and size	Orbicular, covering $\geq 95\%$ of the fruit, from the base upward.	Basal, slightly concave, ca. 1 cm in diameter.	Basal, concave, ca. 1.5 cm in diameter.	- (nut fused to wall).
8. Nut indumentum	Glabrous.	Greyish pubescent.	Sparsely sericeous then dull brown.	- (nut fused to wall).

This species is endemic to Thailand and is currently only known from one location in Ton Pariwat Wildlife Sanctuary. The sanctuary covers a region of low-lying forested mountains with a total area of approximately 100,000 ha at the southern end of the Phuket mountain range. As such it is an integrated part of the Southern Forest Complex of Thailand. The sanctuary is popular for its rich bird- and wildlife (e.g. Blue-banded Kingfisher (*Alcedo euryzona* Temminck, Alcedinidae) and Whitehanded Gibbons (*Hylobates lar* L., Hylobatidae) as well as rare flora, such as *Rafflesia kerrii* Meijer (Rafflesiaceae). Its unique species composition, high diversity and relatively intact forest structure underscore the importance of strengthening ongoing and future conservation measures at Ton Pariwat Wildlife Sanctuary, as a key element of wider conservation efforts in southern Thailand.

Updated key for the species of *Lithocarpus* occurring in Thailand

Following the treatment of Fagaceae for the Flora of Thailand (Phengklai 2008), no further updates have been published. In the updated key we include here, we incorporate the identification of *L. orbicarpus* and add several additional corrections.

Key to the thai species of *Lithocarpus*

(based on vegetative characters and acorns)

- 1 Outer surface of cupules with annular or lamellate markings or markings lacking
- 2 Cupules without lamellae, chartaceous or subcoriaceous, enclosing nearly all of the nut, more or less dehiscent when mature
- 3 Cupules weakly dehiscent from the apex, cupule surface distinctly undulate with vertical and horizontal lines
- 4 Cupule urn-shaped
- 5 Cupule base broadly conical, much broader than apex, skin distinct with many vertical filiform lines or without. Nut conical.....**5. *L. blumeanus***
- 5 Cupule base obconic, much narrower than apex, surface distinct with 3-4 horizontal filiform lines. Nut obconical..... **33. *L. maingayi***
- 4 Cupule top or globe shaped
- 6 Cupule top-shaped, enclosing 4/5 of nut, surface with 2-6 distinct horizontal, filiform lines.....**30. *L. macphailii***
- 6 Cupule globe-shaped, enclosing nut completely, except for a tiny section at the apex, surface with distinct irregularly placed scales along 5-9 horizontal and vertical lines **36. *L. orbicarpus***
- 3 Cupules readily dehiscent into irregular parts from the top, surface with 2-5 filiform, undulate, horizontal lines
- 7 Cupules with 2 or 3 such lines **18. *L. encleisocarpus***
- 7 Cupules with 4 or 5 such lines **37. *L. pattaniensis***
- 2 Cupules with distinct lamellae, coriaceous, enclosing a variable amount of the nut, indehiscent
- 8 Cupule enclosing not less than 1/2 of the nut
- 9 Cupule enclosing about 1/2 of the nut
- 10 Nuts ovoid to conical at apex, scar shallowly concave or flattened.....
.....**24. *L. gracilis***
- 10 Nuts subhemispheric or depressed at apex, scar deeply concave
..... **8. *L. clementianus***
- 9 Cupule enclosing not less than 3/4 of the nut
- 11 Cupules obconic, enclosing nut almost completely except around the umbonate apex
- 12 Nut longer than broad, ca. 1 by 0.7 cm **26. *L. hendersonianus***
- 12 Nut shorter than broad, 1-2.7 by 2-3 cm **32. *L. magnificus***
- 11 Cupules saucer-shaped, enclosing ca. 3/4 of the nut.....**1. *L. aggregatus***
- 8 Cupule enclosing not more than 1/4 of the nut
- 13 Nuts hemispheric or depressed on both sides
- 14 Cupule enclosing 1/5 to 1/4 of the nut **39. *L. platycarpus***
- 14 Cupule enclosed only the base of the nut

- 15 Acorns sessile. Scar deeply concave.....**15. *L. eichleri***
- 15 Acorns with stalk up to 0.5 cm long. Scar slightly concave.....**6. *L. cantleyanus***
- 13 Nuts conical to broadly ovoid, or with a dome-shaped apex
- 16 Cupule enclosing only the base of the nut
- 17 Acorns sessile. Leaves oblanceolate.....**29. *L. lucidus***
- 17 Acorns with fruit-stalk up to 0.5 cm long. Leaves oblong.....**43. *L. reinwardtii***
- 16 Cupule enclosing ca. 1/4 of the nut
- 18 Nut with one horizontal ring around equator. Leaves ensiform to linearlan-
ceolate**28. *L. loratefolius***
- 18 Nut without horizontal ring. Leaves ovate, ovate-oblong or narrowly elliptical
- 19 Nut ovoid or conical. Cupules cup or saucer-shaped. Leaves ovate or ovate-
oblong, apex caudate.....**3. *L. bancanus***
- 19 Nut broadly ovoid. Cupules slightly obconical to saucer-shaped. Leaves nar-
rowly elliptical**41. *L. rassa***
- 1 Outer surface of cupules with alternate lamellae (resembling fish scales) or
pseudospines
- 20 Mature cupules of one infructescence more or less fused together
- 21 Acorns broader than long, depressed both on top and at base. Cupules saucer-
or cupshaped or obconic, some hardly distinct from each other through fusion
- 22 Infructescences with densely arranged cupules
- 23 Cupules barely distinct, resembling a large gall**13. *L. echinophorus***
- 23 Cupules distinct, saucer-shaped
- 24 Nut flattened or apiculate at apex, to 2.2 cm diam. Leaves cuneate at base ...
.....**16. *L. elegans***
- 24 Nut retuse at apex, not less than 3 cm diam. Leaves auriculate at base
.....**2. *L. auriculatus***
- 22 Infructescences with spaces between cupules
- 25 Rachis of infructescence always with sub-branches. Acorns stalked
.....**34. *L. mekongensis***
- 25 Rachis of infructescence without sub-branches
- 26 Acorns sessile**24. *L. finetii***
- 26 Acorns stalked.....**50. *L. tenuinervis****
- 21 Acorns longer than broad, conical, ovoid or turbinate. Cupules cup-shaped
or cylindric
- 27 Rachis of infructescence always with sub-branches. Acorns stalked, nuts shining
- 28 Acorn up to 1 cm high. Rachis up to 4 mm in diam.....**7. *L. ceriferus***
- 28 Acorn not less than 1 cm high (to 2.5 cm). Rachis not less than 4 mm in
diam**40. *L. polystachyus***
- 27 Rachis of infructescence without sub-branches. Acorns sessile, nuts more or
less shining
- 29 Twigs glabrous or sparsely pubescent then glabrous
- 30 Cupules cup-shaped, enclosing up to 1/2 of the nut**12. *L. dealbatus***

- 30 Cupules turbinate, enclosing the whole nut, open only around umbo
.....**53. *L. truncatus***
- 29 Twigs ferruginous or tomentose
- 31 Leaves glabrous except along midrib. Cupules enclosing up to 1/3 of the
nut.....**25. *L. harmandianus***
- 31 Leaves densely tomentose especially on lower surface. Cupules enclosing 1/2
of the nut.....**27. *L. lindleyanus***
- 20 Mature cupules of one infructescence, free, not fused
- 32 Acorn longer than broad, conical, ovoid or obconical. Cupules cup- or sau-
cer-shaped or obconic
- 33 Cupules enclosing nut completely or 2/3 of the nut
- 34 Cupules enclosing ca. 2/3 of the nut
- 35 Cupules slightly obconical-shaped, nuts hairy at style apex (if persistent)
.....**45. *L. rufescens***
- 35 Cupules cup or saucer-shaped.....**16. *L. elegans***
- 34 Cupules enclosing nut completely, or up to the apex of the nut
- 36 Cupules dehiscent, obconic or ovoid
- 37 Cupules obovoid, sessile, surface with dense, long and narrow recurved pseu-
dospines.....**42. *L. recurvatus***
- 37 Cupules ovoid, fruit stalk 2–3 mm long, surface finely ornamented with thin,
triangular lamellae throughout.....**35. *L. neo-robinsonii***
- 36 Cupules indehiscent, ovoid, surface clothed with dense, triangular lamellae
- 38 Infructescences up to 18 cm long. Leaves up to 16 cm long.....**9. *L. craibianus***
- 38 Infructescences not less than 20 cm long. Leaves not less than 20 cm long ...
.....**19. *L. erythrocarpus***
- 33 Cupules enclosing up to 1/2 of the nut
- 39 Acorns stalked
- 40 Cupules slightly obconic. Leaves ovate, ovate-oblong or obovate
.....**48. *L. sootepensis***
- 40 Cupules cup-shaped or saucer-shaped
- 41 Cupules cup-shaped. Leaves lanceolate to lanceolate oblong.....**47. *L. siamensis***
- 41 Cupules saucer-shaped to flattened. Leaves oblong to oblong-lanceolate.....
.....**10. *L. curtissii***
- 39 Acorns sessile
- 42 Acorns (mature) not less than 3.5 by 2.2 cm
- 43 Cupule lamellae bearing pseudo-spined reflexed towards the base. Leaves
acute to obtuse at apex.....**46. *L. scortechinii***
- 43 Lamellae curved towards the cupule apex. Leaves acuminate at apex.....
.....**20. *L. eucalyptifolius***
- 42 Acorns (mature) up to 3 by 2.2 cm
- 44 Infructescence with acorns in clusters, but not fused
- 45 Nuts ovoid. Leaves usually curved to one side.....**54. *L. wallichianus***

- 45 Nuts strongly apically depressed, occasionally conic. Leaves not curved
 **51. *L. thomsonii***
- 44 Infructescence with acorns solitary, with spaces between them
- 46 Cupules saucer or cup-shaped, limb recurved. Leaves not less than 12 cm
 long **21. *L. falconeri***
- 46 Cupules obconical, limb not recurved. Leaves up to 11 cm long
 **4. *L. bennettii***
- 32 Acorns broader than long, hemisphaeric-depressed
- 47 Cupules enclosing the nut completely or up to the apex of the nut
- 48 Cupules more or less up to the apex of the nut, lamellae with erect or reflexed
 pseudospines which are not fused
- 49 Pseudo-spines erect or spreading. Leaves oblanceolate. Scar nearly 1/2 of the
 nut **14. *L. echinops***
- 49 Pseudo-spines reflexed. Leaves oblong or oblanceolate
- 50 Infructescence with acorns packed close together, but not fused. Leaves
 slightly cuneate at base **23. *L. garrettianus***
- 50 Infructescence with acorns solitary, with spaces between them. Leaves obtuse
 at base **54. *L. tubulosus***
- 48 Cupules enclosing the nut completely, except the umbo
- 51 Lamellae pointed, with narrowly pseudospines. Infructescence with acorns
 packed close together, but not fused **57. *L. wrayi***
- 51 Lamellae flattened and imbricate. Infructescence with acorns solitary, with
 spaces between them
- 52 Lamellae fused on lower half, the upper half free and adaxially curved
 **22. *L. fenestratus***
- 52 Lamentas fused almost to apex, only a short free lobe adaxially curved
 **52. *L. trachycarpus***
- 47 Cupules enclosing up to 1/2 of the nut
- 53 Acorns stalked, cupules enclosing only base of the nut
- 54 Stalk up to 1 cm long. Leaves glaucous on lower surface, petiole up to 1 cm
 long **49. *L. sundaicus***
- 54 Stalk not less than 1 cm long. Leaves pale on lower surface, not glaucous,
 petiole not less than 1 cm long **31. *L. magneinii***
- 53 Acorns sessile, cupules enclosing up to 1/2 of the nut
- 55 Acorns not less than 2 by 2.5 cm
- 56 Cupules slightly obconical. Leaves oblong, acute to caudate at apex, margin
 not revolute, petiole not less than 1 cm **11. *L. cyclophorus***
- 56 Cupules saucer-shaped. Leaves obovate, obtuse at apex, margin revolute, pet-
 iole up to 0.6 cm long **44. *L. revolutus***
- 55 Acorns up to 1.5 by 2 cm
- 57 Nuts convex at the apex
- 58 Cupules saucer-shaped to flattened and discoid. Leaves not whorled
- 59 Lamellae usually fused throughout. Leaves up to 15 cm long... **38. *L. pierrei***

- 59 Lamellae fused at base only, apices free. Leaves not less than 18 cm.....
long **17. *L. elephantum***
- 58 Cupules cup-shaped. Leaves usually whorled at the twig tips... **58. *L. xylocarpus***
- 57 Nuts flattened at the apex. Cupule cup-shaped, enclosing 1/5 to 1/2 of the
nut. Leaves with unequal sides, usually curved to one side
- 60 Leaves oblong, elliptic oblong, not less than 10 by 3.5 cm, with 14–20 pairs
of lateral nerves..... **55. *L. vestitus***

Acknowledgements

We are grateful to staff of the Ton Pariwat Wildlife Sanctuary for assistance in the field. We thank R. Pooma and staff of the Forest Herbarium (BKF) for access to collections. Finally, we would like to acknowledge S. Suddee and A. Sinbumroong for advice in identification and the preparation of this manuscript. The first author was supported by a Postdoctoral Research Fellowship of the Chinese Academy of Sciences (2011–2014).

References

- Barnett EC (1940) A survey of the genus *Quercus* and related genera of the Fagaceae in Asia with a more detailed account of the Siamese species of these genera and notes on the use of leaf anatomy in taxonomy. D.Sc. Thesis, University of Aberdeen, Scotland.
- Bunpha K, Masuthon S, Phengkklai C (2011) *Quercus thorelii* Hickel & A. Camus (Fagaceae), a new record for Thailand. Thai Forest Bulletin (Botany) 39: 206–209. http://web3.dnp.go.th/botany/pdf/TFB/TFB9/TFB39_12_Quercus.pdf
- Camus A (1948) Les chênes. Monographie des genres *Quercus* et *Lithocarpus*. Atlas, vol.3. Encyclopédie Économique de Sylviculture 7: 152–165.
- Camus A (1952–54) Les chênes. Monographie des genres *Quercus* et *Lithocarpus* 8. 511–1196.
- Cannon CH (2001) Morphological and molecular diversity in *Lithocarpus* (Fagaceae) of Mount Kinabalu. Sabah Parks Nature Journal 4: 45–69. doi: 10.1080/106351501753462849
- Cannon CH, Manos PS (2001) Combining and comparing morphometric shape descriptors with a molecular phylogeny: the case of fruit type evolution in Bornean *Lithocarpus* (Fagaceae). Systematic Biology 50(6): 860–880.
- Hughes JB, Round PD, Woodruff DS (2003) The Indochinese-Sundaic faunal transition at the Isthmus of Kra: an analysis of resident forest bird species distributions. Journal of Biogeography, 30(4): 569–580. doi: 10.1046/j.1365-2699.2003.00847.x
- Manos Paul S, Cannon CH, Oh S-H (2008) Phylogenetic relationships and taxonomic status of the paleoendemic Fagaceae of western North America: recognition of a new genus, *Notolithocarpus*. Madrono 55.3: 181–190. doi: 10.3120/0024-9637-55.3.181
- Meijaard E (2009) Solving mammalian riddles along the Indochinese-Sundaic zoogeographic transition: new insights from mammalian biogeography. Journal of Biogeography, 36(5):801–802. doi: 10.1111/j.1365-2699.2009.02124.x

- Parnell J (2013) The biogeography of the Isthmus of Kra: a review. *Nordic Journal of Botany* 31: 001–015.
- Phengklai C (2004) Three new species and a new variety of Fagaceae from Thailand. *Thai Forest Bulletin (Botany)* 32: 115–122. http://web3.dnp.go.th/botany/pdf/TFB/TFB32/TFB32_9Fagaceae.pdf
- Phengklai C (2008) Fagaceae. Vol.9 (3). In: Santisuk T, Larsen K, Nielsen I, Chayamarit K, Phengklai C, Pedersen H, Parnell J, Middleton D, Newman M, Simpson DA, van Welzen PC, Hul S, Kato M (Eds) *Flora of Thailand*. The Forest Herbarium, National Parks, Wildlife and Conservation Department, Bangkok.
- Soepadmo E (1972) Fagaceae. *Flora Malesiana Series I, Volume 7* (2). Rijksherbarium /Hortus Botanicus, Leiden University, Leiden, The Netherlands.
- Van Steenis C (1950) The delimitation of Malaysia and its main plant geographical divisions. *Flora Malesiana Series 1*.
- Woodruff DS (2003) The location of the Indochinese-Sundaic biogeographic transition in plants and birds. *Natural History Bulletin of the Siam Society* 51: 97–108.
- Woodruff DS, Turner LM (2009) The Indochinese-Sundaic zoogeographic transition: a description and analysis of terrestrial mammal species distributions. *Journal of Biogeography*, 36(5): 803–821. doi: 10.1111/j.1365-2699.2008.02071.x
- Wu CY, Raven PH, Hong DY (1999) (Eds) *Flora of China (Cycadaceae through Fagaceae)* Vol. 4, Beijing & St. Louis, Science Press & Missouri Botanical Garden Press. <http://flora.huh.harvard.edu/china/mss/volume04/FAGACEAE.published.pdf>