

Primulina pingnanensis, a new species of Gesneriaceae from Guangxi, China

Zheng-Long Li^{1,2,3}, Yan-Yun Kuang^{1,4}, Qing-Qing Xu¹, Wei-Chuen Chou², Xin Hong^{1,3,5}, Li Ding^{5,6}

1 Anhui Provincial Engineering Laboratory of Wetland Ecosystem Protection and Restoration, School of Resources and Environmental Engineering, Anhui University, CN-230601, Hefei City, Anhui Province, China

2 Gesneriad Conservation Center of China (GCCC), Guilin Botanical Garden, CAS, Guilin 541006, China

3 Yunnan Key Laboratory for Integrative Conservation of Plant Species with Extremely Small Populations, Kunming Institute of Botany, Chinese Academy of Sciences, CN-650201 Kunming, Yunnan, China

4 State Key Laboratory of Soil and Sustainable Agriculture, Institute of Soil Science, Chinese Academy of Sciences, CN-210008, Nanjing, China

5 Gesneriad Committee of China Wild Plant Conservation Association, National Gesneriaceae Germplasm Resources Bank of GXIB, CN-541006 Guilin, Guangxi, China

6 Guangxi Key Laboratory of Plant Conservation and Restoration Ecology in Karst Terrain, Guangxi Institute of Botany, Guangxi Zhuang Autonomous Region and Chinese Academy of Sciences, CN-541006, Guilin City, Guangxi Zhuang Autonomous Region, China

Corresponding authors: Xin Hong (hongxin1989@vip.qq.com); Li Ding (dingli@gxib.cn)

Abstract

A new species of *Primulina*, *P. pingnanensis*, from the Guangxi Zhuangzu Autonomous Region, China, is described and illustrated here. It is morphologically similar to *P. orthandra* but has significant differences in the bracts, corolla tube and lobes shape, as well as in the indumentum of the outer surface of the corolla, the filaments, the staminodes and the anthers. Colorful photographs and essential information of this new taxon are also provided, including detailed taxonomic description, distribution, habitat, the comparison table, and the IUCN conservation status. We also discuss a validation of new combination *P. crassifolia* and *Chirita crassifolia*.

Key words: Flora of Guangxi, Gesneriaceae, limestone flora, new taxon, *Primulina orthandra*, taxonomy



Academic editor: Laura Clavijo

Received: 19 March 2023

Accepted: 6 July 2023

Published: 27 July 2023

Citation: Li Z-L, Kuang Y-Y, Xu Q-Q, Chou W-C, Hong X, Ding L (2023)

Primulina pingnanensis, a new species of Gesneriaceae from Guangxi, China. PhytoKeys 229: 157–165. <https://doi.org/10.3897/phytokeys.229.103735>

Copyright: © Zheng-Long Li et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

As of December 2022, about 250 taxa (including varieties) of *Primulina* Hance had been published worldwide, with 215 species and 18 varieties distributed in China and 18 species in Vietnam (GRC 2022; Wei et al. 2022). Based on the quantity and distribution of the *Primulina* species, it is apparent that the Karst limestone areas in Guangxi, China, are the center of *Primulina* diversity and distribution (Hao et al. 2014; Kong et al. 2017; Wei 2018; Li et al. 2019; Xu et al. 2019). This further indicates that more undescribed *Primulina* species may still be found in the Karst ecosystem of South and Southwest China (Möller 2019), especially in karst caves (Xin et al. 2018; Fu et al. 2022), cliffs and gorges.

Averyanov et al. (2020) described a new species of *Chirita* in North Vietnam, which they named *Chirita crassifolia* Aver. & K.S.Nguyen. In 2022, after verification, Möller and collaborators transferred *C. crassifolia* to *Primulina* as *P. crassifolia* (Anh et al. 2022). *Primulina crassifolia* (Aver. & K.S.Nguyen) T.T.P.Anh, F. Wen &

Mich.Möller, comb. nov. Basionym: *Chirita crassifolia* Aver. & K.S.Nguyen, Pl. Diversity Fl. Veg. Bat Dai Son 254 (2021). We emphatically revise the Latin name from '*Chirita crassifolia*' to '*Primulina crassifolia*' here and provide appropriate notes.

In March 2018, when we were on an expedition in Pingnan County, Guangxi, China, we found an unknown *Primulina*-like plant growing at the crevices of rocks on the top of a limestone cliff. According to its leaf arrangement (rosette), fleshy leaves, tubular corolla, and cracked fruits (capsule with parietal placentation), it should be included in *Primulina*. However, it did not match any known species of this genus. In addition, this species distinguishes from the genus *Petrocodon* by its lingulate stigma, curved filaments, and fleshy leaves. We collected specimens in the field and introduced some living plants into cultivation at the nursery of the National Gesneriaceae Germplasm Resources Bank of GXIB and the Gesneriad Conservation Center of China. The morphology of the cultivated plants has not changed from that of the wild plants during the past three years. We compared the collected specimens and living plants with other known type specimens and living plants of *Primulina*, and we found that it is distinctly different from other species of *Primulina*, especially those distributed in Guangxi and recently published species of *Primulina* (e.g., Du et al. 2021; 2022; Xiong et al. 2022; Pan et al. 2022), which are also different from this new taxon. We then confirmed that it was a new species.

Material and method

The description is based on specimens collected in the wild. The measurements and morphological characteristics of the new species were taken from the type specimens processed by the authors. The type specimens of the new species were deposited in IBK. We examined *Primulina* specimens in IBSC, K, IBK, KUN, US, PE, and VNMN. We also searched two online herbaria, Chinese Virtual Herbarium (CVH, <http://www.cvh.ac.cn>) and Chinese Field Herbarium (CFH, <http://www.cfh.ac.cn>), to find species with similar morphology. We determined it to be an undescribed taxon. All morphological characters were studied under a dissecting microscope and were described using the terminology used by Wang et al. (1990, 1998).

Taxonomic treatment

Primulina pingnanensis Xin Hong, Z.L.Li & W.C.Chou., sp.nov.

urn:lsid:ipni.org:names:77324168-1

Fig. 1

Diagnosis. *Primulina pingnanensis* morphologically resembles *P. orthandra* but is distinguished from the latter by bracts lanceolate (vs. ovate), corolla tube funnel-form and no constriction in the middle (vs. tube near tubular with constriction in the middle), outer corolla surface sparsely white puberulent (vs. glabrous), corolla lobes oblong (vs. broadly ovate), filaments strongly curved at the middle (vs. straight), anthers fused from the entire adaxial surface and sparsely barbate (vs. confluent at apex, glabrous), staminodes obvious, 1–1.3 cm long, sparsely pubescent (vs.ca. 1.5 mm long, glabrous). Detailed morphological comparisons with *P. orthandra* are provided in Table 1 and Fig. 2.




Figure 1. *Primulina pingnanensis* Xin Hong, Z.L.Li & W.C.Chou **A** plants in natural habitat **B** cymes **C** opened corolla and dissected calyx lobes with pistil **D** leaves (up: adaxial surface, down: abaxial surface) **E** young fruit **F** bracts **G** stigmas **H** stamens.

Type. CHINA. Guangxi Zhuangzu Autonomous Region: Guigang City, Pingnan County, Shuangma Village, 23°37'N, 110°19'E, growing atop a cliff on a limestone hill, 28 March 2018, Chou Wei-Chuen, CWC180328-01(holotype: IBK!; isotypes: AHU!, IBK!).

Description. Perennial herb. Rhizomatous stem suberete. **Leaves** basal, opposite, and congested at rhizome apex after years of growth; petiole 5–8 cm long, densely pubescent. Leaf blade dark green, thick herbaceous, elliptic to ovate-elliptic, 15–20 × 6–8 cm, apex acute to obtuse, base broadly cuneate, slightly oblique,

Table 1. Diagnostic character differences between *Primulina pingnanensis* and *P. orthandra*.

Character	<i>Primulina pingnanensis</i>	<i>P. orthandra</i>
Bracts	lanceolate	ovate
Tube	funnel-form, no constriction in the middle	tube near tubular, constriction in the middle
Outer corolla surface	sparsely white puberulent	glabrous
Corolla lobes	lobes oblong	lobes broadly ovate
Filaments	strongly bent at the middle	straight
Anthers	fused by the entire adaxial surface and sparsely barbate	confluent at apex, glabrous
Staminodes	obvious, 1–1.3 cm long, sparsely pubescent	ca. 1.5 mm long, glabrous

margin sinuate, densely pubescent on both surfaces; 3–4 pairs of lateral veins on each side of the midrib; adaxially midrib main vein conspicuously sunken and lateral veins inconspicuously sunken, abaxially midrib main vein and lateral vein conspicuously raised. **Cymes** 1–7, axillary, 1–3-branched, 1–8(–16)-flowered; peduncles 18–25 cm long, densely brown puberulent and pubescent; bracts 2, opposite, lanceolate, ca. 7 × 2.5 mm, densely brown pubescent on both surfaces. Pedicels 2.5–4 cm long, densely brown puberulent, and pubescent. Calyx 5-lobed from the base, lobes narrowly lanceolate to triangle, apex acute, ca. 4 × 0.9 mm, outside pubescent, inside glabrous. **Corolla** zygomorphic, lilac, –4.5 cm long, outside sparsely white puberulent, inside glabrous; tube funnel-form, 2.0–3.0 cm long, 1.0–1.3 cm in diameter at the mouth, ca. 5 mm in diameter at the base; limb distinctly 2-lipped; dorsal lip 2-lobed to the middle, lobes ca. 1.0 cm long, apex suborbicular; ventral lip 3-lobed to the middle, lobes oblong, 1.0–1.5 cm long. **Stamens** 2, glabrous, adnate to the corolla tube for ca. 10 mm above the base; filaments 8–10 mm long, extending outwards at the base of the corolla, strongly curved at 90° degrees at the middle, sparsely white pubescent at base; anthers white, ca. 2 mm long, fused by the entire adaxial surface, sparsely barbate; Staminode 3, lateral ones obvious, linear, sparsely pubescent, 1–1.3 cm long, adnate to the corolla for 8 mm above the base; the middle one capitate, ca. 1 mm long, adnate at the corolla base. Disc annular, slightly oblique, ca. 1.5 mm in height, margin repand. **Pistil** 2.8–3.2 cm long; ovary ca. 1.8 cm long, densely white pubescent; style ca. 1 cm long, densely puberulent; stigma trapeziform, ca. 2 mm long, apex slightly 2-lobed, lobes obtuse triangle, ca. 0.4 mm long. **Capsule** 4.5–6 cm long, sparsely brown puberulent, narrowly oblong-ovoid, dehiscing loculicidally into two valves.

Phenology. Flowering from the second half of March to the first half of April, fruiting from June to July.

Etymology. The specific epithet is derived from the type locality, Pingnan County. This county is the birthplace of Mr. Chou's mother. Thus, Mr. Chou strongly suggested using "*pingnanensis*" as the scientific name.

Vernacular name. Píng Nán Bào Chūn Jù Tái (Chinese pronunciation); 平南报春苣苔 (Chinese name).

Distribution and habitat. *Primulina pingnanensis* is only known from the type locality, Shuangma Village, Pingnan County, Guigang City, Guangxi Zhuangzu Autonomous Region, growing on limestone cliffs at an elevation of ca. 50 m. The average temperature in the distribution area is 22.6 °C, while the average annual precipitation is 1050–2100 mm. The habitat is very close to the village with human activities.




Figure 2. Morphological comparisons between *Primulina pingnanensis* and *P. orthandra* **A** habit of *P. pingnanensis* **B** habit of *P. orthandra* **C** lateral view of mature flower of *P. pingnanensis* **D** lateral view of mature flower of *P. orthandra* **E** staminodes of *P. pingnanensis* **F** staminodes of *P. orthandra* **G** filaments of *P. pingnanensis* **H** filaments of *P. orthandra* **I** bracts of *P. pingnanensis* **J** bracts of *P. orthandra*.

Preliminary conservation assessment. The terrain in the North and South of Pingnan County is mountainous, while the central area is flat. Very few limestone hills are in the plains, but some mountains are near Shuangma Village. After finding the new species, we conducted several detailed explorations of the area. The current survey results showed that this species has a small population at the top of a limestone mountain near Shuangma Village, with fewer than 200 individuals. According to the results of our field investigations in the type locality and adjacent regions, the EOO and AOO of *Primulina pingnanensis* are about 2 km² and 0.1 km², respectively. The severe drought in the second half of 2022 has seriously affected the population. According to the field survey, it is preliminarily estimated that the number of individuals in this population has decreased by 40% or more. More in-depth habitat surveys are warranted to determine if there are more populations nearby. For this current study, we temporarily assess the status of this species as Critically Endangered [CR B1+B2ab (iii, v)], according to the IUCN Red List Categories and Criteria (IUCN 2022).

Notes

Validation of new combination in *Primulina*

When the new combination for *Primulina crassifolia* was made by Anh et al. (2022), it was later noticed that the combination was invalid. Therefore, we validate it here.

***Primulina crassifolia* (Aver. & K.S.Nguyen) T.T.P.Anh, F. Wen & Mich.Möller, comb. nov.**

[urn:lsid:ipni.org:names:77324169-1](https://urn.nbn.se/resolve?urn=urn:nbn:se:isid:ipni.org:names:77324169-1)

Chirita crassifolia Aver. & K.S.Nguyen, Pl. Diversity Fl. Veg. Bat Dai Son 254 (2021). Basionym.

Type. Vietnam, Ha Giang Province: Quan Ba district, Tung Vai commune, steep rocky slopes to large and deep cave composed of highly eroded marble-like white limestone, elevation 900–1000 m a.s.l., 17 October 2018, L. Averyanov, Nguyen Sinh Khang, T. Maisak, Truong Duc Thieu, VR 938 (holotype: LE!; isotypes: HN!, LE!).

Discussion

The Karst and Danxia landforms from south & southwest China to the north Indo-China Peninsula are well-known for their high species diversity and endemism levels. Some interesting genera of Gesneriaceae for example, *Primulina*, *Petrocodon* Hance, and *Hemiboea* Clarke have been attracting the attention of botanists and taxonomists in the past decades. *Primulina* speciation is positively associated with changes in past temperatures and East Asian monsoons. Climatic change around the mid-Miocene triggered an early burst (Kong et al. 2017; Xu et al. 2019; Hsieh et al. 2022). There is abundant diversity in the morphology of the vegetative and reproductive organs of *Primulina*. Still, it can be distinguished by its rosette leaves, tubular corolla, parietal placentation, etc. We concluded

that *P. pingnanensis* is an undescribed species of *Primulina* based on extensive investigation in south and southwest China. A comparison of known live plants and specimens of the genus *Primulina*. *P. orthandra* can be regarded as similar to *P. pingnanensis* because they both have similar leaf morphology, calyx, and purple corolla. However, the differing characteristics of the peduncle, bract, stamen, and staminodes can quickly distinguish the two species from each other.

The high species diversity and restricted distributions on limestone habitats have made the calciphilous *Primulina* an ideal study subject for understanding plant radiation on Sino-Vietnamese limestone karsts (Wei et al. 2022). The habitat of *P. pingnanensis* is located in the mountains in the middle of Pingnan County. The rest of the area is a flat plain. The spatial heterogeneity has created a unique species of *Primulina* in Pingnan. This suggests that the karst areas in southern China have acted as both “museums” and “cradles” of plant evolution (Xu et al. 2019). The discovery and publication of this new species supplements the diversity of *Primulina* distributed in Guangxi. It further indicates that there are more undiscovered new species of *Primulina* in the vast karst areas of south China (Hong et al. 2020; Fu et al. 2022).

Acknowledgments

We thank Stephen Maciejewski, the Gesneriad Society, and Michael LoFurno, Associate Professor, Temple University, Philadelphia, USA, for their editorial assistance. We are also grateful to Michael Möller, Royal Botanic Garden Edinburgh, United Kingdom, for the notes on *Primulina crassifolia*.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study was financially supported by the Fund of Yunnan Key Laboratory for Integrative Conservation of Plant Species with Extremely Small Populations (PSESP2021F07).

Author contributions

WC, ZL and LD collected this species. ZL, YK, and QX performed the data analysis. ZL and XH wrote the manuscript. LD and XH revised the manuscript. All authors have read and approved the final manuscript.

Author ORCIDs

- Zheng-Long Li  <https://orcid.org/0009-0009-8988-5725>
Yan-Yun Kuang  <https://orcid.org/0009-0001-4964-3969>
Qing-Qing Xu  <https://orcid.org/0009-0007-8725-9291>
Wei-Chuen Chou  <https://orcid.org/0000-0001-6470-9295>
Xin Hong  <https://orcid.org/0009-0003-9477-2414>
Li Ding  <https://orcid.org/0009-0003-1523-1784>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Anh TT, Averyanov LV, Möller M, Maisak TV, Nguyen KS, Quang BH, Do TN, Li RF, Wen F (2022) One transfer to *Primulina* (Gesneriaceae) and amended descriptions for *P. crassifolia* and *P. quanbaensis* from northern Vietnam. Nordic Journal of Botany 2022(5): e03455. <https://doi.org/10.1111/njb.03455>
- Averyanov LV, Nguyen SK, Tran TH, Averyanova AL, Maisak VT, Nguyen HT (2020) Plant Diversity, Flora and Vegetation of Bat Dai Son Mountain Area, Northern Vietnam. Saint-Petersburg, Strata, 561 pp.
- Du C, Liu J, Ye W, Liao S, Ge BJ, Liu B, Ma JS (2021) Annual report of new taxa and new names for Chinese plants in 2020. Shengwu Duoyangxing 29(8): 1011–1020. <https://doi.org/10.17520/biods.2021122>
- Du C, Liu J, Ye W, Liao S (2022) 2021 annual report on new taxa and nomenclatural changes of Chinese plants. Shengwu Duoyangxing 30(8): 22207. <https://doi.org/10.17520/biods.2022207>
- Fu LF, Monro AK, Wei YG (2022) Cataloguing vascular plant diversity of karst caves in China. Shengwu Duoyangxing 30: e21537. <https://doi.org/10.17520/biods.2021537>
- GRC (2022) Gesneriaceae Resource Centre. Royal Botanic Garden Edinburgh. [continuously updated ; Retrieved/Accessed: 21 November 2022] <https://padme.rbge.org.uk/GRC>
- Hao Z, Kuang YW, Kang M (2014) Untangling the influence of phylogeny, soil and climate on leaf element concentrations in a biodiversity hotspot. Functional Ecology 29(2): 165–176. <https://doi.org/10.1111/1365-2435.12344>
- Hong X, Keene J, Shan WY, Do VT, Wen F (2020) Taxonomic identity of *Primulina swinglei* (Gesneriaceae). Guihaia 40(10): 1393–1401.
- Hsieh CL, Xu WB, Chung KF (2022) Plastomes of limestone karst gesneriad genera *Petrocodon* and *Primulina*, and the comparative plastid phylogenomics of Gesneriaceae. Scientific Reports 12(1): e15800. <https://doi.org/10.1038/s41598-022-19812-2>
- IUCN (Standards and petitions committee) (2022) Guidelines for Using the IUCN Red List Categories and Criteria. Version 15. Prepared by the Standards and petitions committee. <https://www.iucnredlist.org/resources/redlistguidelines> [accessed on 4 Mar. 2022]
- Kong HH, Condamine FL, Harris AJ, Chen JL, Pan B, Möller M, Hoang VS, Kang M (2017) Both temperature fluctuations and East Asian monsoons have driven plant diversification in the karst ecosystems from southern China. Molecular Ecology 26(22): 6414–6429. <https://doi.org/10.3897/phytokeys.127.35445>
- Li S, Xin ZB, Chou WC, Huang Y, Pan B, Maciejewski S, Wen F (2019) Five new species of the genus *Primulina* (Gesneriaceae) from Limestone Areas of Guangxi Zhuangzu Autonomous Region, China. PhytoKeys 127: 77–91. <https://doi.org/10.3897/phytokeys.127.35445>
- Möller M (2019) Species Discovery in Time: An Example from Gesneriaceae in China. Guangxi Sciences 26(1): 1–16.
- Pan B, Wang BM, Yang LH, Lai BD, Li PW (2022) *Primulina fangdingii* (Gesneriaceae), a new species from Guangxi, China. Nordic Journal of Botany 2022(10): e03684. <https://doi.org/10.1111/njb.03684>

- Wang WT, Pan KY, Li KY (1990) Gesneriaceae. In: Wang WT (Ed.) Flora Reipublicae Popularis Sinicae (Vol. 69). Science Press, Beijing, 141–271.
- Wang WT, Pan KY, Li ZY, Weitzman AL, Skog LE (1998) Gesneriaceae. In: Wu ZY, Raven PH (Eds) Flora of China (Vol. 18). Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 254–401.
- Wei YG (2018) The Distribution and Conservation Status of Native Plants in Guangxi, China. China Forestry Publishing House, Beijing, 876 pp.
- Wei YG, Do VT, Wen F (2022) A Checklist to the Plants of Northern Vietnam. China Forestry Publishing House, Beijing, 606 pp.
- Xin ZB, Li S, Zhang RL, Fu LF, Dong J, Wen F (2018) *Primulina zhoui* and *P. huangii* (Gesneriaceae), two new species from limestone areas in Guangxi, China. *Taiwania* 63(1): 54–60. <https://doi.org/10.6165/tai.2018.63.54>
- Xiong C, Chou WC, Huang Y, Wen F (2022) *Primulina nana* (Gesneriaceae), a new species from eastern Guangxi, China. *PhytoKeys* 197: 33–39. <https://doi.org/10.3897/phytokeys.197.83089>
- Xu M, Yang L, Kong H, Wen F, Kang M (2019) Congruent spatial patterns of species richness and phylogenetic diversity in karst flora: Case study of *Primulina* (Gesneriaceae). *Journal of Systematics and Evolution* 59(2): 251–261. <https://doi.org/10.1111/jse.12558>