

A new species of *Typhonium* (Araceae) from Vietnam

Hong Truong Luu^{1,2}, Nga Nguyen-Phi^{3,4}, Quoc Dat Nguyen¹, Hieu Cuong Nguyen¹,
Hong Thien Van⁵, Xuan Bach Nguyen-Le¹

- 1 Southern Institute of Ecology, Institute of Applied Materials Science, Vietnam Academy of Science and Technology, No. 1D, TL 29 Street, Thanh Loc Ward, District 12, Ho Chi Minh City, Vietnam
 - 2 Graduate University of Science and Technology, Vietnam Academy of Science and Technology, No. 18, Hoang Quoc Viet Street, Cau Giay District, Ho Chi Minh City, Vietnam
 - 3 Department of Ecology and Evolutionary Biology, University of Science, Vietnam National University HCMC, 227 Nguyen Van Cu Street, District 5, Ho Chi Minh City, Vietnam
 - 4 Vietnam National University – HCMC, Linh Trung Ward, Thu Duc City, Ho Chi Minh City, Vietnam
 - 5 Institute of Biotechnology and Food Technology, Industrial University of Ho Chi Minh City, No. 12 Nguyen Van Bao Street, Go Vap District, Ho Chi Minh City, Vietnam
- Corresponding author: Xuan Bach Nguyen-Le (nlxbach@gmail.com)

Abstract

Typhonium obtusum is described as a new species endemic to Vietnam. It is unique in the genus in having an oblong-elliptic spathe limb with an obtuse apex and yellowish-greenish filiform staminodes with a down-curved acumen. The ecology, distribution and assessment of the conservation status of the new taxon, as well as a key to all known *Typhonium* species in Vietnam, are provided.

Key words: Araceae, endemic, new species, *Typhonium obtusum*, Vietnam

Introduction

The genus *Typhonium* (Schott 1829) of the Araceae is estimated to have about 80 to 100 species distributed over the world (Sriboonma et al. 1994; Hettterscheid and Boyce 2000; Boyce et al. 2012; Hettterscheid 2013; Low et al. 2021; POWO 2024). A latest checklist of 70 accepted *Typhonium* species names is provided online (POWO 2024). Indochina was proved to be the centre of *Typhonium* diversity with about 40 species described (Low et al. 2021; Pham et al. 2023). The genus was revised several times for Vietnam (Gagnepain 1942a, b; Pham-Hoang 1993, 2003; Nguyen and Vu 2004; Nguyen 2005, 2017). In fact, the last three decades have witnessed many new discoveries which make the total number of *Typhonium* in the country to be 23 (Nguyen and Croat 1997; Hettterscheid and Boyce 2000; Hettterscheid and Nguyen 2001; Nguyen 2005, 2008; Nguyen and Croat 2010; Hettterscheid 2013; Luu et al. 2017; Van et al. 2017, 2021; Nguyen et al. 2021, 2022a, b; Nguyen-Phi et al. 2023; Pham et al. 2023; Serebryanyi et al. 2023).

As part of ongoing study of *Typhonium* in Vietnam, we have collected several putatively new taxa, one of which was found in Phu Yen Province, central Vietnam. At the first glance, the plant looks like *T. rhizomatosum* A.Galloway and P.Schmidt (Galloway 2012) and *T. cordifolium* S.Y.Hu (Hu 1968) as all of them share the following in common: the general appearance of leaves and spathe



Academic editor: Duilio Iamónico
Received: 19 September 2023
Accepted: 31 January 2024
Published: 16 February 2024

Citation: Luu HT, Nguyen-Phi N, Nguyen QD, Nguyen HC, Van HT, Nguyen-Le XB (2024) A new species of *Typhonium* (Araceae) from Vietnam. *PhytoKeys* 238: 119–126. <https://doi.org/10.3897/phytokeys.238.112973>

Copyright: © Hong Truong Luu et al.
This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

limbs and the structure of inflorescences (Nguyen et al. 2022b; Nguyen-Phi et al. 2023). However, after our careful examination of its morphological characteristics, it turns out that our plant is, indeed, a new species that we describe here, based on living collections.

Material and methods

The studied material was collected from Phu Yen Province, Central Vietnam. Specimens were sampled and processed using methods guided by the Royal Botanic Gardens, Kew (Bridson and Forman 1999); the herbarium acronyms follow Thiers (2024). Detailed photographs and description of taxonomically important characters of the new species were taken of fresh materials in the field using a digital camera. Taxonomic identification was based on morphological vegetative and reproductive characters following the aforementioned literature.

Taxonomic treatment

***Typhonium obtusum* Luu, X.B.Nguyen-Le & H.C.Nguyen, sp. nov.**

urn:lsid:ipni.org:names:77336459-1

Fig. 1

Type. VIETNAM. Phu Yen Province, Tay Hoa District, Hoa Thinh Ward; 12°54'30.1"N, 109°14'15.1"E, 30 m elevation; 5 August 2015; Nguyen Le Xuan Bach & Nguyen Hieu Cuong PY495-505 (holotype SGN!; isotypes SGN!, PHH!).

Diagnosis. *Typhonium obtusum* is morphologically similar to *T. rhizomatosum* and *T. cordifolium* in having ovate leaf blades with cordate base, oblong and elongate spathe limb as long as the spadix and an elongated fusiform spadix appendix. However, the novel taxon can be distinguished from *T. rhizomatosum* by its subcylindrical (vs. globose to subglobose) tubers without (vs. with) rhizomatous offsets, stipitate and fusiform (vs. sessile and conical) spadix appendix and oblong-elliptic (vs. narrowly triangular ovate) spathe limb and from *T. cordifolium* by its subcylindrical (vs. subglobose) tubers, leaves without (vs. with) adventitious buds, oblong-elliptic (vs. narrowly triangular ovate) spathe limb and stipitate and fusiform (vs. sessile and cylindrical) spadix appendix.

Description. Herbs, seasonally dormant, 10–17 cm tall; tuber underground, subcylindrical, fleshy, 2–2.5 cm long, 1–1.5 cm in diameter, with many filiform roots. **Leaves** 2–3 together. **Petiole** smooth, slender, 5–9 cm long, 1.8–2.2 mm in diameter, white to brown at base, green towards the apex. **Leaf blade** entire or trilobate, strongly cordate, glabrous, adaxially green, abaxially lighter green, concave, 3.2–4.7 cm long, 3–4 cm wide; entire leaf blade ovate, rounded at apex, venation pinnate with abaxially prominent midrib, lateral veins 5–8, brochiodromous, collective veins at 2–5 mm from margin; anterior lobes ovate, broadly elliptic to obovate, 2.5–3.7 cm long, 2.3–3.0 cm wide, obtuse to rounded at apex with a minute mucro, mid-rib abaxially prominent, lateral veins 4–6, brochiodromous, collective veins at 1–2 mm from margin; lateral lobes elliptic to ovate, 2.2–3.1 cm long, 0.6–1.6 cm wide, obtuse at apex, oblique at base. **Inflorescences** solitary; **peduncle** white, 2.5–6 cm long, ca. 4 mm in diameter; **spathe** 5–6 cm long, tube and limb separated by a constriction, outside green-brown, inside purple-brown; **spathe tube** ovoid, 1.1–1.3 cm long, 7–8 mm in diameter;




Figure 1. *Typhonium obtusum* **A** whole plants **B** stem **C** different forms of leaf blade **D** inflorescence **E** spathe and spadix **F** male, sterile and female zones **G** stamens **H** thecae, opened **I** pistils, cross-dissection **J** fruits.

spathe limb oblong-elliptic, 4–4.5 cm long, 1.1–1.3 cm wide, obtuse at apex, hardly open at base at anthesis; **spadix** equal or slightly shorter than spathe, sessile; **female part** conical, 1.6–2 mm long, 3.9–4.3 mm in diameter at the base, 2–3 mm in diameter at the top, with 3–4 rows of crowded pistils; **ovary** obovate, ca. 1 mm long, 0.6–0.8 mm in diameter, yellowish-greenish, with purple spots, unilocular, uniovulate placentation, transparently white; **ovule** light yellow, on a basal placenta and hold obliquely on a funicle; **stigma** sessile, disciform, 0.3–0.5 mm in diam., 0.1–0.2 mm high, light yellow, penicillate; **interstice** 1–1.3 cm long, lower part ca. 4 mm densely covered with staminodes, upper part naked, smooth, white to yellowish; **staminodes** filiform, subulate, 1–3 mm long, 0.4 mm in diameter, apically curved downward, yellowish-greenish; **male part** cylindrical, ca. 4.5 mm long, ca. 2.5 mm in diameter; **stamens** free, sparsely to (mostly) densely arranged; **thecae** globular, ca. 0.3 mm in diameter, dark purple, opening by apical slit; **pollens** translucent white; **appendix** stipitate, elongated fusiform, pale yellowish, 2.5–3.9 cm long, 2–3 mm in diameter, stipe ca. 3.5 mm long, pale yellow. **Fruits** ovoid to capsule-shaped, 3–4 mm long, 1.5–1.8 mm in diameter, white at base, green towards the apex with many dark purple spots.

Etymology. The species is named for the obtuse apex of its spathe limb.

Vernacular name. Bán hạ mo tù (Vietnamese); Obtuse-spathed typhonium (English, here proposed).

Table 1. Morphological differences between *Typhonium obtusum* and close species.

| | <i>T. obtusum</i> | <i>T. hayatae</i> | <i>T. inopinatum</i> | <i>T. medusae</i> | <i>T. varians</i> | <i>T. rhizomatosum</i> | <i>T. cordifolium</i> |
|------------------------|---|--|---|--|---|---|---|
| Tuber | subcylindrical, without rhizomatous offsets | globose, without rhizomatous offsets | subcylindrical, globose, without rhizomatous offsets | depressed, without rhizomatous offsets | depressed, without rhizomatous offsets | globose to subglobose, with rhizomatous offsets | subglobose, without rhizomatous offsets |
| Leaf blade | ovate, smooth | – | ovate to triangular or hastate, smooth | broadly triangular, 10 cm long, 14 cm wide, hairy | broadly triangular, more or less distinctly trilobate to subpentalobate, smooth | ovate to elliptical-ovate, smooth | narrowly ovate elliptic to narrowly elliptic, acuminate, with a bulbil when mature |
| Spadix appendix | 2.5–3.9 cm long, 0.2–0.3 cm in diameter, pale yellowish, fusiform, stipe ca. 0.35 cm long, base gradually narrowing | 6–14 cm long, 0.5–1.5 cm in diameter, conical, stipe short, base truncate | 4–6 cm long, 0.4–0.5 cm in diameter, yellow, yellowish-brown, elongate conical, subsessile, base truncate | 1 cm long, 0.3 cm in diameter, reddish-brown, elongate conical, stipe 0.15 cm long, base truncate | 4.5–9 cm long, to 1 cm in diameter, dark grey, elongate conical, stipe 0.5 cm long, base truncate | 8 cm long, 0.35 cm in diameter, beige, conical, sessile, base truncate | 3.6–7.7 cm long, 0.1–0.2 cm in diameter, brick orange, cylindrical, sessile, base gradually narrowing |
| Spathe tube | ovoid, 1.1–1.3 cm long, 0.7–0.8 cm in diameter, green with brown dots | oblong-ovoid 2.5–4.5 cm long, 1–2 cm wide, dark brownish-purple | ovoid, 0.8–1.5 cm long, ca. 1 cm in diameter, green | ovoid, 1.8–2 cm long, 1.2–1.5 cm in diameter, white with pinkish or purplish-brown flushing | ovoid, ca. 2.5 cm long, ca. 1.5 cm in diameter, glossy green | subglobose, 1 cm long, 0.8 cm in diameter, outside bright pale green, inside as outside, but with pale pink flush | ovoid, 1.2 cm long, 1 cm in diameter, light brown-green outside, brown or reddish-brown inside |
| Spathe limb | oblong-elliptic, 4–4.5 cm long, 1.1–1.3 cm wide, green with brown mottling | widely ovate in lower part, narrowly triangular in upper part, 12–26 cm long, 7.5–12 (–15) cm wide, dark brownish-purple | narrowly ovate to lanceolate, 5.5–7 cm long, 1.2–2 cm wide, basally brownish, apically green | orbicular, 2.7–3 cm long, 3 cm wide, whitish-greenish with a dense pinkish-brownish or brownish mottling | triangular ovate, 7–14 cm long, 4–7.5 cm wide, dark green flushed with dirty brown | narrowly triangular ovate, 7–9.5 cm long, 0.9–1.5 cm wide, bright pale green with brown longitudinal veins | narrowly triangular ovate, 5–9 cm long, to 3 cm wide, light brown-green |
| Staminode | slender filiform, decurved, yellowish-greenish | unknown | filiform, horizontally spread and slightly curved, yellow | subulate, upper ones straight, lower ones variously curved, mostly downwards, pale yellow or cream | subulate, upper ones straight, lower ones strongly curved downward, pale yellow | cylindrical, mostly perpendicular to axis, creamy white | cylindrical, spreading, dark yellow |
| Ovary | obovoid, yellowish-greenish | unknown | ellipsoid, yellowish-greenish | elongate, tapering to the base, basal half white, upper part spotted reddish-pink | more or less ellipsoid, pale green | elongate obovate, creamy white | elongate, white |
| Stigma | light yellow | unknown | yellow | reddish-pink | dirty whitish-greyish | creamy white | white |
| Distribution | C. Vietnam | S. Vietnam | N. and Central India to Thailand | C. Thailand | N. Thailand | Thailand | Thailand, Myanmar, S. Vietnam |

Ecological notes. The new species was found growing in clumps on basalt soils in open places of rural farms. It appears in September to December and becomes dormant in January to August. Flowering and fruiting were seen in August.

Distribution. *Typhonium obtusum* has been recorded only from the type locality.

Conservation status. Data Deficient (DD) (IUCN Standards and Petitions Subcommittee 2022). The new species has, so far, been found in one location and further inventory should be employed for a certain assessment.

Taxonomic notes. The key morphological similarities and differences of *T. obtusum* versus *T. rhizomatosum* and *T. cordifolium* are presented in the diagnosis. In addition, the two latter are different from the new species by their unique characters in the genus: *T. cordifolium* with adventitious buds

appearing at the mature leaf blade apex or sometimes at the top of the sheath, while *T. rhizomatosum* often forms large colonies thanks to its rhizomatous offshoots (Murata et al. 2010; Boyce et al. 2012; Nguyen-Phi et al. 2023).

The new species may be morphologically close to *T. hayatae* Sribonma & J. Murata (Sriboonma et al. 1994), *T. inopinatum* Prain (King and Prain 1898), *T. medusae* Hett. & Sookch. (Hetterscheid et al. 2001) and *T. varians* Hett. & Sookch. (Hetterscheid et al. 2001), as they have a short spathe limb and similar general shape and structure of the spadix and filiform staminodes. However, these species are easily distinguishable from the new taxon as they have elongate conical, sessile or shortly stipitate spadix appendix with truncate base. Furthermore, *T. hayatae* has globose tuber, long spadix appendix (6–14 cm) and spathe limb (12–26 cm); *T. inopinatum* has globose tuber, ovate-lanceolate spathe limb that is shorter than the spadix appendix and yellow staminodes; *T. medusae* has depressed tuber, velvety petioles, hairy leaf blades, orbicular spathe limb, pale yellow or cream staminodes and elongate conical appendix; *T. varians* has depressed tuber, dark grey spadix appendix 4.5–9 cm long, triangular ovate spathe limb of 7–14 cm length and 4–7.5 cm in diameter and pale pink anthers. Their different morphological characters are summarised in Table 1.

Key to the 24 presently known Vietnamese species of *Typhonium*

- 1 Sterile interstice of spadix entirely covered with staminodes ***T. flagelliforme***
- Only base of sterile interstice of spadix covered with staminodes **2**
- 2 Leaves perfectly trifoliolate **3**
- Not as above **4**
- 3 Plant with 3 leaves; appendix stipitate; female section of spadix with 5–6 rows of pistils ***T. thatsonense***
- Plant with 1 leaf; appendix sessile; female section of spadix with 2–3 rows of pistils ***T. hangiae***
- 4 Inflorescence appearing before the leaves **5**
- Inflorescence appearing together with the leaves **6**
- 5 Spathe 6–9 cm long; staminodes ca. 6 mm long ***T. penicillatum***
- Spathe 14–30 cm long; staminodes \leq 3 mm long ***T. hayatae***
- 6 Spathe limb elongate, narrowly lanceolate-triangular **7**
- Spathe limb wide, oblong-elliptic, ovate to lanceolate **15**
- 7 Staminodes red with a light yellow acumen **8**
- Staminodes unicolourful **9**
- 8 Spathe tube globose, to 1.5 cm long; staminodes 5 mm long, clavate ***T. bachmaense***
- Spathe tube oblong or cylindrical, 2 cm long; staminodes 12 mm long, acute ***T. kbangense***
- 9 Spathe limb white ***T. praelongum***
- Not as above **10**
- 10 Spathe limb corrugated ***T. corrugatum***
- Not as above **11**
- 11 Leaf 3-lobed **12**
- Not as above **13**
- 12 Stigma funnel-shaped and lobed ***T. stigmatilobatum***
- Stigma disciform and unlobed ***T. huense***

- 13 Leaves with 7 leaflets, leaflets linear to or linear-lanceolate ***T. lineare***
- Not as above **14**
- 14 Spathe limb much shorter than spadix appendix ***T. vermiforme***
- Spathe limb as long as spadix appendix ***T. dongnaiense***
- 15 Spadix longer than spathe **16**
- Spadix as long as or shorter than spathe **18**
- 16 Stamines folded 180° apically ***T. phuocbinhense***
- Stamines straight up to parallel to axis **17**
- 17 Male zone cylindrical, stamines cylindrical to conical ***T. khonkaenensis***
- Male zone subglobose, stamines clavate ***T. acetosella***
- 18 Leaves developing bulbils at the top and/or the base, upper surface often grey variegated ***T. cordifolium***
- Not as above **19**
- 19 Spathe limb very strongly circinnately recoiled over the entire length ***T. circinnatum***
- Not as above **20**
- 20 Tuber producing rhizomatous offsets about 5 cm apart ***T. rhizomatosum***
- Not as above **21**
- 21 Stamines ≤ 3 mm long **22**
- Stamines > 5 mm long **23**
- 22 Spathe tube above the ground; spathe limb oblong-elliptic, with obtuse apex ***T. obtusum***
- Spathe tube underground; spathe limb triangular ovate, with acute apex ***T. vietnamense***
- 23 Stamines red with a light yellow acumen, upward straight or slightly curved ***T. blumei***
- Stamines whitish, curly ***T. trilobatum***

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This research is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 106.03-2020.37.

Author contributions

Conceptualization, Hong Truong Luu, Xuan Bach Nguyen-Le; Formal analysis, Hong Truong Luu, Nga Nguyen-Phi, Xuan Bach Nguyen-Le, Hong Thien Van; Funding acquisition, Hong Truong Luu; Investigation, Xuan Bach Nguyen-Le, Hieu Cuong Nguyen; Methodology, Hong Truong Luu, Nga Nguyen-Phi, Quoc Dat Nguyen, Quoc Dat Nguyen; Supervision, Hong Truong Luu; Visualization, Hong Truong Luu, Nga Nguyen-Phi, Xuan Bach Nguyen-Le; Writing – original draft, all authors; Writing – review & editing, Hong Truong Luu, Nga Nguyen-Phi, Xuan Bach Nguyen-Le. All authors have read and agreed to the published version of the manuscript.

Author ORCIDs

Hong Truong Luu  <https://orcid.org/0000-0002-7036-7081>

Nga Nguyen-Phi  <https://orcid.org/0000-0001-7429-3641>

Quoc Dat Nguyen  <https://orcid.org/0000-0002-3687-1567>

Hieu Cuong Nguyen  <https://orcid.org/0000-0001-8707-2944>

Hong Thien Van  <https://orcid.org/0000-0003-0151-5068>

Xuan Bach Nguyen-Le  <https://orcid.org/0000-0001-8391-338X>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Boyce PC, Sookchaloem D, Hettterscheid WL, Gusman G, Jacobsen N, Idei T, Nguyen VD (2012) Araceae. Flora of Thailand 11: 101–321.
- Bridson D, Forman L (1999) Herbarium Handbook (3rd Ed.). Lubrecht & Cramer Ltd, Kew, 346 pp.
- Gagnepain F (1942a) Aracées. In: Lecomte H (Ed.) Flore Générale de l'Indo-Chine 6. Masson, Paris, 1174–1181.
- Gagnepain F (1942b) *Typhonium* (Araceae) nouveaux d'Indochine. Bulletin de la Société botanique de France 89: 10–12. <https://doi.org/10.1080/00378941.1942.10833053>
- Galloway A (2012) New Araceae species from Laos and Thailand. Aroideana 35: 51–64.
- Hettterscheid W (2013) New *Typhonium* species from Asia. Aroideana 36: 93–97.
- Hettterscheid WLA, Boyce PC (2000) A reclassification of *Sauromatum* Schott and new species of *Typhonium* Schott (Araceae). Aroideana 23: 48–55.
- Hettterscheid WLA, Nguyen VD (2001) Three new species of *Typhonium* (Araceae) from Vietnam. Aroideana 24: 24–29.
- Hettterscheid W, Sookchaloem D, Murata J (2001) *Typhonium* (Araceae) of Thailand: New species and a revised key. Aroideana 24: 30–55.
- Hu SY (1968) Studies in the Flora of Thailand 41. Araceae. Dansk Botanisk Arkiv 23: 409–457.
- IUCN Standards and Petitions Subcommittee (2022) Guidelines for using IUCN Red List Categories and Criteria. Version 15. Prepared by the Standards and Petitions Subcommittee. <https://www.iucnredlist.org/documents/RedListGuidelines.pdf> [Accessed 31 January 2024]
- King G, Prain D (1898) Descriptions of some new plants from the north-eastern frontiers of India. Journal of the Asiatic Society of Bengal 67: 284–305.
- Low SL, Yu CC, Ooi IH, Eiadthong W, Galloway A, Zhou ZK, Xing YW (2021) Extensive Miocene speciation in and out of Indochina: The biogeographic history of *Typhonium sensu stricto* (Araceae) and its implication for the assembly of Indochina flora. Journal of Systematics and Evolution 59(3): 419–428. <https://doi.org/10.1111/jse.12689>
- Luu HT, Van HT, Ngo TTD, Nguyen LP, Le HP (2017) *Typhonium thatsonense* (Araceae), a new species from Vietnam. Novon 25(4): 438–441. <https://doi.org/10.3417/D-16-00020>
- Murata J, Ohi-Toma T, Tanaka N (2010) New or noteworthy plant collections from Myanmar (4): *Typhonium cordifolium* and two new species, *T. neogracile* and *T. praecox* (Araceae). Shokubutsu Kenkyu Zasshi 85: 1–7.
- Nguyen VD (2005) Araceae. In: Nguyen TB (Ed.) Checklist of Plant Species of Vietnam (Vol. 3). Agriculture Publishing House, Hanoi, 871–898.

- Nguyen VD (2008) *Typhonium stigmatilobatum* (Araceae tribe Areae), a new species from Vietnam. *Kew Bulletin* 63(3): 491–493. <https://doi.org/10.1007/s12225-008-9044-8>
- Nguyen VD (2017) *Flora of Vietnam 16: Araceae Juss.* Publishing House for Science and Technology, Hanoi, 458 pp.
- Nguyen VD, Croat TB (1997) A new species of *Typhonium* (Araceae) from Vietnam. *Aroideana* 20: 48–52.
- Nguyen VD, Croat TB (2010) A new species of *Typhonium* (Araceae) from Vietnam. *Novon* 20(2): 195–197. <https://doi.org/10.3417/2008091>
- Nguyen VD, Vu XP (2004) The genus *Typhonium* (Araceae) in flora of Vietnam. *Academia Journal of Biology* 26: 25–31.
- Nguyen VD, Le CT, Nguyen VD, Ha MT, Nguyen TVA, Croat T (2021) *Typhonium phuocbinhense* sp. nov. (Araceae: Areae), a new species from central Vietnam. *Phytotaxa* 482: 073–079. <https://doi.org/10.11646/phytotaxa.482.1.8>
- Nguyen VD, Le CT, Dinh QD, Nguyen TVA, Tran VT, Croat TB (2022a) *Typhonium kbangense*, a new species of Araceae (Aroidea–Areae) from Central Vietnam. *Nordic Journal of Botany* 2022(8): e03601. <https://doi.org/10.1111/njb.03601>
- Nguyen VD, Nguyen TVA, Tran VT, Le CT (2022b) Two new records of species and a new synonym of genus *Typhonium* (Araceae) for flora of Indochina. *VNU Journal of Science: Natural Sciences and Technology* 38: 54–60.
- Nguyen-Phi N, Van HT, Luu HT (2023) *Typhonium rhizomatosum* (Araceae), a new record for the flora of Vietnam. *Academia Journal of Biology* 45(3): 125–131. <https://doi.org/10.15625/2615-9023/18612>
- Pham TKT, Nguyen DD, Nguyen VC, Nguyen VD, Tran VT (2023) *Typhonium hangiae* (Araceae: Aroideae: Areae), a new species from Central Vietnam with a key to Vietnamese species of the genus. *Kew Bulletin* 78(4): 1–6. <https://doi.org/10.1007/s12225-023-10134-8>
- Pham-Hoang H (1993) *Cây cỏ Việt Nam [An illustrated flora of Vietnam]*. Vol. 3. Montreal, Canada, 1176 pp.
- Pham-Hoang H (2003) *Cây cỏ Việt Nam [An illustrated flora of Vietnam]*. Vol. 3. Youth Publishing House, Ho Chi Minh, 1020 pp.
- POWO (2024) *Plants of the World Online*. Facilitated by the Royal Botanic Gardens, Kew. <http://www.plantsoftheworldonline.org/> [Accessed 31 January 2024]
- Schott HW (1829) *Wiener Zeitschrift für Kunst, Litteratur, Theater und Mode*. Vienna.
- Serebryanyi M, Trinh T, Hettterscheid W (2023) New tuberous Araceae from Binh Thuan Province (South Vietnam). *Blumea* 68(1): 39–48. <https://doi.org/10.3767/blumea.2023.68.01.03>
- Sriboonma D, Murata J, Iwatsuki K (1994) Revision of *Typhonium* (Araceae). *Journal of the Faculty of Science, University of Tokyo Section III. Botany* 15: 255–313.
- Thiers B (2024) *Index Herbariorum: A global directory of public herbaria and associated staff*. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/> [Accessed 31 January 2024]
- Van HT, Nguyen-Phi N, Vu NL, Galloway A, Luu HT (2017) *Typhonium dongnaiense* (Araceae), a new species from Vietnam. *Annales Botanici Fennici* 54(4–6): 405–408. <https://doi.org/10.5735/085.054.0622>
- Van HT, Van Son L, Nguyen-Phi N, Nguyen QD, Nguyen TQT, Nguyen HC, Luu HT (2021) A new species and a new record of *Typhonium* (Araceae) from southern Vietnam. *Phytotaxa* 527(3): 201–208. <https://doi.org/10.11646/phytotaxa.527.3.4>