

Four new synonyms and a new combination in *Parnassia* (Celastraceae)

Yumin Shu¹, Zhixiang Zhang^{1,2}

1 Laboratory of Systematic Evolution and Biogeography of Woody Plants, College of Nature Conservation, Beijing Forestry University, Beijing 100083, China **2** Museum of Beijing Forestry University, Beijing 100083, China

Corresponding author: Zhixiang Zhang (zxzhang@bjfu.edu.cn)

Academic editor: P. Stoev | Received 16 December 2016 | Accepted 17 February 2017 | Published 15 March 2017

Citation: Shu Y, Zhang Z (2017) Four new synonyms and a new combination in *Parnassia* (Celastraceae). *PhytoKeys* 77: 81–88. <https://doi.org/10.3897/phytokeys.77.11513>

Abstract

Parnassia yunnanensis had been previously described based on mixed specimens containing materials partially belonging to *P. cacuminum*, which makes the application of *P. yunnanensis* ambiguous. Therefore, we lectotypified *P. yunnanensis* and meanwhile synonymized *P. lanceolata* var. *oblongipetala* under it. *P. yunnanensis* var. *longistipitata* was found more similar to *P. cacuminum* rather than *P. yunnanensis*, thus a new combination, *P. cacuminum* var. *longistipitata* **comb. nov.** was proposed. Furthermore, other three names (*P. vevusta*, *P. degeensis* and *P. kangdingensis*) were reduced to synonyms of *P. cacuminum* too.

Keywords

Taxonomy, lectotypification, *Parnassia*, morphology

Introduction

Parnassia L. is a genus containing approximately 70 species (Ku and Hultgård 2001) which are predominantly distributed in arctic and temperate zones of the northern hemisphere and are mostly diverse in China and the Himalayas (Simmons 2004).

Parnassia yunnanensis Franch. (1896: 266) was described from Heqing, Yunnan province in China in 1896 which based on two collections (*Delavay* 710 and *Delavay s.n.*). Four duplicates of *Delavay* 710 were successfully traced from P and K (herbaria acronyms following Thiers 2016). However, one of these duplicates deposited in P (barcode number P00709380, Figure 1a) is a mixed specimen: an individual indicated by a circle bears obovate petals (magnified as Figure 1b, d), while the rest




Figure 1. A mixed type specimen of *Parnassia yunnanensis*. The sheet of Delavay 710 with barcode P 00709380 (a), the individual indicated with a circle and magnified should be determined as *P. cacuminum* (b, d), an individual represents the rest, indicated with a square and magnified (c, e), *P. cacuminum* (f) and *P. yunnanensis* (g) in the wild.

possess lanceolate petals (Figure 1c, e, magnified from the individual indicated by a square). Obviously, they are different taxa, the individual with obovate petals should be *P. cacuminum* Hand.-Mazz. (1931: 433). Franchet used ‘oblonga’ to describe the shape of petals in the protologue of *P. yunnanensis*, that made this name ambiguous, some new names were published to be closely related to *P. yunnanensis* but were actually close to or conspecific with itself or *P. cacuminum*.

Pan described *P. longipetaloides* J.T. Pan (1985: 222) based on *Jinshajing Exped.* 6105 in KUN collected from Eryuan, Yunnan in 1963 and this name was placed in synonymy of *P. yunnanensis* in *Flora Yunnanica* (Chuang 2006). According to T.T. Yu 7429A collected from Muli, Sichuan province in 1937, *P. lanceolata* T.C. Ku (1987: 34) was described by Ku and soon after, she proposed a new variety, *P. lanceolata* var. *oblongipetala* T.C. Ku (1991: 82) based on an isotype of *P. longipetaloides*, *Jinshajing Exped.* 6105 stored in PE. Wu et al. (2008) reduced *P. lanceolata* to a synonym of *P. yunnanensis*, but they did not mention var. *oblongipetala*.

Parnassia yunnanensis var. *longistipitata* Z.P. Jien (1963: 255) was proposed in 1963. This variety has white petals which are significantly different from *P. yunnanensis* but closer to *P. cacuminum*. *Parnassia venusta* Z.P. Jien (1963: 257), *P. degeensis*

T.C. Ku (1987: 30) and *P. kangdingensis* T.C. Ku (1987: 35) are three taxa that are morphologically close to *P. cacuminum*, but the authors compared them to some other taxa in the protologues when published them. Further studies in collections and field expeditions have been carried out to synonymise them in this paper.

Methods

This paper is based on the critical review of the protologues and examination of specimens in herbaria BJFC, CDBI, K, KUN, P, PE, HNWP, SM, SZ and online on Jstor Global Plants (<https://plants.jstor.org/> accessed on 12 December 2016). The measurements provided herein were mostly taken from dried herbarium specimens, and certain features such as colours were supplemented with the information from field observation. By applying Art 9.1 and 9.2 of the ICN (McNeill et al. 2012) strictly, the lectotype was selected. Specimens examined were listed alphabetically.

Results and discussion

Since the syntypes of *P. yunnanensis* belong to more than one taxon, a lectotype needs to be designated for it. Wu et al. (2008) cited the type of *P. yunnanensis* in their taxonomic work as “*China. Yunnan, Hokin, Delavay 710 (holotype K!); Sichuan, near Tatchienlu, Pratt 542 (syntype, BM!)*”. As Franchet did not mention *Pratt 542* in the protologue when he described this new name, *Pratt 542* should not be regarded as a syntype of *P. yunnanensis*. In addition, they did not use the phrase “designated here” or the term “lectotype” in the statement. According to Art. 7.10 and 9.23 of the ICN, their typification of *P. yunnanensis* was not effective. Franchet worked at P (Stafleu and Cowan 1976), so we select a well-presented specimen with barcode P00709378 that contains most individuals as the lectotype.

Ku stated in the protologue of *P. lanceolata* var. *oblongipetala* that it differs from the typical variety in having oblong petals, obtuse at the apex, and *P. lanceolata* differs from *P. longipetaloides* in having 3-lobed staminodes while the latter have obscure 4–6 dentate ones. By taxonomic revision of this variety, we determined that the type specimen of this name was indistinguishable from that of *P. yunnanensis* in terms of taxonomically important morphological characters (shapes of petals and staminodes, Figures 1g, 2a–d) and their general distribution ranges (both type specimens were collected from Mt. Maer between Hoqing and Eryuan). Thus we treated this variety as a new synonym of *P. yunnanensis*.

Parnassia venusta was described based on *T.T. Yu 22666* collected from Gongshan, Yunnan by Jien. As concluded from the protologue, he thought that this species differs from *P. cacuminum* in two characters: leaves reniform, petals lanceolate-obovate, slightly fimbriated at the base (Figure 2i) while *P. cacuminum* has cordata leaves, petals late obovate, erose or slightly fimbriated at the base (Figures 1f, 2e, f). The difference




Figure 2. Morphology of the petals and staminodes in *Parnassia*. *P. lanceolata* var. *oblongipetala* (a, b *Jinshajinag Exped.* 6105, the holotype), *P. yunnanensis* (c, d T.T. Yu 7429), *P. cacuminum* (e T.T. Yu 12254 f Y.M. Shu et al. sz286), *P. kangdingensis* (g, h Y.M. Shu et al. sw365), *P. venusta* (i Y.M. Shu et al. sw365), *P. cacuminum* var. *longistipitata* (j Y.M. Shu et al. sw459), *P. degeensis* (k Y.W. Tsui 4997, the holotype). Scale bar = 1 mm.

in the shape of the leaves was observed and we found it do not held good; reniform to cordate leaves always occur within a population even within a single individual. *Parnassia cacuminum* has obovate petals, but it is variant, 4–9 mm in length by 2–5 mm in width are observed in the sample specimens, and no dividing line can be found amongst lanceolate-obovate, obovate and late obovate. Hence, we treated *P. venusta* as a new synonym of *P. cacuminum*.

In 1987, Ku proposed two new species, *P. degeensis* and *P. kangdingensis* from Dege and Kangding respectively. *P. degeensis* was described based on Y.W. Tsui 4997, which has only one sheet containing one individual with two flowers. Ku described the

staminodes of this name as undulate or 5–7 dentates at the apex and compared it with *P. farreri* W.E. Evans (1921: 174), which differed significantly in the shape of petals. We examined the holotype of *P. degeensis* (Figure 2k); the petals were obovate with distinct veins, staminodes 3 lobed at the apex and that agreed well with *P. cacuminum*. Ku stressed in the protologue of *P. kangdingensis* that this new taxon was closely related to *P. lanceolata*. However, from the description, it could be concluded that this name was quite different from *P. lanceolata* (obovate white petal vs. lanceolata green petal) but close to *P. cacuminum* by the numerical characters which Ku used to distinguish them, such as heights of individuals and length of petals. As we examined the type specimens and observed the populations in Kangding (Figure 2g, h), variations of height from 2–9 cm and length of petals from 3–6 mm were found within a population. Therefore, we propose to recognise *P. degeensis* and *P. kangdingensis* as synonyms of *P. cacuminum*.

Not far from the location where *P. kangdingensis* was described in Kangding, a new variety, *P. yunnanensis* var. *longistipitata* was proposed. This variety has cordate leaves, elliptic to obovate white petals and flat, linear staminodes with entire or obscure 3 dentate apices (Figure 2j). All these characters made it different from *P. yunnanensis* but close to *P. cacuminum*. The only difference between it and *P. cacuminum* is the linear shaped staminodes. The field observation found that these linear shaped staminodes were easily distinguished from those of *P. cacuminum* and they were stable within populations. Thus, it was proposed to recognise this taxon as a variety of *P. cacuminum*.

Taxonomic treatment

***Parnassia yunnanensis* Franch., J. Bot. (Morot) 10(16): 266. 1896.**

Parnassia longipetaloides J.T. Pan, Acta Phytotax. Sin. 23(3): 222. 1985.

Type: China. Yunnan: Eryuan, Mt. Maer, elev. 3600 m, 25 Jul 1963, Jinshajinag Exped. 6105 (holotype: KUN [KUN0437666!]; isotype: KUN!).

Parnassia lanceolata T.C. Ku, Bull. Bot. Res. 7(1): 34. 1987.

Type: China. Sichuan: Muli, elev. 3900 m, 28 Jul 1937, T.T. Yu 7429A (holotype: PE [PE01842934!]; isotypes: PE!, KUN!, E [image!]).

Parnassia lanceolata T.C. Ku var. *oblongipetala* T.C. Ku, Acta Phytotax. Sin. 29(1): 82. 1991.

Type: China. Yunnan: Eryuan, Mt. Maer, elev. 3600 m, 25 Jul 1963, Jinshajinag Exped. 6105, **syn. nov.** (holotype: PE [PE01842922!]).

Type. China. Yunnan: Heqing (Hokin), Kowa-la-po, in the shrub, 26 Aug 1884, Delavay 710 (Lectotype, designated here: P [P00709378!]; isolectotypes: P [P00709379!, P00709380! (exclude the individual with obovate petals)], K [K000739471!]).

Specimens examined. CHINA. Gansu: Lintan, 26 Jul 2013, C. Shang I”-218 (BJFC); Sichuan: Jiulong, elev. 3800 m, 15 Jul 1979, T.C. Wei 20512 (CDBI); Jiulong, elev. 3800 m, 25 Aug 1980, Z.A. Liu 22993 (CDBI); Kangding, elev. 4010 m, 5

Aug 2015, Y.M. Shu et al. sw447 (BJFC); Kangding, elev. 3600 m, 6 Jul 1974, N.Z. Zhang 4801 (PE, CDBI); Kangding, elev. 3080 m, 25 Jul 1934, C.S. Liu 883 (PE, SZ); **Yunnan**: Dali, elev. 3800 m, 31 Jul 2014, Y.M. Shu et al. sz072 (BJFC); Dongchuan, 18 Jul 2009, Huang 1529 (KUN); Lijiang, elev. 4000 m, 17 Aug 2003, D. Wu et al. 3005 (KUN); Lijiang, elev. 4000 m, 26 Aug 2002, D. Wu et al. 2005 (KUN).

***Parnassia cacuminum* Hand.-Mazz., Symb. Sin. 7(2): 433. 1931.**

Parnassia venusta Z.P. Jien, Acta Phytotax. Sin. 8(3): 257. 1963.

Type: China. Yunnan: Gonshan Hsien, Sawalunba, elev. 4000 m, 3 Sep 1938, T.T. Yu 22666, **syn. nov.** (holotype: PE [PE01842930!]; isotypes: PE!, KUN!).

Parnassia degeensis T.C. Ku, Bull. Bot. Res. 7(1): 30. 1987.

Type: China. Sichuan: Dege, Haizikou, 24 Jul 1951, Y.W. Tsui 4997, **syn. nov.** (holotype: PE [PE01896065!]).

Parnassia kangdingensis T.C. Ku, Bull. Bot. Res. 7(1): 35. 1987.

Type: China. Sichuan: Kangding, 28 Jul 1951, W.P. Fang et al. 10632, **syn. nov.** (holotype: PE [PE01842917!]; isotype: SZ!).

Type. China. Sichuan: Muli, elev. 4450–4500 m, 30 Jul 1915, Handel-Mazzetti 7338 (holotype: WU [WU0046641 image!])

Specimens examined. CHINA. Qinghai: Yushu, elev. 4250 m, 23 July 1964, Yushu Exped. 587 (HNWP, PE); **Sichuan:** Baoxing, elev. 4000 m, 1 Aug 2015, Y.M. Shu et al. sw365 et sw370 (BJFC); Baoxing, elev. 4048 m, 2 Jul 2010, C.S. Chang et al. SI0933 (PE); Daofu, elev. 3100 m, 17 Jul 1979, s.n. 1203 (SM); Dege, elev. 4000 m, 2 Jul 1979, s.n. 311 (SM); Dege, elev. 4100 m, 23 Jun 1974, Sichuan Zhibei Exped. 7121(CDBI); Dege, elev. 4300 m, 19 Jun 1974, Qingzang Exped. 63 (PE); Dege, elev. 4100 m, 23 Jun 1974, s.n. 7121 (PE); Jiulong, elev. 4110 m, 19 Jul 2012, L. He et al. PH20120719-03 (BJFC); Jiulong, 30 Jun 1974, Z.G. Liu 4730 (PE, CDBI); Jiulong, elev. 4100 m, 16 Jun 1984, W.L. Chen et al. 6345 (PE); Kangding, elev. 3730 m, 15 Jul 2012, L. He et al. PH20120715-06 (BJFC); Kangding, elev. 3700 m, 20 Jun 1984, W.L. Chen et al. 6541 (PE); Kangding, C.S. Liu 940 (PE); Kangding, 1 Aug 1963, West Sichuan Exped. 1414 (PE); Luhuo, elev. 3900 m, 5 Jul 1974, Q.H. Li et al. 6446 (PE, CDBI); Muli, 18 Aug 1937, T.T. Yu 7795 (KUN); Muli, Jun 1928, J.F. Rock 16564 (IBSC); Muli, elev. 4333 m, May 1932, J.F. Rock 23749 (PE); Muli, elev. 3700 m, 21 Jun 1937, T.T. Yu 6528 (PE, KUN); Muli, elev. 3400 m, 18 Aug 1937, T.T. Yu 7795 (PE); Rangtang, elev. 4300 m, 16 Jul 1979, s.n. 783 (SM); **Xizang:** Bomi, elev. 4300 m, 5 Sep 1982, B.S. Li et al. 00673 (PE); Chayu, elev. 3800 m, 8 Sep 1982, Qingzang Exped. 10118 (PE); Chayu, elev. 4100 m, 9 Sep 1982, Qingzang Exped. 10231 (PE, KUN); Chayu, elev. 3800 m, 19 Jul 2010, X.H. Jin et al. STET0551 (PE); Chayu, elev. 4100 m, 16 Jul 2010, X.H. Jin et al. STET0806 (PE); Chayu, elev. 3700 m, Aug 1935, T.W. Wang 65980 (PE); Cuona, elev. 4500 m, 19 Jul 1975, Z.Y. Wu et al. 75-1119 (KUN); Gongjue, elev. 4220 m,

15 Aug 2010, Kangzang Exped. 10-1966 (PE); Milin, elev. 4300 m, 28 Jul 1983, B.S. Li et al. 5960 (PE); Lasa, elev. 4952 m, 8 Jul 2012, L.M. Gao GLM-123788 (KUN); Linzhi, 23 Jul 2014, L. He PH20140723-10 (BJFC); **Yunnan:** Bijiang, elev. 4300 m, 12 Sep 1964, S.K. Wu 8790 et 8810 (KUN); Deqin, elev. 4300 m, 19 Jul 2014, Y.M. Shu et al. sz 286 (BJFC); Deqin, elev. 4300 m, 31 Jul 2003, H. Wang et al. 3066 (KUN); Deqin, elev. 4300 m, 15 Jul 2004, J. Cai et al. 4209 (KUN); Deqin, elev. 4100 m, 10 Jul 1983, Hengduan Mt. Exped. 4576 (PE); Deqin, elev. 3200 m, Aug 1935, C.W. Wang 64943 et 64974 (PE); Deqin, elev. 3500 m, 6 Jul 1937, T.T. Yu 8758 (PE, KUN); Deqin, elev. 4000 m, 9 Sep 1938, T.T. Yu 22245 (PE, KUN); Deqin, elev. 3600 m, 4 Aug 1940, K.M. Feng 5921 (KUN); Deqin, 15 Aug 1976, J.S. Yang 8573 (KUN); Deqin, elev. 3800 m, 1 Aug 1940, K.M. Feng 6209 (KUN); Eryuan, elev. 3700 m, 20 Jul 1963, Jinshajiang Exped. 63-6118 (KUN); Lijiang, elev. 4100 m, 14 Aug 2005, D. Wu et al. 5012 (KUN); Qiaojia, elev. 3800 m, 17 Jul 1973, B.X. Sun et al. 1032 (KUN); Weixi, elev. 3500 m, Jul 1935, C.W. Wang 64660 (PE); Weixi, elev. 3600 m, Aug 1935, C.W. Wang 68611 (PE); Xiangelila, elev. 4658 m, 3 Sep 2010, Kangzang Exped. 10-3237 (PE); Xiangelila, elev. 3600 m, 18 Jul 1937, T.T. Yu 12254 (PE, KUN); Xiangelila, 25 Jul 1939, K.M. Feng 1802 (PE); Xiangelila, elev. 4000 m, 6 Jul 2006, W.B. Yu et al. 5052 et 5053 (KUN); Xiangelila, elev. 3700 m, 12 Jul 1937, T.T. Yu 12085 (PE, KUN); Xiangelila, elev. 4000 m, 25 Jul 2003, D. Wu et al. 3028 (KUN); Yangbi, elev. 2600 m, 30 Jul 2003, D. Wu et al. 326 (KUN).

***Parnassia cacuminum* Hand.-Mazz. var. *longistipitata* (Z.P. Jien.) Y.M.Shu & Z.X.Zhang, comb. nov.**

urn:lsid:ipni.org:names:77161383-1

Basionym. *Parnassia yunnanensis* Franch. var. *longistipitata* Z.P. Jien, Acta Phytotax. Sin. 8(3): 255. 1963.

Type. China. Sichuan: Kangding, Mt. Che To, 6 Aug 1934, C.S. Liu 1028 (holotype: PE [PE01842918!])

Specimens examined. CHINA. Sichuan: Kangding, elev. ca. 4200 m, 6 Aug 2015, Y.M. Shu et al. sw459 et sw463 (BJFC); Kangding, elev. 4187 m, 12 Aug 2015, Y.M. Shu et al. sw505 (BJFC); Kangding, 7 Aug 2011, S.X. Yu 5008 (PE); Kangding, 10 Aug 2009, WPW 108 (KUN); Kangding, elev. 4000 m, 14 Jul 1981, Z.J. Zhao 114906 (SZ); Qianning, elev. 4300 m, 4 Jul 1974, Sichuan zhibei Exped. 5497 (PE, CDBI).

Acknowledgements

We are indebted to the curator and staff members of BJFC, CDBI, K, KUN, P, PE, HNWP, SM, SZ and WU herbaria for either providing images of specimens on their website or allowing us to examine collections in their care for this study. Lijuan Liu is

thanked for providing magnified photos of type specimens. This study was funded by the Science and Technology Basic Work (grant no. 2013FY112100) and the National Natural Science Foundation of China (grant no. 31110103911).

References

- Chuang H (2006) *Parnassia*. In: Wu ZY (Ed.) *Flora Yunnanica*. vol. 12. Science Press, Beijing, 121–140. [In Chinese]
- Evans WE (1921) Diagnoses specierum novarum in herbario Horti Regii Botanici Edinburgensis cognitarum. Notes from the Royal Botanic Garden, Edinburgh 13(63–64): 172–175. <http://www.biodiversitylibrary.org/item/99837>
- Franchet AR (1896) Saxifragaceae, Crassulaceae, et Combretaceae novae. Journal de Botanique (Morot) 10(16): 266–267. <http://www.biodiversitylibrary.org/item/35707>
- Handel-Mazzetti H (1931) Saxifragaceae, *Parnassia*. Symbolae Sinicae 7(2): 432–435. <http://dx.doi.org/10.5962/bhl.title.878>
- Jien ZP (1963) Some new plants of Chinese *Parnassia*. Acta Phytotaxonomica Sinica 8(3): 253–260.
- Ku TC (1987) A revision of the genus *Parnassia* (Saxifragaceae) in China. Bulletin of Botanical Research 7(1): 1–61.
- Ku TC (1991) New taxa of Saxifragaceae and Rosaceae from the Hengduan mountains. Acta Phytotaxonomica Sinica 29(1): 80–83.
- Ku TC, Hultgård UM (2001) *Parnassia*. In: Wu ZY, Raven PH (Eds) *Flora of China*, vol. 8. Missouri Botanical Garden Press, St. Louis and Science Press, Beijing, 95–118.
- McNeill J, Barrie FR, Buck WR, Demoulin V, Greuter W, Hawksworth DL, Herendeen PS, Knapp S, Marhold K, Prado J, Prud'homme van Reine WF, Smith GF, Wiersema JH, Turland NJ (Eds) (2012) International Code of Nomenclature for algae, fungi and plants (Melbourne Code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011. Regnum Vegetabile 154. Koeltz Scientific Books. <http://www.iapt-taxon.org/nomen/main.php>
- Pan JT (1985) New taxa of the genus *Parnassia* from China. Acta Phytotaxonomica Sinica 23(3): 220–224.
- Simmons MP (2004) Parnassiaceae. In: Kubitzki K (Ed.) *The families and genera of Vascular Plants*. Springer Verlag, Berlin, 291–296. https://doi.org/10.1007/978-3-662-07257-8_33
- Stafleu FA, Cowan RS (1976) *Taxonomic Literature: A Selective Guide to Botanical Publications and Collections with Dates, Commentaries and Types* (2nd edn), Vol. 1. Scheltema & Holkema, Utrecht, Bohn, 869 pp.
- Thiers B (2016) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih/> [Accessed: 15 Sep 2016]
- Wu D, Lu JM, Wang H (2008) Taxonomic Revision on Some Species of *Parnassia* (Parnassiaceae) from China. Acta Botanica Yunnanica 30(6): 657–661.