

# Taxonomic studies of *Glochidion* (Phyllanthaceae) from the Indo-China Peninsula (II): The identities of *G. anamiticum* and *G. annamense*

Gang Yao<sup>1</sup>, Zhuqiu Song<sup>2</sup>, Youheng Wu<sup>2,3</sup>, Nguyen Thi Kim Thanh<sup>4</sup>, Yuling Li<sup>1</sup>, Shi Shi<sup>1</sup>, Shixiao Luo<sup>2</sup>

**1** South China Limestone Plants Research Center, College of Forestry and Landscape Architecture, South China Agricultural University, Guangzhou 510642, China **2** Key Laboratory of Plant Resources Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 516650, China **3** University of Chinese Academy of Sciences, Beijing 1000049, China **4** Department of Botany, Faculty of Biology, HNU Herbarium, VNU University of Science, Vietnam

Corresponding author: Shixiao Luo ([luoshixiao@scbg.ac.cn](mailto:luoshixiao@scbg.ac.cn))

---

Academic editor: A. Sennikov | Received 21 October 2018 | Accepted 21 November 2018 | Published 13 December 2018

---

**Citation:** Yao G, Song Z, Wu Y, Thanh NTK, Li Y, Shi S, Luo S (2018) Taxonomic studies of *Glochidion* (Phyllanthaceae) from the Indo-China Peninsula (II): The identities of *G. anamiticum* and *G. annamense*. *PhytoKeys* 114: 1–9. <https://doi.org/10.3897/phytokeys.114.30725>

---

## Abstract

The names *Glochidion anamiticum* (Kuntze) Kuntze ex Merrill and *G. annamense* Beille were previously reduced to synonyms of *G. eriocarpum* Champ. ex Benth. However, literature examination and morphological comparison suggest that *G. annamense* is conspecific with *G. anamiticum* and the species can be readily distinguished from *G. eriocarpum* by its styles connate into a long cylindric column and up to 3 mm long, ovaries usually 3-locular, stigma usually 3-lobed, capsules pubescent and usually 6-grooved, persistent styles 3–4 mm long. Thus, the specific status of *G. anamiticum* is here reinstated and *G. annamense* is treated as its synonym.

## Keywords

*Glochidion*, Phyllanthaceae, Taxonomy, Vietnam

## Introduction

*Glochidion* J. R. Forster & G. Forster is a large genus within the tribe Phyllanthae Dumortier, family Phyllanthaceae Martinov (Webster 2014, APG 2016). It consists of over 300 species mainly distributed in the Indo-Pacific region, east to southeast Polynesia and south into Australia (Webster 2014, Yao and Zhang 2016). The genus can be distinguished from all the other genera in the tribe by its usually unlobed styles, apiculate anthers and usually fleshy seed coat (Webster 2014).

Molecular phylogenetic studies revealed that *Glochidion*, together with *Breynia* J. R. Forster & G. Forster, *Phyllanthodendron* Hemsl. and *Sauropus* Blume, were all deeply embedded within *Phyllanthus* L. s. str. (Hoffmann et al. 2006, Kathriarachchi et al. 2006). So, some authors suggested the merger of these genera with *Phyllanthus* and accepted the broad concept of *Phyllanthus* s. l. (including *Breynia*, *Glochidion*, *Phyllanthodendron* and *Sauropus*) as a super-genus that includes over 1,200 species (Hoffmann et al. 2006). However, Pruesapan et al. (2012) suggested that the reinstatement of *Breynia*, *Glochidion*, *Phyllanthodendron* and *Sauropus*, as well as the disintegration of the paraphyletic *Phyllanthus* s. str. (over 800 species) into smaller genera, seems to be more reasonable, because these groups can be distinguished easily from each other in morphology and *Phyllanthus* s. str. is actually a large and morphologically heterogeneous genus with wide distribution areas. The latter suggestion is further supported by morphological (van Welzen et al. 2014), palynological (Yao and Zhang 2016) and wood anatomical (Jangid and Gupta 2016) studies. Thus, the acceptance of the generic status of *Glochidion* is widely adopted in most recent literature (e.g. van Welzen et al. 2014, Webster 2014, Duocet Group 2016 onwards, Kato and Kawakita 2017).

The species *Glochidion anamiticum* (Kuntze) Kuntze ex Merrill was originally described by Kuntze (1891) in the genus *Diasperus* L. ex Kuntze based on one collection (*O. Kuntze* 3798) from Vietnam and *Glochidion annamense* Beill was described by Beille (1927) based on five collections (*F. Evrard* 233, *C. Gaudichaud* 161, *M. Krempf* 1567, *E. Poilane* 7838 and *E. Poilane* 10022) from Vietnam, too. The latter was soon reduced to the synonymy of *G. anamiticum* (Merrill 1936). In the taxonomic study of the Thai Euphorbiaceae, Airy Shaw (1972) provisionally placed both names under *G. eriocarpum* Champ. ex Benth., expressing uncertainty by question marks and this questionable treatment was directly followed by van Welzen (2007) in Flora of Thailand. Subsequent taxonomists accepted Airy Shaw's proposal to treat *G. anamiticum* and *G. annamense* as synonyms of *G. eriocarpum* (Govaerts et al. 2000, Li and Gilbert 2008, The Plant List 2013 onwards, Govaerts et al. 2018). However, *G. annamense* was recognised as a separate species by Nguyen (2007) when treating the Vietnamese Euphorbiaceae, but *G. anamiticum* was not considered. In our comprehensive taxonomic studies of *Glochidion*, we found that *G. anamiticum* and *G. annamense* represent the same species, but this species is clearly different from *G. eriocarpum* in morphology. We therefore propose to reinstate the specific status of *G. anamiticum* and treat *G. annamense* as a synonym of *G. anamiticum*.

## Material and methods

Specimens of *Glochidion* deposited in the herbaria HN, IBSC, K, KUN, NY, P and PE were studied carefully in the present study and field investigations of *G. anamiticum* and *G. eriocarpum* were carried out in recent years. Morphological characters of relevant species were also photographed and some of them were measured. Herbarium abbreviations cited here follow the Index Herbariorum (Thiers 2018 onwards).

## Results and discussions

*Glochidion anamiticum* (Figures 1A and D) and *G. annamense* (Figure 1B) represent the same species, which is similar to *G. eriocarpum* (Figures 1C and E) in habit to some extent. However, the Vietnamese endemic species *G. anamiticum* is morphologically distinct and can be readily distinguished from *G. eriocarpum* (Table 1) by its stipules triangular to narrowly triangular and ca. 1–1.5 mm long (Figure 2A) (vs. narrowly triangular to linear, 1–4 mm long), sepals of female flowers ovate-triangular to narrowly triangular and pubescent outside (Figures 2A and C) (vs. oblong and villous outside; Figures 2B and D), styles connate into a long cylindrical column up to 3 mm long (Figures 2A and C) (vs. a cylindrical column 1–1.5 mm long; Figure 2B), ovaries usually 3-locular (vs. usually 4–5-locular), stigma usually 3-lobed (Figure 2A) (vs. usually 4–5-lobed; Figure 2B), capsules pubescent and usually 6-grooved (Figures 2E and F) (vs. villous and usually 8–10-grooved; Figure 2G), persistent style cylindrical column 3–4 mm long (Figures 2E and F) (vs. shortly column-shaped to subconical, less than 1 mm long; Figure 2G). Thus, our results suggest that it may be best to treat *G. anamiticum* and *G. eriocarpum* as separate species. The following taxonomic treatment is therefore necessary.

## Taxonomic treatment


### *Glochidion anamiticum* (Kuntze) Kuntze ex Merrill, *Brittonia* 2: 195. 1936

Figures 1A, B, D, 2A, C, E, F

**Basionym.** *Diasperus anamiticus* Kuntze, Revis. Gen. Pl. 2: 601. 1891.

**Type.** Vietnam. Annam, Turong, March 1875, *O. Kuntze* 3798 (lectotype designated by Merrill (1936: 195): NY-00263420, photo!; isolectotype: K-001081193, photo!).

*Glochidion annamense* Beille (1927, p. 627). Type: Vietnam. Annam, Lien-chien, 25 August 1923, *E. Poilane* 7838 (lectotype designated by Nguyen (2007): HM; isolectotypes: K-001081192, P-00379159, P-00379160, photos!). Remaining syntypes: Vietnam. Annam, Trai-ca, province, Phanrang, 9 March 1924, *E. Poilane* 10022 (P-00379161, photo!); Tourane, January 1837, *C. Gaudichaud* 161 (P-00379151, photo!); Nha-trang, *M. Krempf* 1567 (P-00379152, photo!); Dalat réserve du Calmy, 24 October 1920, *F. Evrard* 233 (P-00379150, photo!).



**Figure 1.** Type (A–C) and field images (D–E) of *Glochidion* species. **A** Lectotype of *G. anamiticum* (Kuntze) Kuntze ex Merrill (O. Kuntze 3798, NY-00263420) **B** Syntype of *G. annamense* Beille (M. Krempf 1567, P-00379152) **C** Type of *G. eriocarpum* Champ. ex Benth. (J.G. Champion 470, K-001081087) **D** *G. anamiticum*; **E**, *G. eriocarpum*.




Shrubs, 1–3 m tall, monoecious; branchlets usually pubescent; leaf blade membranous, 6–8 × 2.5–3 cm, lanceolate, oblong to ovate, sometimes asymmetrical, base acute to round, apex acuminate, adaxially dark brown or blackish, pubescent on the median rib, abaxially light brown, pubescent; lateral veins 6–8 pairs, elevated abaxially; petiole ca. 2–3 mm long, pubescent; stipules triangular to narrowly triangular, 1–1.5 mm long. Male flowers: pedicels ca. 10 mm long, pubescent; sepals 6, oblong to ovate, hairy outside, glabrous inside; stamens 3, connate, surmounted by a conical prolongation. Female flowers: in axillary clusters, subsessile; sepals 6, ca. 1.2 mm long, ovate-triangular or narrowly triangular, hairy outside, glabrous inside; ovary depressed globose, 3 (rarely 4)-locular, pubescent; style cylindrical column, long exceeding the sepals, 2–3 mm long, hairy in the lower quarter, glabrous in the upper part, 3 (rarely 4)-lobed at the top. Capsule depressed globose, 6 (rarely 8)-grooved, 7–9 mm in diameter, ca. 5 mm high, very briefly hairy; persistent style a long cylindrical column up 3–4 mm long; fruit pedicels stout, 3–4 mm long, pubescent.

**Distribution and habitat.** *Glochidion anamiticum* is endemic to Vietnam, mostly from Da Nang to Ninh Thuan (Nguyen 2007).

**Notes.** In the monograph *Revisio Generum Plantarum*, Kuntze (1891) transferred hundreds of *Glochidion* and *Phyllanthus* species into the genus *Diasperus* L. ex Kuntze and he also described many new species, including *Diasperus anamiticus* Kuntze which was based on one of Kuntze's collections (*O. Kuntze* 3798) from Vietnam. At the same time, however, a specific name (*Glochidion anamiticum*), under the generic name *Glochidion*, was also suggested for *D. anamiticus*, but it was listed as a synonym of the new species. According to Art. 34.1 of the ICN (McNeill et al. 2012), the name *G. anamiticum* Kuntze is invalid because it was not accepted by the author and just cited as a synonym of *Diasperus anamiticus*. On the other hand, many of the new species described by Kuntze (1891) were overlooked for a long time, until they were re-evaluated by Merrill (1936) and then the species *Glochidion anamiticum* was accepted formally as a member of *Glochidion* for the first time. So, the author's attribution of the species should be *Glochidion anamiticum* (Kuntze) Kuntze ex Merrill.

In his commentary on Loureiro's (1790) new species described from southern Vietnam, Merrill (1935) transferred the species *Nymphanthus pilosus* Lour. into *Glochidion* and reduced *Glochidion annamense* to a synonym of his new combination *Glochidion pilosum* (Lour.) Merr., although the type of *G. pilosum* was not examined in his study. Merrill's (1935) taxonomic treatment of *Glochidion annamense* was followed by Ho (2003) in An Illustrated Flora of Vietnam. However, as noted by Loureiro (1790) in the protologue of *Nymphanthus pilosus*, the species was characterised by pinnate leaves and bacciformis fruits, which are much different from those of the genus *Glochidion* (characterised by alternate and distichous leaves, fruits capsule), but consistent with those of *Emblica* Gaertn. or *Phyllanthus* sect. *Emblica*, thus the species also had been transferred into these two genera and named as *Emblica pilosa* (Lour.) Spreng. or *Phyllanthus pilosus* (Lour.) Müll. Arg., respectively. In the following year, however, Merrill (1936) treated the name *Glochidion annamense* as a synonym of *G. anamiticum*, based on the study of type specimens, but the name *G. pilosum* was not considered.



**Figure 2.** Morphological comparisons between *Glochidion anamiticum* (Kuntze) Kuntze ex Merrill (**A, C, E, F**) and *G. eriocarpum* Champ. ex Benth. (**B, D, G**). **A–D** Female flower **D–G** Fruit. The female flowers and persistent styles in the fruits are shown by the solid arrowhead and hollow arrowhead, respectively.

**Table 1.** Morphological comparisons between *Glochidion anamiticum* (Kuntze) Kuntze ex Merrill and *G. eriocarpum* Champ. ex Benth.

Traits/Species	<i>Glochidion anamiticum</i>	<i>G. eriocarpum</i>
Stipule	Triangular to narrowly triangular; ca. 1–1.5 mm long;	Narrowly triangular to linear; 1–4 mm long;
Sepals of female flower	Ovate-triangular to narrowly triangular; pubescent outside;	Oblong; villous outside;
Style	Cylindrical column, long exceeding the sepals, 2–3 mm long;	Cylindrical column, slightly exceeding the sepals, 1–1.5 mm long;
Stigma	Deeply 3(4)-lobed;	Deeply 4–5-lobed;
Ovary	Pubescent; 3(4)-locular;	Villous; 4–5-locular;
Capsule	Pubescent; 7–9 mm in diameter; deeply 6(8)-grooved;	Villous; 8–10 mm in diameter; deeply 8–10-grooved;
Persistent style	Cylindrical column; up to 4 mm long;	Shortly column-shaped to subconical; less than 1 mm long.

Morphological comparisons between *Glochidion anamiticum* and *G. eriocarpum* can be seen in Table 1 and Figures 1, 2. The most noticeable traits of *G. anamiticum* are its long style (up to 3 mm long) in the female flowers and long persistent style (up to 4 mm long) in the fruits.

**Additional specimens examined.** *Glochidion anamiticum*: VIETNAM. Detailed locality unknown, 7 April 1978, *Nhan 294* (HN); Annam, Hué, Flower pale yellow; Near river, 11 March 1927, *R. W. Squires 178* (P-00379167); Arbore Broyé (Tuyen Duc), altitude 1500 m, 29 March 1953, *M. Schmid s.n.* (P-00509785); Dalat, ravin au S. du Langbian Palace, 17 May 1924, *F. Evrard 910* (P-00379154); Dalat, lac des Soupirs, altitude 1500 m, 12 January 1964, *M. Schmid s.n.* (P-00509783); Dalat, ravin derrière la gendarmerie, 2 October 1924, *F. Evrard 1293* (P-00379155); Dalat, ravin au Sud du Langbian Palace, 14 November 1924, *F. Evrard 1788* (P-00379156); Région de Dalat, September 1952, *M. Schmid 1288* (P-00509784); Cochinchine, prov. de Bienhoa, 11 October 1931, *E. Poilane 19574* (P-00379164); Env. de Hué, prov. Thua Thiên-Huê, 23 December 1943, *J.E. Vidal 917a* (P-00476551). Hue, Bach Ma National Park, 3 August 2018, *Y.H. Wu 17126* (IBSC). *Glochidion eriocarpum*: CHINA. Guangdong Province, Jiangmen, 21 October 2013, *L.X. Zhou 11542* (KUN); Guangdong Province, Yangchun, Ehuangzhang Nature Reserve, 14 November 2008, *G. Yao 018* (IBSC); Guangdong Province, Zhaoqing, Dinghushan Nature Reserve, 17 December 2008, *G. Yao 058* (IBSC); Guangxi Province, Rongshui, 24 October 2008, *G. Yao 08-008* (IBSC); Hainan Province, Xinglong, 6 March 2009, *G. Yao 101* (IBSC); Hongkong, J.G. Champion 470 (K); Yunan Province, Puer, 1 April 2003, *Y. Chen s.n.* (KUN); Yunan Province, Xishuangbanna, 29 July 1977, *G.D. Tao 15679* (KUN); VIETNAM. Annam, Environs du poste forestier de Bang Trê Lat sur sông Con, prov. de Vinh, 24 July 1929, *E. Poilane 16419* (P-00379163); Hanh Hoa Province, Thach Thanh District, Cuc Phuong NP, Thanh Yen Commune, Sanh Village, altitude 150 m, 27 February 2001, *N.M. Cuong 1320* (P-00520940); Tonkin, Binh Lei, 43 km à l'ouest de Chapa, altitude 1000–1200 m, 12 August 1926, *E. Poilane 12899* (P-00379162); Tonkin, Taai Wong Mo Shan and vicinity, Ton fa market, Ha-coi, September 1939, *W.T. Tsang 29570* (P-00379168).

## Acknowledgements

The authors are grateful to the editor and reviewer for helpful comments on our manuscript and to the curators of the herbaria HN, IBSC, K, KUN, NY, P and PE for providing specimen images or hosting our visits. This study was financially supported by a grant from the Southeast Asia Biodiversity Research Institute, Chinese Academy of Sciences (Y4ZK111B01).

## References

- Airy Shaw HK (1972) The Euphorbiaceae of Siam. *Kew Bulletin* 26(2): 191–363. <https://doi.org/10.2307/4117717>
- Angiosperm Phylogeny Group (APG) (2016) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society* 181(1): 1–20. <https://doi.org/10.1111/boj.12385>
- Beille L (1927) *Glochidion*. In: Lecomte H (Ed.) *Flore générale de l'Indo-Chine* 5. Masson, Paris, 608–628.
- Duocet Group (2016 onwards) Duocet Wiki of Plants. <http://duocet.ibiodiversity.net> [accessed 21 Oct. 2018]
- Govaerts R, Frodin DG, Radcliffe-Smith A, Carter S (2000) World checklist and bibliography of Euphorbiaceae (Pandaceae). Royal Botanic Garden, Kew, 1622 pp.
- Govaerts R, Dransfield J, Zona S, Hodel DR, Henderson A (2018) World Checklist of Phyllanthaceae. Facilitated by the Royal Botanic Gardens, Kew. <http://wccsp.science.kew.org> [accessed 21 Oct. 2018]
- Ho PH (2003) *An Illustrated Flora of Vietnam* 2. Nha Xuat Ban Tre, Tp. Ho Chi Minh, 951 pp.
- Hoffmann P, Kathriarachchi H, Wurdack KJ (2006) A phylogenetic classification of Phyllanthaceae (Malpighiales; Euphorbiaceae sensu lato). *Kew Bulletin* 61: 37–53. <https://www.jstor.org/stable/20443245>
- Jangid PP, Gupta S (2016) Systematic wood anatomy of the tribe Phyllanthae (Phyllanthaceae, Euphorbiaceae s.l.) from India: Implication in reinstatement of *Phyllanthus*, *Glochidion* and allies. *Nordic Journal of Botany* 34(4): 496–512. <https://doi.org/10.1111/njb.01072>
- Kathriarachchi H, Samuel R, Hoffmann P, Mlinarec J, Wurdack KJ, Ralimanana H, Stuessy TF, Chase MW (2006) Phylogenetics of tribe Phyllanthae (Phyllanthaceae; Euphorbiaceae sensu lato) based on nrITS and plastid *matK* DNA sequence data. *American Journal of Botany* 93(4): 637–655. <https://doi.org/10.3732/ajb.93.4.637>
- Kato M, Kawakita A (2017) *Obligate pollination mutualism*. Springer, Tokyo, 309 pp. <https://doi.org/10.1007/978-4-431-56532-1>
- Kuntze O (1891) *Revisio Generum Plantarum* 2. A. Felix, Leipzig, 377–1011.
- Li PT, Gilbert MG (2008) *Glochidion*. In: Wu ZY, Raven PH (Eds) *Flora of China* 11. Science Press & Missouri Botanical Garden Press, Beijing & St. Louis, 193–202.
- Loureiro J (1790) *Flora cochinchinensis* 2. Academicis, Ulyssipone, Lisboa, 744 pp.



- McNeill J, Barrie FR, Buck WR, Demoulin V, Greuter W, Hawksworth DL, Herendeen PS, Knapp S, Marhold K, Prado J, Prud'Homme Van Reine WF, Smith GF, Wiersema JH, Turland NJ (2012) International Code of Nomenclature for algae, fungi, and plants (Melbourne Code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011 [Regnum Vegetabile 154]. Gantner, Ruggell, 240 pp.
- Merrill ED (1935) A commentary on Loureiro's "Flora Cochinchinensis". Transactions of the American Philosophical Society, new series 24: 1–445. <https://doi.org/10.2307/1005470>
- Merrill ED (1936) Otto Kuntze's new genera and new species of Indo-China plants. Brittonia 2(3): 189–200. <https://doi.org/10.2307/2804861>
- Nguyen NT (2007) Taxonomy of Euphorbiaceae in Vietnam. Vietnam National University Publishers, Hanoi, 407 pp.
- Pruesapan K, Telford IR, Bruhl JJ, van Welzen PC (2012) Phylogeny and proposed circumscription of *Breynia*, *Sauropus* and *Synostemon* (Phyllanthaceae), based on chloroplast and nuclear DNA sequences. Australian Systematic Botany 25(5): 313–330. <https://doi.org/10.1071/SB11005>
- The Plant List (2013 onwards) Version 1.1. <http://www.theplantlist.org> [accessed 21 Oct. 2018]
- Thiers B (2018 onwards) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. The New York Botanical Garden, New York. <http://sweetgum.nybg.org/ih> [accessed 21 Oct. 2018]
- van Welzen PC (2007) *Glochidion*. In: Chayamarit K, van Welzen PC (Eds) Flora of Thailand 8(2). Royal Forest Department, Bangkok, 308–331.
- van Welzen PC, Pruesapan K, Telford IRH, Esser HJ, Bruhl JJ (2014) Phylogenetic reconstruction prompts taxonomic changes in *Sauropus*, *Synostemon* and *Breynia* (Phyllanthaceae tribe Phyllanthae). Blumea 59(2): 77–94. <https://doi.org/10.3767/000651914X684484>
- Webster GL (2014) Euphorbiaceae. In: Kubitzki K (Eds) Flowering Plants. Eudicots. The Families and Genera of Vascular Plants 11. Springer, Berlin, Heidelberg, 51–216. [https://doi.org/10.1007/978-3-642-39417-1\\_10](https://doi.org/10.1007/978-3-642-39417-1_10)
- Yao G, Zhang DX (2016) Pollen morphology of Chinese *Glochidion* (Phyllanthaceae) and its taxonomic implications. Nordic Journal of Botany 34(1): 102–110. <https://doi.org/10.1111/njb.00912>