RESEARCH ARTICLE



Two new records of the fern genus Coniogramme (Pteridaceae) from Vietnam

Caihong Wang¹, Wenli Yang², Junwen Zhao¹, Danke Zhang¹, Gangmin Zhang¹

I Laboratory of Systematic Evolution and Biogeography of Woody Plants, College of Nature Conservation, Beijing Forestry University, Beijing 100083, China 2 College of Landscape Architecture and Tourism, Hebei Agricultural University, Baoding 071000, China

Corresponding author: Gangmin Zhang (gary1967@bjfu.edu.cn)

Academic editor: Thais Almeida | Received 21 January 2019 | Accepted 22 March 2019 | Published 11 April 2019

Citation: Wang C, Yang W, Zhao J, Zhang D, Zhang G (2019) Two new records of the fern genus *Coniogramme* (Pteridaceae) from Vietnam. PhytoKeys 119: 137–142. https://doi.org/10.3897/phytokeys.119.33126

Abstract

Two new records of the fern genus *Coniogramme* Fée from Vietnam, *C. japonica* and *C. procera*, are presented. In addition, a key to recognising the species of *Coniogramme* in Vietnam is given in this paper.

Keywords

Cryptogrammoideae, taxonomy, Vietnam

Introduction

Coniogramme Fée, which belongs to the subfamily Cryptogrammoideae in the family Pteridaceae (Christenhusz et al. 2011; Zhang and Ranker 2013; PPG I 2016), is mainly distributed in the tropical and subtropical regions of Asia, extending south to Africa. It is characterised by its large habit with creeping rhizomes and 1–3 pinnate fronds with exindusiate sori borne along the lateral veins.

The genus was first monographed in its modern form by Hieronymus (1916), who accepted 17 species in the world. Dixit and Das (1979) recorded 13 taxa within the genus from India. In the course of the study of pteridophytes in China, some scholars (Ching 1930, 1974, 1982; Shing 1981; Kong 1982; Ching and Liu 1984;

Guo and Chen 2013) have published more than 30 new taxa of *Coniogramme*. Amongst them, Shing (1981) reported 27 new taxa and recognised 39 species and eight varieties in Flora Reipublicae Popularis Sinicae (Shing 1990). In the taxonomic keys, many species are distinguished only by the shape of pinnules, size of serrated teeth from pinnule margins and position of hydathodes at the top of the veins, but these traits are unstable and vary occasionally within normal populations and are difficult to use in practical identification. This taxonomic treatment has been described as the "inclusion of many erroneous new species entirely confusing the variation within species which is easily observable in the field" (Fraser-Jenkins et al. 2015). So far, the genus has long been one of the most problematic fern groups with respect to its specific definition. Fraser-Jenkins (2008) concluded that the taxonomy of *Coniogramme* is very complicated and has been confusing taxonomists in species circumscription.

Although many new taxa have been recorded recently, the fern diversity in Vietnam remains unclear (Lu et al. 2014). In revising the taxonomy of *Coniogramme*, the senior author was fortunate to visit Vietnam and examine the specimens deposited at HNU and HN. Two new records were discovered, i.e. *C. japonica* (Thunberg) Diels and *C. procera* Fée. This work is a contribution to the knowledge of fern flora in Vietnam.

Results

Coniogramme japonica (Thunb.) Diels (1899: 262) Figures 1, 3

Type. Japan. No exact location. C. P. Thunberg s.n. (UPS!).

Specimens examined. VIETNAM. Cao Bang Province: Ha Lang District, Dong Loan municipality, vicinities of Ban Lung and Lung Phuc, 22°46'N, 106°44'E, 500–600 m elev., 25 Nov 1998, *L. Averyanov et al. CBL 656* (HN). Bac Kan Province: Cho Don District, Ban Thi municipality, Phia Khao village, 22°17'03"N, 105°30'34"E, ca. 800 m elev., 5 Mar 2011, *N.Q. Hieu, N.T. Hiep, P.K. Loc, P.V. The, & N.T. Vinh CPC 1240* (HNU 017603, HNU 017604).

Taxonomic notes. This species is very unique in morphology, differing from other species in its anastomosing veins, which form 1–3 rows of areoles along each side of midrib and hydathodes not extending the base of short serrated teeth. Shing (1981, 1990) considered that pinnules of *Coniogramme japonica* are narrowly lanceolate with a cuneate or rounded-cuneate base and published a similar new species, *C. centrochinensis* Ching, whose pinnules were widely lanceolate with a rounded base. After specimen examination, combined with fieldwork, we found that the morphology of the pinnules was not stable and varied occasionally within normal populations. More research work should be undertaken to elucidate their phylogenetic relationship.

Distribution and habitat. *Coniogramme japonica* is distributed in China, Japan (including Ryukyu Islands), Korea and Vietnam (new record). The species usually grows in shady wet places at an elevation of about 100 to 2000 m.



Figure 1. *Coniogramme japonica* (Thunberg) Diels **A** one of the voucher specimens **B** portion of a pinnule, showing anastomosing veins **C** morphology and habitat.

Coniogramme procera Fée (1865: 22)

Figures 2, 3

Type. Nepal. April 1821. Wallich no 3 (K!).

Specimens examined. VIETNAM. Kon Tum Province: NW slopes of Ngoc Link mountain, 2380 m elev., 06 Mar 1995, *L. Averyanov et al. VH 519* (HN); W slope of Ngoc Link mountain, 1950 m elev., 10 May 1995, *L. Averyanov et al. VH 1290* (HN).

Taxonomic notes. This species is large and up to 1.8 m tall, differing from other species in its far more dissect laminae, basal pinnae having more than 10 pairs of pinnules, pinnules with rounded-truncate or truncate (sometimes slightly cordate) base and coarsely serrated margin and sori extending only to 1/2–2/3 of veins. Fraser-Jenkins (2008) reported that the species has a characteristically strong odour when the leaves were crushed or broken, similar to that of *Coniogramme fraxinea* (D.Don) Diels.

Distribution and habitat. *Coniogramme procera* was once recorded being distributed in Vietnam in Flora Reipublicae Popularis Sinicae (Shing 1990) and this was followed by Flora of China (Zhang and Ranker 2013). After specimen examination, we found that there were no accounts of *C. procera* in K, BM, P, PE and other major herbaria and we wondered about the basis of this recognition. In addition, the species has never been recorded in the literature on flora of Vietnam (Tardieu-Blot and Christensen 1941;



Figure 2. *Coniogramme procera* Fée. **A** One of the voucher specimens **B** Portion of a pinnule, showing a truncate base **C** Morphology and habitat.



Figure 3. Distribution of *C. japonica* (dots) and *C. procera* (star) in Vietnam.

Pham 1991; Phan 2010), including the recently updated checklist (Phan 2010). *C. procera* is therefore confirmed to be distributed in central Vietnam for the first time. It is also distributed in Bhutan, China, India, Myanmar, Nepal, Philippines and Thailand. The species usually grows by streams in woodlands at a high elevation, about 1400 to 3600 m.

Based on previous literature (Tardieu-Blot and Christensen 1941; Pham 1991; Phan 2010), along with our specimen identification work at HNU, HN and K, six taxa of *Coniogramme* were recognised in Vietnam, namely *C. fraxinea* (D.Don) Diels, *C. intermedia* Hieron., *C. macrophylla* (Blume) Hieron., *C. petelotii* Tardieu, *C. japonica* (Thunberg) Diels and *C. procera* Fée. Their main differences in character were illustrated in the following key:

Key to the species of Coniogramme in Vietnam

1	Veins anastomosing to form 1 or 2 continuous rows of areoles on each side of
	midrib <i>C. japonica</i>
_	Veins all free
2	Pinnule margins entire
_	Pinnule margins serrate
3	Hydathodes extending to cartilaginous lamina margin C. macrophylla
_	Hydathodes spindle-shaped, not extending to lamina margin4
4	Base of pinnules rounded or slightly cordate
_	Base of pinnules cuneate or rounded-cuneate
5	Basal pinnae having more than 10 pairs of pinnules; pinnules broadly lanceolate,
	base rounded-truncate or truncate (sometimes slightly cordate) C. procera
_	Basal pinnae having 2-3 pairs of pinnules; pinnules lanceolate, base rounded
	to rounded-cuneate C. intermedia

Acknowledgements

We are grateful to Professor Phan Ke Loc and the curator of the HN herbarium for providing access to the material in their care. The study was supported by the National Natural Science Foundation of China (Grant no. 31770221). We thank LetPub (http://www.letpub.com) for its linguistic assistance during the preparation of this manuscript.

References

Ching RC (1930) The studies of Chinese ferns I. Sinensia 1: 43–56. Ching RC (1974) Flora Tsinlingensis (2). Science Press, Beijing, 246 pp. Ching RC (1982) Notes on some Chinese ferns. Acta Phytotaxonomica Sinica 20(2): 235–236. Ching RC, Liu ZY (1984) New ferns from Jinfoshan, Nanchuan, Sichuan (II). Bulletin of

Botanical Research 4(3): 1-32.

- Christenhusz MCM, Schneider H, Zhang XC (2011) A linear sequence of extant lycophytes and ferns. Phytotaxa 19(1): 7–54. https://doi.org/10.11646/phytotaxa.19.1.2
- Diels L (1899) Polypodiaceae. In: Engler A, Prantl K (Eds) Die Natürlichen Pflanzenfamilien, vol. 1(4). Wilhelm Engelmann, Leipzig, 139–339.
- Dixit RD, Das A (1979) The genus *Coniogramme* Fée in India. Proceedings of the Indian Academy of Sciences 88 B, II: 253–268.
- Fée ALA (1865) Mémoires sur la famille des fougères 10. Description de Fougères exotiques rares ou nouvelles. Mémoirs de la Société des Sciences Naturelles de Strasbourg 6(1): 21–22.
- Fraser-Jenkins CR (2008) Taxonomic Revision of three hundred Indian subcontinental Pteridophytes with a revised census-list, a new picture of fern-taxonomy and nomenclature in the Indian subcontinent. Bishen Singh Mahendra Pal Singh, Dehra Dun, 685 pp.
- Fraser-Jenkins CR, Kandel DR, Pariyar S (2015) Ferns and Fern-allies of Nepal (1). Godawari, Lalitpur, Nepal, 492 pp.
- Guo XS, Chen JP (2013) *Coniogramme bashanensis* (Pteridaceae), a new fern species from Shaanxi, China. Novon 22(3): 292–296. https://doi.org/10.3417/2009145
- Hieronymus G (1916) Über die Gattung *Coniogramme* und ihre Arten. Hedwigia 57: 265–328.
- Kong XX (1982) New ferns from Sichuan, China. Acta Botanica Yunnanica 4(4): 337–343.
- Lu NT, Zhang L, Zhang LB (2014) Ten new records of *Polystichum* (Dryopteridaceae) for Vietnam and lectotypification of *Polystichum atroviridissimum* and *P. fimbriatum*. Phytotaxa 164(2): 115–123. https://doi.org/10.11646/phytotaxa.164.2.5
- Pham HH (1991) An illustrated flora of Vietnam, vol. 1. Montreal, 42–261.
- Phan KL (2010) The updated checklist of the fern flora of Vietnam following the classification scheme of A. Smith et al. (2006). Journal of Fairylake Botanical Garden 9(3): 1–13.
- PPG The Pteridophyte Phylogeny Group I (2016) A community-derived classification for extant lycophytes and ferns. Journal of Systematics and Evolution 54(6): 563–603. https:// doi.org/10.1111/jse.12229
- Shing KH (1981) New taxa of the genus *Coniogramme* Fée from China. Acta Botanica Yunnanica 3(2): 217–238.
- Shing KH (1990) Coniogramme. In: Ching RC, Shing KH (Eds) Flora Reipublicae Popularis Sinicae 3 (1). Science Press, Beijing, 228–274.
- Smith AR, Pryer KM, Schuettpelz E, Korall P, Schneider H, Wolf PG (2006) A classification for extant ferns. Taxon 55(3): 705–731. https://doi.org/10.2307/25065646
- Tardieu-Blot ML, Christensen C (1941) Cryptogames Vascularies. In: Lecomte H, Gagnepain F (Eds) Flore générale de l'Indo-Chine, vol. 7(2). Masson & Cie, Paris.
- Zhang GM, Ranker TA (2013) *Coniogramme*. In: Wu ZY, Raven PH (Eds) Flora of China Vol. 2–3. Science Press, Beijing, 171–178.