

# Two new species of *Cylindrolobus* (Orchidaceae) from the eastern Himalayas

Ji-Dong Ya<sup>1</sup>, Xiao-Hua Jin<sup>2</sup>, Cheng Liu<sup>1</sup>

**1** Germplasm Bank of Wild Species, Kunming Institute of Botany, Chinese Academy of Sciences, Lanhei Road 132, Heilongtan, Kunming, Yunnan, 650201, China **2** State Key Laboratory of Systematic and Evolutionary Botany and Herbarium, Institute of Botany, Chinese Academy of Sciences, Nanxincun20, Xiangshan, Beijing, 100093, China

Corresponding author: Xiao-Hua Jin ([xiaohuajin@ibcas.ac.cn](mailto:xiaohuajin@ibcas.ac.cn))

Academic editor: Cai Jie | Received 27 February 2019 | Accepted 18 May 2019 | Published 29 August 2019

**Citation:** Ya J-D, Jin X-H, Liu C (2019) Two new species of *Cylindrolobus* (Orchidaceae) from the eastern Himalayas. In: Cai J, Yu W-B, Zhang T, Li D-Z (Eds) Revealing of the plant diversity in China's biodiversity hotspots. PhytoKeys 130: 107–113. <https://doi.org/10.3897/phytokeys.130.33989>

## Abstract

Two new species, *Cylindrolobus motuoensis* and *C. glabriflorus* (Orchidaceae), from Southwestern China and north of Myanmar are described and illustrated with detailed photos. *Cylindrolobus motuoensis* is morphologically similar to *C. gloensis* and *C. foetidus*, but can be distinguished from them by having amplexicaul sterile bracts, dark red floral bracts, white flowers, falcate-lanceolate lateral sepals and central keel of lip running from base to the tip of mid-lobe. *Cylindrolobus glabriflorus* is similar to *C. hegdei* and *C. tenuicaulis* but differs from them by having longer and wider leaves, obovate bracts, and the reddish brown central papillate keel of lip.

## Keywords

Taxonomy, Orchidaceae, *Cylindrolobus*, new species, China

## Introduction

The genus *Cylindrolobus* Blume consists of 60–70 species, distributed in the tropical region from East Himalaya, China, Southeast Asia to New Guinea (Chen et al. 2009, Ormerod 2014, Ng et al. 2018). *Cylindrolobus* was originally proposed as a section of *Eria* Lindl., Pridgeon et al. (2005) suggested that *Cylindrolobus* should be subsumed into *Callostylis* Blume. Recent molecular and morphological studies suggest that *Cylindrolobus* is a distinct genus, characterized by a multi-noded stem with leaves at apex, short inflorescences with one to several flowers and conspicuous and colorful bracts (Chen et al. 2009, Ng et al. 2018).

## Materials and methods

Living plants were collected from Xizang (Tibet) Autonomous Region of China and north of Myanmar during the botanical expeditions in 2016 and in 2018. The shapes, colors of flowers and other details of the plants observed, measured and photographed, as well as specimens collected, were based on living materials from 2017 to 2019. Morphological photographs of the lip, column and pollinia were taken using an Olympus SZX16. All voucher specimens were deposited in KUN (Herbarium of Kunming Institute of Botany, Chinese Academy of Sciences).

## Taxonomic treatment

***Cylindrolobus motuoensis* X.H.Jin & J.D.Ya, sp. nov.**

urn:lsid:ipni.org:names:60479344-2


Figures 1, 3A

**Diagnosis.** *Cylindrolobus motuoensis* is similar to *C. gloensis* (Ormerod & Agrawala) Schuit., Y.P. Ng & H.A. Pedersen, and *C. foetidus* (Aver.) Schuit., Y.P. Ng & H.A. Pedersen in terms of morphological structure and shape of the flowers (Hu et al. 2010, Agrawala and Ormerod 2014, Ng et al. 2018). The new species can be distinguished from *C. gloensis* by the smaller flowers, elliptic and concave bracts, and ovate lip with three keels, mid-lobe thickened and papillate on margin. The new species can be distinguished from *C. foetidus* with longer and wider leaves, dark red and elliptic floral bracts, white flowers and falcate-lanceolate lateral sepals.

**Type.** CHINA. Xizang Autonomous Region: Motuo, subtropical evergreen broad-leaved forest, alt. 2000 m, 26 Feb 2017, *Ji-Dong Ya, Cheng Liu, Hua-Jie He* 17HT0073 (holotype: KUN!).

**Additional specimen examined.** CHINA. Xizang Autonomous Region: Motuo, subtropical, evergreen broad-leaved forest, 26 Feb 2017, *Xiao-Hua Jin, Ji-Dong Ya*, 17HT1088 (paratype: KUN!)

**Description.** Epiphytic herb. Roots terete, slender, pubescent, ca. 1.0–1.5 mm thick. Rhizome creeping, to 3–4 mm thick. Stem terete, slender, 3(2) leaved apically, covered by close-fitting sheaths, 18–24 cm long, 3–6 mm thick. Leaves ligulate-lanceolate, acuminate, 10–13 cm long, 1.5–2.0 cm wide. Inflorescences axillary, pubescent, borne on near the apical of the stem, 2–3 cm long, 2 flowered; peduncle 1.0–1.5 cm long; 2 sterile bracts, smaller, amplexicaul; rachis 0.2 cm long, floral bracts dark red, elliptic, acute, concave, sparsely tomentum, 7 mm long, 3 mm wide. Flowers white, sepal externally with brown tomentum, peduncle and ovary ca. 1.0–1.5 cm long, densely brown tomentum. Dorsal sepal lanceolate, acute, 5 veined, 11 mm long, 4 mm wide; lateral sepals falcate-lanceolate, acute, 5 veined, 9 mm long, 5 mm wide, base adnate to column foot form a subglobose and obtuse mentum; petals lanceolate, slightly oblique, acute, 3 veined, 10 mm long, 3 mm wide; labellum ovate in outline, 3-lobed, base hinged to the apex of the column foot, apex obtuse and emarginate,



**Figure 1.** *Cylindrolobus motuoensis* X.H.Jin & J.D.Ya. **A** Plant **B** Inflorescence **C** lateral view of flower **D** ventral view of flower **E** front view of flower **F** adaxial sepals and petals **G** abaxial sepals and petals **H** lateral view of column and lip **I** lateral view of column **J** front view of column **K** front view of labellum **L** lateral view of labellum (rip cutting) **M** adaxial bract **N** abaxial bract **O** polar view of pollinarium **P** ventral view of pollinarium **Q** lateral view of pollinarium **R** adaxial anther cap **S** abaxial anther cap (Photographed by J.-D. Ya).

curved, c. 6 mm long, 3 mm wide; lateral lobes suberect, subovate, apex slightly introvert; mid-lobe ligulate, ca. 3 × 3 mm, thickened and papillate on margin, apex emarginate; disk with 3 keels, central keel longitudinal thickened, with orange papilla, running from base to the tip of mid-lobe, lateral keels glabrous, running from base to middle of mid-lobe. Column semiterete, ca. 4 mm long, broad winged at ventrally; foot incurved, ca. 3.5 mm. Anther cap ovate, ca. 1 mm × 1 mm, pollinia 8, yellowish white, compressed rectangular, anterior ca. 0.5 mm × 0.4 mm × 0.2 mm, posterior 4 smaller. Fl. February-March.

**Etymology.** The new species is named after Motuo, Xizang Autonomous Region of China, where it was discovered in a subtropical evergreen broad-leaved forest.


**Vernacular name.** Mo Tuo Zhu Lan (墨脱柱兰).

***Cylindrolobus glabriflorus* X.H.Jin & J.D.Ya, sp. nov.**

urn:lsid:ipni.org:names:60479345-2

Figures 2, 3B

**Diagnosis.** *Cylindrolobus glabriflorus* is similar to *C. hegdei* (Agrawala & H. J. Chowdhery) A. N. Rao and *C. tenuicaulis* (S. C. Chen & Z. H. Tsi) S. C. Chen & J. J. Wood



**Figure 2.** *Cyndrolobus glabriflorus* X.H.Jin & J.D.Ya. **A** Plant **B** inflorescence **C** ventral view of flower **D** lateral view of flower **E** adaxial sepals and petals **F** abaxial sepals and petals **G** lateral view of column and lip **H** front view of labellum **I** lateral view of labellum **J** ventral view of labellum **K** lateral view of column **L** front view of column **M** adaxial bract **N** abaxial bract **O** lateral view of pollinarium **P** polar view of pollinarium **Q** abaxial anther cap **R** adaxial anther cap (Photographed by J.D. Ya).

(Agrawala and Chowdhery 2008, Chen et al. 2009). The new species can be distinguished from *C. hegdei* by the longer and wider leaves, glabrous inflorescence, smaller and yellow flowers, yellowish green and obovate bracts and reddish brown central papillate keel of lip. It differs from *C. tenuicaulis* by having longer and wider leaves, longer inflorescence, yellowish green and obovate bracts, longer lip with three calli, lateral margins of mid-lobe thickened and erect.

**Type.** MYANMAR. Kachin State: Putao Township, Hponkanrazi Wildlife Sanctuary, subtropical, evergreen, broad-leaved, humid montane forest, alt. 2200 m, 12 Apr 2018, *Xiao-Hua Jin, Ji-Dong Ya* 18HT1618 (holotype: KUN!).

**Additional specimen examined.** CHINA. Xizang Autonomous Region: Motuo, subtropical, evergreen broad-leaved forest, alt. 1796 m, 2 Apr 2019, *Ji-Dong Ya, Cheng Liu*, 18HT2586 (paratype: KUN!). CHINA. Xizang Autonomous Region: Motuo, subtropical, evergreen broad-leaved forest, alt. 1750 m, 7 Apr 2018, *Hong Jiang, Weiping Zhang, Zhou-dong Han* 07336 (paratype: YAF!).

**Description.** Epiphytic herb. Roots terete, slender, pubescent, ca. 0.8–1.0 mm thick. Rhizome inconspicuous. Stem clustered, terete, slender, 4-leaved apically, covered by close-fitting sheaths, 13–25 cm long, 3–4 mm thick. Leaves lanceolate, acuminate, 4.5–6.5 cm long, 0.8–1.5 cm wide. Inflorescences axillary, glabrous, arising from the apical of the stem, 1.3 cm long, 2-flowered; peduncle 0.5 cm long; sterile bracts



**Figure 3.** Holotype. **A** *Cylindrolobus motuoensis* X.H.Jin & J.D.Ya. **B** *Cylindrolobus glabriflorus* X.H.Jin & J.D.Ya.

1–2, smaller. Flowers yellow, sepal externally reddish yellow, 7–8mm long; floral bracts yellowish green with red brown edges, obovate, mucronate, concave, glabrous, 4 mm long, 2.5 mm wide; peduncle and ovary ca. 4–6 mm long. Dorsal sepal lanceolate, obtuse, 3 veined, 6 mm long, 2 mm wide; lateral sepals falcate-ovate, obtuse, 3-veined, 5 mm long, 3.5 mm wide, base adnate to column foot form a subglobose and obtuse mentum; petals oblong-ovate, slightly oblique at base, obtuse, 1 veined, 5 mm long, 2 mm wide; labellum oblong in outline, 3-lobed, base hinged to the apex of the column foot, apex emarginate, ca.3.5 mm long, 1.5 mm wide; lateral lobes suberect, subovate, apex slightly introvert, disk with 2 reddish brown calli; mid-lobe sub-square, ca. 3.3 mm × 2 mm, lateral margins thickened and erect, with a central papillate keel, reddish brown, ca. 0.5 mm high; apex emarginate. Column semiterete, ca. 3.5 mm long, narrow winged at ventrally; rostellum triangle, 0.2 mm, a hook-like protrusion under the stigma; foot incurved, ca. 4 mm; cap subrotund, ca. 0.6 mm × 0.6 mm, papillate and protrude in front; pollinia 8, yellowish white, compressed subrotund from the lateral view, anterior 4 ca. 0.2 mm × 0.2mm × 0.1mm, posterior 4 smaller. Fl. April-May.

**Etymology.** The specific epithet “*glabriflorus*” refers to glabrous flowers of this new species.

**Vernacular name.** Zhong Mian Zhu Lan (中缅柱兰).

## Discussion

The generic delimitation of Podochileae Pfitzer has long been confused (Pridgeon et al. 2005, Ng et al. 2018). Recent molecular and morphological evidence reveals that *Eria* s.l. should be split into 21 genera, including *Cylindrolobus* which is morphologically characterized by fleshy and elongate stems, lateral inflorescences, conspicuous and colorful bracts.

These two new species are distributed in a narrow area in the border region between China and Myanmar. We found there are many populations of each species and abundant individuals per population during our botanical expeditions. As the habitats of these two species are in a remote location and the border area has restricted access, the effect of human interference and climate change on them is little known. For the time being, these two species are considered as Least Concern (LC) according to current information on these species and the IUCN Red List category (IUCN 2012).

Located at the margin of Qinghai-Tibet Plateau, Motuo is famous for its vertical vegetation system from tropical forest to permanent glacier with elevation approximately 7000 m, which allow the thriving and diversification of plant diversity. In addition, there are many biodiversity hotspots in the eastern Himalayas, e.g. the north of Myanmar, however the species diversity of this region is poorly known. Hence, the species diversity of these border regions requires solid investigations (Liu et al., 2019) and timely conservation action plans in this region, in order to mitigate increasing anthropogenic disturbance and destruction.

The two new species described here increased the members of *Cylindrolobus* in China to seven species (Chen et al. 2009, Hu et al. 2010, Liu et al. 2013, Ng et al. 2018), the key to Chinese species of *Cylindrolobus* are developed here.

### Key to Chinese species of *Cylindrolobus*

- |   |  |                        |
|---|--|------------------------|
| 1 | Stems stout, clavate.....  | 2                      |
| – | Stems slender, terete.....   | 4                      |
| 2 | Lip yellow.....  | <i>C. cristatus</i>    |
| – | Lip lateral lobes with purple edges, mid-lobe with lighter purple edges and patch..... | 3                      |
| 3 | Inflorescence glabrous, the lateral lobes bigger than mid-lobe.....                    | <i>C. clavicaulis</i>  |
| – | Inflorescence pubescent, the lateral lobes smaller than mid-lobe .....                 | <i>C. marginatus</i>   |
| 4 | Inflorescence glabrous.....  | 5                      |
| – | Inflorescence pubescent .....  | 6                      |
| 5 | lip mid-lobe thickened and papillate on margin.....                                    | <i>C. tenuicaulis</i>  |
| – | lip mid-lobe not thickened and smoothly on margin .....                                | <i>C. glabriflorus</i> |
| 6 | Flowers yellow .....   | <i>C. foetidus</i>     |
| – | Flowers white.....   | <i>C. motuoensis</i>   |



## Acknowledgements

We are grateful to Lian-Yi Li for his help with image processing; we convey our special thanks to Forestry Bureau of Linzhi Prefecture for their kind help in the field. This research was supported by grants from the Chinese Scholarship Council (Grant No. 201704910424), the National Natural Science Foundation of China (Grant No. 31670194), National Science & Technology Infrastructure and the Large-scale Scientific Facilities of the Chinese Academy of Sciences (2017-LSFGBOWS-02), and Southeast Asia Biodiversity Research Institute, Chinese Academy of Sciences (Y4Z-K111B01 to X.H.J.).

## References

- Agrawala DK, Chowdhery HJ (2008) A new species of *Eria* Lindl. (Orchidaceae) from Arunachal Pradesh, India. *Phytotaxonomy* 8: 8–12.
- Agrawala DK, Ormerod P (2014) A new species of *Eria* (Orchidaceae) from India under the section *Cylindrolobus*. *Taiwania* 59(3): 206–209. <https://doi.org/10.6165/tai.2014.59.206>
- Chen SC, Lou YB, Vermeulen JJ (2009) *Cylindrolobus* Blume. In: Wu ZY, Raven PR, Hong DY (Eds) *Flora of China* (Vol. 25). Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 249–250.
- Hu H, Jin XH, Sun Q, Sun XG (2010) A new record of Orchidaceae from China. *Redai Yaredai Zhiwu Xuebao* 18: 401–402.
- IUCN (2012) *IUCN Red List Categories and Criteria*, Version 3.1. (2<sup>nd</sup> edn). IUCN, Gland.
- Liu JJ, Baeten L, Verheyen K (2019) Biodiversity on International Borders Requires Solid Inventories. *Bioscience* 69(5): 322–323. <https://doi.org/10.1093/biosci/biz029>
- Liu Q, Li JW, Yin JT (2013) *Eria clavicaulis*, a new record of Orchidaceae from Yunnan, China. *Guihuia* 33(1): 70–71.
- Ng YP, Schuiteman A, Pedersen HÆ, Petersen G, Watthana S, Seberg O, Pridgeon AM, Cribb PJ, Chase MW, NG YP (2018) Phylogenetics and systematics of *Eria* and related genera (Orchidaceae: Podochileae). *Botanical Journal of the Linnean Society* 186(2): 179–201. <https://doi.org/10.1093/botlinnean/box088>
- Ormerod P (2014) A Synopsis of *Eria* Lindl. Section *Cylindrolobus* (Blume) Lindl. (Orchidaceae: Eriinae) in Malesia. *Harvard Papers in Botany* 19(1): 77–95. <https://doi.org/10.3100/hpib.v19iss1.2014.n6>
- Pridgeon AM, Cribb PJ, Chase MW, Rasmussen FN (2005) Tribe Podochileae. In: *Genera Orchidacearum* 4, Epidendroideae (part one). Oxford University Press, Oxford, 529–596.