

Two new species of *Fargesia* (Poaceae, Bambusoideae) from southwestern China

Xia-Ying Ye^{1,2}, Yu-Xiao Zhang³, De-Zhu Li²

1 Agronomy and Life Science Department, Zhaotong University, Zhaotong, Yunnan 657000, China **2** Germplasm Bank of Wild Species, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, Yunnan 650201, China **3** Yunnan Academy of Biodiversity, Southwest Forestry University, Kunming, Yunnan 650224, China

Corresponding author: De-Zhu Li (dzl@mail.kib.ac.cn)

Academic editor: E. Ruiz-Sanchez | Received 18 September 2020 | Accepted 29 October 2020 | Published 10 December 2020

Citation: Ye X-Y, Zhang Y-X, Li D-Z (2020) Two new species of *Fargesia* (Poaceae, Bambusoideae) from southwestern China. *PhytoKeys* 170: 25–37. <https://doi.org/10.3897/phytokeys.170.58780>

Abstract

Two new species of *Fargesia*, one from Xizang (Tibet) and one from Yunnan, China, are described and illustrated. *Fargesia viridis* D.Z. Li & X.Y. Ye is characterized by its densely white powder, nearly solid internodes, yellow setose sheath scar and culm sheaths, and 4–6 leaves of large size. *Fargesia purpurea* D.Z. Li & X.Y. Ye has thinner culms (0.5–1.4 cm in diameter), a ring of 4–5 mm tall brown setae below nodes, fewer branches, glabrous sheath scar and culm sheaths, differentiated from the related species.

Keywords

Fargesia, new species, southwestern China, taxonomy, temperate woody bamboos

Introduction

Tribe Arundinarieae, i.e. the temperate woody bamboos, is one of the three tribes of the subfamily Bambusoideae (Poaceae), containing approximately 581 species in 31 genera (Bamboo Phylogeny Group 2012; Clark et al. 2015; Clark and Oliveira 2018). These bamboos are distributed primarily in the temperate to subtropical zones of the Northern Hemisphere, with nearly 90% of species distributed in East Asia (Ohrnberger 1999; Li et al. 2006).

Among the 31 genera, *Fargesia* Franchet is the largest one, consisting of more than 90 species (Li et al. 2006; Yi et al. 2008), out of which, 85 species occur in China and

83 taxa are endemic to the country (Vorontsova et al. 2016). The *Fargesia* species are mainly distributed in temperate mountain areas (alt. 800–4300 m) of East Asia (Keng 1987; Yi 1988; Ohrnberger 1999; Li et al. 2006; Vorontsova et al. 2016). This group is especially common and diverse in the high elevation ecosystem of southwest China where they have undergone rapid diversification associated with the orogeny of the Hengduan mountains (Ye et al. 2019).

Fargesia is characterized by the presence of short-necked pachymorph rhizomes (usually < 20 cm), unicaespitose clumps, 7–15 branches at mid-culm nodes, semelaucent inflorescence, racemose to paniculate, compressed or open, with 3 stamens (Li et al. 2006). Although reproductive features are important for bamboo classification, vegetative morphological characters are usually used to distinguish species due to long flowering cycles (Janzen 1976; Zhang and Ren 2016). Based on morphological characters of buds and culm sheaths, Yi (1988) divided the genus *Fargesia* into two sections, *F.* sect. *Ampullares* Yi and *F.* sect. *Fargesia* (Keng and Wang 1996). The section *Ampullares* is distinguished by compound buds consisting of multiple distinct buds and deciduous culm sheaths. The section *Fargesia* is characterized by compound buds composed of several obscure buds and late deciduous or persistent culm sheaths, and contains four series, namely, ser. *Murielae* Yi, ser. *Fargesia* Yi, ser. *Angustissimae* Yi and ser. *Yunnanenses* Yi. The series *Murielae* has oblong or narrowly elliptical culm sheaths, with rounded apex, as wide as the base, while in the latter three series, the shape of culm sheaths is different and featured as narrowly triangular or narrowly orbicular-triangular, apex triangular or linear, much narrower than the base. Moreover, the texture and length of culm sheaths are varied in these three series. For example, the culm sheaths of ser. *Fargesia* and ser. *Angustissimae* are longer than internodes, but shorter or equal in ser. *Yunnanenses*. The culm sheaths of ser. *Fargesia* are apically leathery and narrowed for distal ca. 1/5 of length but apically thickly papery and narrowed for distal ca. 1/3–1/2 of length for species of ser. *Angustissimae*.

Although flowering is not frequent in this genus, it shows considerable diversity in vegetative morphology and many new species continue to be described (Yi 2000a, Yi 2000b, Yi 2000c; Yi et al. 2005, 2006; Yi et al. 2007; Yang and Yi 2013a, Yang and Yi 2013b) from southwest China. During floristic surveys of bamboos between 2015 and 2018, the authors collected vast specimens of *Fargesia* from southwest China. After scrutiny of the data available (Keng and Wang 1996; Li et al. 2006; Yi et al. 2008; Vorontsova et al. 2016), we found that several specimens could not be assigned to any described species. Here, we described two new species of *Fargesia* based on morphological comparison and the phylogenetic results (Ye et al. 2019).

Materials and methods

Observation and measurement of morphological characters were undertaken using living plants in the field and specimens in the lab. Morphological features of related species were obtained from specimens and literature (Keng and Wang 1996; Li et al. 2006; Yi et al. 2008).

Taxonomic treatment

Fargesia viridis D.Z. Li & X.Y. Ye, sp. nov.

urn:lsid:ipni.org:names:77213334-1

Figs 1, 2

Diagnosis. *Fargesia viridis* D.Z. Li & X.Y. Ye resembles *F. frigidis* Yi, *F. zayuensis* Yi and *F. similaris* Hsueh & Yi, but can be distinguished from *F. frigidis* by thinner and glabrous culm, more leaves on the ultimate branch, longer leaf sheath and large leaf blade, from *F. zayuensis* by shorter and thinner culm, solid internode, more leaves on the ultimate branch and broader leaf blade, and from *F. similaris* by solid internode, prominent sheath scar, setose culm sheath, glabrous petiole, more leaf number and larger leaf blade.

Type. CHINA, Yunnan, Gongshan County, along the road to Dulongjiang Town, 27°51'28"N, 98°26'46"E, 2667 m alt., 1 September 2015, X.Y.Ye YXY272 (holotype & isotype: KUN!).

Description. Rhizomes pachymorph, rhizome neck 3–6 cm long, 1–1.6 cm in diameter, solid. Culms 2–3 (4) m tall, pluricaespitose, 0.6–1.2 cm in diameter; internodes terete, 16–22 (30) cm long, densely white powdery and black when culms old, glabrous, nearly solid; nodes with weakly prominent supra-nodal ridge; sheath scar prominent, initially brown setose, with persistent remains of sheath base. Branches 8–10, fascicular, open; buds oblong, margins yellow-brown ciliolate. Culm sheaths persistent or tardily deciduous, leathery, narrowly rounded, 1/3 as long as internodes, yellow setose, densely at base and readily deciduous, longitudinal ribs prominent, margins yellow ciliolate, apex asymmetrical; auricles absent; oral setae absent or 1–2, ca. 2 mm long; ligule concave or truncate, ca. 1 mm tall, glabrous, fissured; blades erect or reflexed, linear-lanceolate, glabrous, narrower than the apex of culm sheath. Foliage leaves 4–6 per ultimate branch; sheath 3–4 cm long, glabrous, purple, margins ciliolate; auricles and oral setae absent; ligule truncate, ca. 1 mm tall; petiole 1–3 mm long, glabrous; blade lanceolate, 4–9 × 0.7–1.4 cm, glabrous, base broadly cuneate, secondary veins 2–3 pairs, transverse veins conspicuous, margins serrate. Inflorescence unknown.

Phenology. New shoots July to August.

Etymology. The specific epithet refers to the beautiful color of leaf blade.

Vernacular name. Cù Lǜ Jiàn Zhú (Chinese pronunciation); 翠绿箭竹 (Chinese name).

Distribution and habitat. *Fargesia viridis* is only known from the type locality, the Dulongjiang Town. It occurs along the stream and grows as pure bamboo forest or under the evergreen broadleaved forest at an elevation of 2600–2800 m alt.

Notes. Morphological comparisons between *Fargesia viridis* and the related species were provided in Table 1. Other four species of this genus were found in the Dulongjiang Town, i.e., *F. declivis* Yi, *F. sagittatinea* Yi, *F. acuticontracta* Yi and *F. praecipua* Yi, with this new species being easily distinguished from the other species in this region by its shorter and thinner culms, solid internodes (except *F. acuticontracta*), and shorter culm sheath (only 1/3 as long as internode).




Figure 1. *Fargesia viridis* D.Z. Li & X.Y. Ye **A** habitat **B** individual **C** rhizome **D** culm showing solid and nearly solid internodes **E** culm bud and sheath scar with yellow setose **F** young culms with culm sheaths **G** culm sheath showing densely setose at base and oral characters **H** branchlet.



Figure 2. *Fargesia viridis* D.Z. Li & X.Y. Ye **A** branchlet **B** rhizome **C** node, showing branches and sheath scar with setose **D** young culm with culm sheaths **E, F** culm leaves showing sheath and densely setose at base **G** culm buds.

***Fargesia purpurea* D.Z. Li & X.Y. Ye, sp. nov.**

urn:lsid:ipni.org:names:77213335-1

Figs 3–5

Diagnosis. *Fargesia purpurea* D.Z. Li & X.Y. Ye resembles *F. pauciflora* (Keng) Yi and *F. brevipedis* Yi, but can be distinguished from the former by thinner and taller culms,

Table 1. Morphological comparison of *Fargesia viridis* and its related species.

Characters	<i>Fargesia viridis</i>	<i>Fargesia frigidis</i>	<i>Fargesia zayuensis</i>	<i>Fargesia similaris</i>
Culm height	2–3 (4) m	1.5–3.5 m	6 m	Shrubby
Culm diameter	0.6–1.2 cm	1–1.7 cm	0.8–1.5 cm	0.8–1.2 cm
Internode	16–22 (30) cm long, densely white powdery, glabrous, nearly solid	22–24 cm long, initially densely white waxy and white-gray setose below nodes, glabrescent, nearly solid	25–35 cm long, initially sparsely white powdery; hollow, wall 1.5–2 mm thick	9.5–18.2 cm long, white or black powdery below nodes, wall 2–3 mm thick, cavity filled with lamellate pith
Branch complement	8–10	4–13	5–10	3–8(15)
Sheath scar	Prominent, initially yellow setose, with persistent remains of sheath base	Very prominent, woody	Prominent	Weakly prominent
Culm sheath	Persistent or tardily deciduous, yellow setose and densely at base, readily deciduous, longitudinal ribs prominent, margins yellow ciliate, apex asymmetrical	Gradually deciduous to persistent, very sparsely appressed light yellow setulose, upper margins yellow–brown ciliate initially, longitudinal ribs conspicuous, apex asymmetrical	Gradually deciduous, abaxially slightly gray-brown setulose, margins brown ciliate or not	Glabrous, margins densely ciliate, apex slightly white powdery
Culm sheath oral setae	Absent or 1–2, 2 mm long	Absent	Readily deciduous	Absent or 1–3
Culm sheath ligule	Concave or truncate, ca. 1 mm	Convex or truncate, 1–1.5 mm, glabrous	Truncate, ca. 1 mm	Truncate, ca. 1 mm
Culm sheath blade	Erect or reflexed, triangular or linear-lanceolate	Reflexed, readily deciduous, triangular to linear-lanceolate	Readily deciduous, reflexed, rarely erect, linear-lanceolate	Erect, triangular-conical, glabrous
Leaf number of the ultimate branch	4–6	1–4	1–3	2–4
Leaf sheath	3–4 cm long, glabrous	1.5–2 cm long, glabrous	3–4 cm, glabrous	Glabrous or with white pubescent margins
Leaf oral setae	Absent	Absent or sometimes few	Absent	2–6, 2–4 mm long, yellow-brown or gray
Leaf ligule	Truncate, ca. 1 mm	Inclined- truncate, ca. 0.4 mm	Truncate, glabrous	Truncate, ca. 1 mm
Petiole	1–3 mm long	1 mm long	1 mm long	Sparely gray-white pubescent
Leaf blade	4–9 × 0.7–1.4 cm, glabrous, secondary veins 2–3 pairs	2.3–5.2 × 0.45–0.7 cm, glabrous, secondary veins 2 or 3 pairs	5–8.5 × 0.4–0.6 cm, glabrous, secondary veins 2 pairs	1.3–6.5 × 0.4–0.6 cm, glabrous or abaxially white-gray pubescent, secondary veins 2- or 3 paired
Habitat	Along the stream or under the evergreen broadleaved forest at the altitude of 2600–2800 m, northwest, Yunnan.	On the shady slope of barren hills at 3100–3700 m, west Yunnan.	Under the <i>Pinus</i> or broadleaved forest, 2500–3000 m, Zayu, Xizang (Tibet).	Unknown, Yunnan

a ring of 4–5 mm tall brown setae below nodes, glabrous sheath scar, fewer branches and more leaf number, from the latter by a ring of 4–5 mm tall brown setae below nodes, less branch number, glabrous sheath scar, oral setae absent and narrower leaf blade.

Type. CHINA, Xizang (Tibet), Zayu County, Xiachayu Town, bamboo mountain of new village, 28°31'14"N, 96°57'59"E, 2705 m alt., 24 August 2015, X.Y.Ye & X.He YXY254-1 (holotype & isotype: KUN!).

Description. Rhizomes pachymorph, rhizome neck 5–10 cm long, 1.2–2 cm in diameter, solid. Culms (3)4–5(6) m tall, unicaespitose, 0.5–1.4 cm in diameter; inter-




Figure 3. *Fargesia purpurea* D.Z. Li & X.Y. Ye **A** habitat **B** young and densely white powdery culm with purple culm sheath **C** individual **D** rhizome.

nodes terete, 30–46 cm long, white powdery and black when culms old, with a ring of 4–5 mm brown setae below nodes, longitudinal ribs prominent; wall 1–4 mm thick, cavity filled with lamellate pith; nodes with weakly prominent supra-nodal ridge; sheath scar prominent, with persistent remains of sheath base. Branches 3–7, open; buds triangular. Shoots purple, or with dark purple spots. Culm sheaths persistent, leathery, narrowly triangular, 1/3 as long as internodes, glabrous, longitudinal ribs prominent, upper margins ciliolate; auricles and oral setae absent; ligule truncate or inclined-truncate, 1–2 mm; blade reflexed, linear-lanceolate, glabrous, narrow than apex of culm sheath, readily deciduous. Foliage leaves 3–5 per ultimate branch; sheaths 2.5–4.5 cm long, glabrous, purple, margins ciliolate; auricles and oral setae absent; ligules truncate, ca. 1 mm; petiole 1–3 mm long; blades lanceolate, 5–12 × 0.5–1.4 cm, abaxially densely white pubescent, base cuneate, secondary veins 3–4 pairs, transverse veins conspicuous, margins serrate. Inflorescence unknown.

Phenology. New shoots July to August.

Etymology. The specific epithet refers to the color of culm sheath and leaf sheath.




Figure 4. *Fargesia purpurea* D.Z. Li & X.Y. Ye **A** internodes, showing branches and persistent culm sheath **B** young culms with culm sheaths **C** culm bud **D** branches **E** node, showing brown setae below node **F** culm sheath, showing details of blade and ligule **G** branchlet **H** leaf sheath **I** abaxial surface of leaf, showing densely pubescence.




Figure 5. *Fargesia purpurea* D.Z. Li & X.Y. Ye **A** internode with branchlet **B** culm sheath abaxial view, showing culm leaf blade **C** node with branches **D** culm bud **E** yong culm with culm sheath **F** abaxial surface of leaf, showing densely pubescence.

Vernacular name. Zǐ Qiào Jiàn Zhú (Chinese pronunciation); 紫鞘箭竹 (Chinese name).

Distribution and habitat. *Fargesia purpurea* is only known from the type locality, bamboo mountain of new village in Zayu county. It grows under the evergreen broad-leaved forest at an elevation of 2700–2800 m alt.

Table 2. Morphological comparison of *Fargesia purpurea* and its related species.

Characters	<i>Fargesia purpurea</i>	<i>Fargesia pauciflora</i>	<i>Fargesia brevistipedis</i>
Culm height	(3)4–5(6) m	2–4 m	4–5 m
Culm diameter	0.5–1.4 cm	1–3 cm	1.2–2 cm
Internode	30–46 cm long, longitudinal ribs prominent, densely white powdery, with a ring of 4–5 mm brown setulose; wall 1–4 mm thick	35–40 cm long, longitudinal ribs prominent, initially densely white powdery, glabrous; wall 2–3 mm thick	28–35 (40) cm long, initially white powdery, glabrous; wall 1.5–2(3) mm thick
Branch complement	3–7	6–10	many
Sheath scar	Prominent, glabrous, with persistent remains of sheath base	Prominent, initially densely yellow-brown setose	Prominent, initially yellow-brown setose
Culm sheath	Persistent, glabrous, upper margins yellow-brown ciliate initially	Persistent or gradually deciduous, glabrous or sparsely yellow-brown setose, margins brown ciliate	Persistent or gradually deciduous, glabrous or sparsely yellow-brown setose, white powdery, margins brown ciliate
Culm sheath ligule	Truncate or inclined-truncate, 1–2 mm	Truncate or arcuate, 1–2.5 mm	Truncate or arcuate, 1–1.5 mm
Culm sheath blade	Reflexed, readily deciduous,	Reflexed, linear-lanceolate, glabrous	Reflexed, linear or linear-triangular,
Leaf number of the ultimate branch	3–5	2 or 3	(3)5(6)
Leaf sheath	Purple, glabrous	Glabrous	Purple or light green, glabrous
Leaf oral setae	Absent	Absent	4–8, 5–6 mm long
Leaf ligule	Truncate, 1 mm tall	Arcuate or truncate, glabrous	0.5 mm tall
Petiole	1–3 mm long	Initially abaxially pubescent	Initially pubescent
Leaf blade	5–12 × 0.5–1.4 cm, secondary veins 3–4 pairs, abaxially densely pubescent	9–14 × 0.7–1.2 cm, secondary veins 2–4 pairs, abaxially pubescent	6.5–11.5 × 0.5–0.85 cm, secondary veins 3–4 pairs, initially abaxially gray pubescent
Habitat	Under the evergreen broadleaved forest at the altitude of 2700–2800 m, Zayu, Xizang (Tibet).	Under the <i>Pinus</i> or broadleaved forest, or under shrubs, 2000–3200 m, southwest Sichuan and northeast Yunnan.	Under shrubs at the elevation of about 1250 m, central Sichuan.

Additional specimens examined (paratype). CHINA, Xizang (Tibet), Zayu County, Xiachayu Town, bamboo mountain of new village, 28°32'04"N, 96°59'07"E, 2724 m alt., 24 August 2015, X.Y.Ye & X.He YXY254-2 (KUN!).

Notes. Morphological comparisons between *Fargesia purpurea* and the related species were provided in Table 2. Two species of this genus were distributed in the Zayu county, namely, *F. zayuensis* Yi and *F. macclureana* (Bor) Stapleton, with this new species being easily distinguished from them in this region by its glabrous culm sheath and abaxially densely white pubescent leaf blade.

Discussion

Both *Fargesia viridis* and *F. purpurea* have persistent culm sheaths and buds containing several obscure buds, making them belong to the section *Fargesia*. The shape of culm sheaths is different from these two species. *F. viridis* is characterized by narrowly rounded culm sheath, with apex nearly as wide as base, which is similar to the species of the series *Murielae*. *F. purpurea* is characterized by triangular culm sheaths, shorter than internodes, with apex narrower than base; these features are similar to those species of

the series *Yunnanenses*. Therefore, *F. viridis* and *F. purpurea* are assigned to the series *Murielae* and series *Yunnanenses*, respectively.

Fargesia is a polyphyletic genus and could be divided into three or four clades based on plastome sequences (Zhang et al. 2018; Zhou et al. 2019) and double-digested restriction enzyme-associated DNA sequencing (ddRAD-seq) data (Ye et al. 2019). *F. viridis* was classified as belonging to V-*Fargesia*4 clade based on the phylogenetic results of ddRAD-seq analyses (Ye et al. 2019), but no conclusion could be made for its position on the plastome phylogeny. Additionally, the phylogenetic relationship of *F. purpurea* in *Fargesia* has not been studied and that may be supplemented in the future.

Fargesia viridis (*F. sp.*2 in Fig. 2 of Ye et al. 2019) is closest to *F. frigidis* not only in morphology but also in phylogenetic relationships (Table 1, Ye et al. 2019), but the altitude distribution range of them are different. Moreover, *F. viridis* can be easily distinguished from *F. frigidis* by several morphologic characters, i.e. thinner culms, glabrous internodes, more leaves on ultimate branch. According to the identification keys, *F. viridis* is also similar to *F. zayuensis* and *F. similaris*; for example, they all have narrowly rounded culm sheath, with apex nearly as wide as the base, branch number usually above 5, auricles absent, glabrous leaf blade. However, a number of subtle features make *F. viridis* distinctive, such as internode nearly solid, densely white powdery culm, culm sheath persistent and densely yellow setose.

Fargesia purpurea resembles *F. pauciflora* and *F. brevistipedis* by its internode length, prominent sheath scar, culm sheath persistent, auricles and oral setae absent, and leaf blade abaxially pubescent, but differs in terms of the habitat, thinner culm, internode with a ring of 4–5 mm brown setulose, less branch number, glabrous culm sheath and sheath scar.

Mountains of Southwest China are the diversity center for *Fargesia* species; 80 out of 85 are distributed in this area and 73 of them are endemic. The two new species established here are also distributed in these mountains, indicating that the species diversity of *Fargesia* in this region may be beyond our knowledge. The species of *Fargesia* have an island-like distribution and allopatric speciation might have great impact on their diversity (Ye et al. 2019). However, the diversification of species could be caused by many reasons, such as heterogeneous environment, fluctuating climatic conditions, and adaptive evolution (Xing and Ree 2017; Ding et al. 2020). This genus with species distributed on a different elevation provides a case to disentangle the extrinsic and intrinsic factors that could promote species divergence. And research in this area may improve our ability to predict the evolutionary tendency and mitigate the threats posed by global warming to species distributed in the mountains of Southwest China.

Acknowledgements

We thank Xie He of Kunming Institute of Botany, Chinese Academy of Science, and the guide of Zayu county, Xizang (Tibet), for their assistance with field work. The study was funded by the National Natural Science Foundation of China (No. 31800315, 31430011), and the Applied and Fundamental Research Foundation of Yunnan Province (2019FD059).

References

- Bamboo Phylogeny Group (2012) An updated tribal and subtribal classification of the bamboos (Poaceae: Bambusoideae). In: Gielis J, Potters G (Eds) The 9th World Bamboo Congress. Bamboo Science Cult, 27 pp.
- Clark LG, Oliveira RP (2018) Diversity and evolution of the new world bamboos. Proceedings World Bamboo Congress, Mexico.
- Clark LG, Londoño X, Ruiz-Sanchez E (2015) Bamboo taxonomy and habitat. In: Liese W, Köhl M (Eds) Bamboo – The Plant and its Uses, Springer, 30 pp.
- Ding WN, Ree RH, Spicer RA, Xing YW (2020) Ancient orogenic and monsoon-driven assembly of the world's richest temperate alpine flora. *Science* 369(6503): 578–581. <https://doi.org/10.1126/science.abb4484>
- Janzen DH (1976) Why bamboos wait so long to flower. *Annual Review of Ecology and Systematics* 7(1): 347–391. <https://doi.org/10.1146/annurev.es.07.110176.002023>
- Keng PC (1987) On the nomenclature of the high-alpine bamboos from China. *Journal of Bamboo Research* 6: 11–17.
- Keng PC, Wang ZP (1996) *Flora Reipublicae Popularis Sinicae*. Science Press, Beijing, 387–480.
- Li DZ, Wang ZP, Zhu ZD, Xia NH, Jia LZ, Guo ZH, Yang GY, Stapleton CMA (2006) Tribe Bambuseae. In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China* (Vol. 22). Beijing and Missouri Botanical Garden Press, St. Louis, 7–180.
- Ohrnberger D (1999) The bamboos of the world: Annotated nomenclature and literature of the species and the higher and lower taxa. [Access Online via Elsevier]
- Vorontsova MS, Clark LG, Dransfield J, Govaerts R, Baker WJ (2016) World checklist of bamboos and rattans. Science Press, Beijing.
- Xing YW, Ree RH (2017) Uplift-driven diversification in the Hengduan Mountains, a temperate biodiversity hotspot. *Proceedings of the National Academy of Sciences of the United States of America* 114(17): E3444–E3451. <https://doi.org/10.1073/pnas.1616063114>
- Yang L, Yi TP (2013a) A new species of *Fargesia* Franch. emend. Yi from northwestern Yunnan of China (Poaceae) and Chinese name of *Gelidocalamus stellatus* Wen. *Journal of Sichuan Forestry Science and Technology* 34: 48–51.
- Yang L, Yi TP (2013b) A new species of *Fargesia* Franch. emend. Yi from western Guizhou of China (Bambusoideae). *Bulletin of Botanical Research* 33: 513–515.
- Ye XY, Ma PF, Yang GQ, Guo C, Zhang YX, Chen YM, Guo ZH, Li DZ (2019) Rapid diversification of alpine bamboos associated with the uplift of the Hengduan Mountains. *Journal of Biogeography* 46(12): 2678–2689. <https://doi.org/10.1111/jbi.13723>
- Yi TP (1988) A study of the genus *Fargesia* from China. *Journal of Bamboo Research* 7: 6–15.
- Yi TP (2000a) New materials of alpine bamboos from southwest China. *Journal of Sichuan Forestry Science and Technology* 21: 1–6.
- Yi TP (2000b) A new species of *Fargesia* from northeastern Sichuan, China. *Yunnan Zhi Wu Yan Jiu* 22: 251–254.
- Yi TP (2000c) Some new taxa of Bambusoideae in western Sichuan, China. *Journal of Sichuan Forestry Science and Technology* 21: 13–23.

- Yi TP, Shi JY, Wang HT, Ma LS, Yang L (2005) Two new species and a new combination of bamboo from Sichuan and Yunnan, China. *Journal of Sichuan Forestry Science and Technology* 26: 33–42.
- Yi TP, Shi JY, Wang HT, Ma LS, Yang L (2006) A new species and a new forma of *Fargesia* Franch. emend. Yi from Yunnan, China. *Journal of Sichuan Forestry Science and Technology* 27: 47–49.
- Yi TP, Shi JY, Yang L (2007) Alpine new bamboos from Sichuan, Tibet, Chongqing China. *Bulletin of Botanical Research* 27: 515–520.
- Yi TP, Shi JY, Ma LS, Wang HT, Yang L (2008) *Iconographia Bambusoidearum Sinicarum*. Science Press, Beijing, 415–505.
- Zhang YQ, Ren Y (2016) Supplementary description of flowers and flowering branches of four *Fargesia* and one *Drepanostachyum* species (Bambusoideae, Poaceae), and notes on their taxonomy. *Nordic Journal of Botany* 34(5): 565–572. <https://doi.org/10.1111/njb.00975>
- Zhang YQ, Zhou Y, Hou XQ, Huang L, Kang JQ, Zhang JQ, Ren Y (2018) Phylogeny of *Fargesia* (Poaceae: Bambusoideae) and infrageneric adaptive divergence inferred from three cpDNA and nrITS sequence data. *Plant Systematics and Evolution* 305(1): 61–75. <https://doi.org/10.1007/s00606-018-1551-y>
- Zhou Y, Zhang YQ, Xing XC, Zhang JQ, Ren Y (2019) Straight From the Plastome: Molecular phylogeny and morphological evolution of *Fargesia* (Bambusoideae: Poaceae). *Frontiers of Plant Science* 10: e981. <https://doi.org/10.3389/fpls.2019.00981>