

On the recognition of the long neglected *Vitis adenoclada* Hand.-Mazz. (Vitaceae) from southern China

Jun Wen¹, Zhi-Yao Ma¹

¹ Department of Botany, National Museum of Natural History, Smithsonian Institution, PO Box 37012, Washington, DC 20013-7012, USA

Corresponding author: Jun Wen (WENJ@si.edu)

Academic editor: A. Trias-Blasi | Received 4 March 2021 | Accepted 14 April 2021 | Published 17 June 2021

Citation: Wen J, Ma Z-Y (2021) On the recognition of the long neglected *Vitis adenoclada* Hand.-Mazz. (Vitaceae) from southern China. *PhytoKeys* 179: 29–33. <https://doi.org/10.3897/phytokeys.179.65519>

Abstract

This study reports the recognition of *Vitis adenoclada* Hand.-Mazz. from southern China. The species was not recognized in the *Flora Reipublicae Popularis Sinicae* and *Flora of China* treatments. Recent field studies and examination of herbarium collections including the type material suggest that *Vitis adenoclada* is morphologically similar to *V. heyneana*, in their densely arachnoid tomentose abaxial leaves, yet it can be easily distinguished from the latter by its red-purple glandular hairs on the young branches (vs. glandular hairs absent in *V. heyneana*) and inflorescences usually subtended by a tendril at the base (vs. only occasionally with a tendril in *V. heyneana*). *Vitis adenoclada* may be a species of hybrid origin, with the highly tomentose *Vitis heyneana* as one of the parental species, and likely the glandular-hair bearing *V. davidii* as the other parental species. *Vitis adenoclada* is recorded from southern China in Guangdong, Guangxi, Guizhou, Hunan and Zhejiang provinces.

Keywords

Grapes, species delimitation, Vitaceae, *Vitis*, *Vitis adenoclada*, *Vitis heyneana*

Introduction

Vitis L. (the grape genus) consists of about 75 species widely distributed in the Northern Hemisphere, with eastern Asia and North America as its current centers of diversity (Wen 2007; Wen et al. 2018a, b; Ma et al. 2020). Li (1998) and Ren and Wen (2007) did not treat *Vitis adenoclada* Hand.-Mazz., which was described from Hunan by Han-

del-Mazzetti (1925), in the *Flora Reipublicae Ropularis Sinicicae* and *Flora of China* treatments. Species delimitation of eastern Asian *Vitis* remains highly problematic (Li et al. 1996; Wan et al. 2008; Ma et al. 2018, 2020). Based on our recent field work and the examination of the type material and additional specimens, we herein propose to recognize the long-neglected *Vitis adenoclada* from southern China.

Systematics

***Vitis adenoclada* Hand.-Mazz., Anz. Akad. Wiss. Wien, Math.-Naturwiss. Kl. 62: 145. 1925.**

Figures 1, 2

Vitis adenoclada Hand.-Mazz., Anz. Akad. Wiss. Wien, Math.-Naturwiss. Kl. 62: 145. 1925. Type: China. Hunan: Xinhua Xian, near mine at Hsikwangschan (Xikuangshan), Hsinhwa (Xinhua), 600 m, 13 May 1918, in fl., *H. Handel-Mazzetti 11819* (holotype: WU, WU0029734; isotype: K, K000574937).

Description. Robust high climber 3–12 m, polygamo-dioecious. Young branches with arachnoid tomentum and red purple glandular hairs, glandular hairs hardened at maturity; tendrils bifurcate to sometimes trifurcate, stout. Leaves simple; stipules brown, narrowly triangular to ovate-triangular, 3.5–5 × 2–2.5 mm, membranaceous, entire; petiole 3–7 cm, arachnoid tomentose; blade oval, 7–16 × 5–12 cm, abaxially densely tomentose, adaxially pubescent, veins 7–10 pairs, base cordate to subcordate, notch obtuse, margin serrate with 20–35 teeth on each side, notch area entire, teeth fine and short, apex acute to acuminate. Inflorescence a thyrses, leaf-opposed, 7–18 cm, paniculate in shape, usually subtended by an unbranched tendril at the base; peduncle 2–7 cm, arachnoid tomentose. Pedicel 1–3 mm, pubescent. Calyx 0.1–0.2 × 0.2–0.3 mm. Petals calyptrate, 1.5–1.8 × 0.4–0.5 mm. Stamens 2–3 mm; anthers 0.6–0.7 mm. Ovary oval; style short. Fruits purple-black, globose, 9–12 mm in diam. Seeds obovoid, chalaza rounded, ventral infolds furrowed upward 1/3–1/4 from base.

Phenology. Fl. May–Jul; fr. Jul–Oct.

Distribution and ecology. In Guangdong, Guangxi, Guizhou, Hunan and Zhejiang provinces of China. 500–1015 m. Edge of forests, forests, often in limestone areas.

Additional specimens examined. CHINA. **Guangdong:** Huaiji Xian, 31 Oct 1958, *Y.G. Liu 02809* (IBSC). **Guangxi:** Jingxiu Xian, Dayaoshan, 16 km, 1015 m, 8 May 2010, floral buds, *J. Wen 11560* (US), *J. Wen 11577* (US). **Guizhou:** Jun 6, 1987, *BJFC Graduate Students 434* (BJFC00010633, BJFC00010634). **Hunan:** Hongjiang Xian, Xuefengshan, 500 m, 12 May 2014, *X.Q. Liu 114* (CCAU), *X.Q. Liu 115* (CCAU); Xinning Xian, Zhiyunshan, 13 Sep 1984, 900 m, *Zhiyunshan Exp. Team 476* (PE). **Zhejiang:** Nishui, 28 Jul 1959, fr, *S.Y. Zhang 6061* (KUN, PE).

Notes. *Vitis adenoclada* is morphologically similar to *V. heyneana* Roem. & Schult, with both having arachnoid tomentose abaxial leaves. It can be easily distinguished



Figure 1. General morphology of *Vitis adenoclada*, especially showing the highly tomentose leaves and the inflorescences subtended by an unbranched tendril (*J. Wen 11560*, US).

from the latter by its red-purple glandular hairs on the young branches (vs. glandular hairs absent on young branches in *V. heyneana*) and inflorescences usually subtended by a tendril at the base (vs. only occasionally with a tendril in *V. heyneana*) (Figures 1, 2). *Vitis adenoclada* may be a species of hybrid origin, with the highly tomentose *Vitis heyneana* as one of the parental species, and a glandular-hair bearing *Vitis* species, e.g., *V. romanetii* Rom.Caill. or *V. davidii* (Rom.Caill.) Foëx as the other potential parental species. Both *V. davidii* and *V. romanetii* share the unique glandular trichomes with *V. adenoclada* (see Ma et al. 2016), and *V. davidii* occurs in southern China in the same geographic area as *V. adenoclada*. We hence regard *Vitis davidii* as a more likely parental species than *V. romanetii* due to their geographic distribution and the highly




Figure 2. Characteristic glandular trichomes on young branches of *Vitis adenoclada* **A** X.Q. Liu 114 (CCAU) **B** X.-Q. Liu 115 (CCAU) (photo credit: Z.Y. Ma).

similar glandular trichome morphology. Ma et al. (2018, 2020) also showed that *Vitis adenoclada* and *V. davidii* grouped in one clade in the nuclear phylogeny, consistent with our hypothesis.

Acknowledgements

We thank Robin Everly and Gil Taylor for assistance with obtaining literature, the following herbaria (BJFC, CCAU, IBSC, K, KUN, PE, US, and WU) for allowing the examination of their specimens, and two reviewers (Dr. Betsy Jackes and Dr. Limin Lu) and the handling editor Dr. Anna Trias-Blasi for their helpful suggestions.

References

- Handel-Mazzetti H (1925) Plantae novae sinenses, diagnosibus brevibus descriptae a Dre. Heinr. Handel-Mazzetti (35. Fortsetzung). Anzeiger der Akademie der Wissenschaften in Wien. Sitzung der Mathematisch-Naturwissenschaftliche Klasse 62: 143–148.
- Li CL (1998) Vitaceae. Flora Reipublicae Ropularis Sinicicae 48(2). Science Press, Beijing, ii–vii. [1–3, 12–208.] [in Chinese]

- Li CL, Cao YL, He YH (1996) Systematics on Chinese *Vitis* L. Chinese Journal of Applied and Environmental Biology 2: 234–253. [in Chinese]
- Ma ZY, Wen J, Ickert-Bond SM, Chen LQ, Liu XQ (2016) Morphology, structure, and ontogeny of trichomes of the grape genus (*Vitis*, Vitaceae). Frontiers of Plant Science 7: e704. <https://doi.org/10.3389/fpls.2016.00704>
- Ma ZY, Wen J, Tian JP, Jamal A, Chen LQ, Liu XQ (2018) Testing reticulate evolution of four *Vitis* species from East Asia using restriction-site associated DNA sequencing. Journal of Systematics and Evolution 56(4): 331–339. <https://doi.org/10.1111/jse.12444>
- Ma ZY, Wen J, Tian JP, Gui LL, Liu XQ (2020) Testing morphological trait evolution and assessing species delimitations in the grape genus using a phylogenomic framework. Molecular Phylogenetics and Evolution 148: e106809. <https://doi.org/10.1016/j.ympev.2020.106809>
- Ren H, Wen J (2007) *Vitis*. In: Wu Z-Y, Hong D-Y, Raven PH (Eds) Flora of China 12. Science Press, Beijing and Missouri Botanical Garden Press, St. Louis, 210–222.
- Wan YZ, Schwaninger H, Li D, Simon CJ, Wang YJ, Zhang CH (2008) A review of taxonomic research on Chinese wild grapes. *Vitis* 47(2): 81–88.
- Wen J (2007) Vitaceae. In: Kubitzki K (Ed.) The Families and Genera of Vascular Plants 9. Springer-Verlag, Berlin, 467–479. https://doi.org/10.1007/978-3-540-32219-1_54
- Wen J, Harris AJ, Kalburgi Y, Zhang N, Xu Y, Zheng W, Ickert-Bond S, Johnson G, Zimmer EA (2018a) Chloroplast phylogenomics of the New World grape species (*Vitis*, Vitaceae). Journal of Systematics and Evolution 56(4): 297–308. <https://doi.org/10.1111/jse.12447>
- Wen J, Lu LM, Nie ZL, Liu XQ, Zhang N, Ickert-Bond S, Gerrath J, Manchester SR, Boggan J, Chen ZD (2018b) A new phylogenetic tribal classification of the grape family (Vitaceae). Journal of Systematics and Evolution 56(4): 262–272. <https://doi.org/10.1111/jse.12427>