

Carex malipoensis (Cyperaceae), a new species from southeast Yunnan, China

Yuan-Yuan Li^{1,2}, Ya-Ping Chen¹, Li-Qiong Jiang³,
En-De Liu¹, Yuan Luo^{1,2}, Hua Peng¹

1 CAS Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China **2** University of Chinese Academy of Sciences, Beijing 100049, China **3** School of Ecology and Environmental Sciences & School of Life Sciences, Yunnan University, Kunming 650091, China

Corresponding author: Hua Peng (hpeng@mail.kib.ac.cn)

Academic editor: P. Jiménez-Mejías | Received 17 September 2021 | Accepted 16 December 2021 | Published 7 January 2022

Citation: Li Y-Y, Chen Y-P, Jiang L-Q, Liu E-D, Luo Y, Peng H (2022) *Carex malipoensis* (Cyperaceae), a new species from southeast Yunnan, China. PhytoKeys 188: 19–29. <https://doi.org/10.3897/phytokeys.18875401>

Abstract

Carex malipoensis, a new species from southeast Yunnan, China, is here described and illustrated. It is morphologically similar to *C. trichophylla* in sect. *Euprepes*, but differs from it by its longer inflorescences and peduncles, pendulous spikes, hispidulous female glumes, densely hispidulous utricles, and longer nutlets.

Keywords

Carex sect. *Euprepes*, *Carex trichophylla*, morphology, taxonomy

Introduction

As one of the largest angiosperm genera, *Carex* Linnaeus (1753) (Cyperaceae) comprises an extremely high diversity of about 2000 species (Roalson et al. 2021; WCSP 2021). The genus has a Cosmopolitan distribution. It was placed within the tribe Cariceae and divided into four subgenera by Kükenthal (1909): subg. *Psyllophora* (Degland 1828) Petermann (1849), subg. *Vigneastra* (Tuckerman 1843) Kükenthal (1899), subg. *Vigneae* (P. Beauv. ex Lestiboudois 1819) Petermann (1849) and subg. *Carex*. This classification of Kükenthal had been widely adopted by subsequent researchers for a long time. Recent molecular phylogenetic studies revealed that Cariceae is a natural group, and the previously recognized genera of this tribe, i.e., *Cymophyllus* Mack. ex Britton & A.Br.,

Kobresia Willdenow, *Schoenoxiphium* Nees von Esenbeck, and *Uncinia* Persoon, should be merged into *Carex* (Yen and Olmstead 2000; Global *Carex* Group 2015; Starr et al. 2015; Lévillé-Bourret et al. 2018; Martín-Bravo et al. 2019; Villaverde et al. 2020; Larridon et al. 2021; Roalson et al. 2021). A framework of the combined giant genus was urgently needed to increase our understanding of *Carex*. In order to solve this problem, Villaverde et al. (2020) conducted a robust phylogeny of *Carex* based on molecular data (308 nuclear exon matrices, 543 nuclear intron matrices and 66 plastid exon matrices) and six clades were recognized. Accordingly, Villaverde et al. (2020) proposed an updated infrageneric classification of *Carex* and classified it into six subgenera: subg. *Siderosticta* M.J. Waterway, subg. *Carex*, subg. *Euthyceras*., subg. *Psyllophorae*, subg. *Uncinia* (Pers.) Peterm. and subg. *Vignea*. However, the classification within the subgenera still remained unresolved, so a more systematic and friendly infrageneric classification system was required. A framework infrageneric classification of *Carex* was proposed recently which divided *Carex* into 62 formally named Linnean sections and 49 informal groups based on the current phylogenetic knowledge of *Carex* (Roalson et al. 2021).

A total of 527 *Carex* were recorded in Flora of China (Dai et al. 2010), the Catalogue of Life China (Chen and Zhang 2018) recorded 593 species, which represents the most complete and update list of the genus in China. The number of species of *Carex* continues growing in China as more new species have been reported in recent years (Lu and Jin 2018; Zhang et al. 2018; Jin et al. 2020; Lu et al. 2020; Yang and Liu 2020).

During our field investigations between 2016 and 2018, we collected specimens of an unknown species of *Carex* in Malipo County, southeast Yunnan Province, China. After careful morphological studies, examination of herbarium specimens and relevant literature, we concluded that it can be assigned to *Carex* sect. *Euprepes* based on a combination of some morphological characters: cauline leaves well-developed; leaf blades and involucre bracts elliptic to linear-elliptic, with prominent transverse veins; complex branched inflorescence; spikes androgynous; and presence of utriculiform cladophylls (Jiménez-Mejías et al. 2016) at the base of spikes. Molecular phylogenetic studies indicated that *C.* sect. *Euprepes* belongs to the core *Carex* clade (Starr et al. 2015; Villaverde et al. 2020), and the most recent infrageneric classification framework placed it with the Indica Clade together with species traditionally placed in sections *Euprepes* and *Mapaniifoliae* (Roalson et al. 2021). Consisting of seven species, sect. *Euprepes* are restricted to South and Southeast Asia (Nelmes 1955). Only one species of the section, *C. zizaniifolia* Raymond (1959), was previously reported from China, distributed in southeast Yunnan Province (Dai et al. 2010; Chen and Zhang 2018). However, after our research on this section, we conclude that our new collections are different from all known species and represent a species new to science. We describe and illustrate it here below.

Materials and methods

The new species was compared morphologically with specimens of other taxa of *Carex* sect. *Euprepes* from the following public herbaria A, BM, E, GH, HNU, IBSC, K, KUN, HUH, MO, MT, NY, P, PE, and US [acronyms follow Theirs (continuously

updated)] as well as our new collections across China (especially with material collected from southeast Yunnan and neighboring area; herbarium specimens kept in KUN). Meanwhile, protologues and other related taxonomic literature were collated and reviewed. The characters' data come from specimens measurements and the prologue (*C. atrivaginata* Nelmes and *C. tricophylla* Nelmes (1955), *C. euprepes* Nelmes and *C. laosensis* Nelmes (1939), *C. tavoyensis* Nelmes (1948), *C. zizaniifolia* Raymond and *C. poilanei* Raymond (1959)). The terminology used by Kükenthal (1909) for the morphological description of *Carex* species was adopted here. The distribution of the new species was compiled from the herbarium specimen records and our own collections, and shown on the distribution map.

Results

A detailed morphological comparison of the potential new species and the seven species of *C. sect. Euprepes* is summarized in Table 1. Morphologically, the new species is most similar to *C. tricophylla* but can be distinguished by the characteristics of culms (70–105 cm long, 2–4 mm thick, sides ribbed in the new species vs. culms 45–65 cm long, 1–1.5 mm thick, sides concave in *C. tricophylla*); leaf sheaths (1.5–7 cm long vs. 1.2–2 cm long); inflorescences (20–45 cm long, peduncles up to 9 cm long vs. 3–15 cm long, more or less exserted), spikes (15–33 mm long, pendulous vs. 7–10 mm long, erect), female glumes (hispidulous vs. glabrous), utricles (densely hispidulous vs. glabrous below, adpressed-hispid above), and nutlets (4–4.5 mm long vs. 2.25–2.5 mm long). The new taxon can be distinguished from all the other seven species in this section by their culms (length, cross-sectional shape, sides and indumentum); leaves (shape, transverse veins), sheath (length, indumentum and the appendage of mouth); inflorescence bigger (20–45 × 3–5 cm in the new taxa vs. 3–25 × 1–5 cm in the other species) or with different shape (oblong in the new taxon vs. oblong, narrowly oblong or triangular-ovate in other species); peduncles longer (up to 9 cm long in the new taxon vs. scarcely or slightly exserted in other species; only *C. atrivaginata* has relative long peduncles not exceeding 5 cm); and the female glumes, utricles and nutlets (differences in shape, size and indumentum).

Taxonomy

***Carex malipoensis* Yuan Y. Li & H. Peng, sp. nov.**

urn:lsid:ipni.org:names:77235050-1

Figures 1, 2

Type. CHINA. Yunnan Province, Malipo County, Mengdong Village, Xiangchunping, 22°54'36.77"N, 104°38'54.09"E, alt. 1850 m, 5 December 2016, E.D. Liu *et al.* LiuED6425 (holotype: KUN! Barcode 1433368; isotype: KUN! Barcode 1347669).




Figure 1. Morphology of *Carex malipoensis* **A** habitat **B** plant **C** rhizome **D** inflorescence **E** spike **F** leaf blade **G** sheath **H** the transverse veins **I** leaf margin **J** male glumes **K** female glume **L** utricles **M** achene. Scale bars: 2 mm. Photographed by X.X. Zhu (**A**, **D**, **E** *LiuED6425*, *LiuED5912* KUN), Y.P. Chen (**B**, **C**, **F**, **I** *Y.P. Chen & L.Q. Jiang MLP01* KUN), and Y. Y. Li (**G**, **H**, **J**–**K** *Y.P. Chen & L.Q. Jiang MLP01* KUN; **L**, **M** *LiuED6425* KUN).

Table 1. Morphological comparisons between *Carex malipoensis* and species in *C.* sect. *Euprepes*.

Characters	<i>C. malipoensis</i>	<i>C. atrivaginata</i>	<i>C. euprepes</i>	<i>C. laosensis</i>	<i>C. poilanei</i>	<i>C. tauoyensis</i>	<i>C. tricophylla</i>	<i>C. sizaniifolia</i>
Culm								
Length	70–105 cm	65 cm	70 cm	unknown	70–105 cm	unknown	45–65 cm	60–70 cm
Thick	2–4 mm	3 mm	3 mm	2 mm	2–3 mm	2–2.5 mm	1–1.5 mm	2 mm
Transverse angle	obtuse	acute	obtuse	obtuse	acute	acute, obtuse to subacute	obtuse	obtuse
Sides	ribbed	concave	concave	concave	concave	ribbed, slightly twisted	concave	concave
Indumentum	hispidulous	glabrous	glabrous	glabrous	glabrous	glabrous	hispidulous upward	glabrous
Leaf								
Number	3–5	7	4–6	1–4	6–8	unknown	6 or more	8
Shape	elliptic	narrowly linear-lanceolate	elliptic	narrowly linear-lanceolate	narrowly linear-lanceolate	Linear-elliptic	elliptic, apex attenuated	narrowly elliptic or lanceolate
Length	12–28 cm	13–16 cm	17–24 cm	20–30 cm	18–30 cm	25–28 cm	12–18 cm	15–25 cm
Width	2–3 cm	1–1.3 cm	3–4.5 cm	8–15 mm	1–2 cm	2.3–3 cm	1.2–2 cm	2–2.5 cm
Indumentum	glabrous, sparsely hispidulous on undersurface midrib	glabrous, apex hispidulous	scabrid along veins undersurface	glabrous	glabrous	scabrid along veins undersurface	scabrid along veins undersurface	scabrid along veins
Leaf sheath								
Length	1.5–7 cm	1.5–5 cm	1–2 cm	1.5–2 cm	2–5 cm	unknown	1.5–2 cm	1.4–2 cm
Indumentum	hispidulous	glabrous	glabrous below, hispidulous above	hispidulous	glabrous	glabrous	hispidulous	glabrous
Appendage of sheath mouth	not developed	not developed	prominent	small	small	small	prominent	small
Inflorescence								
Shape	oblong	oblong	narrowly oblong	narrowly oblong	triangular-ovate	narrowly oblong	narrowly oblong	narrowly oblong
Length	20–45 cm	8–10 cm	10–22 cm	3.5–13.5 cm	12–20 cm	15–25 cm	3–15 cm long	5–6 cm
Width	3–5 cm	2–5 cm	3–4 cm	1–2.5 cm	3–6 cm	1–2 cm	1–2 cm	1–1.5 cm
Peduncles	exserted	exserted	scarcely or slightly exserted	scarcely exserted	exserted	scarcely exserted	exserted	scarcely exserted
Spikes								
Length	15–30 (45) mm	10–25 mm	8–10 mm	5 mm	8–18 mm	5–9 mm	7–10 mm	5–6 mm
Male part vs female part	much longer	much longer	equal	longer	much longer	slightly longer	longer	slightly longer
Utricles								
Shape	ovate-elliptic	ovate-elliptic	elliptic	broadly ellipsoid or ellipsoid-obovoid	ovate or rhomboid-ovate	ellipsoid or obovoid-ellipsoid	obovoid or ellipsoid-obovoid	ovate-elliptic

Characters	<i>C. malipoensis</i>	<i>C. atrivaginata</i>	<i>C. estrepes</i>	<i>C. laosensis</i>	<i>C. poilanei</i>	<i>C. taoyuensis</i>	<i>C. trichophylla</i>	<i>C. zizaniifolia</i>
Length	6–7.5 mm	6–7.5 mm	4–4.5 mm	4–4.5 mm	4–5 mm	4–5 mm	6–6.5 mm	ca. 3 mm
Indumentum	densely hispidulous	densely hispidulous	glabrous, margins hispidulous	glabrous	unknown	glabrous below, adpressed-hispid above	glabrous below, adpressed-hispid above	densely hispidulous
nutlets	ca. 4 mm	ca. 4 mm	2.25–2.5 mm	2.25–2.8 mm	unknown	ca. 2.3 mm	2.25–2.5 mm	immature

The characters marked “unknown” for the missing description of the protologue or the original materials incomplete.

Diagnosis. The new species is most similar to *C. trichophylla* Nelmes (1955), but differs in inflorescences 20–45 cm long (vs. shorter than 15 cm in *C. trichophylla*), peduncles up to 9 cm long (vs. more or less exserted in *C. trichophylla*), spikes pendulous (vs. erect in *C. trichophylla*), female glumes hispidulous (vs. glabrous in *C. trichophylla*), utricles densely hispidulous (vs. glabrous below, adpressed-hispid above in *C. trichophylla*), and nutlets 4–4.5 mm long (vs. 2.25–2.5 mm long in *C. trichophylla*).

Description. Rhizomes elongate. Culms tufted, 70–105 cm long, 2–4 mm in diam, obtusely trigonous, hispidulous, basal sheaths dark brown. Leaves 3–5, basal and cauline, loosely arranged; leaf blade elliptic, 12–28 × 2–3 cm, transverse veins prominent, margin hispidulous-villous, base round to cuneate, apex acute, greyish green when dried; sheaths 1.5–7 cm long, hispidulous, mouth hairy, not developed into prominent appendage. Involucral bracts leaf-like, longer than inflorescence, sheathing; panicles 20–45 × 3–5 cm, with 12–16 branches, single or binate; peduncles up to 9 cm long, reduced toward apex, tenuous, glabrous or slightly pubescent; inflorescence axes sharply trigonous, hairy on edges; bractlets glumiform, ca. 5 mm long, apex awned, awns 4–5 mm long; cladophylls utriculiform, 3–4 mm long; spikes bisexual, androgynous, 15–30 (–45) mm long; male part of spike much longer than female part; male part densely many flowered, ca. 2.5 mm wide; female part fewer flowered, 3–4.5 mm wide; male glumes oblong-lanceolate, 6–9 × 1.5–2 (–3.2) mm, awned; female glumes oblong-lanceolate, 3.8–4.2 mm long, pale brown, green at middle, apex acute, midrib extending into a scabrid awn. Utricles ovate-elliptic, 6–7.5 mm long, green to brown, veined, densely hispidulous, apex attenuating into a long beak, ca. 3 mm long, orifice oblique. Nutlets ca. 4 × 1.8–2 mm, dark brown, obovate-elliptic, trigonous. Stigmas 3.

Phenology. Flowering from November to December, and fruiting in May.

Distribution. The new species is currently known from Malipo County in south-east Yunnan at the Sino-Vietnamese border (Fig. 3).

Habitat. The new species usually grows in evergreen broad-leaved forests at altitudes of 1100–1850 m.

Etymology. The specific epithet refers to Malipo County of Yunnan Province, China, from where the type specimens were collected.

Common name (assigned here). Ma Li Po Tai Cao (麻栗坡臺草; Chinese name).




Figure 2. Line drawing of *Carex malipoensis* **A** habit **B** male glume **C** female glume **D** utricles **E** achene **F** part of blade, show the transverse veins. Scale bars: 6 cm (**A**); 2 mm (**B–E**). Drawn by Yuan Luo from the type specimen.

Additional specimens examined (paratypes). CHINA. Yunnan: Maguan County, Ching-kou Loa-chün-shan, 7 December 1947, *K.M. Feng 13677* (KUN0368409; KUN1263725); Malipo County, Mengdong Village, 4 December 2016, *E.D. Liu et al. LiuED 6336* (KUN1433717), *LiuED6403* (KUN1433131); Malipo County, Laojun-shan, Bailingyan, 13 May 2017, *E.D. Liu et al. LiuED5912* (KUN1340680); Malipo County, Tianbao Town, Bajiaoping Village, 1 December 2018, *Y.P. Chen & L.Q. Jiang MLP01* (KUN).



Figure 3. Distribution map of *Carex malipoensis* (▲) and *C. trichophylla* (●).

Specimens examined of other species. *Carex atrivaginata*: VIETNAM. Chapa: Pételot, 3179 (P00277787, P00277788); E. Poilane, 27084 (MT00072452, MT00072458); *C. euprepes*: LAOS. Tawieng, Chiengkvang: 2 April, 1932, Kerr, 20927 (BM001172101, K000291207, K000291208, K000291210, NY04059693, P00282617); *C. laosensis*: LAOS. Pak Munung, Wieng chan: 22 April, 1932, Kerr, 21202 (K000291209, K000291210, P00284722); *C. poilanei*: LAOS. Phong Saly: 6 September, 1941, E. Poilane, 25984 (MT00117677); E. Poilane, 32994 (MT00072475); *C. tavoyensis*: MYANMAR. Padachaung, Tavoy. 3 April, 1921, P.T. Russell, 1935 (K000999214); *C. trichophylla*: VIETNAM. Chapa (Fig. 3): 1 July, 1930, Pételot, 5325 (GH00027549, P00302178, P00302179, P00302180, US00087306); *C. zizaniifolia*: CHINA. Yunnan, Pingbian: 1934, H. T. Tsai 62809 (A00027543, IBSC0653006, KUN0368701, PE00030290).

Conservation status. The new species is currently known from Maguan and Malipo Counties in Yunnan, China. Only six collections have been recorded since 1947. It may be classified as Endangered (EN) or Vulnerable (VU) according to the IUCN Red List criteria (IUCN 2012). However, collections of *Carex* are often deficient, and a solid suggestion is needed based on a comprehensive investigation about the new species. Therefore, we suggest to characterize the conservation status of *C. malipoensis* as Data Deficient (DD) at present.

Discussion

In comparison with other species of *C.* sect. *Euprepes*, *C. malipoensis* is morphologically most similar to *C. trichophylla*. Both species have obtusely trigonous culms and elliptic and subpetioleate leaf blades, inflorescences big and loose, and utricles longer than 6 mm, all characters that differ from all the remaining species of *C.* sect. *Euprepes*. Despite their shared similarities, *C. malipoensis* can be distinguished from *C. trichophylla* in the length of inflorescences and peduncles, and indumentum of sheath and utricles. Specifically, *C. malipoensis* has much longer culms and spikes, and larger leaves, panicles, and achenes compared with that of *C. trichophylla*. Moreover, the female glumes and utricles are hispidulous in *C. malipoensis*, but glabrous in *C. trichophylla*.

Acknowledgements

We would like to thank Dr. Chun-Lei Xiang and two anonymous reviewers for their constructive suggestions that greatly improved this paper. We are grateful to the staff of herbaria IBSC, KUN and PE for their kind assistance with research facilities. We also thank Dr. Xin-Xin Zhu for providing the photographs of the new species. This work was supported by the National Natural Science Foundation of China (Grant No. 31872649).

References

- Chen WL, Zhang SR (2018) Seed Plants. Catalogue of Life China. Science Press, Beijing, 49–91.
- Dai LK, Liang SY, Zhang SR, Tang YC, Koyama T, Tucker GC (2010) *Carex* Linnaeus. In: Wu ZY, Raven PH, D.Y. H (Eds) Flora of China. Science Press & Missouri Botanical Garden Press, Beijing & St. Louis, 293–294.
- Degland JV (1828) *Carex*. In: Loiseleur-Deslongchamps JLA (Ed.) Flora Gallica 2. 2nd edition. J. B. Baillière, Paris, 281–313.
- Global *Carex* Group (2015) Making *Carex* monophyletic (Cyperaceae, tribe Cariceae): A new broader circumscription. *Botanical Journal of the Linnean Society* 179(1): 1–42. <https://doi.org/10.1111/boj.12298>
- IUCN (2012) IUCN Red List Categories and Criteria, Version 3.1. 2 ed. Gland, Switzerland and Cambridge, UK, 32 pp.
- Jiménez-Mejías P, Luceno M, Wilson KL, Waterway MJ, Roalson EH (2016) Clarification of the Use of the Terms Perigynium and Utricle in *Carex* L. (Cyperaceae). *Systematic Botany* 41(3): 519–528. <https://doi.org/10.1600/036364416X692488>
- Jin XF, Liu YD, Lu YF, Sun WY, Wang H (2020) Notes on *Carex* (Cyperaceae) from China (VI): The identity of *Carex retrofracta* Kuk. (sect. *Confertiflorae*). *Phytotaxa* 429(2): 135–147. <https://doi.org/10.11646/phytotaxa.429.2.4>

- Kükenthal G (1899) Die *Carex* vegetation des außertropischen Südamerika (ausgenommen Paraguay und Südbrasilien). Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie 27: 485–563.
- Kükenthal G (1909) Cyperaceae-Caricoideae. In: Engler HGA (Ed.) Das Pflanzenreich. Wilhelm Engelmann, Leipzig, 824 pp.
- Larridon I, Zuntini AR, Lévillé-Bourret É, Barrett RL, Starr JR, Muasya AM, Villaverde T, Bauters K, Brewer GE, Bruhl JJ, Costa SM, Elliott TL, Epiawalage N, Escudero M, Fairlie I, Goetghebeur P, Hipp AL, Jiménez-Mejías P, Sabino Kikuchi IAB, Luceño M, Márquez-Corro JI, Martín-Bravo S, Maurin O, Pokorny L, Roalson EH, Semmouri I, Simpson DA, Spalink D, Thomas WW, Wilson KL, Xanthos M, Forest F, Baker WJ (2021) A new classification of Cyperaceae (Poales) supported by phylogenomic data. Journal of Systematics and Evolution 59(4): 852–895. <https://doi.org/10.1111/jse.12757>
- Lestiboudois TG (1819) Essai sur la famille des Cypéracées. Didot Jeune, Paris, 46 pp.
- Lévillé-Bourret É, Starr JR, Ford BA (2018) Why are there so many sedges? Sumatroskirpeae, a missing piece in the evolutionary puzzle of the giant genus *Carex* (Cyperaceae). Molecular Phylogenetics and Evolution 119: 93–104. <https://doi.org/10.1016/j.ympev.2017.10.025>
- Linnaeus C (1753) Species Plantarum 2. Impensis Laurentii Salvii, Stockholm, 561–1200.
- Lu YF, Jin XF (2018) Notes on *Carex* (Cyperaceae) from China (V): New species, combination and subspecies from Zhejiang. Phytotaxa 372(3): 203–211. <https://doi.org/10.11646/phytotaxa.372.3.3>
- Lu ZC, Su YL, Lu YF, Jin XF (2020) *Carex pingleensis* (*Carex* sect. *Mitratae*), a new species of Cyperaceae from Guangxi, China. Taiwania 65: 391–395. <https://doi.org/10.6165/tai.2020.65.391>
- Martín-Bravo S, Jiménez-Mejías P, Villaverde T, Escudero M, Hahn M, Spalink D, Roalson EH, Hipp AL, Global Carex Group (2019) A tale of worldwide success: Behind the scenes of *Carex* (Cyperaceae) biogeography and diversification. Journal of Systematics and Evolution 57: 695–718. <https://doi.org/10.1111/jse.12549>
- Nelmes E (1939) Notes on *Carex*: VIII. Bulletin of Miscellaneous Information (Royal Botanic Gardens, Kew) 6: 305–306 <https://doi.org/10.2307/4111742>
- Nelmes E (1948) A New Species of *Carex* from Burma. Kew Bulletin 3(1): 67–68. <https://doi.org/10.2307/4118923>
- Nelmes E (1955) The Genus *Carex* in Indo-China, including Thailand and Lower Burma. Mémoires du Muséum National d'Histoire Naturelle Nouvelle Série Série B, Botanique. Editions Du Muséum, Paris, 103–108.
- Petermann WL (1849) Deutschlands Flora. Wigand, Leipzig, 668 pp.
- Raymond M (1959) Carices Indochinenses necnon Siamenses. Memoires de Jardin Botanique de Montreal 53: 36–37.
- Roalson EH, Jiménez-Mejías P, Hipp AL, Benítez-Benítez C, Bruederle LP, Chung K-S, Escudero M, Ford BA, Ford K, Gebauer S, Gehrke B, Hahn M, Hayat MQ, Hoffmann MH, Jin X, Kim S, Larridon I, Lévillé-Bourret É, Lu Y, Luceño M, Maguilla E, Márquez-Corro JI, Martín-Bravo S, Masaki T, Míguez M, Naczi RFC, Reznicek AA, Spalink D, Starr JR (2021) A framework infrageneric classification of *Carex* (Cyperaceae) and its organizing principles. Journal of Systematics and Evolution 59(4): 726–762. <https://doi.org/10.1111/jse.12722>

- Starr JR, Janzen FH, Ford BA (2015) Three new, early diverging *Carex* (Cariceae, Cyperaceae) lineages from East and Southeast Asia with important evolutionary and biogeographic implications. *Molecular Phylogenetics and Evolution* 88: 105–120. <https://doi.org/10.1016/j.ympev.2015.04.001>
- Thiers B (continuously updated) Index Herbariorum: A global directory of public herbaria and associated staff. <http://sweetgum.nybg.org/ih> [accessed 12 August 2021]
- Tuckerman E (1843) *Enumeratio methodica Caricum quarundam I. Riggs, Schenectadae*, 27 pp.
- Villaverde T, Jiménez-Mejías P, Luceño M, Waterway MJ, Kim S, Lee B, Rincón-Barrado M, Hahn M, Maguilla E, Roalson EH, Hipp AL (2020) A new classification of *Carex* (Cyperaceae) subgenera supported by a HybSeq backbone phylogenetic tree. *Botanical Journal of the Linnean Society* 194(2): 141–163. <https://doi.org/10.1093/botlinnean/boaa042>
- WCSP (2021) World checklist of selected plant families. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet. <http://wcsp.science.kew.org/> [accessed 12 August 2021]
- Yang HB, Liu GD (2020) *Carex ledongensis* (Cyperaceae), a new sedge from Hainan Isl. Of South China. *Phytotaxa* 461(2): 72–78. <https://doi.org/10.11646/phytotaxa.461.2.1>
- Yen AC, Olmstead RG (2000) Molecular systematics of Cyperaceae tribe Cariceae based on two chloroplast DNA regions: *ndhF* and *trnL* intron-intergenic spacer. *Systematic Botany* 25(3): 479–494. <https://doi.org/10.2307/2666691>
- Zhang SR, Zhang J, Liu ZY, Qu S, Han RG (2018) *Carex nodosa* (Cyperaceae), a new species of *C.* sect. *Occlusae* from southwest China. *Nordic Journal of Botany* 36(8): e01900. <https://doi.org/10.1111/njb.01900>