

Pollen morphology of the genera *Hidalgoa* and *Dahlia* (Coreoapseidae, Asteraceae): implications for taxonomy

Erandi Sánchez-Chávez¹, Andrew Vovides¹, Victoria Sosa¹

¹ *Biología Evolutiva, Instituto de Ecología, A.C. Carretera Antigua a Coatepec 351, 91073 El Haya, Xalapa, Veracruz, Mexico*

Corresponding author: Erandi Sánchez-Chávez (erandi.sanchez4@gmail.com)

Academic editor: Alexander Sennikov | Received 17 December 2021 | Accepted 18 May 2022 | Published 13 June 2022

Citation: Sánchez-Chávez E, Vovides A, Sosa V (2022) Pollen morphology of the genera *Hidalgoa* and *Dahlia* (Coreoapseidae, Asteraceae): implications for taxonomy. *PhytoKeys* 199: 187–202. <https://doi.org/10.3897/phytokeys.199.79501>

Abstract

Hidalgoa and *Dahlia* are two closely related genera in Asteraceae, tribe Coreoapseidae whose limits need to be clarified. Pollen morphology has been useful for delimitation at the genus level in this family. To better define these genera, the morphology of pollen grains was observed and measured using light and scanning electron microscopy. The pollen grains of 25 species of *Dahlia* and *Hidalgoa* were acetolyzed and analyzed. Pollen is tricolporate in most of the species studied, although in a few species in *Dahlia*, grains were found to be hexacolporate. The most outstanding differentiating characters among species of *Dahlia* and *Hidalgoa* are colpus length (greater in *Hidalgoa*) and shape of spines (conical in *Hidalgoa*). In addition, lalongate ora are larger in *Hidalgoa* than in *Dahlia*. A PCA analysis of thirteen pollen characters, identified species of *Hidalgoa* in a discrete group and *Dahlia cuspidata* as an outlier. These distinctive attributes in pollen morphology support the idea that pollen morphology is useful for delimitation at the generic level in the *Dahlia* clade. Further evidence from other sources, genetic or anatomical, might contribute to demarcating *Dahlia* and *Hidalgoa*, and provide insight into the family's evolutionary history.

Keywords

Asteraceae, *Dahlia*, hexacolporate, *Hidalgoa*, pollen morphology, tricolporate

Introduction

Hidalgoa La Llave and *Dahlia* Cav. are two closely related genera in tribe Coreopsideae of the Asteraceae (Sørensen 1969; Turner 2010; Sánchez-Chávez et al. 2019). *Hidalgoa* comprises four accepted species (Panero 2007; Turner 2010), with *H. ternata* La Llave having the most widespread distribution, from Mexico to northern South America. The rest have restricted distributions. *H. pentamera* Sherff and *H. uspanapa* B.L. Turner are endemic to southeastern Mexico, and *H. werklei* Hook.f. is distributed in Costa Rica and in the Andean region of Colombia. Habitats for the species in *Hidalgoa* are cloud forests, mainly in microhabitats associated with rivers and very humid places. Remarkably, plants of *Hidalgoa* are vines, climbing onto vegetation by twisting petioles. The heads of *Hidalgoa* have five to twelve pistillate, fertile ray florets, and functionally staminate disc florets (Fig. 1A, B). Cypselae are compressed, with two apical lateral cusps (Panero 2007; Crawford et al. 2009; Turner 2010; Pruski and Robinson 2015).

Dahlia includes 40 species, of which 37 are endemic to Mexico (Villaseñor and Redonda-Martínez 2018; Carrasco-Ortiz et al. 2019; Reyes-Santiago et al. 2019), distributed mostly in pine and oak forests (Carrasco-Ortiz et al. 2019). Based on their chromosome numbers and morphological descriptors such as life form, size, shape, and segmentation of the compound leaves, four sections were recognized in *Dahlia* (Sørensen 1969): sect. *Epiphytum* with a single epiphytic species (*D. macdougallii* Sherff), sect. *Pseudodendron* with three suffrutescent species, sect. *Entemophyllon* with eight suffrutescent species with solid petioles, and sect. *Dahlia* with 28 herbaceous species with hollow petioles (Suppl. material 1: Table S1). With the exception of the epiphytic species, the rest possess tubers, and a few are rupicolous. Some of the species in sect. *Entemophyllon* that live for more than one season, have stems that become quite woody and give the full-grown plants a shrubby aspect. In *Dahlia* the ray florets can be neutral, pistillate fertile or sterile, while the disk florets are hermaphroditic, and vary in number from 15 to 170 (Fig. 1C, D) (Sørensen 1969; Pruski and Robinson 2015). Cypselae are compressed, linear to spatulate, sometimes shallowly tuberculate, with pappus absent or consisting of 2(5) small teeth, weak filiform, sometimes elongated to 1 mm caducous bristles (Sørensen 1969; Panero 2007; Pruski and Robinson 2015).

Hidalgoa was thought to be closely related to the genera *Fitchia* Hook. f., *Moonia* Arn., *Oparanthus* Sherff, and *Petrobium* R. Br., for sharing the character of functional male disc florets (Ryding and Bremer 1992). The similar floral morphology of *Hidalgoa* and *Dahlia* was suggested to be the result of a close relationship (Sørensen 1969; Turner 2010). Furthermore, a previous phylogenetic molecular study that included species of *Dahlia* and *Hidalgoa* found that the latter was embedded in a large clade with *Dahlia*; species of the two genera formed a well-supported monophyletic group (Sánchez-Chávez et al. 2019). However, taxonomic decisions will not be made until additional anatomical and palynological characters, as well as further molecular data, can be analyzed.

In spite of the uniform pollen morphology in the genera of Asteraceae, some pollen characters have been shown to be useful for supporting recognition and delimitation in numerous genera (El-Ghazaly and Anderberg 1995). Some examples



Figure 1. Morphological variation in *Dahlia* and *Hidalgoa* species **A** *Hidalgoa pentamera* **B** *Hidalgoa uspanapa* **C** *Dahlia moorei* **D** *Dahlia mixtecana*. Photos by E. Sánchez-Chávez (**A, C, D**) and Andrés Ortiz (**B**).

of differentiating characters in the taxa of this family are the polar diameter and dimensions of the colpus and endoaperture in *Viguiera* Kunth (Magenta et al. 2010) and in *Xanthium* L., the length and number of spines and the number of columellae (Coutinho et al. 2020). Sexine thickness, the type of aperture, and spine dimensions are the differentiating traits for species of *Stilpnopappus* and *Strophopappus* (Carrijo et al. 2013), as is pollen surface ornamentation in the complex *Phaeostigma* of the genus *Ajania* (Huang et al. 2017). Pollen type and pollen grain shape are taxonomically useful for distinguishing species and genera of the subtribe Lepidaploinae (Marques et al. 2021).

The pollen in tribe Coreopsidae is helianthoid (exine with columellae containing internal foramina and clearly caveate), predominantly spheroidal, tricolporate, echinate, with internal foramina, full cavea present, and endexine much thicker than the foot layer (Blackmore et al. 2009).

Pollen morphology has not been analyzed in detail for the species of *Hidalgoa* or *Dahlia*. Previous palynological research by Wodehouse (1929) on four *Dahlia* species identified the presence of six apertures (hexacolporate) in pollen grains. This attribute constitutes a notable exception within the family. Further studies examined and de-

scribed the meiotic stages of the pollen mother cell in *Dahlia* to determine the development of these apertures (Wodehouse 1930).

The aims of this study are to compile and compare pollen morphology of the species of *Hidalgoa* and *Dahlia* to identify informative characters and understand the relationships and limits of these taxa.

Materials and methods

Twenty-five samples of pollen grains were obtained from herbarium specimens deposited in the IBUG (Instituto de Botánica de la Universidad de Guadalajara) and XAL (Instituto de Ecología, A. C.) herbaria. Vouchers of specimens are included in Table 1.

Pollen grains were acetolyzed according to the methodology of Erdtman (1960), and for difficult material in which compounds formed thin coats on the grains that interfered during the scanning process, the suggestions of Fonnegra (1989) were implemented. The grains were immersed in glacial acetic acid for 24 hours before acetolysis and then transferred to the acetolysis mixture for 1 to 6 hours and the temperature of

Table 1. Studied species of *Hidalgoa* and *Dahlia* for analyzing pollen grains, indicating their voucher and the herbarium in which they were deposited. Herbarium acronyms are according to Index Herbariorum.

Species	Locality	Collector	Herbarium
<i>Dahlia atropurpurea</i> P.D. Sørensen	Guerrero	A. Castro C. 2251	IBUG
<i>Dahlia australis</i> (Sherff) P.D. Sørensen	Puebla	A. Rodríguez C. 6491	IBUG
<i>Dahlia barkenae</i> Knowles & Westc.	Jalisco	A. Castro C. 2304	IBUG
<i>Dahlia brevis</i> P.D. Sørensen	México	A. Rodríguez C. 5869	IBUG
<i>Dahlia campanulata</i> Saar, P.D. Sørensen & Hjert.	Oaxaca	A. Rodríguez C. 6495	IBUG
<i>Dahlia coccinea</i> Cav.	Jalisco	A. Rodríguez C. 7490	IBUG
<i>Dahlia cordifolia</i> (Sessé & Moc.) McVaugh	Guerrero	A. Rodríguez C. 5224	IBUG
<i>Dahlia cuspidata</i> Saar, P.D. Sørensen & Hjert.	Guanajuato	E. Ventura 9581	IBUG
<i>Dahlia dissecta</i> S. Watson		A. Rodríguez C. 6412	IBUG
<i>Dahlia imperialis</i> Roehl ex Ortgies	Chiapas	A. Rodríguez C. 6983	IBUG
<i>Dahlia linearis</i> Sherff	Guanajuato	E. Ventura 6143	IBUG
<i>Dahlia merckii</i> Lehm.		L. Gutierrez s/n	IBUG
<i>Dahlia mollis</i> P.D. Sørensen	Hidalgo	A. Rodríguez C. 6414	IBUG
<i>Dahlia neglecta</i> Saar	Hidalgo	A. Rodríguez C. 6466	IBUG
<i>Dahlia parvibracteata</i> Saar & P.D. Sørensen	Guerrero	A. Rodríguez C. 6092	IBUG
<i>Dahlia pugana</i> Aarón Rodr. & Art. Castro	Jalisco	A. Rodríguez C. 7731	IBUG
<i>Dahlia rudis</i> P.D. Sørensen		A. Ma. Hernández 12	XAL
<i>Dahlia rupicola</i> P.D. Sørensen	Durango	A. Rodríguez C. 6133	IBUG
<i>Dahlia scapigera</i> Knowles & Westc.	Queretaro	E. Gonzalez P. 560	IBUG
<i>Dahlia sorenseii</i> H.V. Hansen & Hjert.		J. Suárez J. 584	IBUG
<i>Dahlia spectabilis</i> Saar & P.D. Sørensen	San Luis Potosí	A. Rodríguez C. 6352	IBUG
<i>Dahlia tenuicaulis</i> P.D. Sørensen	Jalisco	M. Chazaro B. 5736	IBUG
<i>Dahlia wixarika</i> Art. Castro, Carr.-Ortiz & Aarón Rodr.	Jalisco	A. Castro C. 2983	IBUG
<i>Hidalgoa pentamera</i> Sherff	Veracruz	E. Sánchez-Chávez 28	XAL
<i>Hidalgoa ternata</i> La Llave	Veracruz	T. B. Croat 25505	XAL

the water bath was raised to 96 °C. For light microscopy (LM), the pollen grains were mounted in glycerol jelly, sealed, and then examined with a Carl Zeiss Fomi III Optical Microscope, equipped with a Cannon Power Shot G9 digital camera. Permanent slides were deposited in the Palynological Laboratory of the Instituto de Ecología, A. C. The following pollen measurements were obtained from 25 grains per sample: polar axis, equatorial diameter, exine thickness, colpus length, colpus width, ora width, ora length, spinae length, spine width at base and number of apertures.

To observe the pollen with a scanning electron microscope (SEM), acetolyzed pollen grains were washed in ethanol and later in water. Grains were sputter-coated with gold and observed using a Carl Zeiss EVO-50 scanning electron microscope. The terminology of Halbritter et al. (2018) was used, and for pollen structure the terminology of Erdtman (1969) was followed. Number of spines/100 μm^2 , colpus end, base of spine and pollen surface ornamentation were described for five grains per sample. Final morphological data are presented in Table 2.

A matrix based on thirteen pollen characters was constructed. To estimate the quantitative variation within *Dahlia* and *Hidalgoa*, each character was measured and the average for each species was estimated. Qualitative characters were coded with

Table 2. Pollen attributes analyzed on the studied species of *Hidalgoa* and *Dahlia*. The values given in exine, colpus, ora, spine are averages. Cl colpus length, Cw colpus width, Ow Os width, Ol Os length, Swab Spine Width at base.

Specie	Polar axis (P)(μm)			Equatorial diameter (E)(μm)			P/E	Pollen shape	Number of apertures
	Min	Max	Mean	Min	Max	Mean			
<i>Dahlia atropurpurea</i>	28.18	32.70	30.43	28.85	32.46	30.92	0.98	Oblate-spheroidal	Tricolporate
<i>Dahlia australis</i>	25.38	29.98	28.43	23.88	30.49	28.32	1.00	Spheroidal	Tricolporate
<i>Dahlia barkerae</i>	30.62	36.15	33.78	30.86	34.82	33.34	1.01	Prolate-spheroidal	Tricolporate
<i>Dahlia brevis</i>	26.41	32.33	29.04	25.53	32.58	29.41	0.99	Oblate-spheroidal	Tricolporate
<i>Dahlia campanulata</i>	30.61	34.72	32.84	30.50	34.48	32.96	1.00	Spheroidal	Tricolporate
<i>Dahlia coccinea</i>	29.42	39.47	34.74	31.66	39.83	34.79	1.00	Spheroidal	Tricolporate
<i>Dahlia cordifolia</i>	27.80	32.03	29.64	28.00	33.46	30.61	0.97	Oblate-spheroidal	Tricolporate
<i>Dahlia cuspidata</i>	31.21	35.48	33.16	31.63	37.05	34.44	0.96	Oblate-spheroidal	Hexacolporate/ Tricolporate
<i>Dahlia dissecta</i>	28.34	32.50	30.55	27.72	32.21	30.31	1.01	Prolate-spheroidal	Hexacolporate/ Tricolporate
<i>Dahlia imperialis</i>	25.95	30.64	28.04	25.78	31.50	28.94	0.97	Oblate-spheroidal	Hexacolporate/ Tricolporate
<i>Dahlia linearis</i>	29.83	33.79	31.82	29.85	34.99	32.68	0.97	Oblate-spheroidal	Tricolporate
<i>Dahlia merckii</i>	26.19	31.65	28.54	28.64	34.78	31.61	0.90	Oblate-spheroidal	Hexacolporate/ Tricolporate
<i>Dahlia mollis</i>	26.59	30.26	28.38	28.14	32.36	29.93	0.95	Oblate-spheroidal	Tricolporate
<i>Dahlia neglecta</i>	32.72	37.48	35.06	30.75	37.77	35.77	0.98	Oblate-spheroidal	Tricolporate
<i>Dahlia parvibracteata</i>	29.58	32.88	31.29	28.73	32.67	31.57	0.99	Oblate-spheroidal	Hexacolporate/ Tricolporate
<i>Dahlia pugana</i>	29.11	33.23	31.09	29.56	33.94	31.78	0.98	Oblate-spheroidal	Tricolporate
<i>Dahlia rudis</i>	30.08	35.73	33.29	33.40	36.78	34.94	0.95	Oblate-spheroidal	Hexacolporate/ Tricolporate
<i>Dahlia rupicola</i>	28.20	35.37	31.16	26.92	32.40	30.08	1.04	Prolate-spheroidal	Tricolporate
<i>Dahlia scapigera</i>	26.68	32.66	30.56	27.01	31.57	29.38	1.04	Prolate-spheroidal	Tricolporate
<i>Dahlia sorensenii</i>	28.87	34.80	31.90	30.73	36.42	33.90	0.94	Oblate-spheroidal	Hexacolporate/ Tricolporate
<i>Dahlia spectabilis</i>	27.71	34.07	30.33	25.20	32.12	30.11	1.01	Prolate-spheroidal	Tricolporate
<i>Dahlia tenuicaulis</i>	28.54	32.84	31.03	31.58	35.37	33.01	0.94	Oblate-spheroidal	Tricolporate
<i>Dahlia wixarika</i>	27.85	33.23	30.12	27.24	32.07	30.15	1.00	Spheroidal	Tricolporate
<i>Hidalgoa pentamera</i>	25.73	30.35	27.97	25.10	30.27	26.94	1.04	Prolate-spheroidal	Tricolporate
<i>Hidalgoa ternata</i>	28.01	34.07	30.36	26.45	32.08	29.63	1.02	Prolate-spheroidal	Tricolporate

Table 2. Continued.

Specie	Exine (μm)	Colpus (μm)			Os (μm)		Spine		Number of spines/100 μm^2	Base of spine
		Cl	Cw	Copus ends	Ol	Ow	Length (μm)	Swab (μm)		
<i>Dahlia atropurpurea</i>	2.88	3.79	3.12	acute	2.28	2.55	7.80	5.99	7–8	distended
<i>Dahlia australis</i>	2.63	3.01	3.08	acute	2.56	2.05	6.01	4.51	6–8	distended
<i>Dahlia barkerae</i>	3.58	6.38	5.63	acute	2.20	3.09	7.55	7.21	4–5	distended
<i>Dahlia brevis</i>	1.81	4.83	4.74	obtuse	2.06	2.93	7.37	5.79	7–10	distended
<i>Dahlia campanulata</i>	2.24	4.34	1.54	obtuse	1.44	1.92	7.35	6.99	5–7	smooth
<i>Dahlia coccinea</i>	3.05	5.42	4.53	obtuse	3.18	2.71	9.81	8.17	4–5	distended
<i>Dahlia cordifolia</i>	2.96	5.75	4.26	obtuse	2.32	2.74	7.87	6.64	6–7	smooth
<i>Dahlia cuspidata</i>	1.64	8.61	6.22	obtuse	4.75	5.24	9.07	6.94	4–6	distended
<i>Dahlia dissecta</i>	2.88	3.80	2.95	obtuse	2.18	2.35	7.33	6.57	5–7	smooth
<i>Dahlia imperialis</i>	1.98	3.85	2.93	obtuse	2.17	2.93	7.46	5.95	6–8	smooth
<i>Dahlia linearis</i>	2.19	3.98	4.44	obtuse	2.42	3.70	5.29	5.59	8–10	smooth
<i>Dahlia merckii</i>	3.05	4.59	3.16	obtuse	1.45	2.75	6.97	6.25	7–8	distended
<i>Dahlia mollis</i>	1.88	4.14	2.42	obtuse	2.20	2.42	7.79	6.27	5–7	distended
<i>Dahlia neglecta</i>	4.20	4.81	3.34	obtuse	2.35	3.03	6.82	6.80	4–5	distended
<i>Dahlia parvibracteata</i>	3.11	4.82	2.97	obtuse	2.05	2.97	8.97	6.86	6–7	distended
<i>Dahlia pugana</i>	3.42	4.41	4.08	obtuse	2.68	2.23	6.22	6.35	6–7	distended
<i>Dahlia rudis</i>	3.06	4.19	2.50	acute	2.41	2.50	8.19	7.74	6–7	narrower
<i>Dahlia rupicola</i>	1.71	3.36	2.76	obtuse	2.41	2.76	8.36	5.99	6–7	distended
<i>Dahlia scapigera</i>	2.15	6.58	2.65	acute	2.68	2.65	7.90	6.09	4–5	narrower
<i>Dahlia sorensenii</i>	1.68	5.17	2.40	obtuse	1.73	2.40	9.07	6.12	4–5	narrower
<i>Dahlia spectabilis</i>	2.46	4.72	2.05	obtuse	1.54	2.05	8.57	6.00	6–8	distended
<i>Dahlia tenuicaulis</i>	2.82	4.19	4.94	obtuse	2.56	2.79	9.22	6.43	7	distended
<i>Dahlia wixarika</i>	2.42	3.36	2.63	obtuse	2.95	2.63	7.44	6.01	7–9	narrower
<i>Hidalgoa pentamera</i>	1.78	14.97	3.29	acute	2.06	9.98	5.47	6.09	4–5	narrower
<i>Hidalgoa ternata</i>	1.71	14.55	3.34	acute	2.89	8.21	5.80	4.63	4	narrower

the following states: number of apertures (tricolporate:0/ hexacolporate:1), colpus end (obtuse:0/ acute:1), base of spine (narrower:0/ distended:1) alongate ora (absent:0/ present:1; present/absent:2) (Suppl. material 1: Table S2). A principal component analysis (PCA) was run in R (R Core Team, 2019) to evaluate the contribution of each pollen variable to the affiliation of species (Table 3). Graphical representation displayed distribution of thirteen pollen characters.

Results

Pollen grains from a total of 25 species belonging to *Dahlia* (23 species) and *Hidalgoa* (2 species) were analyzed. Table 2 summarizes measurements and character states and Figs 2, 3 and 4 show the diversity in their pollen morphology.

The majority of species analyzed in *Dahlia* are tricolporate. However, two out of ten pollen grains are hexacolporate in *D. cuspidata*, *D. dissecta*, *D. imperialis*, *D. merckii*, *D. parvibracteata*, *D. rudis* and *D. sorensenii*, with three apertures on one hemisphere and three on the other hemisphere (Fig. 2U). The shape is spheroidal-oblate, spheroidal or spheroidal-prolate (P/E = 0.90–1.04) and radially symmetric. Pollen size

Table 3. Palynological characters used in the multivariate analysis of *Hidalgoa* and *Dahlia* species. The contribution of every character for Axis 1 and Axis 2 is indicated (see Fig. 4).

		Character	Axis 1	Axis 2
1	Pa	Polar axis (μm)	1.63	23.53
2	Et	Exine thickness (μm)	6.59	2.30
3	Cl	Colpus length (μm)	25.18	1.19
4	Cw	Colpus width (μm)	8.88	10.75
5	Ow	Os width (μm)	22.82	1.34
6	Ol	Os length (μm)	3.75	11.48
7	Sl	Spinae length (μm)	7.98	10.97
8	Swab	Spine width at base (μm)	5.48	20.32
9	Na	Number of apertures	1.06	2.37
10	Ns	Number of spines/100 μm^2	4.49	11.70
11	Ce	Colpus ends	8.81	0.25
12	Sb	Base of spine	3.14	1.55
13	Osl	Os lalongate	0.20	2.23

is $P = 25.3$ (31) $39.4 \mu\text{m}$, $E = 23.8$ (31.6) $39.8 \mu\text{m}$ (Fig. 6A), and corresponds to a medium grain (Erdtman 1969). The ora are rarely lalongate, and situated distally from the equator, length 1.04 (2.51) $5.41 \mu\text{m}$, and width 1.27 (3.60) $8.61 \mu\text{m}$ (Figs 2B, 2F, 6B), rarely acute. Colpus usually short, almost equal to ora length, more or less oval to oblong, length 2.24 (4.81) $9.2 \mu\text{m}$, and width 1.27 (2.85) $5.89 \mu\text{m}$ (Figs 3E, 3H, 3K, 3N, 4I, 4M, 6C), apices obtuse to acute. Exine thickness thin, excluding spines, ranging from 1.1 (2.59) to $5.6 \mu\text{m}$ (Fig. 2). Ornamentation echinate; spines 4 (6–7) $10/100 \mu\text{m}^2$, spine length from 4.2 (7.75) to $12.26 \mu\text{m}$, and width at base from 3.06 (6.40) to $10.35 \mu\text{m}$, shape of spines more deltate than conical (Fig. 6D), with a distended or narrower base and with acuminate apex (Figs 3, 4). Tectum with the base of the spine always microperforate.

Pollen grains in the species of *Hidalgoa* analyzed are tricolporate and spheroidal-prolate ($P/E = 1.02$ – 1.04), radially symmetric. Pollen size is $P = 25.7$ (29.1) $34 \mu\text{m}$, $E = 25.1$ (28.5) $32 \mu\text{m}$ (Fig. 6A), and corresponds to a medium grain (Erdtman 1969). The lalongate ora length 1.71 (2.47) to $4.87 \mu\text{m}$, and width 6.32 (9.39) to $13.02 \mu\text{m}$ (Figs 2W, 6B), usually wider than longer and with acute apices. The colpus is elliptical, length 11.28 (14.76) to $16.83 \mu\text{m}$, and width 2.69 (3.31) to $4.21 \mu\text{m}$ (Fig. 6C), apex always acute (Figs 3A, 3B, 4N). Exine is thinner, 1.12 (1.74) $2.95 \mu\text{m}$ excluding the spines. Ornamentation is echinate; spines 4–5/100 μm^2 , spine length ranging from 4 (5.6) to $6.77 \mu\text{m}$ and width at base 3.6 (5.3) to $7.3 \mu\text{m}$, shape of spines conical (Figs 3C, 4O, 6D), with apex acute. Tectum with base of spine always microperforate. Palynological characters of the two studied species of *Hidalgoa* are similar, only a slight variation in pollen grain size was detected.

Results of the PCA indicate that the first two components explain 46.04% of the observed variation (see Table 3 to for the contribution of each variable to Dim 1 and Dim 2). A bidimensional projection of the axes of the two first components is displayed in Fig. 5. The first principal component explains 25.19% of the variation and is associated with colpus length (Cl) and os width (Ow). The second principal component explains




Figure 2. Pollen grains of *Dahlia* and *Hidalgoa* observed with light microscopy (LM) **A** *D. australis* **B** *D. barkerae* **C** *D. brevis* **D** *campanulata* **E** *D. coccinea* **F** *D. cordifolia* **G** *D. cuspidata* **H** *D. dissecta* **I** *D. imperialis* **J** *D. linearis* **K** *D. merckii* **L** *D. mollis* **M** *D. neglecta* **N** *D. parvibracteata* **O** *D. pugana* **P** *D. rudis* **Q** *D. rupicola* **R** *D. scapigera* **S** *D. sorensenii* **T** *D. spectabilis* **U** *D. rudis*, pollen grain hexacolporate with three apertures on one hemisphere and three on the other hemisphere **V** *H. ternata* **W** *H. pentamera*. Scale bars: 10 μ m.




Figure 3. Scanning Electron Microscope (SEM) images of *Dahlia* and *Hidalgoa* pollen grains **A–C** *Hidalgoa ternata* **A** equatorial view **B** detail of colpus **C** detail of spine **D–F** *Dahlia australis* **D** polar view **E** detail of colpus **F** detail of spine **G–I** *Dahlia cuspidata* **G** equatorial view **H** detail of colpus **I** detail of spine **J–L** *Dahlia neglecta* **J** equatorial view **K** detail of colpus **L** detail of spine **M–O** *Dahlia coccinea* **M** equatorial view **N** detail of colpus **O** detail of spine.




Figure 4. Scanning Electron Microscope (SEM) images of *Dablia* and *Hidalgoa* pollen grains **A, B** *Dablia atropurpurea* **C** *Dablia barkerae* **D, E** *Dablia brevis* **F** *Dablia dissecta* **H** *Dablia imperialis* **I** *Dablia linearis* **J** *Dablia merckii* **K** *Dablia scapigera* **L** *Dablia tenuicaulis* **M** *Dablia wixarika* **N–O** *Hidalgoa pentamera*.

20.85% and variables that contributed the most are polar axis (Pa), spine width at base (Swab), spine length (Sl), colpus width (Cw), os length (Ol), number of spines/100 μm^2 (Ns). Length of arrows in Fig. 5 suggests adequate sampling for all characters, except for alongate os (Osl), number of apertures (Ap) and base of spine (Sb).




Figure 6. Comparison of *Dahlia* and *Hidalgoa* pollen grains **A** box plot of Polar diameter **B** os length/width **C** colpus length/width **D** spine length/width at base. **1** *Dahlia atropurpurea*. **2** *Dahlia australis*. **3** *Dahlia barkerae*. **4** *Dahlia brevis*. **5** *Dahlia campanulata*. **6** *Dahlia coccinea*. **7** *Dahlia cordifolia*. **8** *Dahlia cuspidata*. **9** *Dahlia dissecta*. **10** *Dahlia imperialis*. **11** *Dahlia linearis*. **12** *Dahlia merckii*. **13** *Dahlia mollis*. **14** *Dahlia neglecta*. **15** *Dahlia parvibracteata*. **16** *Dahlia pugana*. **17** *Dahlia rudis*. **18** *Dahlia rupicola*. **19** *Dahlia scapigera*. **20** *Dahlia sorensonii*. **21** *Dahlia spectabilis*. **22** *Dahlia tenuicaulis*. **23** *Dahlia wixarika*. **24** *Hidalgoa pentamera*. **25** *Hidalgoa ternata*.

aperture varies from tricolporate to hexacolporate. Hexacolporate species observed here were: *D. cuspidata*, *D. dissecta*, *D. imperialis*, *D. merckii*, *D. parvibracteata*, *D. rudis* and *D. sorensonii*. Wodehouse (1930) recognized the same pattern in *D. brevis*, *D. coccinea*, *D. pinnata* and *D. imperialis*. However, Wodehouse (1930) described these six apertures as uniform in all pollen grains; the observations were made on species belonging to the San Francisco *Dahlia* Society, plants that are probably of hybrid origin. The specimens collected from the herbariums for this study do not display characters of hybrid origin. Furthermore, hexacolporate grains have been indeed reported in Old World Vernoniae and *Adenanthemum* (Blackmore et al 2009). Other members of Coreopsideae such as *Bidens* also vary in the number of colpi from 3–4 (Tadesse et al. 1995) or polypentaporate (Younis et al. 2020).

The os and colpus displayed more variation in *Dahlia* than in *Hidalgoa*. In *Hidalgoa* the os is alongate, the widest is up to 13 µm width, and the longest colpus is approximately 17 µm, with apices always acute. These traits of *Hidalgoa* are similar to those observed in *Bidens* (Tadesse et al. 1995). In *Dahlia* the ora are either alongate or longate. The os is slightly wider than larger and apices are obtuse, rarely acute. The

widest and largest colpus was observed in *D. cuspidata* (5.89 μm and 9.2 μm respectively) and *D. scapigera* (ca. 7 μm), and the smallest in *D. australis* (2.9 μm). Variation in the apertures like those of *Dahlia* has not been reported in other genera of the tribe Coreopsidae (Blackmore et al. 2009).

Spines are variable between *Dahlia* and *Hidalgoa*. In *Hidalgoa* they are conical and smaller (4.08 to 6.77 μm) while in *Dahlia* they are deltate and larger (4.29 to 12.26 μm), with exception of *D. linearis* (smaller). Spines in *Dahlia* varied more in shape and size. They are commonly triangular or deltate with a broadened base (distended base), as described by Tellería (2017) for tribe Coreopsidae. Sometimes spines emerge abruptly from the exine surface, e.g. *D. campanulata*, *D. imperialis*, *D. parvibracteata*, *D. scapigera*. These spines are similar to those described in *Coreopsis* (Tadesse et al. 1995). The transition between the microperforate basal portion of the spine and the unperforated apical portion is abrupt in almost all species, except in *D. cordifolia*, *D. linearis* and *D. sorensenii*. Exine thickness did not vary among *Hidalgoa* and *Dahlia* species. The thinnest was observed in *D. cuspidata* and *D. sorensenii* (1.6 μm) and the thickest in *D. neglecta* (4.2 μm).

Multivariate analyses did not reveal a clear clustering among species of *Dahlia* according to the sections proposed by Sørensen (1969), based mainly on life form and in the phylogeny of Saar et al. (2003). Nevertheless, *D. cuspidata* and the *Hidalgoa* species are significantly different from the other *Dahlia* species included in this study. *Hidalgoa* species have long colpi and wide ora, and *D. cuspidata* has long ora. *Dahlia cuspidata* possesses unusual morphological characters such as large involucrel outer bracts and cuspidate leaf shape. Thus, further research might decide the position of this species.

The most recent phylogeny that included *Dahlia* and *Hidalgoa* (Sánchez-Chávez et al. 2019) identified *Hidalgoa* within the *Dahlia* clade. However, both genera are morphologically complex (Sørensen 1969; Turner 2010), and contrasting characters such as life form, number and arrangement of fertile and sterile flowers have been used to separate these two groups. A further phylogeny, including all species, may help us to better understand pollen evolution in the genus.

Conclusions

The palynological descriptions for *Hidalgoa* presented here are the first, and despite the similarities in its floral morphology to that of *Dahlia*, its pollen is remarkably different, mostly in colpus length and shape of their spines. *Hidalgoa* has pollen grains with large colpi and small, conical spines. In addition, the length of the elongate ora differ. Hexacolporate grains with a distended base, were found in a number of *Dahlia* species but have not been identified in *Hidalgoa*. Likewise, morphological characters such as pistillate fertile ray florets, cypselae with two apical lateral cusps and twisting petioles in *Hidalgoa* contrast with the ray florets, which can be fertile, pistillate or sterile, cypselae with pappus absent or present with 2(5) small teeth or of two weak filiform, caducous bristles of *Dahlia*; characters that have been utilized to tell these

two genera apart. The results obtained in this palynological study support the idea that pollen morphology is useful for delimitation at the generic level in the *Dahlia* clade. These differentiating attributes in pollen morphology in the species of *Dahlia* and *Hidalgoa* indicate that they should be recognized as separate genera. However, as indicated above, additional anatomical and molecular characters are needed to make the taxonomic decision and help us understand evolution in the genera, and their relationship to other genera in Coreopsideae.

Acknowledgements

We thank Tadesse Mesfin, Arturo Castro-Castro and an anonymous reviewer for comments and suggestions that greatly improved this manuscript. We thank the curators of the IBUG and XAL herbaria for kindly allowing us to sample the pollen of *Hidalgoa* and *Dahlia*. We are grateful to Bianca Delfosse for editing our English. Sonia Galicia provided invaluable help in the anatomy lab and with optical microscopy, and Ana Lucía Tovar Álvarez produced the SEM photomicrographs of LANIVEG-QRO. E. Sánchez-Chávez was supported by a Doctorate in Science scholarship from CONA-CyT, Mexico (780481).

References

- Blackmore S, Wortley AH, Skvarla JJ, Robinson H (2009) Evolution of pollen in Compositae. In: Funk, VA, Susanna A, Stuessy TF, Bayer RJ (Eds) Systematics, evolution and biogeography of Compositae. International Association for Plant Taxonomy, Vienna, 101–130.
- Carrasco-Ortiz M, Munguía-Lino G, Castro-Castro A, Vargas-Amado G, Harker M, Rodríguez A (2019) Riqueza, distribución geográfica y estado de conservación del género *Dahlia* (Asteraceae) en México. Acta Botánica Mexicana 126: 1–24. <https://doi.org/10.21829/abm126.2019.1354>
- Carrijo TT, Garbin ML, Picanço Leite W, Mendonça CBF, Esteves RL, Gonçalves-Esteves V (2013) Pollen morphology of some related genera of Vernonieae (Asteraceae) and its taxonomic significance. Plant Systematics and Evolution 299(7): 1275–1283. <https://doi.org/10.1007/s00606-013-0795-9>
- Coutinho AP, Silveira P, Pita C, Santos MJ, Saraiva C, Perpétuo NC (2020) Pollen morphology of *Xanthium* L. (s.l.) (Asteraceae, Asteroideae, Heliantheae, Ambrosiinae) in the Iberian Peninsula – a palynotaxonomic approach to a poisonous, allergenic and invasive genus. Grana 60(1): 35–56. <https://doi.org/10.1080/00173134.2020.1737729>
- Crawford DJ, Tadesse M, Mort M, Kimball RT, Randle CP (2009) Coreopsideae. In: Funk VA, Susanna A, Stuessy TF, Bayer RJ (Eds) Systematics, evolution and biogeography of Compositae. International Association for Plant Taxonomy, Vienna, 713–727.
- El-Ghazaly G, Anderberg AA (1995) Pollen morphology of *Phagnalon* and *Aliella* (Asteraceae, Gnaphalieae) and its taxonomical implications. Grana 34(2): 89–99. <https://doi.org/10.1080/00173139509429999>

- Erdtman G (1960) The acetolysis method, a revised description. *Svensk Botanisk Tidskrift* 54(4): 561–564.
- Erdtman G (1969) Handbook of palynology: morphology, taxonomy, ecology; an introduction to the study of pollen grains and spores. Munksgaard, 486 pp.
- Fonnegra R (1989) Métodos de Estudio Palinológico. Centro de Investigaciones de la Facultad de Ciencias Naturales y Exactas Universidad de Antioquia, Medellín, 57 pp.
- Halbritter H, Ulrich S, Grímsson F, Weber M, Zetter R, Hesse M, Buchner R, Svojtka M, Frosch-Radivo A (2018) Illustrated Pollen Terminology. Illustrated Pollen Terminology. Springer International Publishing, 487 pp. <https://doi.org/10.1007/978-3-319-71365-6>
- Huang Y, An YM, Meng SY, Guo YP, Rao GY (2017) Taxonomic status and phylogenetic position of *Phaeostigma* in the subtribe Artemisiinae (Asteraceae). *Journal of Systematics and Evolution* 55(5): 426–436. <https://doi.org/10.1111/jse.12257>
- Magenta MG, Nunes ADD, Mendonça CBF, Gonçalves-Esteves V (2010) Palynotaxonomy of Brazilian *Viguiera* (Asteraceae) species. *Boletín de La Sociedad Argentina de Botánica* 45(3–4): 285–299.
- Marques D, do Pico GMV, Nakajima JN, Dematteis M (2021) Pollen morphology and its systematic value to southern South American species of *Lepidaploa* (Vernoniaeae: Asteraceae). *Rodriguésia* 72: 1–18. <https://doi.org/10.1590/2175-7860202172017>
- Panero JL (2007) Coreopsidae. In: Kadereit JW, Jeffrey C (Eds) The families and genera of vascular plants VIII. Flowering Plants. Eudicots, Asterales. The families and genera of vascular plants. Springer, Berlin, 406–428.
- Pruski JF, Robinson H (2015) Asteraceae. In: Davidse G, Sousa-Sánchez M (Eds) *Flora Mesoamericana* 5(2): 1–1362.
- R Core Team (2019) R: A language and environment for statistical computing. R Foundation for Statistical Computing. <https://www.r-project.org/>
- Reyes-Santiago J, Islas-Luna MA, Munguía-Lino G, Castro-Castro A (2019) *Dahlia mixtecana* (Asteraceae, Coreopsidae), a striking new species from Oaxaca, Mexico. *Phytotaxa* 394(3): 209–218. <https://doi.org/10.11646/phytotaxa.394.3.2>
- Ryding O, Bremer K (1992) Phylogeny, distribution, and classification of the Coreopsidae (Asteraceae). *Systematic Botany* 17(4): 649–659. <https://doi.org/10.2307/2419733>
- Saar DE, Polans NO, Sørensen PD (2003) A phylogenetic analysis of the genus *Dahlia* (Asteraceae) based on internal and external transcribed spacer regions of nuclear ribosomal DNA. *Systematic Botany* 28(3): 627–639. <http://www.jstor.org/stable/25063902>
- Sánchez-Chávez E, Rodríguez A, Castro-Castro A, Pérez-Farrera MA, Sosa V (2019) Spatio-temporal evolution of climbing habit in the *Dahlia-Hidalgoa* group (Coreopsidae, Asteraceae). *Molecular Phylogenetics and Evolution* 135: 166–176. <https://doi.org/10.1016/j.ympev.2019.03.012>
- Sherff EE (1947) Addition to the genera *Scalesia* Arn and *Hidalgoa* Llave and Lex. Publications of the Field Museum of Natural History. Botanical Series 23 (7): 334–335.
- Sørensen PD (1969) Revision of the genus *Dahlia* (Compositae, Heliantheae– Coreopsidinae). *Rhodora* 71: 309–365.
- Tadesse M, Crawford DJ, Smith EB (1995) Pollen morphology of North American *Coreopsis* (Compositae – Heliantheae). *Grana* 34(1): 21–27. <https://doi.org/10.1080/00173139509429029>

- Tellería MC (2017) Spines vs. microspines: An overview of the sculpture exine in selected basal and derived Asteraceae with focus on Asteroideae. *Journal of Plant Research* 130(6): 1023–1033. <https://doi.org/10.1007/s10265-017-0956-y>
- Turner BL (2010) The Comps of Mexico. A systematic account of the family Asteraceae (Chapter 10: Subfamily Coreopsideae). *Phytologia Memoirs* 1: 1–224.
- Villaseñor JL, Redonda-Martínez R (2018) A new species of *Dahlia* (Asteraceae, Coreopsideae) from the state of Oaxaca, Mexico. *Phytotaxa* 362(2): 239–243. <https://doi.org/10.11646/phytotaxa.362.2.11>
- Wodehouse RP (1929) The origin of symmetry patterns of pollen grains. *Bulletin of the Torrey Botanical Club* 56(7): 339–350. <https://doi.org/10.2307/2480750>
- Wodehouse RP (1930) The origin of the six-furrowed configuration of *Dahlia* pollen grains. *Bulletin of the Torrey Botanical Club* 57(6): 371–380. <https://doi.org/10.2307/2480640>
- Younis S, Shaheen S, Harun N, Abbas M, Ahmad M, Hussain K, Khan F (2020) Application of multimicroscopic techniques (LM and SEM) in comparative palynological study of Asteroideae members, inhabited in Pakistan. *Microscopy Research and Technique* 84(5): 1063–1077. <https://doi.org/10.1002/jemt.23667>

Supplementary material I

Tables S1, S2

Authors: Erandi Sánchez-Chávez, Andrew Vovides, Victoria Sosa

Data type: Tables (docx. file)

Explanation note: **Table S1.** *Dahlia* diversity. Sections proposed by Sorensen (1969).

Table S2. Data matrix showing the values found for the 13 characters (columns) in the 25 species (rows) of *Hidalgoa* and *Dahlia* included in this study. (Characters designated according to Table 3).

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/phytokeys.199.79501.suppl1>