

Cephalaria anamurensis (Caprifoliaceae), a new species from south Anatolia, Turkey

Ramazan Süleyman Göktürk¹, Hüseyin Sümbül¹

¹ Department of Biology, Faculty of Science, Akdeniz University, 07058, Antalya, Turkey

Corresponding author: Ramazan Süleyman Göktürk (gokturk@akdeniz.edu.tr)

Academic editor: C. Morden | Received 21 March 2016 | Accepted 10 June 2016 | Published 15 June 2016

Citation: Göktürk RS, Sümbül H (2016) *Cephalaria anamurensis* (Caprifoliaceae), a new species from south Anatolia, Turkey. *PhytoKeys* 65: 25–33. doi: 10.3897/phytokeys.65.8571

Abstract

A new species, *Cephalaria anamurensis* (Caprifoliaceae) is described and illustrated from south Anatolia, Turkey. The species grows on steppe and stony places in Anamur (C4 Mersin province) in south Anatolia. Diagnostic morphological characters from closely similar taxa are discussed. The geographical distribution of the new species and two closely related species in Turkey are mapped.

Keywords

Cephalaria, New species, Taxonomy, Turkey

Introduction

The genus *Cephalaria* Schrad. ex Roem. & Schult. was first described by J.J. Roemer and J.A. Schultes (Roemer and Schultes 1818). It is distributed from Mediterranean area to west China, and some of the species are also found in southern Africa (Szabó 1940). The genus *Cephalaria* has long been regarded as belonging to the Dipsacaceae, whereas according to APG III it is included within the larger family Caprifoliaceae (Dipsacales) and consist of 100 species (Reveal and Chase 2011). In Turkey, the family Caprifoliaceae is represented by 12 genera, namely *Centranthus* DC. (3 spp.), *Cephalaria* (39 spp.), *Dipsacus* L. (5 spp.), *Knautia* L. (9 spp.), *Lonicera* L. (11 spp.), *Morina* L. (2 spp.), *Pterocaphalus* Adans. (9 spp.), *Scabiosa* L. (32 spp.), *Succisa* Haller (1 sp.), *Tremastelma* Rafin. (1 sp.), *Valeriana* L. (14 spp.) and *Valerianella* Mill. (31 spp.) (Güner et al. 2012).

In July 2015, the authors collected a specimen of *Cephalaria* from the Taurus Mountains (Mersin province) during fieldwork for Expo Antalya 2016 natural plant supply project. Fruiting material was gathered in the same area in September 2015 by the authors. This specimen has been compared to many specimens including two supposedly closely related species in the Herbaria of Akdeniz University Herbarium, ANK, GAZI and HUB, records in the literature (Szabó 1940, Göktürk et al. 2012, Göktürk and Sümbül 2014) and consulting floras of Turkey and neighboring countries (Bobrov 1957, Matthews 1972, Ferguson 1976, Feinbrun-Dothan 1978, Matthews 1988, Lack 1991). Successful efforts have also been made to find additional locations in this vicinity where this novel plant may be located. After comparison with material of morphologically similar taxa, we concluded that these specimens represent a species new to science.

Materials and methods

In a total, five specimens (23 individuals) of the new species were collected from three adjacent localities. The illustrations of the species were made from dry materials by using Adobe Photoshop CS4. Herbarium studies were made in Akdeniz University Herbarium, which has the richest *Cephalaria* collections in Turkey, and ANK, GAZI and HUB.

Taxonomic treatment

Cephalaria anamurensis Göktürk & Sümbül, sp. nov.

urn:lsid:ipni.org:names:77155495-1

Figs 1, 2

Diagnosis. *Cephalaria anamurensis* is similar to *C. elmaliensis* Hub.-Mor. & V. A. Matthews and *C. speciosa* Boiss. & Kotschy. It can be distinguished from them by its rhizomatous growth, the lower stem leaves 35–55 × 3.5–4.5 cm, a globose capitula, involucre bracts that are ovate-orbicular to triangular-ovate, 3–7 × 2.5–6 mm, and completely blackish or blackish on the dorsal side and at the acute or subobtusate apex, receptacular bracts that are oblanceolate and blackish at an acuminate apex, and the involucre with long teeth 3–4 mm long and short teeth 1–1.5 mm long.

Type. TURKEY, Mersin, Anamur, Anamur to Kazancı, Kırkkuyu, Bıçkıcı boğazı, 36°28'35"N; 032°44'11"E, 1784 m, steppe and stony places, 24 July 2015 Göktürk 8018, Sümbül & Çingay (holotype: Akdeniz University Herbarium 3446!; isotypes: ANK!, GAZI!, HUB!, NGBB!).

Description. Plant stout, erect, perennial, rhizomatous herbs, up to 1.5 m, simple, striate, covered with densely stellate hairy and retrorse hairy in lower part. Leaves coriaceous, densely stellate hairy; lower stem leaves simple, lanceolate, 35–55 × 3.5–4.5




Figure 1. Photographs and distribution map of *Cephalaria anamurensis*. **A** habit and habitat of the type plant **B** Close-up of flowering capitula **C** Close-up of fruiting capitula **D** Distribution of *Cephalaria anamurensis* (★), *C. elmaliensis* (●) and *C. speciosa* (■) in Turkey. Photos: Ramazan Süleyman Göktürk.



Figure 2. **A** Holotype specimen of *Cephalaria anamurensis* Göktürk & Sümbül **B** Reproductive parts of *Cephalaria anamurensis* (Göktürk 8018) (1), *C. elmaliensis* (Göktürk 3532) (2) and *C. speciosa* (Göktürk 4727) (3): **a** corolla **b** involucre bract **c** receptacular bract **d** involucre **e** calyx.

cm, crenate-serrate, acuminate; cauline leaves simple or lyrate; simple leaves lanceolate or broadly lanceolate, $10\text{--}30 \times 1.5\text{--}5.5$ cm, entire, acuminate; lyrate leaves lanceolate or broadly lanceolate in outline, $10\text{--}25 \times 2\text{--}5$ cm, with 2–3 segments, lateral segments lanceolate, $0.3\text{--}1.5 \times 0.1\text{--}0.4$ cm, entire, acute, terminal segment larger than lateral ones, lanceolate or broadly lanceolate, $8\text{--}22 \times 2\text{--}5$ cm, margins entire or crenate-serrate, acute; upper stem leaves simple, sessile, linear or linear-lanceolate, $0.5\text{--}6 \times 0.2\text{--}0.8$ cm, entire, acuminate. Capitula globose, 25–50-flowered, 2–4 cm in diameter in flower, 2–3 cm in diameter in fruit. Involucre bracts ovate-orbicular to triangular-ovate, $3\text{--}7 \times 2.5\text{--}6$ mm, completely blackish or blackish in dorsal side and at apex, pubescent or adpressed pilose, margins ciliate, acute or subobtusate at apex; receptacular bracts oblanceolate, $8\text{--}13 \times 2\text{--}4$ mm, straw-coloured on dorsal side and base, blackish at apex, pubescent and adpressed pilose in dorsal side and apex, margins ciliate, acuminate at apex. Calyx cupuliform, 1–2 mm in diameter, with irregular teeth. Corolla cream or pale yellow, 8–15 mm long, densely adpressed hairy outside. Involucre 4-angled, 7–13 mm long in fruit, pilose, 4 long and 4 short teeth at apex; long teeth 3–4 mm long, short teeth 1–1.5 mm long. Flowering from July to August, fruiting from August to September.

Distribution and ecology. This species is endemic to South Anatolia, Turkey, and east Mediterranean (mountain) element (Fig. 1D). According to EUNIS (2007),

habitat type of this new species is E2.5 (Meadows of the steppe zone). *Cephalaria anamurensis* grows in steppe and stony places at an elevation of 1784–1800 m. It is associated with endemics such as *Marrubium lutescens* Boiss. & Heldr. subsp. *micranthum* (Boiss. & Heldr.) P. H. Davis, *Nepeta nuda* L. subsp. *glandulifera* Hub.-Mor. & P. H. Davis, *Papaver pilosum* Sibth. & Sm. subsp. *pilosum*, *Sideritis libanotica* Labill. subsp. *violascens* (P. H. Davis) P. H. Davis, *Verbascum cucullatibracteum* Hub.-Mor. and non-endemic plants such as *Digitalis ferruginea* L. subsp. *ferruginea*, *Euphorbia kotschyana* Fenzl, *Berberis crataegina* DC., *Genista albida* Willd., *Hordeum bulbosum* L., *Onopordum sibthorpiatum* Boiss. & Heldr., *Phlomis armeniaca* Willd., *Scutellaria orientalis* L. subsp. *pinnatifida* J. R. Edm. and *Thymus sipyleus* Boiss.

Conservation status. This species is known only from three adjacent localities with small populations in Kırkkuyu collected by the authors. It is suggested that this new species should be placed under the IUCN threat category “Critically Endangered (CR)” (IUCN 2014) because the estimated area of occupancy is less than 10 km², the population size of the new species is estimated to be less than 50 mature individuals, and the population size of the new species could be reduced in the near future based on heavily grazing pressure [CR B2; C2a(i)].

Etymology. The specific epithet is derived from the name of Anamur district in Mersin province, where the holotype of *C. anamurensis* was collected.

Results

The new species is included in a group of *Cephalaria* species that are covered with stellate hairs. There are nine species in this group in Turkey, and eight of them are endemic to the country. Only *Cephalaria stellipilis* Boiss. has a distribution extending out of Turkey to Lebanon. *Cephalaria anamurensis* is morphologically closest to *C. elmaliensis* and *C. speciosa*. *Cephalaria elmaliensis* is endemic to Çıglikara nature protection area and grows on stony ground and in openings in cedar forests (*Cedrus libani* A. Rich. var. *libani*) in the Elmalı district/Antalya. *Cephalaria speciosa* is endemic to east Anatolia and grows in rocky places and roadsides (Fig. 1D). A comparison of *Cephalaria anamurensis*, *C. elmaliensis* and *C. speciosa* is given in Table 1.

The genus *Cephalaria* was represented by 29 species in the *Flora of Turkey and the East Aegean Islands* (Matthews 1972). Since then, 11 new species, one new subspecies and one variety have been described from Turkey (Matthews 1988, Sümbül 1991, Göktürk and Sümbül 1997, Göktürk et al. 2003, Göktürk and Sümbül 2003, Kuş and Göktürk 2005, Aksoy et al. 2007, Parolly and Eren 2007, Göktürk et al. 2012). Göktürk and Sümbül (2014) placed *C. amana* Rech. f. as a synonym of *C. taurica* Szabó. According to Göktürk and Sümbül (2014), the total number of species of *Cephalaria* reported from Turkey is 39 and the total number of taxa of *Cephalaria* is 41 in Turkey. With the description here of *C. anamurensis* the number of species in Turkey is now 40 and the total number of taxa of *Cephalaria* is also 42 in Turkey.

Table 1. Morphological comparison of *Cephalaria anamurensis*, *C. elmaliensis* and *C. speciosa*.

Characters	<i>C. anamurensis</i>	<i>C. elmaliensis</i>	<i>C. speciosa</i>
Stem	rhizomatous, up to 1.5 m, retrorse hairy in lower part	not rhizomatous, up to 1 m, antrorse hairy in lower part	not rhizomatous, up to 1.5 m, retrorse hairy in lower part
Lower stem leaves	lanceolate, 35–55 × 3.5–4.5 cm, crenate-serrate, acuminate	lanceolate, 10–26 × 1.3–4 cm, entire or crenate-serrate, acute	oblong-lanceolate, 10–40 × 2.5–6.5 cm, entire or crenate-serrate, acute or acuminate
Lyrate cauline leaves	lanceolate or broadly lanceolate in outline, 10–25 × 2–5 cm, with 2–3 segments	narrowly ovate-lanceolate in outline, 6–15.5 × 0.8–1.6 cm, with 2–6 segments	lanceolate or oblong-lanceolate in outline, 8–20 × 3–6 cm, with 2–6 segments
Capitula	globose, 2–4 cm in diameter in flower	subglobose, 1–2 cm in diameter in flower	ovate to globose, 2.5–4.5 cm in diameter in flower
Involucral bracts	ovate-orbicular to triangular-ovate, 3–7 × 2.5–6 mm, completely blackish or blackish in dorsal side and at apex, acute or subobtuse	ovate, 4–6 × 2.5–3 mm, completely straw-coloured or only brown at apex, acute	ovate to triangular- lanceolate, 7–15 × 3–7 mm, completely straw-coloured, acuminate or subacuminate
Receptacular bracts	oblanceolate, 8–13 × 2–4 mm, blackish at apex, acuminate	ovate or lanceolate, 8–12 × 2–3 mm, completely straw-coloured, acute	triangular-lanceolate, 12–20 × 3.5–6 mm, brown at apex, pungent
Involucel	long teeth 3–4 mm long, short teeth 1–1.5 mm long	long teeth 3 mm long, short teeth 1 mm long	long teeth 4 mm long, short teeth 2 mm long

A morphological key of *Cephalaria* species in Turkey with stellate hairs.

- 1 Lower stem leaves lyrate.....2
- Lower stem leaves simple4
- 2 Lower stem leaves only with sparsely stellate hairs, stem hollow ... *C. demirizii*
- Lower stem and cauline leaves with dense stellate hairs, stem not hollow3
- 3 Bracts blackish at apex; involucral bracts acute or subacute; receptacular bracts ovate-oblong..... *C. davisiana*
- Bracts straw-coloured at apex; involucral bracts obtuse; receptacular bracts oblong or narrowly oblanceolate *C. sumbuliana*
- 4 Petiole of lower stem leaves deflexed.....*C. duzceënsis*
- Petiole of lower stem leaves not deflexed5
- 5 Capitula ovoid; involucel sericeous *C. elazigensis*
- Capitula not ovoid; involucel pilose.....6
- 6 Plant stout, greater than 1 m high.....7
- Plant slender, up to 1 m high.....8
- 7 Capitula ovate-globose; involucral bracts ovate to triangular-lanceolate.....
..... *C. speciosa*
- Capitula globose; involucral bracts ovate-orbicular to triangular-ovate.....
..... *C. anamurensis*
- 8 Lower and cauline stem leaves lanceolate*C. elmaliensis*
- Lower and cauline stem leaves oblong-spathulate.....*C. stellipilis*

Specimens examined

Cephalaria anamurensis (Paratypes), TURKEY- C4 Konya: Ermenek, Kazancı district plateau, Kırkkuyu, 1800 m, 19.07.1984, *Sümbül 3217* (HUB!, ANK!); C4 Mersin: Anamur, Anamur to Kazancı, Kırkkuyu, Bıçkıcı boğazı, 36°28'31"N; 032°44'53"E, 1800 m, steppe and stony places, 24.07.2015, *Göktürk 8020, Sümbül & Çingay* (Akdeniz University Herbarium!); C4 Mersin: Anamur, Anamur to Kazancı, Kırkkuyu, Bıçkıcı boğazı, 36°28'20"N; 032°45'05"E, 1800 m, steppe and stony places, 24.07.2015, *Göktürk 8024, Sümbül & Çingay* (Akdeniz University Herbarium!); C4 Mersin: Anamur, Anamur to Kazancı, Kırkkuyu, Bıçkıcı boğazı, 36°28'20"N; 032°45'05"E, 1800 m, steppe and stony places, 29.09.2015, *Göktürk 8066 & Sümbül* (fruiting) (Akdeniz University Herbarium!).

Cephalaria elmaliensis TURKEY- C2 Antalya: Elmalı, Çıglıkara, near security building, in openings in *Cedrus libani* var. *libani*, 1700–1900 m, 25.08.1993, *H. Duman et al. 5345* (GAZI!, Akdeniz University Herbarium!); C2 Antalya: Elmalı, Çıglıkara, near security building, in openings in *Cedrus libani* var. *libani*, 1700–1900 m, 12.08.1995, *Göktürk 3532* (Akdeniz University Herbarium!); Elmalı, Çıglıkara, between Ayıngediği-Kaş gediği, in openings in *Cedrus libani* var. *libani*, stony ground, 1750 m, 17.08.2007, *Göktürk 6111* (Akdeniz University Herbarium!).

Cephalaria speciosa TURKEY- B7 Erzincan: Keşiş mountain, Cimin, rocky slopes, c. 2300 m, 28.08.1957, *P. H. Davis 31828* (ANK!); Kemah, above Kömürköy, steppe, 1850 m, 31.07.1996, *Dönmez 5367* (HUB!); Tunceli: Ovacık, Munzur Mountain, Aksu stream, c. 1700 m, 21.07.1957, *P. H. Davis 31462 & I. C. Hedge* (ANK!); B8 Erzincan: Aşkale to Tercan, dry rocky places, c. 1700 m, 25.08.1957, *P. H. Davis 32657 & I. C. Hedge* (ANK!); between Tercan-Aşkale, slopes, 1765 m, 17.08.2000, *Göktürk 4528 & F. Göktürk* (Akdeniz University Herbarium!); B8 Muş: Varto, Muş to Varto, rocky slopes, 1350 m, 02.08.2001, *Göktürk 4727 & M. Göktürk* (Akdeniz University Herbarium!).

Acknowledgement

We wish to thank the subject editor and reviewers for their comments which helped us to improve the manuscript, TÜBİTAK (Project Number TBAG-2074) and Expo Antalya 2016 natural plant supply project in Taurus Mountains for financial support and the Curators of the Herbaria Ankara University (ANK), Gazi University (GAZI) and Hacettepe University (HUB), who allowed us to study their *Cephalaria* specimens. This paper was produced from project (number: 21.01.0121.24) supported by Scientific Research Projects Unit of Akdeniz University.

References

- Aksoy N, Göktürk RS, Açık L, Çelebi A (2007) *Cephalaria duzceënsis* (Dipsacaceae), a new species from the western Black Sea Region, Turkey. *Nordic Journal of Botany* 25(1): 64–69. doi: 10.1111/j.0107-055X.2007.00100_11.x
- Bobrov EG (1957) Dipsacaceae. In: Shishkin BK (Ed.) *Flora of the USSR*, Vol. 24. Akademiya Nauk SSSR Press, Moscow, 20–37.
- EUNIS (2007) EUNIS Habitat Classification. http://www.eea.europa.eu/themes/biodiversity/eunis/ds_resolveuid/9a179b2f8dd74f7d91bcf4f3a284e55e [accessed 05 June 2016]
- Feinbrun-Dothan N (1978) *Cephalaria* Schrad. ex Roem. & Schult. In: Feinbrun-Dothan N (Ed.) *Flora Palaestina*, Vol. 3. Israel Academy of Sciences Press, Jerusalem, 263–266.
- Ferguson IK (1976) *Cephalaria* Schrad. ex Roem. & Schult. In: Tutin GT, Heywood VH, Burges NA, Moore DM, Valentine DH, Walters SM, Webb DA (Eds) *Flora Europaea*, Vol. 4. Cambridge University Press, Cambridge, 57–58.
- Göktürk RS, Sümbül H (1997) A new species of *Cephalaria* (Dipsacaceae) from South Anatolia, Turkey. *Annales Botanici Fennici* 34(3): 153–155. <http://www.sekj.org/AnnBot.html>
- Göktürk RS, Sümbül H (2003) *Cephalaria aytachii* (Dipsacaceae), a new species from Central Anatolia, Turkey. *Annales Botanici Fennici* 40(2): 123–127. <http://www.sekj.org/AnnBot.html>
- Göktürk RS, Sümbül H, Açık L (2003) A new species of *Cephalaria* Schrad. ex Roem. & Schult. (Dipsacaceae), including a new variety from East Anatolia, Turkey. *Israel Journal of Plant Sciences* 51(1): 59–65. doi: 10.1560/332B-JGEF-R6RC-DHL5
- Göktürk RS, Sümbül H, Çelebi A, Açık L (2012) Two new species of *Cephalaria* (Caprifoliaceae) from Turkey. *Turkish Journal of Botany* 36(4): 311–321. doi: 10.3906/bot-1101-4
- Göktürk RS, Sümbül H (2014) A taxonomic revision of the genus *Cephalaria* (Caprifoliaceae) in Turkey. *Turkish Journal of Botany* 38(5): 927–968. doi: 10.3906/bot-1310-6
- Güner A, Aslan S, Ekim T, Vural M, Babuş MT (2012) *Türkiye Bitkileri Listesi (Damarlı Bitkiler)*. Nezahat Gökyiğit Botanik Bahçesi ve Flora Araştırmaları Derneği Yayını, 1290 pp.
- IUCN (2014) *Guidelines for Using the IUCN Red List Categories and Criteria*. Version 11. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf> [accessed 2 June 2016]
- Kuş S, Göktürk RS (2005) A new *Cephalaria* species (Dipsacaceae) from the European part of Turkey. *Nordic Journal of Botany* 23(4): 427–430. doi: 10.1111/j.1756-1051.2003.tb00415x
- Lack HW (1991) *Cephalaria* Schrad. ex Roem. & Schult. In: Rechinger KH (Ed.) *Flora Iranica*, Dipsacaceae, Number 168. Akademische Druck-u Verlagsanstalt, Graz, 8–18.
- Matthews VA (1972) *Cephalaria* Schrad. ex Roem. & Schult. In: Davis PH (Ed.) *Flora of Turkey and the east Aegean islands*, Vol. 4. Edinburgh University Press, Edinburgh, 585–597.
- Matthews VA (1988) *Cephalaria* Schrad. ex Roem. & Schult. In: Davis PH, Mill RR, Tan K (Eds) *Flora of Turkey and the East Aegean islands*, Vol. 10, suppl. Edinburgh University Press, Edinburgh, 156 pp.
- Parolly G, Eren Ö (2007) Contributions to the Flora of Turkey, 2. *Willdenowia* 37(1): 243–271. doi: 10.3372/wi.37.37114
- Reveal JL, Chase MW (2011) APG III: Bibliographical information and synonymy of Magnoliidae. *Phytotaxa* 19: 71–134. <http://mapress.com/phytotaxa/backissues.htm>

- Roemer JJ, Schultes JA (1818) *Cephalaria* Schrad. ex Roem. & Schult. In: Linne C (Ed.) *Systema vegetabilium* 3. Sumtibus JG Cottae, Stuttgartiae, 43–54.
- Sümbül H (1991) Ten new species from Anatolia and two new records for the flora of Turkey. *Edinburgh Journal of Botany* 48: 27–40.
- Szabó Z (1940) *Cephalaria*-génusz monográfiája. Kiadja A Magyar Tudományos Akadémia Press, 352 pp.