

# *Lysimachia coriacea* (Primulaceae, Myrsinoideae), a new species from Chongqing, China

Hai-Fei Yan<sup>1,2</sup>, Ya Huang<sup>3</sup>, Hong-Jing Zhang<sup>3</sup>, Si-Rong Yi<sup>3</sup>

**1** Key Laboratory of Plant Resources Conservation and Sustainable Utilization, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China **2** South China National Botanical Garden, Guangzhou 510650, China **3** Chongqing Three Gorges Medical College, Wanzhou, Chongqing 404120, China

Corresponding author: Si-Rong Yi ([yisirong123@aliyun.com](mailto:yisirong123@aliyun.com))

Academic editor: W.A. Mustaqim | Received 10 August 2022 | Accepted 27 November 2022 | Published 16 December 2022

**Citation:** Yan H-F, Huang Y, Zhang H-J, Yi S-R (2022) *Lysimachia coriacea* (Primulaceae, Myrsinoideae), a new species from Chongqing, China. PhytoKeys 215: 87–94. <https://doi.org/10.3897/phytokeys.215.91488>

## Abstract

A new species, *Lysimachia coriacea*, from Chongqing, China, is described and illustrated. It is assigned to subgen. *Lysimachia* sect. *Nummularia* ser. *Paridiformes* and resembles *L. paridiformis* var. *stenophylla*, but is characterised by smaller leathery leaves with black glandular striations near the margin. It is also similar to *L. nanpingensis* in its two to three pairs of leaves sub-whorled at the stem apices, but differs by smaller leathery leaves and densely glandular stem, petiole and pedicel, and calyx lobes with sparse black glandular stripes.

## Keywords

field expedition, *Lysimachia*, morphology, Nanchuan County, taxonomy

## Introduction

*Lysimachia* L. is the largest genus in the tribe Lysimachieae (Primulaceae), which consists of approximately 200 species worldwide (Chen and Hu 1979; Chen and Hu 1989; Hu and Kelso 1996; Marr and Bohm 1997; Hao et al. 2004; Julius et al. 2016). Traditionally, the classification of Lysimachieae includes six genera (*Anagallis* L., *Asterolinon* Hoffmanns. & Link, *Glaux* Tourn. ex L., *Lysimachia*, *Pelletiera* A.St.-Hil., and *Trientalis* Ruppius ex L.) based on several characters, such as the dehiscing pattern of the capsule and the number of corolla lobes. Molecular evidence even suggested expanding the delimitation of *Lysimachia* by including all satellite genera in the tribe (e.g. Anderberg et

al. 2007). China has been considered as the diversity centre of this genus with approximately 130 species recorded in Flora of China (Hu and Kelso 1996). Subsequently, at least 20 new species have been discovered in the last two decades (Peng and Hu 1999; Shao et al. 2004; Shao et al. 2006; Yan and Hao 2012; Liu et al. 2014a, b; Estes et al. 2015; Zhou et al. 2015; Baskose et al. 2016; Julius et al. 2016; Wang et al. 2016; Yan et al. 2017; Wang et al. 2018; Huang et al. 2019; Liu et al. 2020; Mou et al. 2020; Yi 2020; Ju et al. 2021; Lu et al. 2021). In recent years, multiple field expeditions have been conducted in the Municipality of Chongqing, China. During several fieldworks, a species new to science of *Lysimachia* was discovered and is described below.

## Materials and methods

Multiple field investigations were conducted between April 2019 and July 2022 to collect the specimens of the putative new species. The morphological descriptions are based on both living and dried materials, which are deposited in the Herbaria of South China Botanical Garden, Chinese Academy of Sciences (IBSC) and Kunming Institute of Botany, Chinese Academy of Sciences (KUN). This study was based on an examination of herbarium specimens at IBSC, PE, KUN and digital images from the Chinese Virtual Herbarium (CVH: <https://www.cvh.ac.cn/index.php>), Global Plants JSTOR (<http://plants.jstor.org/>) and Plants of the World Online (POWO: <http://www.plantsoftheworldonline.org/>). In addition, the descriptions of its most similar species (*L. paridiformis* var. *stenophylla* Franch. and *L. nanpingensis* F.H. Chen & C.M. Hu) from the relevant taxonomic literature (Chen and Hu 1979; Chen and Hu 1989; Hu and Kelso 1996) were also consulted. The conservation assessment of the putative new species was undertaken using IUCN categories of threat (IUCN Standards and Petitions Committee 2022).

## Taxonomic treatment

***Lysimachia coriacea* S.R. Yi & H.F. Yan, sp. nov.**

urn:lsid:ipni.org:names:77310050-1

Figs 1, 2

**Type.** CHINA. Chongqing: Nanchuan District, Nanping Town, Shangengqiao, 29°02'N, 107°08'E, 740 m a.s.l., 28 April 2019, *Si-Rong Yi* YSR8174 (holotype IBSC!; isotypes KUN!).

**Diagnosis.** The new species belongs to subgen. *Lysimachia* sect. *Nummularia* ser. *Paridiformes* Chen & Hu (1979: 36) characterized by verticillate upper leaves, scale-like lower leaves and umbellate inflorescence. It is similar to *L. paridiformis* var. *stenophylla* in having scale-like lower leaves, upper leaves in a terminal whorl and terminal umbels, but it is easily distinguished by its smaller and leathery leaves (vs. papery) only with black

glandular stripes near the margin (vs. black glandular stripes on the whole leaf blade), pedicels with dense stalked glands (vs. glabrous), and corollas without black glandular stripes (vs. occasionally with black glandular stripes) (Hu and Kelso 1996). At first glance, it also looks similar to *L. nanpingensis*, but differs by its smaller leathery leaves (vs. papery) and densely glandular stem, petiole and pedicel (vs. densely fulvous pubescent), and glabrous calyx lobes with sparse black glandular stripes (vs. pubescent and sparsely reddish glandular punctate). A more detailed comparison of the three species is provided in Table 1.

**Description.** Herbs perennial, 5–15 cm tall. **Rootstock** with numerous fibrous roots. **Stem** terete, erect or lower part procumbent, rooting at nodes, simple or branched, with dense stalked glands when young. **Leaves** opposite, lowest 1–2 pairs scale-like, upper 2–3 pairs closely crowded; blades elliptic or ovate-elliptic, 1.8–3.5 × 1.0–1.8 cm, leathery, adaxially dark green, smooth and glabrous, lustrous; abaxially light green, glabrous, black glandular striate near entire margin, base cuneate, apex acute, mid-vein impressed above, raised below, lateral veins and veinlets inconspicuous; petiole 5–15 mm long, with dense stalked glands. **Flowers** 1–5, crowded at apex of stems, bracts oblanceolate, 6 mm long, green, sparsely glandular; pedicel 6–15 mm long, with dense stalked glands, erect in fruit; calyx lobes narrowly lanceolate, 4.5–6.8 × 0.9–1.5 mm, divided to base, apex acuminate, abaxially sparsely black glandular striate, margin glandular, membranous; corolla yellow, tube 2–3 mm high, lobes elliptic, 6–9 × 5.5–6.8 mm, apex obtuse or slightly emarginate; filaments connate basally into a tube, ca. 2 mm high, with dense granular glandular spots, free parts 2–3 mm long; anthers oblong, 0.8–1.2 mm long, dorsifixed; ovary subglobose, 0.9–1.2 mm in diam., glabrous; style 4.5–6.0 mm long, glabrous. **Capsule** subglobose, 2.5–4.5 mm in diam.

**Distribution and habitat.** *Lysimachia coriacea* is only known from the western slopes of Jinfo Mountain, Nanchuan, Chongqing, China. It grows on damp rocks or cliffs in forests, along roadsides and in mountain valleys at elevations of 740–1,300 m.


**Conservation status.** This new species has only been found on Jinfo Mountain with at least four populations, where its area of occupancy is less than 10 km<sup>2</sup>. Fortunately, the current distribution area of the species is under the protection of the Jinfo Mountain National Natural Reserve. Thus, it is assigned the status of “Least Concern” (LC) according to the IUCN Red List Categories and Criteria (IUCN Standards and Petitions Committee 2022).

**Phenology.** Flowering from April to May and fruiting from May to June.

**Etymology.** Latin *coriacea*, leathery, alluding to texture of leaves.


**Chinese name.** 革叶过路黄 (Gé Yè Guò Lù Huáng).

**Additional specimens examined.** CHINA. **Chongqing:** Nanchuan District, Nanping Town, Shangenqiao, 29°02'N, 107°08'E, 740 m a.s.l., in shady fissures of wet rocks, 7 June 2019, *Si-Rong Yi* YSR8004 (KUN1510469!); Nanchuan District, Nanping Town, Huangniya, on a damp rocky roadside, 1,300 m a.s.l., 28 May 1986, *Z. Y. Liu* 0663 (PE01895529!); Nanchuan District, Dutouma (from Lanba to Sanhui), on the roadside, 860 m a.s.l., 7 May 1957, *J.H. Xiong & Z.L. Zhou* 90705 (IBSC0019656!); Nanchuan District, Sanhui Dianchanggou, on a damp rock in forests, 970 m a.s.l., 5 July 1957, *J.H. Xiong & Z.L. Zhou* 91820 (IBSC0019657!).



**Figure 1.** *Lysimachia coriacea* **A** habit **B** leaf **C** flower **D** calyx lobes, abaxial view (right) and adaxial view (left) **E, F** opened corolla, showing filaments connate at base **G** anthers, front view (left) and back view (right) **H** pistil **I** flower with corolla removed **J** young fruit with persistent calyx. Drawn by Yun-Xiao Liu.





**Figure 2.** *Lysimachia coriacea* **A** flowering plants **B** inflorescence **C** fruiting plants **D** flowers **E, F** leaves **G** flower with corolla removed **H** calyx, abaxial view **I** calyx, adaxial view **J** corolla, abaxial view **K** corolla, adaxial view **L–M** stamens **N** pistil **O** young fruits. Photographed by Si-Rong Yi.

**Table 1.** Comparison of diagnostic characters of *Lysimachia coriacea*, *L. paridiformis* var. *stenophylla* and *L. nanpingensis*.

	<i>L. coriacea</i>	<i>L. paridiformis</i> var. <i>stenophylla</i>	<i>L. nanpingensis</i>
Stem	with dense stalked glands when young	glabrous	with dense fulvous multicellular hairs
Leaf blade	elliptic or ovate-elliptic, 1.8–3.5 × 1.0–1.8 cm, leathery; base cuneate, apex acute; with black glandular stripes near the margin; veins inconspicuous, except mid-vein	narrowly elliptic to broadly lanceolate or linear-lanceolate, 4–16 × 1.2–5 cm, papery; base cuneate, apex short acuminate; with or without black glandular stripes; veins 4 or 5 pairs, conspicuous	elliptic to ovate-elliptic, 3.5–5.5 × 2–4.5 cm, papery; base subrounded, apex acute; indistinctly glandular punctate; veins inconspicuous, except mid-vein
Petiole	5–15 mm long; with dense stalked glands	sessile or subsessile; glabrous	3–12 mm long; densely fulvous pubescent
Calyx lobes	4.5–6.8 × 0.9–1.5 mm; glabrous on both sides, sparsely ciliate on the margin, with sparse black glandular stripes	8–13 × 2.5–3.5 mm; glabrous on both sides, occasionally ciliate on the margin, occasionally with black glandular stripes	6–7.5 × 1.2–1.9 mm; pubescent on the abaxial surface, obscurely glandular punctate
Pedice	6–15 mm long, erect in fruit, with dense stalked glands	3–15 mm long; erect in fruit, glabrous	4–9 mm long, recurved in fruit, densely fulvous pubescent
Corolla lobes	6–9 × 5.5–6.8 mm, without glandular stripes	9–11 × 4–4.5 mm, with or without black glandular stripes	9–11 × 3.5–4 mm, sparsely reddish glandular punctate

**Notes.** Specimens of the new species were first collected by Ji-Hua Xiong and Zi-Lin Zhou in 1957 and deposited in IBSC (IBSC0019656 and IBSC0019657). The two specimens were identified as “*Lysimachia paridiformis* var. *stenophylla* Franch.” by Chi-Ming Hu. However, Hu should also notice the differences between this specific taxon and *L. paridiformis* var. *stenophylla*, because he wrote a temporary name “*Lysimachia nitida* Chen et C.M. Hu” on the annotation labels of the specimens. Later, specimens of the new species were collected by some other collectors (see above).

Acknowledgements

This research was supported by the Science & Technology Basic Resources Investigation Program of China (2017FY100100), the National Natural Foundation of China (31870192), and the Biological Resources Programme, Chinese Academy of Sciences (KFJ-BRP-017-104). We thank Ms. Yun-Xiao Liu for the line drawings of the new species.

References

Anderberg AA, Manns U, Källersjö M (2007) Phylogeny and floral evolution of the Lysimachieae (Ericales, Myrsinaceae): Evidence from *ndhF* sequence data. Willdenowia 37(2): 407–421. <https://doi.org/10.3372/wi.37.37202>

Baskose I, Keskin A, Gurbanov K (2016) *Lysimachia savranii* (Primulaceae), a new species from the eastern Taurus in Turkey. Phytotaxa 267(3): 228–232. <https://doi.org/10.11646/phytotaxa.267.3.6>

- Chen FH, Hu CM (1979) Taxonomic and phytogeographic studies on Chinese species of *Lysimachia*. *Acta Phytotaxonomica Sinica* 17: 21–53.
- Chen FH, Hu CM (1989) *Lysimachia*. In: Chen FH, Hu CM (Eds) *Flora Reipublicae Popularis Sinicae*. Vol 59. Science Press, Beijing, 3–133.
- Estes D, Shaw J, Mausert-Mooney C (2015) *Lysimachia lewisii* (Primulaceae): A new species from Tennessee and Alabama. *Phytoneuron* 17: 1–15.
- Hao G, Yuan YM, Hu CM, Ge XJ, Zhao NX (2004) Molecular phylogeny of *Lysimachia* (Myrsinaceae) based on chloroplast trnL–F and nuclear ribosomal ITS sequences. *Molecular Phylogenetics and Evolution* 31(1): 323–339. [https://doi.org/10.1016/S1055-7903\(03\)00286-0](https://doi.org/10.1016/S1055-7903(03)00286-0)
- Hu CM, Kelso S (1996) Primulaceae. In: Wu ZY, Raven PH (Eds) *Flora of China*. Vol 15. Science Press, Beijing & Missouri Botanical Garden Press, St Louis, 39–189.
- Huang YF, Dong LN, Xu WB (2019) *Lysimachia fanii*, a new species of Primulaceae from limestone area of Guangxi, China. *PhytoKeys* 130: 75–84. <https://doi.org/10.3897/phytokeys.130.34655>
- IUCN Standards and Petitions Committee (2022) Guidelines for using the IUCN Red List Categories and Criteria. Version 15. Prepared by the Standards and Petitions Committee. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Ju WB, Deng HN, Xu B, He XJ, Gao XF (2021) *Lysimachia xuyongensis* (Primulaceae), a new species from Sichuan, China. *Phytotaxa* 525(1): 59–64. <https://doi.org/10.11646/phytotaxa.525.1.7>
- Julius A, Tagane S, Naiki A, Gutiérrez-Ortega JS, Suddee S, Rueangruea S, Yahara T, Utteridge T (2016) *Lysimachia kraduensis* (Primulaceae), a new species from northeastern Thailand. *Phytotaxa* 289(1): 69–76. <https://doi.org/10.11646/phytotaxa.289.1.5>
- Liu K, Hong X, Zhou SB, Cheng YS, Tang CF, Xu HJ (2014a) A new species of *Lysimachia* (Myrsinaceae) from Dabieshan Mountain, China. *Plant Systematics and Evolution* 300(7): 1615–1620. <https://doi.org/10.1007/s00606-014-0986-z>
- Liu K, Zhou SB, Chen YS, Hong X (2014b) *Lysimachia dabieshanensis* sp. nov. (Primulaceae), a new species from Dabieshan Mountain, China. *Phytotaxa* 174(2): 119–122. <https://doi.org/10.11646/phytotaxa.174.2.8>
- Liu A, Yan HF, Yu XL, Hao G (2020) *Lysimachia caloneura* (Primulaceae), a new species from Hunan, China. *Annales Botanici Fennici* 58(1–3): 45–48. <https://doi.org/10.5735/085.058.0108>
- Lu ZC, Yuan Q, Wei SJ, Dong LN, Xu WB (2021) *Lysimachia lijiangensis* (Primulaceae), a new species from limestone area of central Guangxi, China. *Taiwania* 66(1): 65–72.
- Marr KL, Bohm BA (1997) A taxonomic revision of the endemic Hawaiian *Lysimachia* (Primulaceae) including three new species. *Pacific Science* 51: 254–287.
- Mou C, Wu Y, Xiang L, Xiang XM, Zhang DG (2020) *Lysimachia xiangxiensis* (Primulaceae), a new species from limestone area in Hunan Province, central China. *PhytoKeys* 140: 23–32. <https://doi.org/10.3897/phytokeys.140.47995>
- Peng CI, Hu CM (1999) *Lysimachia chingshuiensis* (Primulaceae), a new species from eastern Taiwan. *Botanical Bulletin of Academia Sinica* 40: 49–52.
- Shao JW, Zhang XP, Guo XH (2004) A new species of *Lysimachia* in Primulaceae. *Bulletin of Botanical Research* 24: 389–391.

- Shao JW, Zhang XP, Guo XH (2006) *Lysimachia dextrorsiflora* X.P. Zhang, X.H. Guo & J.W. Shao, a new species of Primulaceae from China. *Zhiwu Fenlei Xuebao* 44(5): 589–594. <https://doi.org/10.1360/aps050135>
- Wang ZH, Li XW, Liu ED, Shangguan FZ, Chang XK (2016) *Lysimachia septemfida* (Primulaceae), a new species from Yunnan, China. *Plant Diversity* 38(4): 201–206. <https://doi.org/10.1016/j.pld.2016.08.002>
- Wang Y, Ma W, Zhou SB, Liu K (2018) *Lysimachia tianmaensis* (Primulaceae), a new species from Anhui, China. *PhytoKeys* 98: 117–124. <https://doi.org/10.3897/phytokeys.98.23751>
- Yan HF, Hao G (2012) *Lysimachia huchimingii* sp. nov. (Primulaceae) from China. *Nordic Journal of Botany* 30(4): 443–445. <https://doi.org/10.1111/j.1756-1051.2012.01401.x>
- Yan HF, Xu Y, Zhu ZM, Hu CM, Hao G (2017) *Lysimachia sinopilosa* (Primulaceae), a new species from Yunnan, China. *Annales Botanici Fennici* 54(1–3): 45–48. <https://doi.org/10.5735/085.054.0308>
- Yi SR (2020) *Lysimachia porcatiseipala*, a new species of *Lysimachia* (Primulaceae) from Chongqing, China. *Phytotaxa* 434(1): 118–122. <https://doi.org/10.11646/phytotaxa.434.1.9>
- Zhou JJ, Yu XL, Deng YF, Yan HF, Lin ZL (2015) *Lysimachia huangsangensis* (Primulaceae), a new species from Hunan, China. *PLoS ONE* 10(7): e0132713. <https://doi.org/10.1371/journal.pone.0132713>