

Quercus mangdenensis, a new species of *Quercus* (Fagaceae) from Kon Tum Province, Vietnam

Nguyen Van Ngoc¹, Hoang Thi Binh¹

¹ Faculty of Biology, Dalat University, 01- Phu Dong Thien Vuong, Dalat, Vietnam

Corresponding author: Hoang Thi Binh (binhht@dlu.edu.vn)

Academic editor: Norbert Holstein | Received 19 August 2022 | Accepted 28 November 2022 | Published 15 December 2022

Citation: Ngoc NV, Binh HT (2022) *Quercus mangdenensis*, a new species of *Quercus* (Fagaceae) from Kon Tum Province, Vietnam. *PhytoKeys* 215: 73–79. <https://doi.org/10.3897/phytokeys.215.93684>

Abstract

Quercus mangdenensis Binh & Ngoc, **sp. nov.** (Fagaceae) is newly described from Mang Den Town in the central highland of Vietnam. The new species is characterized by lanceolate to oblong-lanceolate leaves with entire margin, 1–5-fruited infructescence, larger fruit size 6–10.5 cm long, broadly bowl-shaped cupules enclosing 1/5 of the nut, bracts of cupule entire and arranged in 5–7 rings, and cylindrical-ellipsoid and basally flat nuts 4.5–7.5 cm long. *Quercus mangdenensis* is morphologically similar to *Q. bidoupensis* Binh & Ngoc and *Q. kontumensis* A.Camus in having similar leaf shape, cuneate leaf base, and bracts arrangement in cupules. However, it differs from *Q. bidoupensis* and *Q. kontumensis* by cupules broadly bowl-shaped, much larger fruits, cylindrical-ellipsoid nut shape, and cupule enclosing 1/5 of the nuts. A description, photographs, and preliminary species conservation status of the new species are also provided.

Keywords

Fagales, flora, Kon Plong, Mang Den, taxonomy

Introduction

Quercus L. is the biggest genus of the family Fagaceae, comprising more than 500 species widely distributed in Europe, North America, the Mediterranean, temperate deciduous forests in East Asia, and tropical montane forests in Southeast Asia (Nixon 1993;

Huang et al. 1999; Phengkklai 2008; Hubert et al. 2014; Valencia-A et al. 2016). In Vietnam, a total of 52 *Quercus* species have been reported (Ho 2003; Ban 2005; Li et al. 2016; Binh et al. 2018a, b, c; Binh et al. 2021; Ngoc et al. 2022), of which, recent efforts in taxonomic works of the genus *Quercus* have resulted in the description of seven new species from Vietnam (Binh et al. 2018a, b, c; Binh et al. 2021; Ngoc et al. 2022), indicating that the species diversity of *Quercus* of the country is significant. Therefore, taxonomic studies of the Vietnamese *Quercus* are still required.

Mang Den town belongs to Kon Plong District, Kon Tum Province of Vietnam with a total area of 148.07 km² (Kon Tum Province 2022) (Fig. 1). In this area, the annual temperature range is 18.7–24.9 °C and the average annual rainfall is around 1780–2200 mm with the rainy season from August to February (Tri et al. 2022). In the region, 273 woody plants have been recorded (265 angiosperms and 8 gymnosperm species). As for Fagaceae, 20 species have been recorded from Kon Plong District (Thanh et al. 2001).

From April to July 2022, we conducted botanical surveys in Mang Den and collected several individuals of the genus *Quercus* that could not be placed with any previously described species. After careful morphological examination and taxonomic review, here we describe those individuals as *Quercus mangdenensis* Binh & Ngoc along with morphological comparisons with the most similar species, photographs, and a preliminary conservation assessment of the new species.

Materials and methods

Plant materials

In this study, six specimens of the new species (Binh et al. QC204, 205, 206, 207, 208, 209) were collected from Mang Den, Kon Plong District, Kon Tum Province (Fig. 1). In addition, we used type specimens and specimens that were collected from the type locality of the following species: *Q. bidoupensis* (Tagane et al. V4328 [DLU!], Binh et al. QC27, 29, 30, 45, 57 [DLU!]) and *Q. kontumensis* (Poilane 18381 [P00754004] and [P00754005]), which are morphologically most similar to the new species in having a lanceolate to oblong-lanceolate leaf shape, cuneate leaf base, and bracts of cupule fusing in 5–7 rings.

Morphological observations

To confirm that the species was undescribed, the relevant literature including Camus (1934), Soepadmo (1972), Huang et al. (1999), Ho (2003), Ban (2005), and Phengkklai (2008), Binh et al. 2018a, b, c; Binh et al. 2021; Ngoc et al. 2022 were consulted. Then, we examined specimens in the herbaria at DLU, K, P, and VNM as well as digital images on the websites of JSTOR Global Plants (<https://plants.jstor.org/>) and the Chinese Virtual Herbarium (<http://www.cvh.ac.cn/>).




Figure 1. Type locality of *Quercus mangdenensis* Binh & Ngoc **A** map of Vietnam **B** map of Kon Tum province **C** map of Kon Plong District, the red star indicated the type locality: Mang Den Town, Dak Long Commune.

Taxonomic treatment

Quercus mangdenensis H.T.Binh & Ngoc, sp. nov.

urn:lsid:ipni.org:names:77309987-1

Fig. 2

Diagnosis. *Quercus mangdenensis* is morphologically most similar to *Q. bidoupensis* and *Q. kontumensis* with a lanceolate to oblong-lanceolate leaf shape, cuneate leaf base, and bracts of cupule fusing in 5–7 rings. However, *Q. mangdenensis* is distinguished from *Q. bidoupensis* by its entire leaf margin (vs. undulate, distinctly serrate in the upper 1/2), longer petioles 2.5–4 cm long (vs. 1.3–2 cm long), broadly bowl-shaped cupules (vs. obconical-shaped), larger cupule (2.6–3.5 cm high × 4–5 cm in diam. vs. 1.3–1.5 cm high × 1.3–1.7 cm in diam.), enclosure of cupule (1/5 vs. 1/3 of the nut), cylindrical-ellipsoid nut (vs. ovoid), larger nut (4.5–7.5 cm high × 3–4 cm in diam. vs. 2.2 cm high × 1.4 cm in diam.), and flat basal scar of the nut (vs. convex). It differs from *Q. kontumensis* in having larger cupule enclosing 1/5 of the nut (2.6–3.5 cm high × 4–5 cm in diam. vs. 1–1.2 cm long × 1.5–2 cm in diam., enclosing 1/2 of the nut), larger cylindrical-ellipsoid nut 4.5–7.5 cm high × 3–4 cm in diam. (vs. cylin-

dricar-ovoid, 2 cm high × 1.5 cm in diam.), and scales of cupule fusing into 5–7 entire ridges (vs. undulate) (Table 1).

Type. VIETNAM. Kontum Province, Kon Plong District, Mang Den Town, alt. 1179 m, 14°36'39.1"N, 108°15'00.5"E, 23 July 2022, *H. T. Binh & N. V. Ngoc QC204* (holotype DLU!; isotypes HNI, VNM!)

Description. Tree, evergreen, 20–25 m tall, 60–80 cm girth. Bark whitish gray. Buds perulate, globose to oblate, 2.5–4 mm high, 2.5–3.5 mm in diam., bud scales imbricate, in 2–3 rows, ovate-triangular, 2.0–3.0 × 1.5–2.0 mm, apex obtuse, margin ciliate, covered with appressed white hairy outside, glabrous inside. Young twigs dark pale green, hairy, old twigs grayish-brown, glabrous, sometimes sulcate, lenticellate. Leaf alternate; blades thickly coriaceous, lanceolate to oblong-lanceolate, (5.5–)11–16.5(–19.5) × (2.6–)3.5–5.5 cm, acute at apex, broadly cuneate at base, margin entire, glossy green adaxially, bright green abaxially, conspicuously pale creamy brown to yellowish brown *in sicco*, glabrous on both surfaces, midrib slightly raised on the upper surface, prominently raised on the lower surface, lateral veins (5–)7–9 pairs, prominent on the lower surface, at an angle of 45–55 degree from the midrib, straight and then curved near margin, tertiary veins scalariform-reticulate, faintly visible on both surfaces; petioles (1.3–)2.5–4 cm long, yellowish brown when dry, glabrous. Male and female inflorescences not seen. Infructescences axillary or pseudo-terminal (sometimes in upper leaf-scars), erect, rachis 1–1.5 cm long, 1–1.3 cm in diam., woody, glabrous, dark yellow *in vivo*, yellowish brown *in sicco*, lenticellate. Mature fruits 6–10.5 cm high (including cupule), solitary, sessile on woody rachis; nuts cylindrical-ellipsoid, 4.5–7.5 cm high, 3–4 cm in diam., rounded at the top, white tomentose around stylopodia, stylopodia 1.5–2.5 mm long, basal scar 2–2.5 cm in diam., flat; cupules broadly bowl-shaped, 2.6–3.5 cm high, 4–5 cm in diam., enclosing 1/5 of the nut when mature, outside tomentose with reddish hairs to glabrous, inside with appressed densely reddish hairs, wall

Table 1. Morphological comparison amongst *Quercus mangdenensis* Binh & Ngoc, sp. nov., *Quercus bidoupensis* Binh & Ngoc and *Quercus kontumensis* A. Camus.

Characters	<i>Q. mangdenensis</i>	<i>Q. bidoupensis</i> ^(1,2)	<i>Q. kontumensis</i> ^(3,4)
Buds shape	gobose to oblate	oblong to ellipsoid	ovoid to ellipsoid
Leaf shape	lanceolate to oblong-lanceolate	oblong-lanceolate	lanceolate to oblong-lanceolate
Leaf margin	entire	margins undulate, distinctly serrate in the upper 1/2	entire or distinctly serrate in the upper 1/5
Length of petioles	(1.3–)2.5–4 cm long	1.3–2 cm long	2.5–4.2 cm long
Number of secondary veins	(5–)7–9 pairs	10–13 pairs	5–8 pairs
Cupule shape and size	broadly bowl-shaped, 2.6–3.5 cm high, 4–5 cm in diam.	obconical-shaped, 1.3–1.5 cm high, 1.3–1.7 cm in diam.	cup-shaped, 1–1.2 cm long, 1.5–2 cm in diam.
Number of rings on cupule	5–6 rings	5–6 rings	6–7 rings
Cupule bract margin	entire	entire	undulate
Cupule coverage	enclosing 1/5 of the nut	enclosing 1/3 of the nut	enclosing 1/2 of the nut
Nut shape and size	cylindrical-ellipsoid, 4.5–7.5 cm high, 3–4 cm in diam.	ovoid, 2.2 cm high, 1.4 cm in diam.	cylindrical-ovoid, 2 cm high, 1.5 cm in diam.
Base of the nut	Flat	Convex	Flat

⁽¹⁾ From the protologue (Binh et al. 2018b); ⁽²⁾ From *Binh et al. QC27* (DLU); ⁽³⁾ From Camus (1934); ⁽⁴⁾ From the material *E. Poilane 18381* (P [P00754005, P00754005]).

0.5–1.3 cm thick, woody, bracts fusing and arranged in 5–7 ridges (rings), the rings' margin completely entire (without scale-like structure).

Distribution. Vietnam. Kon Tum Province, Kon Plong District, Mang Den Town (Fig. 1).

Habitat. *Quercus mangdenensis* were found in the scattered evergreen forest, from 1050 to 1200 m elevation.

Etymology. The specific epithet is derived from its type locality, Mang Den Town, Kon Plong District, Kon Tum Province, Vietnam.

Vernacular name. Sồi Mãng Đen (suggested here).

Phenology. Fruiting specimens and fallen mature fruits were collected in July.

Additional specimens examined. VIETNAM. Kontum Province, Kon Plong District, Mang Den, alt. 1168 m, 14°36'37.4"N, 108°15'11.6"E, 23 July 2022, *H. T. Binh & N. V. Ngoc QC205* [fr.] (DLU!, HN!, VNM!); *ibid.*, 1180 m elev., 14°36'32.5"N, 108°15'18.2"E, 23 July 2022, *H. T. Binh & N. V. Ngoc QC207* [fr.] (DLU!, HN!, VNM!); *ibid.*, 1130 m elev., 14°36'36.5"N, 108°15'31.3"E, 23 July 2022, *H. T. Binh & N. V. Ngoc QC208* [fr.] (DLU!, HN!, VNM!); *ibid.*, 1065 m elev., 14°36'36.5"N, 108°15'31.3"E, 23 July 2022, *H. T. Binh & N. V. Ngoc QC209* [fr.] (DLU!, HN!, VNM!).

Preliminary conservation status. Our botanical inventories were conducted at Mang Den and the surrounding areas from April to July 2022 and we found five sub-populations of 5–7 mature individuals of *Quercus mangdenensis* in the scattered ever-



Figure 2. *Quercus mangdenensis* Binh & Ngoc: **A** twigs with young fruit **B** terminal buds **C** infructescences **D** fallen mature fruit **E**, **F** adaxial and abaxial surface of the leaves **G** nuts **H** outside of cupule **I** densely reddish hairs inside of cupules **J** basal scar of the nut **K**, **L** inside and outside of bud scale. Materials from *H. T. Binh and N. V. Ngoc QC204*.

green forests. In addition, the forest in this area is severely fragmented and continuing to decline caused by human activities, such as farming, logging, and harvesting of non-timber forest products, etc., while almost all of the individuals of *Q. mangdenensis* are distributed along the boundary between the evergreen forest and local people's farms. According to criterion D of the IUCN Red List criteria (IUCN 2019), the species classifies as Critically Endangered (CR). The area of occupancy (AOO) and the extent of occurrence (EOO) were calculated are 0.415 km² and 8.0 km², respectively. Based on criterion B of the IUCN Red List criteria (IUCN 2019), the new species is qualified as Critically Endangered [CR B1ab(i,ii,iii,iv,v) B2ab(i,ii,iii,iv,v)].

Acknowledgements

The authors would like to thank Mr. Nguyen Van Duc and Mrs. My from Kon Tum city who kindly supported us during our fieldwork in Mang Den Town. We wish to thank the curators and staff of the following herbaria DLU, K, P, and VNM for making their materials accessible. This research is funded by the Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 106.03-2019.19.

References

- Ban NT (2005) Vietnam plant checklist, vol. 2, Agriculture Publishers, Hanoi National University. [In Vietnamese]
- Binh HT, Ngoc NV, Bon TN, Tagane S, Yahara T (2018a) A new species and two new records of *Quercus* (Fagaceae) from northern Vietnam. *PhytoKeys* 92: 1–15. <https://doi.org/10.3897/phytokeys.92.21831>
- Binh HT, Ngoc NV, Tagane S, Toyama H, Mase K, Mitsuyuki C, Strijk JS, Suyama Y, Yahara T (2018b) A taxonomic study of *Quercus langbianensis* complex based on morphology, and DNA barcodes of classic and next generation sequences. *PhytoKeys* 95: 37–70. <https://doi.org/10.3897/phytokeys.95.21126>
- Binh HT, Ngoc NV, Tai VA, Son HT, Tagane S, Yahara T (2018c) *Quercus trungkhanhensis* (Fagaceae), a new species from Cao Vit Gibbon Conservation Area, Cao Bang Province, north-eastern Vietnam. *Acta Phytotaxonomica et Geobotanica* 69(1): 53–61. <https://doi.org/10.18942/apg.201713>
- Binh HT, Ngoc NV, Son HT, Tagane S, Yahara T (2021) *Quercus ngochoaensis* (Fagaceae), a new species from Ba Vi National Park, northern Vietnam. *Phytotaxa* 516(3): 283–288. <https://doi.org/10.11646/phytotaxa.516.3.7>
- Camus A (1934) Les Chênes. Monographie du Genre *Quercus* Tome 1. Paul Lechevalier. Paris, 190–293.
- Ho PH (2003) An Illustrated Flora of Vietnam, vol. 2. Young Publishers, Ho Chi Minh City. [In Vietnamese]

- Huang CC, Chang YT, Bartholomew B (1999) Fagaceae. In: Wu ZY, Raven PH (Eds) Flora of China 4. Science Press, Beijing & Missouri Botanical Garden Press, Saint Louis, 314–400.
- Hubert F, Grimm GW, Joussetin E, Berry V, Franc A, Kremer A (2014) Multiple nuclear genes stabilize the phylogenetic backbone of the genus *Quercus*. *Systematics and Biodiversity* 12(4): 405–423. <https://doi.org/10.1080/14772000.2014.941037>
- IUCN (2019) Guidelines for using the IUCN red list categories and criteria, version 14. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf> [Accessed 5 August 2022]
- Kon Tum Province (2022) Established Mang Den Town in Kon Plong District. <https://www.kontum.gov.vn/pages/detail/33247/Thanh-lap-thi-tran-Mang-Den-thuoc-huyen-Kon-Plong.html> [Accessed 5 August 2022]
- Li Q, Zhang J, Coombes A (2016) *Quercus lineata* (Fagaceae): New distribution records from China and Vietnam and its leaf anatomical features. *Phytotaxa* 266(3): 226–230. <https://doi.org/10.11646/phytotaxa.266.3.7>
- Ngoc NV, Binh HT, Son HT, Suyama Y, Yahara T (2022) A new species of *Quercus* genus (Fagaceae) from Son Tra Peninsula, Central Vietnam. *PhytoKeys* 206: 61–73. <https://doi.org/10.3897/phytokeys.206.85635>
- Nixon KC (1993) Infrageneric classification of *Quercus* (Fagaceae) and typification of sectional names. *Annales des Sciences Forestieres* 50(Supplement): 25s–34s. <https://doi.org/10.1051/forest:19930701>
- Phengkhai C (2008) Fagaceae. In: Santisuk T, Larsen K (Eds) Flora of Thailand 9(3). The Forest Herbarium, Bangkok, 179–410.
- Soepadmo E (1972) Fagaceae. *Flora Malesiana Series I, Volume 7(2)*. Noordhoff-Kolff N. V., Djakarta, 339 pp.
- Thanh VV, Nhuan TD, Hoi PG (2001) The results of the research and survey on forest plant resources of Konplong. <http://fipiqn.vn/9-tin-moi/100-ket-qua-dieu-tra-he-thuc-vat-va-tham-thuc-vat-rung-huyen-konplong-tinh-kon-tum> [Accessed 5 August 2022]
- Tri CL, Tuan TA, Kiem TN (2022) Application of AHP-GIS Model to assess the ecological suitability of *Codonopsis javanicain* Kon Plong District, Kon Tum Province, Viet Nam. *Journal of Ecological Engineering* 23(7): 276–283. <https://doi.org/10.12911/22998993/150027>
- Valencia-A S, Rosales JLS, Arellano OJS (2016) A new species of *Quercus*, section Lobatae (Fagaceae) from the Sierra Madre Oriental, Mexico. *Phytotaxa* 269(2): 120–126. <https://doi.org/10.11646/phytotaxa.269.2.5>