

# *Diplycosia platyphylla* (Ericaceae), a new species from Mindanao, Philippines

Peter W. Fritsch<sup>1</sup>, Victor B. Amoroso<sup>2</sup>

**1** Botanical Research Institute of Texas, 1700 University Drive, Fort Worth, Texas 76107, USA **2** Center for Biodiversity Research and Extension in Mindanao, Central Mindanao University, University Town, Musuan 8710, Bukidnon, Philippines

Corresponding author: Peter W. Fritsch (pfritsch@brit.org)

---

Academic editor: Peter de Lange | Received 7 June 2016 | Accepted 11 July 2016 | Published 18 August 2016

---

**Citation:** Fritsch PW, Amoroso VB (2016) *Diplycosia platyphylla* (Ericaceae), a new species from Mindanao, Philippines. PhytoKeys 69: 31–38. doi: 10.3897/phytokeys.69.9466

---

## Abstract

*Diplycosia platyphylla* P.W.Fritsch, a new species from Mindanao Island, Philippines, is described and illustrated. This species is most similar to the Bornean *D. urceolata* but differs by its green or slightly flushed pink petioles 4–7 mm long, wider leaf blades, acute calyx lobe apices, and lavender mature fruiting calyx. The new species is known only from a single collection made from Mount Apo in North Cotabato Province, southern Mindanao.

## Keywords

*Diplycosia*, Ericaceae, Gaultherieae, Mindanao, Mount Apo, new species, Philippines

## Introduction

*Diplycosia* Blume (Ericaceae: Vaccinioideae: Gaultherieae) comprises about 116 species distributed throughout Southeast Asia and New Guinea (Sleumer 1967; Argent 1982, 1989, 2002; Ferreras and Argent 2011; Argent 2013, 2014; Fritsch and Amoroso in press; Fritsch and Bush in press). The genus can be delimited from other genera of the tribe by a base chromosome number of  $x = 18$  (vs 11), and a combination of (usually) entire leaf margins, fasciculate inflorescences, paired bracteoles borne at the apex of the pedicel, anthers with terminal tubules but with neither spurs nor disintegration tissue, and a capsule surrounded by an accrescent, fleshy calyx, or rarely a berry. Molecular studies strongly support the monophyly of the genus, and place it as phylogenetically nested within *Gaultheria* Kalm ex L. in the Wintergreen Group clade (Powell and Kron 2001; Bush et al. 2009; Fritsch et al. 2011).

Nine species of *Diplycosia* are currently recognized in the Philippines, three having been described since the taxonomic treatment of Ericaceae for the *Flora Malesiana* (Sleumer 1967; Ferreras and Argent 2011; Argent 2013; Fritsch and Amoroso in press). Because species of *Diplycosia* are often restricted to a single mountain or very few locations, more Philippine species of this genus likely await discovery (Argent 2013). Five species of *Diplycosia* have been documented from the island of Mindanao (Pelser et al. 2011 onwards; Fritsch and Amoroso in press), the southernmost major island of the Philippines. In April–May 2014 a joint botanical expedition was undertaken by the California Academy of Sciences and the CEBREM Office of Central Mindanao University to mountain peaks in the central part of Mindanao. Several individuals of *Diplycosia* were observed on Mt. Apo, the highest peak in the Philippines at 2954 m a.s.l. These plants appeared not to match any other Philippine species of the genus, according to the key in Argent (2013). Further study of a collection and photographic images made from these plants confirmed that these individuals represent a species new to science, which is here described and illustrated.

## Methods

Plants here described as a new species were dried as herbarium specimen vouchers. Macromorphological characters derived from both the specimens and photographic images of the living material were compared with those from the relevant literature sources (Sleumer 1967; Argent 1982, 1989, 2002, 2013, 2014; Ferreras and Argent 2011), specimens that are available at BRIT, CAS, and E, and online type images accessed from JSTOR Global Plants (<https://plants.jstor.org/>). We focused in particular on the characters used for the most recent key to Philippine *Diplycosia* in Argent (2013). Acronyms of herbaria follow Thiers (2016). Conservation status was assessed in accordance with IUCN Standards and Petitions Subcommittee (2016) criteria. The mature fruit was lost in transit between collection and deposition, and thus was not preserved; its description is based on a poor-quality photograph where only the general features were documented.

## Taxonomy

### *Diplycosia platyphylla* P.W.Fritsch, sp. nov.


urn:lsid:ipni.org:names:60472845-2

Figs 1, 2

**Diagnosis.** Haec species *Diplycosiae urceolatae* simillima, sed ab eo petiolo viridi vel viridi ex roseo 4–7 mm longo, lamina 7–7.5 cm lata, lobis calycis acutis, calyce fructus maturo lavandulo differt.

**Type.** PHILIPPINES. Mindanao Island, North Cotabato Province, Barangay Ilomavis, Mount Apo Natural Park, Matingao River Watershed, road from Site C to Site H at Energy Development Corporation (EDC), 1718 m, 6°59.88498'N, 125°14.86668'E, 29 Apr 2014, *D.S. Penneys 2302* (holotype: PNH!; isotypes: CAS-1196249!, CMUH-00010804!).

**Description.** Terrestrial erect shrublet to 0.6 m tall with ascending-erect branchlets. Young branchlets not observed; old branchlets gray, stout (3.5–4.5 mm wide), subterete, with non-peeling epidermis, glabrous. Leaves spirally arranged, with distinct wintergreen odor when crushed (fresh leaves), ascending; petiole green or slightly flushed pink, 4–7 × 2.5–2.9 mm, grooved above, glabrous; lamina broadly obovate to subrotund, the larger 8.3–9.7 × 7–7.5 cm, 1.2–1.7 times as long as wide, coriaceous, abaxially glossy and evenly dark red-dotted over the surface, adaxially shiny and glabrous; major veins prominent abaxially, sulcate adaxially, major secondary veins originating on proximal half, 1 to 3 on each side of midvein, arching-ascending, alternate, occasionally 3 on one side and only 2 on the other; base broadly cuneate, margin entire, revolute, glabrous, apex rounded or strongly emarginate, the very tip with a thick protruding dark gland to 0.8 mm long. Inflorescences axillary or on older sections of defoliated branchlets proximal to the leaves, fasciculate, 3- to 6-flowered, often with up to 7 additional old fruiting pedicels; bracts ovate-deltoid, 1.5–2 × 1.2–1.7 mm, appressed-puberulent. Pedicels slightly dilated distally, 4–7 × 0.5–0.8 mm, muriculate with short (to 0.16 mm long) ferruginous erect or ascending trichomes, and also pubescent with pale ferruginous irregularly oriented straight to slightly undulate nonglandular trichomes to 0.3 mm long; bracteoles ovate-deltoid, 1–1.5 mm long, muriculate, rarely also sparsely puberulent along midvein, midvein planar to prominent, margin ciliolate with a mix of ferruginous glandular trichomes and pale ferruginous nonglandular trichomes. Calyx green strongly flushed deep pink, 2.2–2.7 × 3.4–3.7 mm, sparsely muriculate or glabrous; limb ca. 2 mm long; lobes broadly deltoid-ovate, 1.1–1.2 × 1.8–2 mm, margins with a mixture of ferruginous glandular and pale ferruginous nonglandular trichomes, the latter more prevalent distally, apex acute. Corolla 5-lobed, white slightly flushed with dull pink distally, broadly urceolate, widest at middle, ca. 4.5 × 2.9 mm, glabrous both outside and within; lobes recurved, ovate, ca. 1 × 1 mm, margins eciliolate, apices obtuse. Stamens 10, included within corolla, ca. 4.5 mm long; filaments ± S-curved, ca. 3 mm long, glabrous; anthers 1.7–1.9 mm long, thecae 1.4–1.5 mm long, echinulate, tubes parallel, 0.3–0.4 mm long, smooth, pores strongly




**Figure 1.** *Diplycosia platyphylla* P.W.Fritsch. **A** Habitat **B** Habit **C** Branchlet with inflorescences **D** Inflorescence **E** Immature fruit. (D.S. Penneys 2302). Scale bars: **C** = 10 cm, **D**, **E** = 1 cm. Photos P.W.F.

oblique. Nectary 10-lobed; lobes emarginate. Ovary cylindrical-hemispherical, ca. 1.5 × 1.5 mm. Style 2.5–3 mm long, glabrous. Immature fruit green flushed red, 5–6 × 5–6 mm, style persistent; mature fruiting calyx lavender, accrescent, fleshy.

**Etymology.** The species is named for its notably wide leaves relative to those of most other species in the genus.

**Phenology.** Flowering and fruiting in April.

**Distribution and habitat.** *Diplycosia platyphylla* is known only from a single location on Mt. Apo, Mindanao Island, North Cotabato Province, Philippines. Several



**Figure 2.** *Diplycosia platyphylla* P.W.Fritsch. **A** Fertile branchlet **B** Pedicel and flower **C** Calyx lobe apex **D** Flower with calyx and corolla partly cut away to show androecium and gynoecium **E** Stamen, ventral view **F** Stamen, lateral view **G** Immature fruit **H** Fruiting pedicel. Drawing by L. Heagy from D.S. Penneys 2302 (CAS) and photographic images of the living plant.

individuals were found growing in gravel on an ultramafic open sunny rockslide area in the Tropical Lower Montane Rain Forest biome at 1718 m a.s.l. with *Nepenthes* L., *Huperzia* Bernh., and *Vaccinium* L. on a general NW-facing 30–60% slope but in a flat microhabitat along a roadside.

**Conservation status.** *Diplycosia platyphylla* is known from a single population and single collection, with only several plants seen. Although the species is afforded pro-

tection by its occurrence in Mount Apo National Park, it is at risk through apparent extreme rarity. We categorize this species as Critically Endangered (CR): D.

**Discussion.** The new species is similar to *Diplycosia urceolata* Stapf from Mount Kinabalu in Borneo, Sabah, Malaysia by its stout glabrous branchlets, coriaceous glabrous leaf blades with rounded apices, and 4–7 mm-long pedicels with a mixture of muriculate trichomes and pale ferruginous nonglandular trichomes. It differs from this species most readily by its green or slightly flushed pink petioles 4–7 mm long (vs vivid red and 10–13 mm), wider leaf blades [7–7.5 cm vs (2.5–)3.5–5(–7.5) cm], acute calyx lobe apices (vs obtuse), and lavender mature fruiting calyx (vs black). It is also similar to *D. sanguinolenta* Sleumer from Mount Kinabalu in its large coriaceous leaf blades [9–14(–17) × 5–8(–9.5) cm for *D. sanguinolenta*; Sleumer 1967] and generally acute calyx lobes, but is easily distinguished by a petiole 4–7 mm long (vs 12–15 mm) and a white corolla ca. 4.5 mm long (vs bright red and 13–15 mm long), among other characters. The new species would key to couplet 3 of the key to the Philippine *Diplycosia* in Argent (2013). The two leads at that point in the key identify the varieties of *D. luzonica* (A.Gray) Merrill, i.e., *D. l.* var. *caelalanensis* (Elmer) Sleumer and *D. l.* var. *merrittii* (Merrill) Sleumer, distinguished by “Leaves broadly pointed to rounded with the terminal gland forming a short mucronate point the largest leaves up to 5 cm long” vs “Leaves acuminate, with a narrowly acute apex, the largest leaves more than 6 cm long,” respectively. Because the new species possesses leaves that are rounded (or strongly emarginate) at the apex but are 8.3–9.7 cm long (the larger ones), the new species does not match either of these leads. Thus, it can be distinguished from all other Philippine *Diplycosia* by the combination of glabrous branchlets and (larger) leaf blades 8.3–9.7 mm long with an apex that is rounded or strongly emarginate, and with a prominent terminal gland.

## Acknowledgements

We thank the Protected Area Management Board of Mount Apo Natural Park and the Department of Environment and Natural Resources (DENR) of Region XII, Republic of the Philippines for the gratuitous permit, the Energy Development Corporation (EDC) for field access and logistical help, Central Mindanao University, the staff of BRIT, CAS, and E for access to their collections, Fulgent P. Coritico for field logistics, George Argent for insights into *Diplocosia* taxonomy, and Linda Heagy for the illustration. This work was supported in part by U.S. National Science Foundation grant DEB-1146409 (Darin S. Penneys, PI, P.W.F., co-PI) and the Philippines Commission on Higher Education.

## References

Argent GCG (1982) New taxa and combinations and comments on Ericaceae. *Botanical Journal of the Linnean Society* 85(1): 2–17.

- Argent GCG (1989) *Diplycosia* in Borneo. Two new species and a provisional key to *Diplycosia* and close allies. Notes from the Royal Botanic Garden, Edinburgh 46(1): 17–26.
- Argent GCG (2002) New taxa and new combinations in the genus *Diplycosia* (Ericaceae) of Borneo and Peninsular Malaysia. Gardens' Bulletin (Singapore) 54(2): 217–238.
- Argent G (2013) A new species of *Diplycosia*: *D. coii* Argent (Ericaceae) is described in honour of Leonardo Co. Philippine Journal of Science 142(special issue): 51–55.
- Argent G (2014) A contribution to the study of the genus *Diplycosia* (Ericaceae) in Sulawesi, Indonesia. Edinburgh Journal of Botany 71(1): 83–115. doi: 10.1017/S0960428613000309
- Bush CM, Lu L, Fritsch PW, Li D-Z, Kron KA (2009) Phylogeny of Gaultherieae (Ericaceae: Vaccinioideae) based on DNA sequence data from *matK*, *ndhF*, and nrITS. International Journal of Plant Sciences 170(3): 355–364. doi: 10.1086/596330
- Ferreras UF, Argent GCG (2011) *Diplycosia bartolomei* (Ericaceae), a new species from the Philippines. Edinburgh Journal of Botany 68(1): 39–42. doi: 10.1017/S0960428610000260
- Fritsch PW, Lu L, Bush CM, Cruz BC, Kron KA, Li D-Z (2011) Phylogenetic analysis of the Wintergreen Group (Ericaceae) based on six genic regions. Systematic Botany 36(4): 990–1003. doi: 10.1600/036364411X604994
- Fritsch PW, Amoroso VB (in press) *Diplycosia kitangladensis* sp. nov. from Mindanao, The Philippines. Natural History Bulletin of the Siam Society.
- Fritsch PW, Bush CM (in press) *Diplycosia rigidifolia* sp. nov. (Ericaceae) from Borneo, Sabah, Malaysia. Nordic Journal of Botany.
- IUCN Standards and Petitions Subcommittee (2016) Guidelines for using the IUCN Red List categories and criteria. Version 12. Prepared by the Standards and Petitions Subcommittee. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf> [accessed 6.6.2016]
- Pelser PB, Barcelona JF, Nickrent DL (Eds) (2011 onwards) Co's Digital Flora of the Philippines. <http://www.philippineplants.org> [accessed 6.6.2016]
- Powell EA, Kron KA (2001) An analysis of the phylogenetic relationships in the wintergreen group (*Diplycosia*, *Gaultheria*, *Pernettya*, *Tepuia*; Ericaceae). Systematic Botany 26(4): 808–817. doi: 10.1043/0363-6445-26.4.808
- Sleumer H (1967) Ericaceae. Flora Malesiana Ser. 1, 6(5): 669–914.
- Thiers B (2016) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/> [continuously updated; accessed 6.6.2016]