

# The correct name for a section of *Ludwigia* L. (Onagraceae)

Peter C. Hoch<sup>1</sup>, Warren L. Wagner<sup>2</sup>, Peter H. Raven<sup>1</sup>

**1** Missouri Botanical Garden, P.O. Box 299, St. Louis, MO 63166-0299, USA **2** Department of Botany, MRC-166, National Museum of Natural History, Smithsonian Institution, P.O. Box 37012, Washington, DC 20013-7012, USA

Corresponding author: Peter C. Hoch (peter.hoch@mobot.org)

---

Academic editor: Peter de Lange | Received 13 March 2015 | Accepted 30 April 2015 | Published 27 May 2015

---

**Citation:** Hoch PC, Wagner WL, Raven PH (2015) The correct name for a section of *Ludwigia* L. (Onagraceae). *PhytoKeys* 50: 31–34. doi: 10.3897/phytokeys.50.4887

---

## Abstract

In 1953, Hara provided new combinations for many sectional and species names when he combined *Jussiaea* L. with *Ludwigia* L., and at the time, *Ludwigia* sect. *Oligospermum* (Micheli) H.Hara was the correct name for one well-defined section. However, subsequent changes to/clarifications of the botanical code have necessitated a change for that name in that now an autonym is treated as having priority over the name or names of the same date and rank that established it. Since Hara's combination was based on *Jussiaea* sect. *Oligospermum* Micheli, the correct name for this section is *Ludwigia* sect. *Jussiaea* (L.) Hoch, W.L.Wagner, & P.H.Raven.

## Keywords

*Ludwigia*, *Jussiaea*, sect. *Oligospermum*, nomenclature

## Introduction

The cosmopolitan group of species known since 1953 as *Ludwigia* sect. *Oligospermum* is one of the most distinctive in the genus (Raven 1963, Wagner et al. 2007). Among the diplostemonous sections of *Ludwigia* – which formerly were segregated as the genus *Jussiaea* L. – this section differs by having woody, subcylindrical capsules with uniseriate seeds firmly embedded in woody pieces of endosperm and pollen released singly

rather than as tetrads or polyads. Most species of the section, which grow in warm-temperate to subtropical moist or wet habitats worldwide, are vigorously aquatic and some (*L. peploides* (Kunth) P.H.Raven, *L. hexapetala* (Hook. & Arn.) Zardini, H.Y.Gu & P.H.Raven) can be invasive weeds in wetlands and other wet areas, most recently so in California (Wood 2006, Hoch and Grewell 2012). This polyploid section comprises a group of nine highly variable species that includes three diploid species ( $n = 8$ ), four tetraploid species ( $n = 16$ ), one hexaploid species ( $n = 24$ , *L. grandiflora* (Michx.) Greuter & Burdet), and one decaploid species ( $n = 40$ , *L. hexapetala*; see also Nesom and Kartesz 2000). Most species in this section have native distributions restricted to the New World, but two species are restricted to the Old World, *L. stolonifera* (Guill. & Perr.) P.H.Raven throughout Africa and Madagascar, extending to Turkey and Iraq, and *L. adscendens* (L.) H.Hara across tropical Asia from India to New Guinea, and from southern Japan to northern Australia, and probably naturalized in Madagascar (Raven 1963, Wagner et al. 2007).

While editing the treatment of *Ludwigia* for the Flora of North America, Jim Zarucchi noticed a problem with the name used for this section, and after consultation with Kanchi Gandhi informed us that a change was necessary. We are grateful to Zarucchi and Gandhi for pointing out this problem for us. We are making this change now so that the correct combination can be available for FNA.

In his treatment for the Flora Brasiliensis (Martius 1875), Micheli divided the genus *Jussiaea* into three sections: sect. *Eujussiaea*, sect. *Oligospermum*, and sect. *Macrocarpon*. This division of the genus has been widely followed in subsequent treatments. Munz (1942) provided a treatment for New World species of *Jussiaea* in which he recognized the same three sections, but provided different names for two of them (no change for sect. *Macrocarpon* Micheli). For sect. *Eujussiaea* Micheli, he used the new name sect. *Myrtocarpus* Munz, and for sect. *Oligospermum* Micheli, he used sect. *Eujussiaea* Munz. His rationale was that the section that included the type species of the genus had to retain the generic name, and since Hitchcock and Greene (1929) effectively lectotypified *Jussiaea* with *J. repens* L. [= *Ludwigia adscendens* (L.) Hara], he proposed the name sect. *Eujussiaea* for the section that included *J. repens* (this lectotypification has been attributed incorrectly in the past to Britton and Brown (1913)).

Hara (1953), following the conclusion by Brenan (1953) and others that *Ludwigia*, *Jussiaea*, and *Isnardia* (a group sometimes segregated) formed a single genus (as *Ludwigia*, as established by Baillon 1877), recognized the sections in question as *Ludwigia* sect. *Oligospermum* (Micheli) H.Hara, sect. *Myrtocarpus* (Munz) H.Hara, and sect. *Macrocarpon* (Micheli) H.Hara. He noted that he was “strictly following the Code” (Hara 1953: 290). This treatment was widely accepted, including by Raven (1963) as well as Munz (1965). Most recently Wagner et al. (2007), in a synopsis of Onagraceae, included all three sections as proposed by Hara.

However, changes made to the ICBN in 1981 and retained in subsequent editions (McNeill et al. 2012), specifically as Article 11.6, invalidated part of this treatment. Article 11.6 states that “an autonym is treated as having priority over the name or

names of the same date and rank that established it.” So when the transfer of names from *Jussiaea* to *Ludwigia* was made, the correct sectional name combination for *Jussiaea* sect. *Oligospermum* should have been *Ludwigia* sect. *Jussiaea*, since this is the section that includes the type of the genus. Therefore, we make the following change in compliance with the ICBN.

## Nomenclature

***Ludwigia* L. sect. *Jussiaea* (L.) Hoch, W.L.Wagner, & P.H.Raven, comb. nov.**

urn:lsid:ipni.org:names:77147317-1

*Jussiaea* L., Sp. pl. 1: 388. 1753. [*Jussia* Adans., Fam. 2: 85, 565. 1763, orth. var.];

*Jussiaea* L. sect. *Eujussiaea* Munz, Darwiniana 4: 184. 1944.

Type. *Jussiaea repens* L. [= *Ludwigia adscendens* (L.) H.Hara] (Lectotype, designated by Hitchcock & Greene, Prop. Brit. Bot. 153. 1929).

*Jussiaea* L. sect. *Oligospermum* Micheli in Martius, Fl. bras. 13(2): 149, 162. 1875.

*Ludwigia* L. sect. *Oligospermum* (Micheli) H.Hara, J. Jap. Bot. 28: 290. 1953.

Type. *Jussiaea hookeri* Micheli [= *Ludwigia hookeri* (Micheli) H.Hara] (Lectotype, designated by Raven, Reinwardtia 6: 335. 1963).

*Cubospermum* Lour., Fl. Cochinch. 258, 275. 1790.

Type. *Cubospermum palustre* (L.) Lour. [= *Ludwigia adscendens* (L.) H.Hara].

*Adenola* Raf., Aut. Bot. 36. 1840.

Type. *Adenola grandiflora* (Michx.) Raf. [= *Ludwigia grandiflora* (Michx.) Greuter & Burdet]. (Lectotype, designated by Pennell, Bull. Torrey Bot. Club 48: 93. 1921).

*Oocarpon* Micheli, Flora 57: 303. 1874. *Ludwigia* sect. *Oocarpon* (Micheli) P.H.Raven, Reinwardtia 6: 336. 1963.

Type: *Oocarpon jussiaeoides* Micheli [= *Ludwigia torulosa* (Arn.) H.Hara].

**Description.** Perennial herbs, stems creeping, floating, or emergent and ascending to erect, rooting at nodes, when floating often forming spongy white pneumatophores at nodes, when erect with spongy base, terete. Leaves alternate; blades with one submarginal vein. Flowers 5(6)-merous; petals present, yellow or white; stamens twice as many (rarely as many) as sepals, pollen shed in monads. Capsules cylindrical, terete, often curved up, woody with thick walls, irregularly and tardily dehiscent. Seeds in one row per locule, pendulous, and firmly embedded in a woody coherent segment of endocarp, with inconspicuous raphe.  $2n = 16, 32, 48, 80, 96$ .

**Taxa included.** *Ludwigia adscendens* (L.) H.Hara, *L. grandiflora* (Michx.) Greuter & Burdet, *L. helminthorrhiza* (Mart.) H.Hara, *L. hexapetala* (Hook. & Arn.) Zardini, H.Y.Gu & P.H.Raven, *L. hookeri* (Micheli) H.Hara, *L. peduncularis* (C.Wright ex Griseb.) M.Gómez, *L. peploides* (Kunth) P.H.Raven subsp. *glabrescens* (O. Kuntze) P.H.Raven, *L. peploides* subsp. *montevideensis* (Sprengel) P.H.Raven, *L. peploides* sub-

sp. *peploides*, *L. peploides* subsp. *stipulacea* (Ohwi) P.H.Raven, *L. stolonifera* (Guill. & Perr.) P.H.Raven, *L. torulosa* (Arn.) H.Hara.

## Acknowledgments

Thanks to Jim Zarucchi for bringing this problem to our attention, and to Kanchi Gandhi for guidance on the nomenclature.

## References

- Baillon HE (1877) Onagraceae. *Histoire des plantes* 6: 305–516.
- Brenan JPM (1953) Notes on African Onagraceae and Trapaceae. *Kew Bulletin* 8: 163–172. doi: 10.2307/4117169
- Britton NL, Brown A (1913) *An illustrated flora of the northern United States, Canada and the British possessions*, Ed 2, 2: 1–735.
- Hara H (1953) *Ludwigia* versus *Jussiaea*. *Journal of Japanese Botany* 28: 289–294.
- Hitchcock AS, Greene ML (1929) *Standard species of Linnean genera of Phanerogamae (1753–54)*. International Botanical Congress. Cambridge (England), 1930: Nomenclature. Proposals by British Botanists, H.M.S.O., London, 110–199.
- Hoch PH, Grewell BJ (2012) *Ludwigia*. In: Baldwin BG, Goldman DH, Keil DJ, Patterson R, Rosatti TJ, Wilkin DH (Eds) *The Jepson Manual: Vascular Plants of California*, Second Edition. Univ. California Press, Berkeley, 948–949.
- Martius CFP de (1875) Onagraceae. *Flora Brasiliensis* 13(2): 146–182.
- McNeill J, Barrie FR, Buck WR, Demoulin V, Greuter W, Hawksworth DL, Herendeen PS, Knapp S, Marhold K, Prado J, PrudPhomme van Reine WF, Smith GF, Wiersema JH, Turland NJ (2012) *International Code of Nomenclature for algae, fungi, and plants (Melbourne Code)* adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011. Koeltz Scientific Books, Königstein, Germany. [Regnum Veg. 154]
- Munz PA (1942) *Studies in Onagraceae XII. A revision of the new world species of Jussiaea*. *Darwiniana* 4: 179–284.
- Munz PA (1965) Onagraceae. *North American Flora*, ser. 2, 5: 1–278.
- Nesom GL, Kartesz JT (2000) Observations on the *Ludwigia uruguayensis* complex (Onagraceae) in the United States. *Castanea* 65: 123–125.
- Raven PH (1963) The Old World species of *Ludwigia* (including *Jussiaea*), with a synopsis of the genus (Onagraceae). *Reinwardtia* 6: 327–427.
- Wagner WL, Hoch PC, Raven PH (2007) Revised classification of the Onagraceae. *Systematic Botany Monographs* 83: 1–240.
- Wood M (2006) Squelching water primrose. *Agricultural Research Magazine* 54(5): 13.