

Nomenclatural changes in Onagraceae

Peter C. Hoch¹, Kanchi Gandhi²

1 *Missouri Botanical Garden, 4344 Shaw Blvd., St. Louis, Missouri 63110-2291, USA* **2** *Harvard University Herbaria, 22 Divinity Ave., Cambridge, MA 02138, USA*

Corresponding author: Peter C. Hoch (peter.hoch@mobot.org)

Academic editor: R. Kriebel | Received 14 February 2020 | Accepted 28 February 2020 | Published 10 April 2020

Citation: Hoch PC, Gandhi K (2020) Nomenclatural changes in Onagraceae. *PhytoKeys* 145: 57–62. <https://doi.org/10.3897/phytokeys.145.51139>

Abstract

A new subspecies and two new combinations are proposed in Onagraceae. *Ludwigia glandulosa* Walter subsp. *brachycarpa* C.-I Peng, **subsp. nov.** is morphologically distinct from the typical subspecies, with smaller capsules and leaves, different seed coat, and a restricted distribution. *Epilobium* sect. *Pachydium* (Fischer & C. A. Meyer) Hoch & K. Gandhi, **comb. nov.** refers to a distinctive group of species formerly known as *Boisduvalia* Spach and as *Epilobium* sect. *Boisduvalia* (Spach) Hoch & P. H. Raven. And *Chamaenerion speciosum* (Decaisne) Hoch & K. Gandhi, **comb. nov.** is proposed for a distinctive Himalayan species originally described in *Epilobium*.

Keywords

Boisduvalia, Chamaenerion, fireweeds, Ludwigia, nomenclature

Introduction

The plant family Onagraceae is known in considerable detail as a result of modern monographic studies of almost the entire family and numerous comparative morphological analyses, summarized in Wagner et al. (2007). Recent phylogenetic analyses (especially Levin et al. 2003, 2004) provided much insight into the relationships in the family and necessitated many changes in the classification. Most of these changes were included in Wagner et al., but recent work revealed the need for several additional nomenclatural changes.

Methods

This contribution is the result of careful nomenclatural review by K. Gandhi of the treatment of Onagraceae by WL Wagner and PC Hoch for the Flora of North America. Gandhi detected several nomenclatural problems, which are corrected by the following changes.

Taxonomic treatment

Ludwigia L., a pan-subtropical genus of 82 species, forms a strongly monophyletic lineage sister to the rest of Onagraceae (e.g., Levin et al. 2003, 2004). Recent molecular analysis (Liu et al. 2017) has challenged the complex sectional classification of *Ludwigia* (22 sections), but full resolution awaits more detailed analysis. One strongly monophyletic clade is section *Isnardia* (L.) W. L. Wagner & Hoch, a group of 19 species centered in southeastern North America (Liu et al. in press) and formerly treated as sections *Dantia* (DeCandolle) Munz (Peng et al. 2005) and *Microcarpium* Munz (Peng 1989; Wagner et al. 2007). While reviewing the treatment of Onagraceae for the Flora of North America, one of us (KG) noticed an error in the treatment of *L. glandulosa* Walter, a widespread species of section *Isnardia*. Peng (1986) initially published *L. glandulosa* subsp. *brachycarpa* (Torrey & A. Gray) C. Peng as a comb. nov., based on *L. cylindrica* Elliott β . *brachycarpa* Torrey & A. Gray, but that latter name was a comb. nov. based on *Jussiaea brachycarpa* Lam., which Peng considered to be a synonym of *L. glandulosa* subsp. *glandulosa*. Therefore, *L. glandulosa* subsp. *brachycarpa* was not a comb. nov., as Peng proposed, but instead a subsp. nov. However, it was invalid since it lacked a diagnosis/description in Latin, as required at that time, and designation of a type specimen. Here we correct the mistake, and validate the name with a type and description. Unfortunately, our colleague Ching-I Peng died in 2018, but since his original intention was clear and he provided the name and description (Peng 1986, 1989), with this authorship we honor his enormous contributions to our understanding of *Ludwigia*.

Ludwigia glandulosa Walter subsp. *brachycarpa* C.-I Peng, subsp. nov.

urn:lsid:ipni.org:names:77209334-1

Diagnosis. *Ludwigia glandulosa* subsp. *brachycarpa* differs from typical *L. glandulosa* in its smaller stature, 10–55(–90) cm (vs. (20–)40–80(–100) cm); narrower leaf blades, 0.3–0.5(–1) cm (vs. 0.3–2.1 cm); shorter sepals, 1.1–1.9 mm (vs. 1.3–2.3 cm); smaller capsules, 2–5 × 1.3–2 mm (vs. (4–)5–8(–9) × 1.6–2(–3) mm); and seed surface cells elongate transversely to length (vs. surface cells elongate parallel to length).

Type material. USA, Louisiana, Cameron Parish, 3.2 km W of junction State Highways 82 and 27, 29°48.62'N, 93°08.25'W, 1 m, 16 August 1980, C.-I Peng, W Peng and J. Chen 4367 (holotype: MO 2806683); see fig. 42 in Peng (1989).

Description. *Stems* rarely reddish green, 10–55(–90) cm. *Leaves:* petiole 0–1 cm, blades linear-elliptic to linear, sometimes very narrowly elliptic, those on main axis 3–5(–7) × 0.3–0.5(–1) cm, those on branches 0.8–3.6 × 0.2–0.3(–0.8) cm. *Inflorescences:* bracteoles attached at base of ovary, 0.4–0.8 × 0.1–0.2 mm. *Flowers:* sepals 1.1–1.9 × 1–1.8 mm, apex acute or short-acuminate; nectary disc obscurely, minutely papillose; style 0.4–0.8 mm, stigma 0.2–0.3 mm diam. *Capsules* obscurely 4-angled, 2–5 × 1.3–2 mm, pedicel 0–0.2 mm. *Seeds* 0.6–0.8 × 0.3–0.4 mm, surface cells elongate transversely to seed length. *Chromosome number:* $n = 16$.

Phenology. Flowering and fruiting April to November.

Etymology. The subspecific epithet '*brachycarpum*' refers to the short capsules.

Distribution and habitat. *Ludwigia glandulosa* subsp. *brachycarpa* is endemic to the US Gulf Coast from southwestern Louisiana to Nueces County, Texas, and more sporadically northward in eastern Texas to south-central Oklahoma. This distribution is at the extreme southwestern edge of that for subsp. *glandulosum*, which grows from Texas and Oklahoma east to Virginia and Florida and north to southern Missouri, Illinois, and Indiana (Peng 1989). Although they overlap in part, the two taxa are only rarely locally sympatric. *Ludwigia glandulosa* subsp. *brachycarpa* grows in ditches, low meadows, coastal prairies, seeps in sandy woods, moist sinkholes in granite outcrops, old clay fields at an elevation of 0–200 m.

New combinations

Epilobium L. is the largest genus in the family Onagraceae; its 165 species are widely distributed in cool or cold regions of the world, with a center of diversity in western North America (Wagner et al. 2007). A group of annual species with affinities to *Epilobium* but considered distinct because they lack seed comas was historically segregated as *Boisduvalia* Spach (Raven 1976). However, molecular and other evidence (Hoch and Raven 1992; Baum et al. 1994; Wagner et al. 2007) unequivocally place *Boisduvalia* within *Epilobium* as two non-monophyletic sections. *Epilobium* sect. *Epilobiopsis* (Spegazzini) Lievens, Hoch & PH Raven is a group of two species characterized by tough, tardily dehiscent capsules, seeds in two rows per locule, and a chromosome number of $n = 15$; and *E.* sect. *Boisduvalia* (Spach) Hoch & P. H. Raven is a group of four species characterized by friable, readily dehiscent capsules, seeds in one row per locule, and chromosome numbers of $n = 9, 10, 19$. In naming this latter section, however, we overlooked an earlier name at the sectional level, one of two proposed for this group by Fischer and Meyer (1836), who treated the group as part of *Oenothera* L. due to the absence of seed comas. Therefore, a new combination is required.

***Epilobium* Linnaeus sect. *Pachydium* (Fischer & C.A. Meyer) Hoch & K. Gandhi, comb. nov.**

urn:lsid:ipni.org:names:77209335-1

Basionym. *Oenothera* sect. *Pachydium* Fischer & C.A. Meyer, Ind. sem. hort. petrop. 2: 45. 1836 [“1835”]. *Boisduvalia* [unranked] *Pachydium* (Fischer & C.A. Meyer) Endlicher, Gen. pl. 1191. 1840. *B.* subg. *Pachydium* (Fischer & C.A. Meyer) Reichenbach, Deut. Bot. Herb.-Buch. 170. 1841; *Boisduvalia* sect. *Pachydium* (Fischer & C.A. Meyer) Munz, N. Amer. Flora 5, II: 228.

Type. *Oenothera densiflora* Lindley [= *Epilobium densiflorum* (Lindley) Hoch & P.H. Raven].

The distinctive, circumboreal/ circumpolar group commonly known as fireweeds has been treated either as a section of *Epilobium* (Haussknecht 1884; Raven 1976; Chen et al. 1992) or as the separate genus *Chamaenerion* Séguier. Although the two groups share the distinctive comose seeds, several floral features, and a base chromosome number of $x = 18$, *Chamaenerion* differs from *Epilobium* in having leaves nearly always spirally arranged, rarely subopposite or verticillate near stem base (vs. opposite at least on proximal stem); lack of a floral tube (vs. more or less distinct floral tube); flowers slightly zygomorphic with subequal stamens that are erect, then deflexed, and styles that are deflexed, then erect (vs. actinomorphic with erect stamens in two series and erect styles); petals entire (vs. emarginate); and pollen shed in monads (vs. tetrads) (Wagner et al. 2007). Recent molecular analyses (Baum et al. 1994; Levin et al. 2004) also demonstrated that the fireweeds form a strongly supported clade sister to the rest of *Epilobium* (Wagner et al. 2007).

Holub (1972) argued that the correct name for the fireweeds at the generic level should be *Chamerion* (Raf.) Raf. ex Holub, not *Chamaenerion*, which he argued was illegitimate. However, as noted in personal correspondence between KG and Ulf Eliasson in 2009, and summarized by Sennikov (2011), clarifications in the botanical code and in the lectotypification of *Chamaenerion* and *Epilobium* negate Holub’s analysis, and Sennikov concluded that the correct and valid name for the fireweeds at the generic level is *Chamaenerion* Séguier.

All but one of the eight species recognized in this genus have been treated at some point as species in *Chamaenerion*, the only exception being a species described in *Epilobium* that is endemic to the Himalayan region from Kashmir to Nepal and Xizang (Tibet), China (Chen et al. 1992), for which the following new combination is provided:

***Chamaenerion speciosum* (Decaisne) Hoch & K. Gandhi, comb. nov.**

urn:lsid:ipni.org:names:77209336-1

Basionym. *Epilobium speciosum* Decaisne, Voy. Ind. (Jacquemont) 4: 57, t. 69 [Apr 1835 – Dec 1844]. *Epilobium latifolium* L. subsp. *speciosum* (Decne.) P. H. Raven, Bull. Brit. Mus. (Nat. Hist.), Bot. 2(12): 349. 1962. *Chamerion speciosum* (Decne.)

Holub, Folia Geobot. Phytotax. 7: 86. 1972; non *Chamaenerion speciosum* Lodd. ex Steudel, Nomencl. Bot. ed. 2, 1: 343. 1840, pro syn.

Type material. India. Himachal Pradesh, near Yurpo, 3800m 1830–1831, *V. Jacquemont 1739* (**holotype**: P; **isotypes**: G, K).

Acknowledgements

We thank Werner Greuter and John Wiersema for useful discussions on the nomenclature of *Ludwigia*, and David E. Boufford, Warren L. Wagner, and Peter H. Raven for comments on the manuscript.

References

- Baum DA, Sytsma KJ, Hoch PC (1994) The phylogeny of *Epilobium* (Onagraceae) based on nuclear ribosomal DNA sequences. *Systematic Botany* 19(3): 363–388. <https://doi.org/10.2307/2419763>
- Chen CJ, Hoch PC, Raven PH (1992) Systematics of *Epilobium* (Onagraceae) in China. *Systematic Botany Monographs* 34: 1–209. <https://doi.org/10.2307/25027806>
- Fischer FEL, Meyer CA (1836) *Index Seminum* [St. Petersburg (Petropolitanus)] 2: 45.
- Hausknecht C (1884) *Monographie der Gattung Epilobium*. Gustav Fischer, Jena.
- Hoch PC, Raven PH (1992) *Boisduvalia*, a coma-less *Epilobium* (Onagraceae). *Phytologia* 73: 456–459. <https://doi.org/10.5962/bhl.part.16727>
- Holub J (1972) Taxonomic and nomenclatural remarks on *Chamaenerion* auct. *Folia Geobotanica et Phytotaxonomica* 7(1): 81–90. <https://doi.org/10.1007/BF02856384>
- Levin RA, Wagner WL, Hoch PC, Nepokroeff M, Pires JC, Zimmer EA, Sytsma KJ (2003) Family-level relationships of Onagraceae based on chloroplast *rbcL* and *ndhF* data. *American Journal of Botany* 90(1): 107–115. <https://doi.org/10.3732/ajb.90.1.107>
- Levin RA, Wagner WL, Hoch PC, Hahn WJ, Rodriguez A, Baum DA, Katinas L, Zimmer EA, Sytsma KJ (2004) Paraphyly in Tribe Onagreae: Insights into phylogenetic relationships of Onagraceae based on nuclear and chloroplast sequence data. *Systematic Botany* 29(1): 147–164. <https://doi.org/10.1600/036364404772974293>
- Liu SH, Hoch PC, Raven PH, Barber JC (2017) Multi-locus phylogeny of *Ludwigia* L. (Onagraceae): Insights on infra-generic relationships and the current classification of the genus. *Taxon* 66(5): 1112–1127. <https://doi.org/10.12705/665.7>
- Liu SH, Yang HA, Kono Y, Hoch PC, Barber JC, Peng CI, Chung KF (in press) Disentangling reticulate evolution of North Temperate haplostemonous *Ludwigia* (Onagraceae). *Annals of the Missouri Botanical Garden*.
- Peng CI (1986) A new combination in *Ludwigia* sect. *Microcarpium* (Onagraceae). *Annals of the Missouri Botanical Garden* 73(2): 490. <https://doi.org/10.2307/2399128>
- Peng CI (1989) The systematics and evolution of *Ludwigia* sect. *Microcarpium* (Onagraceae). *Annals of the Missouri Botanical Garden* 76(1): 221–302. <https://doi.org/10.2307/2399348>

- Peng CI, Schmidt CL, Hoch PC, Raven PH (2005) Systematics and evolution of *Ludwigia* sect. *Dantia* (Onagraceae). *Annals of the Missouri Botanical Garden* 92: 307–359.
- Raven PH (1976) Generic and sectional delimitation in Onagraceae, tribe Epilobieae. *Annals of the Missouri Botanical Garden* 63(2): 326–340. <https://doi.org/10.2307/2395311>
- Sennikov AN (2011) *Chamerion* or *Chamaenerion* (Onagraceae)? The old story in new words. *Taxon* 60(5): 1485–1488. <https://doi.org/10.1002/tax.605028>
- Wagner WL, Hoch PC, Raven PH (2007) Revised classification of the Onagraceae. *Systematic Botany Monographs* 83: 1–240.